

# On Possible Approaches to Differentiation of Rough Real Functions

Zoltán Ernő Csajbók

Department of Health Informatics, Faculty of Health, University of Debrecen  
Sóstói út 2-4, H-4406 Nyíregyháza, Hungary  
[csajbok.zoltan@foh.unideb.hu](mailto:csajbok.zoltan@foh.unideb.hu)

## Abstract

In the mid 1990s Z. Pawlak relying on the rough set theory initiated the study of rough calculus in his many papers. He invented the investigation of its different subfields such as rough continuity, rough derivatives–integrals, rough differential equations, etc. Some authors have systematically investigated the rough continuity of rough real functions in Pawlak’s sense. The following reasonable step would be to define the derivative of rough functions. However, it does not seem clear how it could be carried through this important step. In the paper, a possible approach will be outlined.

*Keywords:* Rough functions, discrete calculus, digital calculus, rough calculus

*MSC:* 03E70, 26A24

## 1. Introduction

In the mid 1990s Z. Pawlak relying on the rough set theory (RST) [6, 7, 15] initiated the study of rough calculus in his many papers [9, 11, 13, 14]. He invented the investigation of its different subfields such as rough continuity–discontinuity, rough derivatives–integrals, rough differential equations, etc.

The paper [3] systematically investigates the rough continuity–discontinuity of rough real functions in Pawlak’s sense. The next reasonable step would be to define the derivative of rough functions.

Pawlak defined the rough derivatives based on discrete calculus. This paper basically, but not completely follows Pawlak’s method.

The rest of paper is organized as follows. After the introduction, Section 1, the rough real numbers are defined in Section 2. Section 3 surveys different possible approximations of rough functions. Section 4 defines the rough derivatives and discusses some special features of this approach.

---

*Copyright © 2020 for this paper by its authors. Use permitted under Creative Commons License Attribution 4.0 International (CC BY 4.0).*

## 2. Rough real numbers

Let  $U, V$  be two classical nonempty sets. A *function*  $f$  with domain  $U$  and co-domain  $V$  is denoted by  $f : U \rightarrow V, u \mapsto f(u)$ .  $V^U$  denotes the set of all functions with domain  $U$ , in notation  $\text{Dom}f = U$ , and co-domain  $V$ , in notation  $\text{Im}f = V$ .

If  $f, g \in V^U$ , the operation  $f \odot g, \odot \in \{+, -, \cdot, /\}$  and the relation  $f \boxtimes g, \boxtimes \in \{=, \neq, \leq, <, \geq, >\}$  are understood by pointwise.

For any  $S \subseteq U, f(S) = \{f(u) \mid u \in S\} \subseteq V$  is the direct image of  $S$ . Especially,  $f(U) \subseteq V$  is the *range* of  $f$ .

If  $a, b \in \mathbb{R} (a \leq b), [a, b] = \{x \in \mathbb{R} \mid a \leq x \leq b\}$  and  $]a, b[ = \{x \in \mathbb{R} \mid a < x < b\}$  denote *closed* and *open* intervals.  $[a, a] = \{a\}$  is identified with the real number  $a \in \mathbb{R}$ . It is easy to interpret the open-closed  $]a, b]$  and closed-open  $[a, b[$  intervals.

$(a, b)$  means an ordered pair of real numbers  $a$  and  $b$ .

Let  $\mathbb{R}^{\geq 0}$  denote the set of nonnegative real numbers. Let  $[n] = \{0, 1, \dots, n\} \subseteq \mathbb{N}$  be a finite set of natural numbers. Accordingly,  $]n] = 1, \dots, n, [n[ = 1, \dots, n - 1,$  and  $]n[ = 1, \dots, n - 1$ .

Definition of rough real numbers can be found in Pawlak's different papers such as [13, 14, 9, 8, 11, 12]. Here, it is briefly summarized.

Let  $I$  denote a closed interval  $I = [0, a] (a \in \mathbb{R}^{\geq 0}, a > 0)$ .

**Definition 2.1.** A *categorization* of  $I$  is a sequence  $S_I = \{x_i\}_{i \in [n]} \subseteq \mathbb{R}^{\geq 0}$ , where  $n \geq 1$  and  $0 = x_0 < x_1 < \dots < x_n = a$ .  $S_I$  is also called the *discretization* of  $I$ .

Let  $I_S$  denote an equivalence relation generated by the categorization  $S_I$ . Let  $x, y \in I$ .  $xI_Sy$  if  $x = y = x_i \in S_I$  for some  $i \in [n]$ , or  $x, y \in ]x_i, x_{i+1}[$  for some  $i \in [n[$ . Hence, the partition  $I/I_S$  associated with the equivalence relation  $I_S$  is:

$$I/I_S = \{\{x_0\}, ]x_0, x_1[, \{x_1\}, \dots, \{x_{n-1}\}, ]x_{n-1}, x_n[, \{x_n\}\}.$$

It should be noted that in classical analysis, the term "partition of  $I$ " is used in a slightly different sense: Two compact real intervals *nonoverlapping* if either they are disjoint or their intersection contains at most one point, which necessarily an endpoint of both intervals ([2], p. 4). In the classical analysis context, a *partition* of  $I$  is a collection of nonoverlapping closed intervals whose union is  $I$  ([1], p. 149).

The block of the partition  $I/I_S$  containing  $x \in I$  is denoted by  $\llbracket x \rrbracket_{I_S}$ . In particular, if  $x \in S_I, \llbracket x \rrbracket_{I_S} = \{x\}$ . If  $x \in ]x_i, x_{i+1}[, \overline{\llbracket x \rrbracket}_{I_S} = [x_i, x_{i+1}]$  is the *closure* of  $\llbracket x \rrbracket_{I_S}$ . Of course, when  $x \in S_I, \llbracket x \rrbracket_{I_S} = \overline{\llbracket x \rrbracket}_{I_S} = \{x\}$ .

In terms of RST terminology,  $I_S$  is an indiscernibility relation on  $I$ . Hence, the naming of the following notions is consistent with the standard terminology of RST.

The members of  $I/I_S$  are called *elementary* or *base* sets. Any union of base sets are referred to as *definable* sets. By definition,  $\emptyset$  is definable. Their collection is denoted by  $\mathcal{D}_{I/I_S}$ .

The principal notions of RST are the lower and upper approximation functions,  $\text{ls}$  and  $\text{us}$ , respectively. Most commonly, their domain and co-domain are the power

set of  $I$ . In the following, however, the closed intervals of the form  $[0, x]$  ( $x \in I$ ) will only be approximated. Therefore,

$$\begin{aligned} l_S([0, x]) &= \{x' \in I \mid \llbracket x' \rrbracket_{I_S} \subseteq [0, x]\} = \cup\{\llbracket x' \rrbracket_{I_S} \in I/I_S \mid \llbracket x' \rrbracket_{I_S} \subseteq [0, x]\}, \\ u_S([0, x]) &= \{x' \in I \mid \llbracket x' \rrbracket_{I_S} \cap [0, x] \neq \emptyset\} = \cup\{\llbracket x' \rrbracket_{I_S} \in I/I_S \mid \llbracket x' \rrbracket_{I_S} \cap [0, x] \neq \emptyset\}. \end{aligned}$$

$\text{PAS}(I) = (I, I/I_S, \mathcal{D}_{I/I_S}, l_S, u_S)$  is called *Pawlak approximation space*.

With a slight abuse of notation, let us define the following numbers:

$$l_S(x) = \max\{x' \in S_I \mid x' \leq x\} \text{ and } u_S(x) = \min\{x' \in S_I \mid x' \geq x\}.$$

Of course,  $l(x) \leq x \leq u_S(x)$ , and  $l_S(x) = u_S(x) = x$  iff  $x \in S_I$ . Moreover,

- $l_S([0, x]) = [0, l_S(x)] = [0, x]$  and  $u_S([0, x]) = [0, u_S(x)] = [0, x]$  (if  $x \in S_I$ );
- $l_S([0, x]) = [0, l_S(x)] \subsetneq [0, x]$  and  $u_S([0, x]) = [0, u_S(x)] \supsetneq [0, x]$  (if  $x \notin S_I$ ).

It is said that the number  $x \in I$  is *exact* with respect to  $\text{PAS}(I)$  if  $l_S(x) = u_S(x)$ , otherwise  $x$  is *inexact* or *rough* [13]. Of course,  $x \in I$  is exact iff  $x \in S_I$ .

Members of  $I/I_S$  are called *rough numbers* with respect to  $\text{PAS}(I)$ . They can be represented as  $\llbracket x \rrbracket_{I_S} = [l_S(x), u_S(x)] = \{x\}$  if  $x \in S_I$ ,  $\llbracket x \rrbracket_{I_S} = ]l_S(x), u_S(x)[$  if  $x \notin S_I$ .

### 3. Approximation of rough real functions

Let  $I = [0, a_I]$ ,  $J = [0, a_J]$  be two closed intervals with  $a_I, a_J \in \mathbb{R}^{\geq 0}$ ,  $a_I, a_J > 0$ . Let  $S_I, P_J$  be categorizations of  $I$  and  $J$ , where  $S_I = \{x_i\}_{i \in [n]}$ ,  $P_J = \{y_j\}_{j \in [m]} \subseteq \mathbb{R}^{\geq 0}$  with  $m, n \geq 1$ ,  $0 = x_0 < x_1 < \dots < x_n = a_I$  and  $0 = y_0 < y_1 < \dots < y_m = a_J$ . The corresponding approximation spaces are  $\text{PAS}(I)$  and  $\text{PAS}(J)$ .

A function  $J^I$  is called a *rough real function* with respect to  $\text{PAS}(I)$  and  $\text{PAS}(J)$ .

To make the blocks of  $I/I_S$  easier to handle technically, they are enumerated as follows.

$$N_I : I/I_S \rightarrow [2n], \llbracket x \rrbracket_{I_S} \mapsto \begin{cases} B_{2i} = 2i, & \text{if } \exists i \in [n] (\llbracket x \rrbracket_{I_S} = \{x_i\} \subset S_I), \\ B_{2i+1} = 2i + 1, & \text{if } \exists i \in [n] (\llbracket x \rrbracket_{I_S} = ]x_i, x_{i+1}[). \end{cases}$$

The inverse of  $N_I$  is:

$$N_I^{-1} : [2n] \rightarrow I/I_S, \quad B_i \mapsto \begin{cases} \{x_{i/2}\}, & \text{if } i \equiv 0 \pmod{2}, \\ ]x_{\frac{i-1}{2}}, x_{\frac{i+1}{2}}[, & \text{if } i \equiv 1 \pmod{2}. \end{cases}$$

The equivalence classes of  $J/J_P$  can be enumerated in the same way by the help of an enumeration function  $N_J$ . Its values are referred to as  $C_j$ 's ( $j \in [2m]$ ).

**Example 3.1.** In the running example, let  $I = [0, x_5]$  with  $S_I = \{x_0, x_1, x_2, x_3, x_4, x_5\}$ , and  $J = [0, y_4]$  with  $P_J = \{y_0, y_1, y_2, y_3, y_4\}$ .

Figure 1 (a) shows the rough coordinate system with respect to  $\text{PAS}(I)$  and  $\text{PAS}(J)$ , and the enumeration of  $I/I_S$  and  $J/J_P$ . Figure 1 (b) presents some rough real functions in this rough coordinate system.

The purpose of this section is to show how rough real functions can be represented taken into account the features of approximation spaces  $\text{PAS}(I)$  and  $\text{PAS}(J)$ . The ideas of these representations mainly rely on Pawlak's paper [9, 10, 11, 12, 13].




Figure 1: Rough coordinate system and rough real functions

### 3.1. Pointwise approximation of rough real functions

**Definition 3.2** ([9, 13]). Let  $f \in J^I$ . The *pointwise*  $(S_I, P_J)$ -lower and  $(S_I, P_J)$ -upper approximations of  $f$  are the functions

$$\underline{f} : I \rightarrow P_J, x \mapsto \text{l}_P(f(x)), \quad \overline{f} : I \rightarrow P_J, x \mapsto \text{u}_P(f(x)).$$

$f$  is *exact* at  $x$ , if  $\underline{f}(x) = \overline{f}(x)$ , otherwise  $f$  is *inexact (rough)* at  $x$ .

$f$  is *pointwise exact* on  $I' \subseteq I$ , if  $\underline{f}(x) = \overline{f}(x)$  for all  $x \in I'$ , otherwise  $f$  is *pointwise inexact (rough)* on  $I'$ .

*Remark 3.3.* Rough real functions are *ab ovo* treated in  $(S_I, P_J)$ -coordinate systems. Hence, it seems reasonable to define the pointwise approximation in this context, too. Nevertheless, a “pointwise” feature much better fits to the whole interval  $I = [a, b]$ . But with the choice  $S = \{x_0 = 0, x_1 = a\}$ , this case is also included in the above definition.

$f \in J^I$  is exact at  $x \in I$  iff  $f(x) = y_j \in P_J$  for some  $j \in [m]$ . Geometrically it means that  $f$  is exact at a point in  $I$  iff in this point  $f$  touches or intersects a line segment  $y = y_j$ , where  $y_j \in P_J$ .

**Example 3.4.** Figure 2 (a) shows the *pointwise*  $(S_I, P_J)$ -lower and  $(S_I, P_J)$ -upper approximations,  $\underline{f}$  and  $\overline{f}$ , of a function  $f \in J^I$ .  $f$  is exact at points  $x^i, x^{ii}, x_2, x^{iii}$  and rough at all other points.

### 3.2. Blockwise approximation of rough real functions

**Definition 3.5.** Let  $f \in J^I$ . The *blockwise*  $(S_I, P_J)$ -lower and  $(S_I, P_J)$ -upper approximations of  $f$  are the functions

$$\underline{f} : I/I_S \rightarrow P_J, \llbracket x \rrbracket_{I_S} \mapsto \text{l}_P(\inf f(\llbracket x \rrbracket_{I_S})), \quad \overleftrightarrow{f} : I/I_S \rightarrow P_J, \llbracket x \rrbracket_{I_S} \mapsto \text{u}_P(\sup f(\llbracket x \rrbracket_{I_S})).$$




Figure 2: Pointwise and blockwise approximations of  $f$

$f$  is *blockwise exact* on  $B_i$  for some  $i \in [2n]$  if  $\overleftarrow{f}(B_i) = \overrightarrow{f}(B_i)$ , otherwise  $f$  is *blockwise inexact (rough)* on  $B_i$ .

Let  $f \in J^I$ ,  $i \in [2n]$ .  $f$  is blockwise exact on  $B_i$  iff  $f(B_i) = \{y_j\} \subset P_J$  for some  $j \in [m]$ . Geometrically it means that  $f$  is exact on  $B_i \in I/I_S$  iff  $f$  touches or intersects a line segment  $y = y_j$  for some  $y_j \in P_J$  at the point  $x_{i/2}$  if  $i \equiv 0 \pmod{2}$ , or  $f$  coincides with a line segment  $y = y_j$  for some  $y_j \in P_J$  on  $]x_{\frac{i-1}{2}}, x_{\frac{i+1}{2}}[$  when  $i \equiv 1 \pmod{2}$ .

**Example 3.6.** Figure 2 (b) shows the *blockwise*  $(S_I, P_J)$ -lower and  $(S_I, P_J)$ -upper approximations,  $\overleftarrow{f}$  and  $\overrightarrow{f}$ , of  $f$ .  $f$  is blockwise exact on  $B_4 = \{x_2\}$  only, and blockwise rough on all other blocks.

It should be noted that the pointwise and blockwise approximations defined above substantially differ from those considered in function approximation theory. Essentially because the lower and upper approximations (pointwise or blockwise) delimit a family of functions.

For instance, in the case of blockwise approximation, the domain and co-domain of every function  $f$  in this family, in their most general form, are  $\text{Dom} f = I$  and  $\text{Im} f = \{y \in J \mid \overleftarrow{f} \leq y \leq \overrightarrow{f}\}$ , *without any assignment rules*. Certain functions in this family can also be interpreted as partially specified ones with *unknown values*. With the help of getting finer and finer rough coordinates systems, actual values of these unknown values are becoming more and more recognizable.

### 3.3. Finite sequence approximations

**Definition 3.7.** Let  $f \in J^I$ . The *finite sequence*  $(S_I, P_J)$ -lower and  $(S_I, P_J)$ -upper approximations of  $f$  are the functions

$$f_\circ : [n] \rightarrow P_J, i \mapsto \text{l}_P(f(x_i)), \quad f^\circ : [n] \rightarrow P_J, i \mapsto \text{u}_P(f(x_i)).$$

This approximation characterizes the rough functions at the categorization points. However, it does not say anything about how a rough function behaves on the open intervals  $]x_i, x_{i+1}[$  ( $i \in [n]$ ).

**Definition 3.8.** Let  $f \in J^I$ . The *extended finite sequence*  $(S_I, P_J)$ -lower and  $(S_I, P_J)$ -upper approximations of  $f$  are the functions

$$f_\bullet : [2n] \rightarrow P_J, \quad i \mapsto \text{l}_P(\inf f(B_i)), \quad f^\bullet : [2n] \rightarrow P_J, \quad i \mapsto \text{u}_P(\sup f(B_i)).$$

Of course,  $f_\circ(i) = f_\bullet(2i)$ ,  $f^\circ(i) = f^\bullet(2i)$  ( $i \in [n]$ ), i.e., they are equal at the categorization points.

**Example 3.9.** Figure 3(a) and Figure 3(b) illustrate the finite sequence and extended finite sequence approximations of the function  $f$  (see Figure 2).




Figure 3: Finite sequence approximations of  $f$

### 3.4. Discrete sequence approximations

The most abstract approximations of rough functions is the discrete sequence one. In the definitions, the following function will be needed.

Let  $I$  and  $J$  two intervals with categorizations  $S_I$  and  $P_J$  be given as above.

$$\text{l}_S^{\text{enum}} : J \rightarrow [m], \quad y \mapsto \max\{i \in [m] \mid y_i \leq y\},$$

$$\text{u}_S^{\text{enum}} : J \rightarrow [m], \quad y \mapsto \min\{i \in [m] \mid y_i \geq y\}.$$

**Definition 3.10.** Let  $f \in J^I$ . The *discrete sequence*  $(S_I, P_J)$ -lower and  $(S_I, P_J)$ -upper approximations of  $f$  are the functions

$$f_\star : [n] \rightarrow [m], \quad i \mapsto \text{l}_S^{\text{enum}}(f(x_i)), \quad f^\star : [n] \rightarrow [m], \quad i \mapsto \text{u}_S^{\text{enum}}(f(x_i)).$$

This approximation also characterizes the rough functions at the categorization points only. It will be extended on the whole partition  $I/I_S$  of  $I$ .

**Definition 3.11.** Let  $f \in J^I$ . The *extended discrete sequence*  $(S_I, P_J)$ -lower and  $(S_I, P_J)$ -upper approximations of  $f$  are the functions

$$f_\star : [2n] \rightarrow [m], \quad i \mapsto \text{l}_S^{\text{enum}}(\inf f(B_i)), \quad f^\star : [2n] \rightarrow [m], \quad i \mapsto \text{u}_S^{\text{enum}}(\sup f(B_i)).$$

Of course,  $f_*(i) = f_*(2i)$ ,  $f^*(i) = f^*(2i)$  ( $i \in [n]$ ), i.e., they are equal at the categorization points.

**Example 3.12.** Figure 4 (a) and Figure 4 (b) illustrate the discrete sequence and extended discrete sequence approximations of  $f$ .




Figure 4: Discrete sequence approximations of  $f$

## 4. Rough derivatives

Classically, discrete derivatives, or more general, discrete calculus of functions was studied in the theory of “calculus of finite differences”, see, e.g., [4], one of the classical treatments of this subject. Traditionally, the subject of the discrete calculus is the functions of the form  $f: \mathbb{N} \rightarrow \mathbb{R}$ , i.e.,  $f$ 's are finite or infinite sequences.

In his papers, however, Pawlak defined the rough derivatives of such functions whose both domain and co-domain are finite set of natural numbers which he called *discrete functions*. In the following, under the discrete functions it is meant a function of the form  $f: \mathbb{N} \rightarrow \mathbb{Z}$ . Obviously,  $f: [n] \rightarrow [m]$  is discrete function as well. The calculus of such functions is called a rough calculus [12], or digital calculus [5].

### 4.1. Pawlak’s approach

Pawlak’s approach is based on the discrete representation of rough functions. This section mainly relies on Pawlak’s papers [10, 11, 12], in addition Nakamura, Rosenfeld’s paper [5]. Although, Nakamura and Rosenfeld defined derivatives in a slightly more general context, their results can be applied here.

**Definition 4.1.** Let  $f: [n] \rightarrow [m]$  be a discrete function. The *rough derivative*  $f'$  of  $f$  is the function

$$f': [n[ = [n - 1] \rightarrow \mathbb{Z}, \quad i \mapsto f(i + 1) - f(i).$$

The relationship between derivation and function operations slightly differ from the classical rules.

**Proposition 4.2** ([5], Theorem 1.4). *Let  $f, g : [n] \rightarrow [m]$  be two discrete functions. Then, for  $f \pm g, k \cdot f, f \cdot g, f/g$ , we have*

- $(f \pm g)'(i) = f'(i) \pm g'(i)$ ;
- $(k \cdot f)'(i) = k \cdot f'(i)$  ( $k \in \mathbb{Z}$ );
- $(f \cdot g)'(i) = f'(i) \cdot g(i) + f(i) \cdot g'(i) + f'(i) \cdot g'(i) = f(i) \cdot g'(i) + g(i+1) \cdot f'(i)$ ;
- $\left(\frac{f}{g}\right)'(i) = \frac{f'(i) \cdot g(i) - f(i) \cdot g'(i)}{g^2(i) + g(i) \cdot g'(i)}$ , provided that  $g^2(i) + g(i) \cdot g'(i) \neq 0$ ,

where  $(f \pm g)', (k \cdot f)', (f \cdot g)', (f/g)' : [n-1] \rightarrow \mathbb{Z}$ .

Higher order derivatives of  $f$  can be defined as usual: the second order derivative of  $f$  is the derivative of  $f'$ , etc. The following notations are used commonly for higher order derivatives:  $f = f^{(0)}$ ,  $f' = f^{(1)}$ ,  $f^{(2)} = f^{(1)'}$ ,  $f^{(3)} = f^{(2)'}$ , etc.

One can observe that  $\text{Dom}f^{(0)} = [n]$ ,  $\text{Dom}f^{(1)} = [n-1]$ ,  $\text{Dom}f^{(2)} = [n-1] - [n-2]$ , and, in general,  $\text{Dom}f^{(k)} = [n-k]$ . It also means that the discrete function  $f : [n] \rightarrow [m]$  has at most derivatives up to  $n$ -th order.

## 4.2. Discussion of rough derivatives

Digital calculus is applied to digital image processing [5, 16]. In this context, direct interpretation of rough derivatives is simple. Let  $f : [n] \rightarrow [m]$  be a discrete function, and  $f' : [n-1] \rightarrow \mathbb{Z}$  its derivative. Then,  $f$  can be interpreted as a piecewise linear function with slopes  $f'(i)$ 's ( $i \in [n-1]$ ). Of course, when  $f'(i) = 0$ ,  $f$  is constant between  $f(i)$  and  $f(i+1)$ , i.e.,  $f(i) = f(i+1)$ . In this interpretation, the domain of  $f$  can be considered as bounded compact real interval  $[0, m] \subset \mathbb{R}$ .

**Example 4.3.** Figures 5 (a) and 5 (b) show the discrete derivatives of  $f_\star$  and  $f^\star$ . Figures 5 (c) and 5 (d) depict their piecewise linear function interpretations.

On the other hand, rough calculus was motivated by the setting up a possible calculus of rough functions. Pawlak used the rough calculus, which is also known as digital calculus or discrete calculus, to achieve this goal. However, Pawlak did not establish a connection between the rough derivatives and rough functions. Indeed, it is hard to interpret rough derivatives as rough functions.

Different approximations of rough functions lead to different *sets* of rough functions. More specifically, let  $\text{PAS}(I)$  and  $\text{PAS}(J)$  be two approximation spaces as defined above. In addition, let  $f \in J^I$  be a fixed rough function. Then, all different approximations of  $f$  set up *sets of rough functions*. In regard to Pawlak's approach, here we will focus on the discrete sequence approximation of rough functions only.

The discrete sequence approximation determines the following set of rough functions:  $\mathcal{R}_f^{ds} = \{g \in J^I \mid \forall i \in [n](y_{f_\star(i)} \leq g(x_i) \leq y_{f^\star(i)})\}$ . That is,  $\mathcal{R}_f^{ds}$  consists of such the rough functions  $g$ 's which are bounded by  $y_{f_\star(i)}$  and  $y_{f^\star(i)}$  ( $i \in [n]$ ,  $y_{f_\star(i)}, y_{f^\star(i)} \in P_J$ ) at the categorization points of  $I$ , but they are not constrained on the open intervals formed by the categorization points of  $I$ .






Figure 5: Discrete derivatives of  $f_*$ ,  $f^*$ , and their interpretations

**Example 4.4.** Figure 6 depicts the discrete approximation of  $f$  (Figure 6 (a)), and the set of rough functions determined by it (Figure 6 (b)). One can observe that the discrete approximation constrains the rough functions at the categorizations points, but it does not say anything about how they behave on the open intervals.




Figure 6: Discrete approximation of  $f$  and its interpretation  $\mathcal{R}_f^{ds}$

Finding out rough derivatives of a set of rough functions, e.g.,  $\mathcal{R}_f^{ds}$ , may be approached, for instance, by differentiating both lower and upper discrete sequence approximations. Some special difficulties of this approach are the following:

(i) Of course,  $f_* \leq f^*$  does not imply  $f_*^{(1)} \leq f^{*(1)}$ .

At an  $i \in [n[$  may occur  $f_*^{(1)}(i) < f^{*(1)}(i)$ ,  $f_*^{(1)}(i) > f^{*(1)}(i)$ , or  $f_*^{(1)}(i) = f^{*(1)}(i)$ . Their interpretations are simple.  $f_*^{(1)}(i) < f^{*(1)}(i)$  means that the lower

approximation changes at  $i$  to a lesser extent than the upper approximation; whereas  $f_{\star}^{(1)}(i) > f^{\star(1)}(i)$  indicates that the lower approximation changes at  $i$  to a greater extent than the upper approximation. If  $f_{\star}^{(1)}(i) = f^{\star(1)}(i)$ , the lower and upper approximations change at  $i$  to the same extent.

(ii)  $f_{\star}^{(1)}(i) < 0$  and/or  $f^{\star(1)}(j) < 0$  for some  $i, j \in [n[$ .

A possible solution is the following. If  $f_{\star}^{(1)}(i) < 0$  for some  $i \in [n[$ , then  $\text{lm}f_{\star}^{(1)}$  may/should be extended to the extent necessary. This can be done, of course, analogously for  $f^{\star(1)}$ , too.

## 5. Concluding remarks

The paper, first, has presented four different rough approximation methods representing the rough real functions. The chances are that a definition of the rough differentiation should rely on one of these representations. In this paper, basically but not completely following Pawlak's method, the rough differentiation based on discrete sequence approximation has been considered only. Of course, additional differentiation definitions should also be studied based on the other representations which may be the subject of many subsequent papers in the future.

**Acknowledgements.** The author would like to thank the anonymous referees for their useful comments and suggestions.

## References

- [1] BARTLE, R., SHERBERT, D.: Introduction to Real Analysis. John Wiley & Sons, Incorporated, 4th edn. (2011)
- [2] BARTLE, R.G.: A Modern Theory of Integration, Graduate Studies in Mathematics, vol. 32. American Mathematical Society (2001)
- [3] CSAJBÓK, Z.E.: On the roughly continuous real functions. In: Mihálydeák, T., Min, F., Wang, G., Banerjee, M., Düntsch, I., Suraj, Z., Ciucci, D. (eds.) Rough Sets. pp. 52–65. Springer International Publishing, Cham (2019)
- [4] JORDAN, C.: Calculus of Finite Differences. Chelsea Publishing Company, New York, Second edn. (1950)
- [5] NAKAMURA, A., ROSENFELD, A.: Digital calculus. Information Sciences 98(1), 83–98 (1997)
- [6] PAWLAK, Z.: Rough sets. International Journal of Computer and Information Sciences 11(5), 341–356 (1982)
- [7] PAWLAK, Z.: Rough Sets: Theoretical Aspects of Reasoning about Data. Kluwer Academic Publishers, Dordrecht (1991)
- [8] PAWLAK, Z.: Rough calculus,. In: Proc. 2nd Annu. Joint Conf. on Inform. Sci., pp. 251–271. John Wiley & Sons, Inc., New York, NY, USA (1995)

- [9] PAWLAK, Z.: Rough real functions. vol. 50. Institute of Computer Science Report, Warsaw University of Technology, Warsaw (12 1994)
- [10] PAWLAK, Z.: On rough derivatives, rough integrals, and rough differential equations. vol. 41. Institute of Computer Science Report, Warsaw University of Technology, Nowowiejska 1519, 00-665 Warsaw, Poland (1995)
- [11] PAWLAK, Z.: On some issues connected with roughly continuous functions. vol. 21. Institute of Computer Science Report, Warsaw University of Technology, Warsaw (1995)
- [12] PAWLAK, Z.: Rough calculus. vol. 58. Institute of Computer Science Report, Warsaw University of Technology, Warsaw (1995)
- [13] PAWLAK, Z.: Rough sets, rough relations and rough functions. *Fundamenta Informaticae* 27(2/3), 103–108 (1996)
- [14] PAWLAK, Z.: Rough real functions and rough controllers. In: Lin, T., Cercone, N. (eds.) *Rough Sets and Data Mining: Analysis of Imprecise Data*. pp. 139–147. Kluwer Academic Publishers, Boston, MA (1997)
- [15] PAWLAK, Z., Skowron, A.: Rudiments of rough sets. *Information Sciences* 177(1), 3–27 (2007)
- [16] ROSENFELD, A.: ‘Continuous’ functions on digital pictures. *Pattern Recognition Letters* 4(3), 177–184 (1986)