

A Note on Parabolic and Linear One-Factorizations of the Complete Graph K_{p+1}

György Kiss^{1,2}, Gábor Korchmáros³, Federico Romaniello⁴ and Valentino Smaldore^{5,*}

¹Department of Geometry and HUN-REN-ELTE Geometric and Algebraic Combinatorics Research Group, Eötvös Loránd University, Pázmány s. 1/c, Budapest, 1117, Hungary

²FAMNIT, University of Primorska, Glagoljaška 8, Koper, 6000, Slovenia

³Dipartimento di Matematica, Informatica ed Economia, Università degli Studi della Basilicata, Viale dell'Ateneo Lucano 10, Potenza, 85100, Italy

⁴Dipartimento delle Culture Europee e del Mediterraneo, Università degli Studi della Basilicata, Via Lanera 20, Matera, 75100, Italy

⁵Dipartimento di Tecnica e Gestione dei Sistemi Industriali, Università degli Studi di Padova, Stradella S. Nicola 3, Vicenza, 361000, Italy

Abstract

One-factorizations of the complete graph K_n have wide applications, as an example they are often used for scheduling round-robin tournaments with n teams. In this note, we characterize parabolic and linear one-factorizations of complete graphs K_{p+1} , when p is an odd prime. This class of one-factorizations arises from the geometry of conics and lines in the affine plane $AG(2, p)$. We also include Magma computations for the cases $p \leq 19$.

Keywords

One-factorization, parabolas, affine lines

Declaration

This work is written in memory of Angelo Sonnino. His ideas on one-factorizations were very inspiring for this note. We hope that the present work provides something he would find interesting to read. The fourth author is also grateful for his mentorship during the writing of Master's thesis, in 2019.

1. Introduction

Some of the most important sport competitions have a final classification which depends on a round-robin phase. In such tournaments, each participant plays at least once against any other team, and the final classification considers all results. Then tournaments as *Serie A* or *NBA* need efficient algorithms to compute all the possible match schedules, see [1, 2]. For example, some tournament uses *Berger's algorithm* [3] (developed by the chess player *Johann Berger*) which divides the n players into two equal sides, from 1 to $\frac{n}{2}$ and from $\frac{n}{2} + 1$ to n ; starts from the first pairing $\{1, n\}, \{2, n-1\}, \dots, \{\frac{n}{2}, \frac{n}{2} + 1\}$; and

ends by giving some combinatorial argument to obtain all the other pairings. In our approach, we use one-factorizations of complete graphs. More precisely, a one-factorization of the complete graph K_n corresponds to a pairing in a round-robin tournament with n teams playing. In this note, we characterize parabolic and linear one-factorizations of complete graphs K_{p+1} , with p an odd prime number. In Section 2, we give preliminaries on graph theory and one-factorizations. In Sections 3 and 4, we give characterization results for parabolic and linear one-factorizations, where a one-factor is said to be *parabolic* or *linear* if it is represented by a parabola or a line, respectively. This note concludes with an Appendix containing computational results for the parabolic one-factorizations of K_{p+1} , $p = 13, 17, 19$, and the Magma code used by the authors.

2. Preliminaries

A graph $G = (V, E)$ is an incidence structure consisting of a set V of objects called vertices and a set E of object called edges. An edge $e \in E$ is denoted in the form $e = \{x, y\}$, where the vertices $x, y \in V$. Two vertices x and y connected by the edge $e = \{x, y\}$, are said to be adjacent. In what follows, neither multiple edges between the same pair of vertices nor loops, i.e. edges of the form $\{x, x\}$, are admitted. When every vertex has an equal number of edges incident to it, the graph is said to be regular. In particular, we will deal with a special class of regular graphs, in which each pair of vertices is connected by an edge. They are called complete graphs

ITAT'24: Information technologies – Applications and Theory, September 20–22, 2024, Drienica Čergovské vrchy, Slovakia

*Corresponding author.

✉ kissgy@cs.elte.hu (Gy. Kiss); gabor.korchmaros@unibas.it (G. Korchmáros); federico.romaniello@unibas.it (F. Romaniello); valentino.smaldore@unipd.it (V. Smaldore)

🆔 0000-0003-3312-9575 (Gy. Kiss); 0000-0002-2776-5754 (G. Korchmáros); 0000-0003-1166-3179 (F. Romaniello); 0000-0003-3433-3164 (V. Smaldore)

© 2024 Copyright for this paper by its authors. Use permitted under Creative Commons License Attribution 4.0 International (CC BY 4.0).



and are usually denoted by K_n , where n is the number of vertices. Let K_n be the complete graph with an even number n of vertices. A *one-factor* is a partition of its vertex set into $\frac{n}{2}$ disjoint edges. A *one-factorization* of K_n is a partition of its edge set into $n-1$ disjoint one-factors. For other notation or definitions on graphs not stated here, we refer the reader to [4]. Our approach to the problem of constructing one-factorizations of complete graphs is geometric, as in [5, 6, 7, 8, 9], and is based on techniques that have also been used for multigraphs, see [10, 11, 12, 13].

3. Parabolic one-factorizations

We adopt the same notation and terminology introduced in a paper by Korchmáros, Pace and Sonnino [5], and then used in subsequent paper by Kiss, Pace and Sonnino [6]. The reader may refer to [14] for notation and terminology on finite geometry not explicitly stated here. Let p be an odd prime. Fix a projective frame in $PG(2, p)$ with homogeneous coordinates $(X_0 : X_1 : X_2)$, and consider $PG(2, p)$ as $AG(2, p) \cup \ell_\infty$ where ℓ_∞ has equation $X_0 = 0$. As usual, the points of $AG(2, p)$ are written as (X, Y) with $X = \frac{X_1}{X_0}$ and $Y = \frac{X_2}{X_0}$. In $AG(2, p)$, let \mathcal{P}_a be the parabola with affine equation $Y = X^2 + a$, where a varies in \mathbb{Z}_p , and $V_\infty = (0 : 0 : 1)$ the point at infinity of the line $X_1 = 0$. We remark that, in the projective closure of $AG(2, p)$, any two parabolas \mathcal{P}_a and \mathcal{P}_b , with $a \neq b$, meet only at the point V_∞ . Let P_i^k denote the affine point $(i + \frac{k}{2}, i^2 + ik)$ and P_i^∞ be the point $(0 : 1 : 2i)$ on the line at infinity ℓ_∞ . The following result is easy to check:

Lemma 3.1. *For a fixed k , the points $P_0^k, P_1^k, \dots, P_{p-1}^k$ are on the parabola $\mathcal{P}_{-\frac{k^2}{4}}$.*

Definition 3.2. *Let $R = (u, v)$ be a point. The symptome of R is defined as $\mu_R = u^2 - v$.*

Let ℓ be a non-vertical line with equation $Y = mX + b$. The symptome of ℓ is defined as $\lambda_\ell = m^2 + 4b$.

We recall some basic properties of the parabolas \mathcal{P}_a . It is straightforward to check the following lemma:

Lemma 3.3. *Let $R = (u, v)$ be a point and $\ell: Y = mX + b$ be a non-vertical line. Then*

- R is an external (internal) point of \mathcal{P}_0 if and only if μ_R is a square (non-square) element in $\text{GF}(p)$.
- ℓ is a tangent (secant or external line) to \mathcal{P}_a if and only if $\lambda_\ell - 4a$ is 0 (a square or a non-square element in $\text{GF}(p)$).
- The pole of ℓ with respect to \mathcal{P}_0 is the point $L = (\frac{m}{2}, -b)$.

The vertices of the complete graph K_{p+1} correspond to the points of $\mathcal{P}_0 \cup V_\infty$, while the edges of K_{p+1} correspond to the points of type P_i^k , with $k = 1, 2, \dots, \frac{p-1}{2}, \infty$. Thus the set of edges of K_{p+1} corresponds to the set of points

$$\mathcal{E} = \left(\bigcup_{k=1}^{\frac{p-1}{2}} \mathcal{P}_{-\frac{k^2}{4}} \right) \cup (\ell_\infty \setminus V_\infty).$$

These points are called *external* w.r.t. \mathcal{P}_0 .

Definition 3.4. *A one-factor represented by a parabola \mathcal{P}_a is a set of $\frac{p-1}{2}$ points of type P_j^k on \mathcal{P}_a , together with a suitable point at infinity. A one-factor so defined is referred to as a parabolic one-factor.*

Definition 3.5. *A one-factor represented by a secant line ℓ of \mathcal{P}_0 is a set consisting of $\frac{p-1}{2}$ points of \mathcal{E} on ℓ , plus the pole of ℓ with respect to \mathcal{P}_0 .*

A one-factor represented by an external line ℓ of \mathcal{P}_0 is a set consisting of $\frac{p+1}{2}$ points of \mathcal{E} on ℓ .

Definition 3.6. *A one-factorization of K_{p+1} is called a parabolic one-factorization if $p-1$ of its one-factors are represented by parabolas and one of its one-factors is represented by a line.*

In [6] the authors proved the existence of an infinite family of parabolic one-factorization.

Theorem 3.7. [6, Theorem 3.4] *Let p be an odd prime. Then the complete graph K_{p+1} has a parabolic one-factorization.*

Proof. The proof is constructive. The set

$$F_0 = \left\{ P_{-\frac{k}{2}}^k : k = 1, 2, \dots, \frac{p-1}{2} \right\} \cup \{P_0^\infty\}$$

is a one-factor represented by the line secant line of \mathcal{P}_0 of equation $X = 0$, and P_0 is the pole. Now we define the sets

$$G_k = \left\{ P_{\frac{k}{2}+2jk}^k : j = 0, 1, \dots, \frac{p-3}{2} \right\} \cup \left\{ P_{-\frac{k}{2}}^\infty \right\},$$

$$H_k = \left\{ P_{\frac{k}{2}+(2j+1)k}^k : j = 0, 1, \dots, \frac{p-3}{2} \right\} \cup \left\{ P_{\frac{k}{2}}^\infty \right\}.$$

By Lemma 3.1 $G_k \setminus \{P_{-\frac{k}{2}}^\infty\}$ and $H_k \setminus \{P_{\frac{k}{2}}^\infty\}$ are disjoint subsets of the parabola $\mathcal{P}_{-\frac{k^2}{4}}$, and both G_k and H_k are one-factors represented by the parabola $\mathcal{P}_{-\frac{k^2}{4}}$. \square

Definition 3.8. *A one-factorization of K_{p+1} is called an almost parabolic one-factorization if at least one of its one-factors is represented by $\mathcal{P}_{-\frac{k^2}{4}}$ for all $k \in \{1, 2, \dots, \frac{p-1}{2}\}$, and all other of its one-factors are represented by lines.*

For $p < 11$ exhaustive computer search shows that each almost parabolic one-factorization of K_{p+1} is parabolic. In the rest of the paper we will assume $p \geq 11$.

Lemma 3.9. *The number of one-factors represented by lines in an almost parabolic one-factorization is either one, or at least $\lceil \frac{p+1}{4} \rceil$.*

Proof. If more than one one-factors are represented by lines, then there exist parabolas $\mathcal{P}_{-\frac{k^2}{4}}$ which represent only one one-factor. Hence $p - \frac{p-1}{2} = \frac{p+1}{2}$ of its points are covered by the lines represented the other one-factors. Any line meets $\mathcal{P}_{-\frac{k^2}{4}}$ in at most two points, so we need at least $\lceil \frac{p+1}{4} \rceil$ lines to cover these points. \square

The following Lemma is a straightforward corollary of [6, Theorem 3.5].

Lemma 3.10. *Let \mathcal{L} denote the set of points of type P_i^k belonging to the one-factors represented by lines in an almost parabolic one-factorization. Suppose that a one-factor is represented by the line $X = 0$. Then for each $k \in \{1, 2, \dots, \frac{p-1}{2}\}$ either $\mathcal{L} \subset G_k$ or $\mathcal{L} \subset H_k$.*

Proposition 3.11. *An almost parabolic one-factorization contains at most two one-factors which are represented by vertical lines.*

Proof. Suppose to the contrary that it contains at least three one-factors which are represented by vertical lines. We may assume that the equations of the corresponding lines are $X = 0$, $X = u$, $X = v$ and $u - v = k' \in \{1, 2, \dots, \frac{p-1}{2}\}$.

We also may assume that the line $X = v$ intersects $G_{k'}$. Then $v = \frac{k'}{2} + 2jk'$ where $j = 0, 1, \dots, \frac{p-3}{2}$. Hence

$$u = v + k' = \frac{k'}{2} + (2j+1)k' \text{ where } j = 0, 1, \dots, \frac{p-3}{2}.$$

So the line $X = u$ intersects $H_{k'}$ contradicting Lemma 3.10. \square

Lemma 3.12. *In $\text{GF}(p)$ let T denote the set*

$$T = \left\{ (4j^2 + 4j)k^2 : j = 0, 1, \dots, \frac{p-1}{2}, \right. \\ \left. k = 1, 2, \dots, \frac{p-1}{2} \right\}.$$

Then $|T| = p$.

Proof. First, we show that the cardinality of the set

$$U = \left\{ 4j^2 + 4j : j = 0, 1, \dots, \frac{p-1}{2} \right\}$$

is $\frac{p+1}{2}$. If $4j_1^2 + 4j_1 = 4j_2^2 + 4j_2$, then $(j_1 - j_2)(j_1 + j_2 + 1) = 0$. So for $j_1 \neq j_2$ we have $4j_1^2 + 4j_1 = 4j_2^2 + 4j_2$, because the sum in the second factor is not 0.

The set U obviously contains 0. Moreover, we claim that U contains both square and non-square elements. Suppose to the contrary, that all of the non-zero products $j(j+1)$ are either squares or non-squares. If 2 is a square, then $1 \cdot 2$ is a square, hence $2 \cdot 3$ is also a square which implies that 3 is a square. In the same way, step by step, we get that all of the elements $4, \dots, \frac{p-1}{2}, \frac{p+1}{2}$ are squares. Hence there are at least $\frac{p+1}{2}$ square elements, a contradiction. If 2 is a non-square, then $1 \cdot 2$ is a non-square, hence $2 \cdot 3$ is also a non-square, so 3 is a square. But 4 is also a square, hence $3 \cdot 4$ is a square, a contradiction again.

Now we show that the set

$$S = \left\{ k^2 : k = 1, 2, \dots, \frac{p-1}{2} \right\}$$

equals to the set of square elements of $\text{GF}(p)$. The cardinality of S is $\frac{p-1}{2}$, because $k_1^2 = k_2^2$ implies $(k_1 - k_2)(k_1 + k_2) = 0$, and the second factor is never 0, because $k_1 + k_2 \leq p - 1$.

Choose elements $u_1, u_2 \in U$ such that u_1 is a square and u_2 is a non-square. Then $u_1U \cap u_2U = \emptyset$, hence

$$|\{0\} \cup u_1U \cup u_2U| = 1 + \frac{p-1}{2} + \frac{p-1}{2} = p,$$

the statement is proved. \square

Proposition 3.13. *If an almost parabolic one-factorization contains a one-factors which is represented by a vertical line, then it cannot contain one-factors represented by non-vertical lines.*

Proof. We may assume that the equation of the corresponding vertical line is $X = 0$. Suppose to the contrary that it contains the line $\ell: Y = mx + b$. We claim that there exists at least one $k \in \{1, 2, \dots, \frac{p-1}{2}\}$ such that $\ell \cap G_k \neq \emptyset \neq \ell \cap H_k$. By Lemma 3.12, there exist j and k such that $\lambda_\ell = (4j^2 + 4j)k^2$. Then

$$\lambda_\ell + k^2 = (4j^2 + 4j)k^2 + k^2 = ((2j+1)k)^2,$$

so, by Lemma 3.3, ℓ intersects $\mathcal{P}_{-\frac{k^2}{4}}$ and the difference of the first coordinates of the two intersections is $(2j+1)k$. Hence one of the two points belongs to G_k and the other one belongs to H_k . The statement follows from Lemma 3.10. \square

The main result of this section is the following theorem, which derives from Propositions 3.9, 3.11 and 3.13.

Theorem 3.14. *Let p be an odd prime. If an almost parabolic one-factorization \mathcal{F} of K_{p+1} contains a one-factor which is represented by a vertical line, then the one-factorization is parabolic.*

Proof. For $p < 11$ the statement follows from an exhaustive computer search, see [6].

If $p \geq 11$, then $\lceil \frac{p+1}{4} \rceil \geq 3$. Hence, by Lemma 3.9, \mathcal{F} contains either one, or at least three one-factors that are represented by lines. In the former case, we are done. The latter case leads to a contradiction, since, by Lemma 3.11, the number of vertical lines is at most two, and by Lemma 3.13, there is no non-vertical line among the lines representing the one-factors. \square

4. Linear one-factorizations

In this section, we consider one-factorizations whose all one-factors are represented by lines.

Let Ω be the set of points of an irreducible conic in $PG(2, q)$ with $q \geq 5$ odd.

Theorem 4.1. *Let K_{q+1} be a one-factorization on Ω whose one-factors are represented by lines. Then some of those lines are a chord of Ω .*

Proof. Let K_{q+1} be represented by the lines ℓ_1, \dots, ℓ_q . Assume on the contrary that each those lines is an external line to Ω . Let L_1, \dots, L_q be the poles of ℓ_1, \dots, ℓ_q with respect to the orthogonal polarity π associated with Ω . For $i = 1, \dots, q$, let φ_i be the involutory perspectivity with center L_i and axis ℓ_i which preserves Ω . Let $G \cong PGL(2, q)$ be the orthogonal group of Ω , i.e. the subgroup of $PGL(3, q)$ which commutes with π . Then G preserves Ω and acts on its points as $PGL(2, q)$ on the projective line over \mathbb{F}_q . Furthermore, $\varphi_i \in G$. Let \mathcal{F} be the set consisting of $\varphi_i, i = 1, \dots, q$ together with the identity of G . Since K_{q+1} is a one-factorization, for any two points $P, Q \in \Omega$ there exists a unique $\varphi \in \mathcal{F}$ such that $\varphi(P) = Q$. Therefore, \mathcal{F} is a sharply transitive permutation set on Ω containing the identity. From the classification of sharply transitive subsets of $PGL(2, q)$ [15], it turns out \mathcal{F} is a subgroup of $PGL(2, q)$ of order $q + 1$. On the other hand, from Dickson's classification of subgroups of $PGL(2, q)$, the subgroups of $PGL(2, q)$ entirely consisting of involutions together with the identity, have order either 2 or 4. But then $q \leq 3$, a contradiction. \square

We conclude by reporting a conjecture that is supported by computer-aided searches. With the aid of Magma [16] we verified that the conjecture holds for $p \leq 23$.

Conjecture 4.2. [6, Conjecture 3.6] *Let $p > 7$ be an odd prime, \mathcal{F} be a one-factorization of the complete graph K_{p+1} such that each one-factor of \mathcal{F} is either represented by a line or a parabola. Then \mathcal{F} is either a parabolic one-factorization or each one-factor of \mathcal{F} is represented by a line.*

Acknowledgments

The research of the first author was partially supported by the Hungarian National Research, Development and Innovation Office OTKA grant no. SNN 132625. The research of the fourth author was partially supported by the Italian National Group for Algebraic and Geometric Structures and their Applications (GNSAGA - INdAM) and by the INdAM - GNSAGA Project *Tensors over finite fields and their applications*, number E53C23001670001.

References

- [1] Lega Calcio Serie A, 2010. URL: <https://www.legaseriea.it/it>.
- [2] National Basketball Association, 1996. URL: <https://www.nba.com/>.
- [3] J. Berger, Schach-Jahrbuch für 1892/93, Verlag Veit, Leipzig, 1893.
- [4] J. A. Bondy, U. S. R. Murty, Graph theory with applications, volume 29, Macmillan London, 1976.
- [5] G. Korchmáros, N. Pace, A. Sonnino, One-factorisations of complete graphs arising from ovals in finite planes, J. Combin. Theory Ser. A 160 (2018) 62–83. doi:10.1016/j.jcta.2018.06.006.
- [6] Gy. Kiss, N. Pace, A. Sonnino, One-factorizations of the complete graph K_{p+1} arising from parabolas, Theory Appl. Graphs 9 (2022) 2. doi:10.20429/tag.2022.090202.
- [7] Gy. Kiss, N. Pace, A. Sonnino, On circular-linear one-factorizations of the complete graph K_n , Discrete Math. 342 (2019). doi:10.1016/j.disc.2019.111622.
- [8] N. Pace, A. Sonnino, One-factorisations of complete graphs arising from hyperbolae in the desarguesian affine plane, J. Geom. 110 (2019) 19 pp. doi:10.1007/s00022-019-0470-6.
- [9] N. Pace, A. Sonnino, One-factorisations of complete graphs constructed in desarguesian planes of certain odd square orders, Electron. J. Combin. 27 (2020) 19 pp. doi:10.37236/8378.
- [10] Gy. Kiss, One-factorizations of complete multigraphs and quadrics in $PG(n, q)$, J. Combin. Des. 10 (2002) 139–143. doi:10.1002/jcd.997.
- [11] Gy. Kiss, C. Rubio-Montiel, A note on m -factorizations of complete multigraphs arising from designs, Ars Math. Contemp. 8 (2015) 163–175. doi:10.26493/1855-3974.537.6bf.
- [12] G. Korchmáros, A. Siciliano, A. Sonnino, 1-factorizations of complete multigraphs arising from finite geometry, J. Combin. Theory Ser. A 93 (2001) 385–390. doi:10.1006/jcta.2000.3115.
- [13] A. Sonnino, One-factorizations of complete multigraphs arising from maximal (k, n) -arcs in

- $PG(2, 2^h)$, Discrete Math. 231 (2001) 447–451. doi:10.1016/S0012-365X(00)00337-X.
- [14] Gy. Kiss, T. Szőnyi, Finite Geometries, CRC Press, Taylor & Francis Group, Boca Raton, FL, 2019.
- [15] S. Eberhard, Sharply transitive sets in $pgl_2(k)$, Adv. Geom. 21 (2021) 611–612. doi:10.1515/advgeom-2021-0029.
- [16] W. Bosma, J. Cannon, C. Playoust, The magma algebra system. i. the user language., J. Symbolic Comput. 24 (1997) 235–265. doi:10.1006/j.sco.1996.0125.

A. Tables

Corollary 3.14 states that almost parabolic one-factorization of K_{p+1} , p odd prime, containing a vertical line are parabolic. In this Appendix, we report examples of such parabolic one-factorizations of the complete graphs K_{14} , K_{18} and K_{20} , found by computations on Magma, [16]. K_4 admits only 1 one-factorization, and are well-known the 6 examples of 1-factorizations of K_6 . We refer to [6] for the cases K_8 and K_{12} .

A.1. $p = 13$

- $F_0 : \{P_{0,\infty} : (0 : 1 : 0), P_{1,6} : (1 : 0 : 3), P_{2,12} : (1 : 0 : 12), P_{3,5} : (1 : 0 : 1), P_{4,11} : (1 : 0 : 9), P_{5,4} : (1 : 0 : 10), P_{6,10} : (1 : 0 : 4)\}$
- $G_1 : \{P_{6,\infty} : (0 : 1 : 12), P_{1,7} : (1 : 1 : 4), P_{1,9} : (1 : 3 : 12), P_{1,11} : (1 : 5 : 2), P_{1,0} : (1 : 7 : 0), P_{1,2} : (1 : 9 : 6), P_{1,4} : (1 : 11 : 7)\}$
- $H_1 : \{P_{7,\infty} : (0 : 1 : 1), P_{1,8} : (1 : 2 : 7), P_{1,10} : (1 : 4 : 6), P_{1,12} : (1 : 6 : 0), P_{1,1} : (1 : 8 : 2), P_{1,3} : (1 : 10 : 12), P_{1,5} : (1 : 12 : 4)\}$
- $G_2 : \{P_{12,\infty} : (0 : 1 : 11), P_{2,1} : (1 : 2 : 3), P_{2,5} : (1 : 6 : 9), P_{2,9} : (1 : 10 : 8), P_{2,0} : (1 : 1 : 0), P_{2,4} : (1 : 5 : 11), P_{2,8} : (1 : 9 : 2)\}$
- $H_2 : \{P_{1,\infty} : (0 : 1 : 2), P_{2,3} : (1 : 4 : 2), P_{2,7} : (1 : 8 : 11), P_{2,11} : (1 : 12 : 0), P_{2,2} : (1 : 3 : 8), P_{2,6} : (1 : 7 : 9), P_{2,10} : (1 : 11 : 3)\}$
- $G_3 : \{P_{5,\infty} : (0 : 1 : 10), P_{3,8} : (1 : 3 : 10), P_{3,1} : (1 : 9 : 4), P_{3,7} : (1 : 2 : 5), P_{3,0} : (1 : 8 : 0), P_{3,6} : (1 : 1 : 2), P_{3,12} : (1 : 7 : 11)\}$
- $H_3 : \{P_{8,\infty} : (0 : 1 : 3), P_{3,11} : (1 : 6 : 11), P_{3,4} : (1 : 12 : 2), P_{3,10} : (1 : 5 : 0), P_{3,3} : (1 : 11 : 5), P_{3,9} : (1 : 4 : 4), P_{3,2} : (1 : 10 : 10)\}$
- $G_4 : \{P_{11,\infty} : (0 : 1 : 9), P_{4,2} : (1 : 4 : 12), P_{4,10} : (1 : 12 : 10), P_{4,5} : (1 : 7 : 6), P_{4,0} : (1 : 2 : 0), P_{4,8} : (1 : 10 : 5), P_{4,3} : (1 : 5 : 8)\}$
- $H_4 : \{P_{2,\infty} : (0 : 1 : 4), P_{4,6} : (1 : 8 : 8), P_{4,1} : (1 : 3 : 5),$

- $P_{4,9} : (1 : 11 : 0), P_{4,4} : (1 : 6 : 6), P_{4,12} : (1 : 1 : 10), P_{4,7} : (1 : 9 : 12)\}$
- $G_5 : \{P_{4,\infty} : (0 : 1 : 8), P_{5,9} : (1 : 5 : 9), P_{5,6} : (1 : 2 : 1), P_{5,3} : (1 : 12 : 11), P_{5,0} : (1 : 9 : 0), P_{5,10} : (1 : 6 : 7), P_{5,7} : (1 : 3 : 6)\}$
- $H_5 : \{P_{9,\infty} : (0 : 1 : 5), P_{5,1} : (1 : 10 : 6), P_{5,11} : (1 : 7 : 7), P_{5,8} : (1 : 4 : 0), P_{5,5} : (1 : 1 : 11), P_{5,2} : (1 : 11 : 1), P_{5,12} : (1 : 8 : 9)\}$
- $G_6 : \{P_{10,\infty} : (0 : 1 : 7), P_{6,3} : (1 : 6 : 1), P_{6,2} : (1 : 5 : 3), P_{6,1} : (1 : 4 : 7), P_{6,0} : (1 : 3 : 0), P_{6,12} : (1 : 2 : 8), P_{6,11} : (1 : 1 : 5)\}$
- $H_6 : \{P_{3,\infty} : (0 : 1 : 6), P_{6,9} : (1 : 12 : 5), P_{6,8} : (1 : 11 : 8), P_{6,7} : (1 : 10 : 0), P_{6,6} : (1 : 9 : 7), P_{6,5} : (1 : 8 : 3), P_{6,4} : (1 : 7 : 1)\}$

A.2. $p = 17$

- $F_0 : \{P_{0,\infty} : (0 : 1 : 0), P_{1,8} : (1 : 0 : 4), P_{2,16} : (1 : 0 : 16), P_{3,7} : (1 : 0 : 2), P_{4,15} : (1 : 0 : 13), P_{5,6} : (1 : 0 : 15), P_{6,14} : (1 : 0 : 8), P_{7,5} : (1 : 0 : 9), P_{8,13} : (1 : 0 : 1)\}$
- $G_1 : \{P_{8,\infty} : (0 : 1 : 16), P_{1,9} : (1 : 1 : 5), P_{1,11} : (1 : 3 : 13), P_{1,13} : (1 : 5 : 12), P_{1,15} : (1 : 7 : 2), P_{1,0} : (1 : 9 : 0), P_{1,2} : (1 : 11 : 6), P_{1,4} : (1 : 13 : 3), P_{1,6} : (1 : 15 : 8)\}$
- $H_1 : \{P_{9,\infty} : (0 : 1 : 1), P_{1,10} : (1 : 2 : 8), P_{1,12} : (1 : 4 : 3), P_{1,14} : (1 : 6 : 6), P_{1,16} : (1 : 8 : 0), P_{1,1} : (1 : 10 : 2), P_{1,3} : (1 : 12 : 12), P_{1,5} : (1 : 14 : 13), P_{1,7} : (1 : 16 : 5)\}$
- $G_2 : \{P_{16,\infty} : (0 : 1 : 15), P_{2,1} : (1 : 2 : 3), P_{2,5} : (1 : 6 : 1), P_{2,9} : (1 : 10 : 14), P_{2,13} : (1 : 14 : 8), P_{2,0} : (1 : 1 : 0), P_{2,4} : (1 : 5 : 7), P_{2,8} : (1 : 9 : 12), P_{2,12} : (1 : 13 : 15)\}$
- $H_2 : \{P_{1,\infty} : (0 : 1 : 2), P_{2,3} : (1 : 4 : 15), P_{2,7} : (1 : 8 : 12), P_{2,11} : (1 : 12 : 7), P_{2,15} : (1 : 16 : 0), P_{2,2} : (1 : 3 : 8), P_{2,6} : (1 : 7 : 14), P_{2,10} : (1 : 11 : 1), P_{2,14} : (1 : 15 : 3)\}$
- $G_3 : \{P_{7,\infty} : (0 : 1 : 14), P_{3,10} : (1 : 3 : 11), P_{3,16} : (1 : 9 : 15), P_{3,5} : (1 : 15 : 6), P_{3,11} : (1 : 4 : 1), P_{3,0} : (1 : 10 : 0), P_{3,6} : (1 : 16 : 3), P_{3,12} : (1 : 5 : 10), P_{3,1} : (1 : 11 : 4)\}$
- $H_3 : \{P_{10,\infty} : (0 : 1 : 3), P_{3,13} : (1 : 6 : 4), P_{3,2} : (1 : 12 : 10), P_{3,8} : (1 : 1 : 3), P_{3,14} : (1 : 7 : 0), P_{3,3} : (1 : 13 : 1), P_{3,9} : (1 : 2 : 6), P_{3,15} : (1 : 8 : 15), P_{3,4} : (1 : 14 : 11)\}$
- $G_4 : \{P_{15,\infty} : (0 : 1 : 13), P_{4,2} : (1 : 4 : 12), P_{4,10} : (1 : 12 : 4), P_{4,1} : (1 : 3 : 5), P_{4,9} : (1 : 11 : 15), P_{4,0} : (1 : 2 : 0), P_{4,8} : (1 : 10 : 11), P_{4,16} : (1 : 1 : 14), P_{4,7} : (1 : 9 : 9)\}$
- $H_4 : \{P_{2,\infty} : (0 : 1 : 4), P_{4,6} : (1 : 8 : 9), P_{4,14} : (1 : 16 : 14), P_{4,5} : (1 : 7 : 11), P_{4,13} : (1 : 15 : 0), P_{4,4} : (1 : 6 : 15), P_{4,12} : (1 : 14 : 5), P_{4,3} : (1 : 5 : 4), P_{4,11} : (1 : 13 : 12)\}$
- $G_5 : \{P_{6,\infty} : (0 : 1 : 12), P_{5,11} : (1 : 5 : 6), P_{5,4} : (1 : 15 : 2), P_{5,14} : (1 : 8 : 11), P_{5,7} : (1 : 1 : 16), P_{5,0} : (1 : 11 : 0), P_{5,10} : (1 : 4 : 14), P_{5,3} : (1 : 14 : 7), P_{5,13} : (1 : 7 : 13)\}$
- $H_5 : \{P_{11,\infty} : (0 : 1 : 5), P_{5,16} : (1 : 10 : 13), P_{5,9} : (1 : 3 : 7), P_{5,2} : (1 : 13 : 14), P_{5,12} : (1 : 6 : 0), P_{5,5} : (1 : 16 : 16),$

$$\begin{aligned}
& P_{5,15} : (1 : 9 : 11), P_{5,8} : (1 : 2 : 2), P_{5,1} : (1 : 12 : 6) \} & P_{3,5} : (1 : 16 : 2) \} \\
G_6 : \{ P_{14,\infty} : (0 : 1 : 11), P_{6,3} : (1 : 6 : 10), P_{6,15} : (1 : 1 : 9) & G_4 : \{ P_{17,\infty} : (0 : 1 : 15), P_{4,2} : (1 : 4 : 12), P_{4,10} : (1 : 12 : 7), \\
& P_{6,10} : (1 : 13 : 7), P_{6,5} : (1 : 8 : 4), P_{6,0} : (1 : 3 : 0), & P_{4,18} : (1 : 1 : 16), P_{4,7} : (1 : 9 : 1), P_{4,15} : (1 : 17 : 0), \\
& P_{6,12} : (1 : 15 : 12), P_{6,7} : (1 : 10 : 6), P_{6,2} : (1 : 5 : 16) \} & P_{4,4} : (1 : 6 : 13), P_{4,12} : (1 : 14 : 2), P_{4,1} : (1 : 3 : 5), \\
H_6 : \{ P_{3,\infty} : (0 : 1 : 6), P_{6,9} : (1 : 12 : 16), P_{6,4} : (1 : 7 : 6), & P_{4,9} : (1 : 11 : 3) \} \\
& P_{6,16} : (1 : 2 : 12), P_{6,11} : (1 : 14 : 0), P_{6,6} : (1 : 9 : 4), H_4 : \{ P_{2,\infty} : (0 : 1 : 4), P_{4,6} : (1 : 8 : 3), P_{4,14} : (1 : 16 : 5), \\
& P_{6,1} : (1 : 4 : 7), P_{6,13} : (1 : 16 : 9), P_{6,8} : (1 : 11 : 10) \} & P_{4,3} : (1 : 5 : 2), P_{4,11} : (1 : 13 : 13), P_{4,0} : (1 : 2 : 0), \\
G_7 : \{ P_{5,\infty} : (0 : 1 : 1PZ0), P_{7,12} : (1 : 7 : 7), P_{7,9} : (1 : 4 : 8), & P_{4,8} : (1 : 10 : 1), P_{4,16} : (1 : 18 : 16), P_{4,5} : (1 : 7 : 7), \\
& P_{7,6} : (1 : 1 : 10), P_{7,3} : (1 : 15 : 13), P_{7,0} : (1 : 12 : 0), & P_{4,13} : (1 : 15 : 12) \} \\
& P_{7,14} : (1 : 9 : 5), P_{7,11} : (1 : 6 : 11), P_{7,8} : (1 : 3 : 1) \} G_5 : \{ P_{7,\infty} : (0 : 1 : 14), P_{5,12} : (1 : 5 : 14), P_{5,3} : (1 : 15 : 5), \\
H_7 : \{ P_{12,\infty} : (0 : 1 : 7), P_{7,2} : (1 : 14 : 1), P_{7,16} : (1 : 11 : 11), & P_{5,13} : (1 : 6 : 6), P_{5,4} : (1 : 16 : 17), P_{5,14} : (1 : 7 : 0), \\
& P_{7,13} : (1 : 8 : 5), P_{7,10} : (1 : 5 : 0), P_{7,7} : (1 : 2 : 13), & P_{5,5} : (1 : 17 : 12), P_{5,15} : (1 : 8 : 15), P_{5,6} : (1 : 18 : 9), \\
& P_{7,4} : (1 : 16 : 10), P_{7,1} : (1 : 13 : 8), P_{7,15} : (1 : 10 : 7) \} & P_{5,16} : (1 : 9 : 13) \} \\
G_8 : \{ P_{13,\infty} : (0 : 1 : 9), P_{8,4} : (1 : 8 : 14), P_{8,3} : (1 : 7 : 16), H_5 : \{ P_{12,\infty} : (0 : 1 : 5), P_{5,17} : (1 : 10 : 13), P_{5,8} : (1 : 1 : 9), \\
& P_{8,2} : (1 : 6 : 3), P_{8,1} : (1 : 5 : 9), P_{8,0} : (1 : 4 : 0), & P_{5,18} : (1 : 11 : 15), P_{5,9} : (1 : 2 : 12), P_{5,0} : (1 : 12 : 0), \\
& P_{8,16} : (1 : 3 : 10), P_{8,15} : (1 : 2 : 5), P_{8,14} : (1 : 1 : 2) \} & P_{5,10} : (1 : 3 : 17), P_{5,1} : (1 : 13 : 6), P_{5,11} : (1 : 4 : 5), \\
H_8 : \{ P_{4,\infty} : (0 : 1 : 8), P_{8,12} : (1 : 16 : 2), P_{8,11} : (1 : 15 : 5), & P_{5,2} : (1 : 14 : 14) \} \\
& P_{8,10} : (1 : 14 : 10), P_{8,9} : (1 : 13 : 0), P_{8,8} : (1 : 12 : 9) G_6 : \{ P_{16,\infty} : (0 : 1 : 13), P_{6,3} : (1 : 6 : 8), P_{6,15} : (1 : 18 : 11), \\
& P_{8,7} : (1 : 11 : 3), P_{8,6} : (1 : 10 : 16), P_{8,5} : (1 : 9 : 14) \} & P_{6,8} : (1 : 11 : 17), P_{6,1} : (1 : 4 : 7), P_{6,13} : (1 : 16 : 0), \\
& & P_{6,6} : (1 : 9 : 15), P_{6,18} : (1 : 2 : 14), P_{6,11} : (1 : 14 : 16), \\
& & P_{6,4} : (1 : 7 : 2) \}
\end{aligned}$$

A.3. $p = 19$

$$\begin{aligned}
& & H_6 : \{ P_{3,\infty} : (0 : 1 : 6), P_{6,9} : (1 : 12 : 2), P_{6,2} : (1 : 5 : 16), \\
& & P_{6,14} : (1 : 17 : 14), P_{6,7} : (1 : 10 : 15), P_{6,0} : (1 : 3 : 0), \\
F_0 : \{ P_{0,\infty} : (0 : 1 : 0), P_{1,9} : (1 : 0 : 14), P_{2,18} : (1 : 0 : 18), & P_{6,12} : (1 : 15 : 7), P_{6,5} : (1 : 8 : 17), P_{6,17} : (1 : 1 : 11), \\
& P_{3,8} : (1 : 0 : 12), P_{4,17} : (1 : 0 : 15), P_{5,7} : (1 : 0 : 8), & P_{6,10} : (1 : 13 : 8) \} \\
& P_{6,16} : (1 : 0 : 10), P_{7,6} : (1 : 0 : 2), P_{8,15} : (1 : 0 : 3), & G_7 : \{ P_{6,\infty} : (0 : 1 : 12), P_{7,13} : (1 : 7 : 13), P_{7,8} : (1 : 2 : 6), \\
& P_{9,5} : (1 : 0 : 13) \} & P_{7,3} : (1 : 16 : 11), P_{7,17} : (1 : 11 : 9), P_{7,12} : (1 : 6 : 0), \\
G_1 : \{ P_{9,\infty} : (0 : 1 : 18), P_{1,10} : (1 : 1 : 15), P_{1,12} : (1 : 3 : 4), & P_{7,7} : (1 : 1 : 3), P_{7,2} : (1 : 15 : 18), P_{7,16} : (1 : 10 : 7), \\
& P_{1,14} : (1 : 5 : 1), P_{1,16} : (1 : 7 : 6), P_{1,18} : (1 : 9 : 0), & P_{7,11} : (1 : 5 : 8) \} \\
& P_{1,1} : (1 : 11 : 2), P_{1,3} : (1 : 13 : 12), P_{1,5} : (1 : 15 : 11) & H_7 : \{ P_{13,\infty} : (0 : 1 : 7), P_{7,1} : (1 : 14 : 8), P_{7,15} : (1 : 9 : 7), \\
& P_{1,7} : (1 : 17 : 18) \} & P_{7,10} : (1 : 4 : 18), P_{7,5} : (1 : 18 : 3), P_{7,0} : (1 : 13 : 0), \\
H_1 : \{ P_{10,\infty} : (0 : 1 : 1), P_{1,11} : (1 : 2 : 18), P_{1,13} : (1 : 4 : 11), & P_{7,14} : (1 : 8 : 9), P_{7,9} : (1 : 3 : 11), P_{7,4} : (1 : 17 : 6), \\
& P_{1,15} : (1 : 6 : 12), P_{1,17} : (1 : 8 : 2), P_{1,0} : (1 : 10 : 0), & P_{7,18} : (1 : 12 : 13) \} \\
& P_{1,2} : (1 : 12 : 6), P_{1,4} : (1 : 14 : 1), P_{1,6} : (1 : 16 : 4), & G_8 : \{ P_{15,\infty} : (0 : 1 : 11), P_{8,4} : (1 : 8 : 10), P_{8,1} : (1 : 5 : 9), \\
& P_{1,8} : (1 : 18 : 15) \} & P_{8,17} : (1 : 2 : 7), P_{8,14} : (1 : 18 : 4), P_{8,11} : (1 : 15 : 0), \\
G_2 : \{ P_{18,\infty} : (0 : 1 : 17), P_{2,1} : (1 : 2 : 3), P_{2,5} : (1 : 6 : 16), & P_{8,8} : (1 : 12 : 14), P_{8,5} : (1 : 9 : 8), P_{8,2} : (1 : 6 : 1), \\
& P_{2,9} : (1 : 10 : 4), P_{2,13} : (1 : 14 : 5), P_{2,17} : (1 : 18 : 0), & P_{8,18} : (1 : 3 : 12) \} \\
& P_{2,2} : (1 : 3 : 8), P_{2,6} : (1 : 7 : 10), P_{2,10} : (1 : 11 : 6), & H_8 : \{ P_{4,\infty} : (0 : 1 : 8), P_{8,12} : (1 : 16 : 12), P_{8,9} : (1 : 13 : 1), \\
& P_{2,14} : (1 : 15 : 15) \} & P_{8,6} : (1 : 10 : 8), P_{8,3} : (1 : 7 : 14), P_{8,0} : (1 : 4 : 0), \\
H_2 : \{ P_{1,\infty} : (0 : 1 : 2), P_{2,3} : (1 : 4 : 15), P_{2,7} : (1 : 8 : 6), & P_{8,16} : (1 : 1 : 4), P_{8,13} : (1 : 17 : 7), P_{8,10} : (1 : 14 : 9), \\
& P_{2,11} : (1 : 12 : 10), P_{2,15} : (1 : 16 : 8), P_{2,0} : (1 : 1 : 0), & P_{8,7} : (1 : 11 : 10) \} \\
& P_{2,4} : (1 : 5 : 5), P_{2,8} : (1 : 9 : 4), P_{2,12} : (1 : 13 : 16), & G_9 : \{ P_{5,\infty} : (0 : 1 : 10), P_{9,14} : (1 : 9 : 18), P_{9,13} : (1 : 8 : 1), \\
& P_{2,16} : (1 : 17 : 3) \} & P_{9,12} : (1 : 7 : 5), P_{9,11} : (1 : 6 : 11), P_{9,10} : (1 : 5 : 0), \\
G_3 : \{ P_{8,\infty} : (0 : 1 : 16), P_{3,11} : (1 : 3 : 2), P_{3,17} : (1 : 9 : 17), & P_{9,9} : (1 : 4 : 10), P_{9,8} : (1 : 3 : 3), P_{9,7} : (1 : 2 : 17), \\
& P_{3,4} : (1 : 15 : 9), P_{3,10} : (1 : 2 : 16), P_{3,16} : (1 : 8 : 0), & P_{9,6} : (1 : 1 : 14) \} \\
& P_{3,3} : (1 : 14 : 18), P_{3,9} : (1 : 1 : 13), P_{3,15} : (1 : 7 : 4), & H_9 : \{ P_{14,\infty} : (0 : 1 : 9), P_{9,4} : (1 : 18 : 14), P_{9,3} : (1 : 17 : 17), \\
& P_{3,2} : (1 : 13 : 10) \} & P_{9,2} : (1 : 16 : 3), P_{9,1} : (1 : 15 : 10), P_{9,0} : (1 : 14 : 0), \\
H_3 : \{ P_{11,\infty} : (0 : 1 : 3), P_{3,14} : (1 : 6 : 10), P_{3,1} : (1 : 12 : 4), & P_{9,18} : (1 : 13 : 11), P_{9,17} : (1 : 12 : 5), P_{9,16} : (1 : 11 : 1), \\
& P_{3,7} : (1 : 18 : 13), P_{3,13} : (1 : 5 : 18), P_{3,0} : (1 : 11 : 0), & P_{9,15} : (1 : 10 : 18) \} \\
& P_{3,6} : (1 : 17 : 16), P_{3,12} : (1 : 4 : 9), P_{3,18} : (1 : 10 : 17), &
\end{aligned}$$

B. Magma code

Below, we report the code used to find the parabolic one-factorizations.

```
p:=19;
F:=GF(p);
AG:=AffineSpace(F,3);
r:=(p-1)/2;
rr:=(p-3)/2;
F0:={AG![0,1,0]};
for k in [1..r] do
kk:=F!k;
l:=AG![1,0,(-kk/2)^2+(-kk/2)*kk];
F0:=F0 join {l};
end for;
printf "F 0 : %o\n" ,F0;
for k in [1..r] do
kk:=F!k;
G:={AG![0,1,-kk]};
H:={AG![0,1,kk]};
for j in [0..rr] do
jj:=F!j;
i:=F!(kk/2+2*jj*kk);
ii:=F!(kk/2+(2*jj+1)*kk);
m:=AG![1,i+kk/2,i^2+i*kk];
n:=AG![1,ii+kk/2,ii^2+ii*kk];
G:=G join {m};
H:=H join {n};
end for;
printf "G %o : %o\n" ,kk,G;
printf "H %o : %o\n" ,kk,H;
end for;
```