

# Effect of Black Point Infection on Germination of different varieties of Wheat Seed

Poonam Rani<sup>1</sup> and Anita Singh<sup>2</sup>

<sup>1</sup>Career Point University, Kota,  
Rajasthan, India

<sup>2</sup>Guru Kashi University, Talwandi Sabo,  
Bathinda, 151302, India

## Abstract

The present investigation for recording the effect of black point on germination of wheat seed were conducted during 2016-17. The seven different varieties of wheat (HD2733, DBW17, PBW725, PBW386, PBW502, HD3086 and HD2967) were selected. The infestation of seed by black point was categorized on 0 to 5 grade scale. In which healthy seeds were consider as 0 grade. Further germination of seeds were categorized into normal seedling, abnormal seedling and un-germinated seeds. The result shows that the influence of infestation on seed germination appeared to be depend upon the wheat variety. In all seven wheat varieties, germination of BP infected wheat seeds was significantly lower for grade 5 (infected seed) as compared to other infestation grades. At 0 grade maximum normal seedling (68-96%) were recorded. Whereas at 1, 2, 3, 4, 5 grades normal seedling ranges from 65-96%, 45-88%, 55.6- 92%, 53.34-72% and 25-50% respectively. Further the abnormal seedling (4-46.67%) and un-germinated seeds (0-75%) indicated that therewasseverity of black point infection on germination of seeds.

**Keywords:** Black Point, Wheat, Varieties, Germination

## 1. Introduction

Black point is common in all wheat growing regions of the world, including China, Australia, Canada, Serbia and others (Conner and Thomas, 1985; Lorenz, 1986; Mathur and Cunfer, 1993; Wang *et al.*, 2006). Several species of fungi are associated with black point grains. In which *Alternariaalternate* (Fr.) and *Drechslerasorokiniana* (Sacc.) is most dominating fungi species (Huguelet and Kiesling, 1973). The disease seeds are brown to black in colour, usually this discolouration restricted

to the embryonic end of the grain. In case of severe infection, the whole grain may be discoloured and shriveled (Hanson and Christensen, 1953; Adlakha and Joshi, 1974; Pathak and Zaidi, 2013). Thus, it can cause significant economic losses by affecting the market price of wheat as well as the flour quality (Fernandez and Conner, 2011; Barkar *et al.*, 2008). Hence the present study was conducted on recording the effect of black point infection on germination of different wheat seeds varieties.

## 2. Materials and Methods

### Dry Inspection of Seeds

The experiment was conducted during 2016-17. Seeds of 7 popular wheat varieties (HD2733, DBW17, PBW725, PBW386, PBW502, HD3086 and HD2967) were obtained from the different grain markets of Punjab for further experimental work. 100 seeds per varieties of wheat were examined under four replicate. The seeds were subjected to visual observation under stereo-microscope. Seeds that showed distinct symptoms and abnormalities were selected and categorized in to six different grades from 0-5 rating scale:

Grade-0 = Grains free from any discolouration (apparently healthy), Grade-1 = Tip of the embryo brown to blackish, Grade-2 = Discolouration covering the whole embryo, Grade-3 = Embryo with 1/4 of the grain discoloured, Grade-4 = Embryo with 1/2 of the grain discoloured, Grade-5 = Embryo with more than 1/2 of the grain discoloured and shriveled

### Plating and Incubation of Seeds:

Infected seeds of each varieties were surface sterilized in 2% HgCl<sub>2</sub> for 15 min and rinsed for 2 min in three changes of sterile distilled water prior to plating. Twenty seeds were plated in each petri dish

containing moist blotter papers. For each of the categories, a total of 400 seed were plated in four replicates of 100 seeds per variety. These petri dishes were incubated at 22°C ± 2°C under alternating cycles of 12 h light and 12 h darkness.

On the 8th day, incubated seeds were observed for germination. After 8 days, the number of normal seedlings, abnormal seedlings and ungerminated seeds were determined. The seeds classified as normal must fall into one of the following categories:

- 1 Intact seedlings, with all essential structures well developed, complete in proportion, and healthy.
- 2 Seedlings with slight defects of their essential structures, provided they show an otherwise satisfactory and balanced development comparable to that intact seedlings in the same test.
- 3 Seedlings with secondary infection that would have fallen into categories 1 or 2 but for infection by fungi or bacteria from sources other than the parent seed.

Germinated seeds were counted and expressed as a measure of seed viability using the formula.

$$Sv = n/N \times 100$$

Where Sv is % seed viability, n is the number of seeds germinated from each normal or abnormal seed type and N is the total number of seeds plated on blotter.

### 3. Results and Discussion

In seven wheat varieties the effect of black point on germination were recorded. Healthy seeds considered to be in 0 grade. However, germination of grade-5 were significantly lower than that of other infection grades. In case of PBW502, HD3086 and HD2967 wheat varieties more variability in healthy seed germination across the different infection grades were recorded. Poor germination and production of abnormal seedling indicated that there was severity of black point infection. The continuous increase in percentage of non-germination was recorded from grade point 0-5 in only two wheat varieties (DBW17 and PBW725). The wheat varieties HD2733, PBW386, HD3086 and HD2967 were most susceptible to black point (Table 4.1). Previous studies disclosed that HD3086, HD2733, HD2967, HD2329 and WH1105 were most susceptible varieties for black point in South Western Punjab (Rani and Singh, 2017). Mihaela *et. al.*, 2013, also recorded higher germination value in black-point-free seeds (BPF) when compared to affected seeds (BP). Previous studies revealed that the germination rate varied with the variety and health of seeds (Sikder and Paul, 2010). Many authors were also reported that the black point decreases the germination rate, number of embryonic roots,

coleoptile length, delayed seedling emergence and seedling vigour (Ozer, 2005; Toklu *et. al.*, 2008).

Tables 4.1 Effect of (Black Point) on germination of wheat seed as determined by the Standard Blotter Test

Infestation Grade	Germination		
	1	2	3
<b>HD2733</b>			
0	84	12	4
1	80	12	8
2	60	8	32
3	60	12	28
4	56	16	28
5	35	25	37.5
<b>DBW17</b>			
0	96	4	0
1	92	4	4
2	88	8	4
3	88	8	4
4	72	16	12
5	40	20	40
<b>PBW725</b>			
0	96	4	0
1	96	4	0
2	96	4	0
3	92	8	0
4	72	12	16
5	25	0	75
<b>PBW386</b>			
0	82.36	5.88	11.76
1	76.47	11.76	11.76
2	76.47	5.88	18.75
3	58.82	17.65	23.53
4	58.82	29.41	11.76
5	35.29	29.41	35.29
<b>PBW502</b>			
0	90	5	5
1	65	15	
2	45	10	25
3	55.6	22.3	33
4	60	20	20
5	50	25	25
<b>HD3086</b>			
0	73.34	20	6.67
1	73.34	20	6.67
2	53.34	26.67	20
3	73.34	6.67	20
4	53.34	26.67	6.67
5	33.34	46.67	20
<b>HD2967</b>			
0	68	28	4
1	84	12	1
2	64	32	1
3	68	12	20
4	35	5	48
5	33.3	16.67	50

1 = Normal germination, 2 = Abnormal germination, 3 = Ungerminated seeds

These seed-borne diseases had been found to affect the growth and productivity of crop plants (Kubiak and Korbas, 1999; Dawson *et. al.*, 2001; Weber *et. al.*, 2001). Similarly, Rena and Gupta (1982) also

reported localized discoloured areas, usually around the embryo end of seeds was often responsible for reduced germinability. Hence the present study revealed that the black point reduced the germination of seed.

### Acknowledgments

Authors are thankful to the Dr. Jagtar Singh Dhiman, former Additional Director Research of PAU and Senior Plant pathologist for identification of diseases from wheat sample.

### References

- [1] Adlakha K L and Joshi L M, Black point of wheat. *Indian Phytopathology*, 27: 41- 44, (1974).
- [2] Borkar S G, Chaudhary K N, Patil P D and Sonawane R B, Distribution and association of fungal pathogens with black point of wheat disease in Maharashtra state. *Journal of Pl Dis Sci*, 3: 13-16, (2008).
- [3] Conner RL and Thomas JB, Genetic variation and screening techniques for resistance to black point in soft white spring wheat. *Canadian Journal of Plant Pathology*, 74: 402-407,(1985).
- [4] Dawson W A J M and Bateman G L, Fungal communities on roots of wheat and barley and effects of seed treatments containing fluquinconazole applied to control take-all. *Plant Pathology*, 50: 5-82, (2001).
- [5] Fernandez M R and Conner R L, Black Point and Smudge in Wheat. *Prairie Soils and Crops*, 4: 158-164,(2011).
- [6] Hanson E W and Christensen J J, The black point disease of wheat in the United States. *The University of Minnesota Agricultural Experiment Station technical bulletin*, 206:30p, (1953).
- [7] Huguélet J E and Kiesling R L, Influence of inoculum composition on the black point disease of durum wheat. *Phytopathology*, 63: 1220-1225,(1973).
- [8] Kubiak K and Korbas M, Occurrence of fungal diseases on selected winter wheat cultivars. *Postępy w Ochronie Roslin*, 39 (2): 801-804,(1999).
- [9] Lorenz K, Effects of black point on grain composition and baking quality of New Zealand wheat. *New Zealand journal of agricultural research*, 29: 711 -718, (1986).
- [10] Mathur SB and Cunfer B, Black point. In: *Seed-Borne Diseases and Seed Health Testing of Wheat*. Copenhagen, Denmark: Danish Government Institute of Seed Pathology for Developing Countries, 13-21,(1993).
- [11] Mihaela C C, Elena D, and Mihai B, Particularities of the wheat varieties seeds germination under the blackpoint attack incidence. *Romanian Biotechnological Letters*, 18(4):8441-8446, (2013).
- [12] Ozer N, Determination of the fungi responsible for black point in bread wheat and effects of the disease on emergence and seedling vigour. *Trakya Univ J Sci*, 61: 35-40,(2005).
- [13] Pathak N and Zaidi R, Fungi associated with wheat seed discoloration and abnormalities in in vitro study. *Agricultural Sciences*, 4 (9): 516-520,(2013).
- [14] Rani P and Singh A, Wheat Abnormalities assessment in South Western Punjab, *International Journal of Current Research*, 9(8): 55377-55383, (2017).
- [15] Rena J P and Gupta P K S, Occurrence of Black point disease of wheat in West Bengal. *Indian Phytopathology*, 35: 700-702,(1982).
- [16] Sikder S and Paul N K, Study of influence of temperature regimes on germination characteristics and seed reserves mobilisation in wheat. *African Journal of Plant Science*, 4: 401- 408, (2010).
- [17] Toklu F, Akgul DS, Biçici M and Karaköy T, The relationship between black point and fungi species and effects of black point on seed germination properties in bread wheat. *Turkish Journal of Agriculture and Forestry*, 32: 267-272, (2008).
- [18] Wang HW, Xing XW, Yuan HX, Sun BJ, Yu QL and Li HL, Evaluation on the resistance of wheat varieties (lines) to black point. *Journal of Triticeae Crops*, 263: 132-135, (2006).
- [19] Weber R, Hrynczuk B, Runowska-Hrynczuk B and Kita W, Influence of the mode of tillage on disease of culm base in some winter wheat varieties, oat and spring wheat. *Journal of Phytopathology*, 149:185-188, (2001).