

Effects of the Transient Outward Potassium Current on Action Potential Upstroke Velocities Tested using the Dynamic Clamp Technique

Arie O Verkerk, Christiaan C Veerman, Jan G Zegers, Ronald Wilders

Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

Abstract

The voltage-gated transient outward potassium current (I_{to1}) plays a prominent role in the early repolarization phase of the cardiac action potential (AP) and thereby contributes to the refractory period and inotropic state of the myocardium. The current is largely responsible for differences in AP repolarization between species, between left and right ventricle, and transmurally, and it is affected by various pathophysiological conditions, such as heart failure.

In voltage clamp experiments, I_{to1} already activates during depolarization to potentials near -50 to -30 mV, suggesting that I_{to1} may be active during the AP upstroke, but whether it modulates the maximal AP upstroke velocity (dV/dt_{max}) is unknown. In the present study, we addressed this issue using the dynamic clamp configuration of the patch-clamp technique.

Experimental data were acquired from HEK-293 cells expressing the fast sodium current that is responsible for the rapid cardiac AP upstroke. The fast component of I_{to1} ($I_{to,f}$) was computed in real time and injected into the HEK-293 cell during its upstroke. $I_{to,f}$ density, activation rate, and voltage dependence were varied.

We conclude that $I_{to,f}$ may modulate dV/dt_{max} and AP overshoot, but only if its activation is fast and its activation threshold is near -50 to -40 mV.

1. Introduction

The transient outward potassium current (I_{to1}) is responsible for the phase-1 repolarization of the cardiac action potential (AP) and thereby contributes to the refractory period and inotropic state of the myocardium. It has become increasingly clear that I_{to1} also modulates conduction velocity through the associated changes in membrane potential (V_m) during the early action potential plateau phase. However, in voltage clamp experiments, I_{to1} already rapidly activates during depolarizing pulses to potentials around -50 to -30 mV [1]. This suggests that I_{to1} may also modulate the maximal AP upstroke velocity

(dV/dt_{max}). However, data on the effects of I_{to1} on the AP upstroke are lacking.

Presently, various knock-out mouse models exist for $K_v4.2$ and $K_v4.3$, pore-forming α -subunits of the potassium ion channels underlying the fast component of I_{to1} ($I_{to,f}$) [1,2]. These mouse models clearly demonstrated the effects of $I_{to,f}$ on early AP repolarization, but data on dV/dt_{max} were not included. This is also the case for various studies where I_{to1} was blocked by drugs. The I_{to1} activator NS5806 decreased dV/dt_{max} , suggesting that I_{to1} indeed modulates dV/dt_{max} [3]. However, NS5806 also reduced the fast sodium current (I_{Na}) [3], which is responsible for the rapid cardiac AP upstroke [4].

In the present study, we evaluated the effects of $I_{to,f}$ on depolarization and repolarization velocity as well as AP overshoot using the dynamic clamp technique [5]. All three characteristics were measured in HEK-293 cells transfected with I_{Na} channels to avoid interference of other depolarizing and repolarizing membrane currents.

2. Methods

2.1. Patch-clamp experiments

HEK-293 cells were transiently transfected with $0.3 \mu\text{g}$ *SCN5A* cDNA and $0.3 \mu\text{g}$ $\beta 1$ cDNA (both wild-type), encoding the α - and β -subunits of the cardiac I_{Na} channel, respectively.

The alternating voltage clamp/current clamp (VC/CC) technique was used to measure I_{Na} -driven upstrokes at physiological Na^+ concentrations and temperature (37°C) in HEK-293 cells, as described previously [4]. In short, cells were voltage clamped at a holding potential of -85 mV, similar to the resting membrane potential of ventricular myocytes, and upstrokes, overshoot, and early repolarization were elucidated by switching to the current clamp mode of the patch clamp technique for 20 ms.

2.2. Dynamic clamp

$I_{to,f}$ of the Bondarenko et al. mouse ventricular AP model [6] was computed in real time and injected, in a




Figure 1. Effect of $I_{to,f}$ density ($g_{to,f}$) on fast depolarization, repolarization, and overshoot. (A) Typical superimposed membrane potential recordings without $I_{to,f}$ ('0% $g_{to,f}$ '), with the default model $I_{to,f}$ settings ('100% $g_{to,f}$ '), and with doubled $I_{to,f}$ density ('200% $g_{to,f}$ '). (B) Injected current (I_{inj}), demonstrating activation of $I_{to,f}$, during the recordings of panel A. (C) Phase plane plot of I_{inj} versus V_m . (D) First derivative of the membrane potential recordings of panel A (dV/dt). Inset: maximal dV/dt during depolarization. (E) Effect of $I_{to,f}$ density on dV/dt during repolarization. (F) Changes in depolarizing dV/dt , repolarizing dV/dt , and overshoot upon an increase in $g_{to,f}$ ($n = 4$). * $P < 0.05$.

continuous feedback loop, into the real cell during its upstroke. The $I_{to,f}$ reversal potential was set to -86 mV, equal to the potassium equilibrium potential. $I_{to,f}$ density, activation rate and voltage dependence were varied.

2.3. Statistics

Data are presented as mean \pm SEM. One-way repeated measures ANOVA was used for comparing the effects of $I_{to,f}$ density, activation rate, and voltage dependence. $P < 0.05$ was considered statistically significant.

3. Results

Fig. 1 shows the effect of $I_{to,f}$ density ($g_{to,f}$) on depolarization and repolarization velocity and overshoot.

As illustrated in Fig. 1, A and D–F, $I_{to,f}$ modulates repolarization velocity without affecting dV/dt_{max} of depolarization and overshoot. The effect on repolarization increases with increasing $g_{to,f}$ (Fig. 1F). The lack of effect on depolarization and overshoot is likely due to the relatively slow activation of $I_{to,f}$ and the resulting virtual absence of $I_{to,f}$ during these AP phases (Fig. 1, B and C).

The $I_{to,f}$ equations of Bondarenko et al. [6] are based on experimental data obtained at room temperature and the Bondarenko et al. AP model "is nominally adjusted for room temperature of 25°C " [6]. Fig. 2 shows the effect of an increase in $I_{to,f}$ activation rate, thereby representing a more close-to-physiological temperature, on depolarization and repolarization velocity and overshoot. As illustrated in Fig. 2, A and D–F, the increase in $I_{to,f}$ activation rate results in a minor, but significant, decrease in both dV/dt_{max} of depolarization




Figure 2. Effect of $I_{to,f}$ activation rate on fast depolarization, repolarization, and overshoot. (A) Typical superimposed membrane potential recordings without $I_{to,f}$ ('no $I_{to,f}$ '), with the default model $I_{to,f}$ activation rate ('default $I_{to,f}$ '), and with 5 and 10 times faster activation ('5× faster' and '10× faster', respectively). (B) Injected current (I_{inj}), demonstrating activation of $I_{to,f}$, during the recordings of panel A. (C) Phase plane plot of I_{inj} versus V_m . (D) First derivative of the membrane potential recordings of panel A (dV/dt). Inset: maximal dV/dt during depolarization. (E) Effect of $I_{to,f}$ activation rate on dV/dt during repolarization. (F) Changes in depolarizing dV/dt , repolarizing dV/dt , and overshoot upon an increase in $I_{to,f}$ activation rate ($n = 5$). * $P < 0.05$.

and overshoot, likely because $I_{to,f}$ is now activated during these AP phases (Fig. 2, B and C).

Electrophysiological properties of I_{to1} are frequently measured in the presence of Cd^{2+} or Co^{2+} to block contaminating Ca^{2+} currents. Unfortunately, these divalent ions induce a positive shift in voltage dependence of I_{to1} . Accordingly, Bondarenko et al. [6] have taken into account an ≈ 10 mV shift in their $I_{to,f}$ equations. However, the shift in voltage dependence of $I_{to,f}$ is not exactly known. Therefore, we performed a final series of dynamic clamp experiments with a 5 times faster $I_{to,f}$ activation and shifts in $I_{to,f}$ voltage dependence ranging from 0 to -50 mV (Fig. 3). Negative shifts result in a decreased dV/dt_{max} of depolarization, faster repolarization, and a smaller overshoot. The effects are more pronounced with larger shifts (Fig. 3F).

4. Conclusion

We conclude that (1) $I_{to,f}$ clearly affects repolarization, and (2) $I_{to,f}$ may modulate dV/dt_{max} of depolarization and overshoot, but only if its activation threshold is fast and its activation velocity is rather negative. The minor effect on upstroke velocity suggests a limited role of $I_{to,f}$ in AP depolarization and overshoot.

References

- [1] Guo W, Li H, London B, Nerbonne JM. Functional consequences of elimination of $I_{to,f}$ and $I_{to,s}$: early afterdepolarizations, atrioventricular block, and ventricular arrhythmias in mice lacking Kv1.4 and expressing a dominant-negative Kv4 α subunit. *Circ Res* 2000;87:73–9.




Figure 3. Effect of a negative shift in voltage dependence of $I_{to,f}$, in addition to a 5 times faster activation, on fast depolarization, repolarization, and overshoot. (A) Typical superimposed membrane potential recordings without $I_{to,f}$ ('no $I_{to,f}$ '), without a shift in voltage dependence ('0 mV shift'), and with a -10 or -20 mV shift in voltage dependence of $I_{to,f}$ (' -10 mV shift' and ' -20 mV shift', respectively). (B) Injected current (I_{inj}), demonstrating activation of $I_{to,f}$, during the recordings of panel A. (C) Phase plane plot of I_{inj} versus V_m . (D) First derivative of the membrane potential recordings of panel A (dV/dt). Inset: maximal dV/dt during depolarization. (E) Effect of $I_{to,f}$ voltage dependence on dV/dt during repolarization. (F) Changes in depolarizing dV/dt , repolarizing dV/dt , and overshoot upon a negative shift in voltage dependence of $I_{to,f}$ ($n = 5$). $*P < 0.05$.

[2] Liu J, Kim K-H, Morales MJ, Heximer SP, Hui C-C, Backx PH. Kv4.3-encoded fast transient outward current is presented in Kv4.2 knockout mouse cardiomyocytes. *PLoS One* 2015;10:e0133274.

[3] Calloe K, Nof E, Jespersen T, Di Diego JM, Chlus N, Olesen S-P, Antzelevitch C, Cordeiro JM. Comparison of the effects of a transient outward potassium channel activator on currents recorded from atrial and ventricular cardiomyocytes. *J Cardiovasc Electrophysiol* 2011;22:1057–66.

[4] Berecki G, Wilders R, de Jonge B, van Ginneken ACG, Verkerk AO. Re-evaluation of the action potential upstroke velocity as a measure of the Na^+ current in cardiac myocytes at physiological conditions. *PLoS One* 2010; 5:e15772.

[5] Wilders R. Dynamic clamp: a powerful tool in cardiac electrophysiology. *J Physiol* 2006;576:349–59.

[6] Bondarenko VE, Szigeti GP, Bett GCL, Kim S-J, Rasmusson RL. Computer model of action potential of mouse ventricular myocytes. *Am J Physiol Heart Circ Physiol* 2004;287:H1378–403.

Address for correspondence:

Arie O Verkerk, PhD
 Department of Anatomy, Embryology and Physiology
 Academic Medical Center, University of Amsterdam
 Meibergdreef 15, 1105 AZ Amsterdam, The Netherlands
 Phone: +31-20-5664644, E-mail: a.o.verkerk@amc.uva.nl