

Synergistic Anti-arrhythmic Effects of Combined Blockade of Sodium and Ultra-rapid Delayed Rectifier Potassium Channels in Human Atria

Haibo Ni, Dominic G Whittaker, Wei Wang, Henggui Zhang

Biological Physics Group, School of Physics and Astronomy
The University of Manchester, Manchester, UK

Abstract

Atrial fibrillation (AF) is the commonest cardiac arrhythmia. Developing effective and safe anti-AF drugs remains unmet. Atrial-selective blocking the ultra-rapid delayed rectifier potassium current (I_{Kur}) and sodium channel (I_{Na}) was recently reported to be potentially effective in the treatment of AF. We used computational models of human atria to assess the antiarrhythmic effects of I_{Kur} and I_{Na} blocks on human atria. A mathematical model of I_{Kur} channel block was developed to describe the blocking effects of acacetin, an I_{Kur} -selective blocker. I_{Na} blockade was simulated using the guarded-receptor model. The effects of these channel blocks were simulated using mathematical models of human atrial myocytes and tissue. Applying acacetin rendered significant APD prolongation (by 29.3 ms at 1 Hz). The I_{Na} blocker significantly reduced peak I_{Na} at fast pacing rates, but produced no effect at 1 Hz in both atria and ventricles. Combined actions of the two drugs further decreased peak I_{Na} and prolonged APD at a fast pacing rate. These effects were also demonstrated in two-dimensional simulations, where combined blockers showed reduced the life span of re-entrant excitation waves, exerting synergistic antiarrhythmic effects in the human atria, indicating that combined blocks of I_{Kur} and I_{Na} may be a valuable strategy for the treatment of AF.

1. Introduction

Atrial fibrillation (AF) is the commonest cardiac arrhythmia [1]. Developing effective and safe anti-arrhythmic drugs remains challenging [1] and fraught with potential complications including promoting ventricular arrhythmias by prolonging action potential durations (APD). The atrial and ventricular fast Na^+ currents (I_{Na}) show different properties in the voltage-dependent inactivation of the channel, mediating the possibility for atrial-selective sodium channel block [2]. The ultra-rapid delayed rectifier potassium current (I_{Kur}) carried by $Kv1.5$ channel contributes to repolarization in

the atria but not in the ventricles [3,4]. Recent studies suggest that the atrial-selective blockade of I_{Kur} is a potentially effective strategy for the pharmacological treatment of AF [5]. Among the selective I_{Kur} blockers, Acacetin, a natural flavone initially isolated from a traditional Chinese medicine *Xuelianhua*, was demonstrated to be a promising atrial-selective agent for the treatment of AF [3,5].

Multiple-channel blockade is receiving growing interest as empirical observations suggest that multi-channel blockers generally offer more effective antiarrhythmic effects [1]. A recent study with mathematical simulations and experimental validations on (presumably healthy) canine hearts suggest that adding potassium channel block enhances the AF-selective anti-AF effects obtainable from optimised I_{Na} blockade.

In this study we use computational models to assess the hypothesis that blocking I_{Kur} and I_{Na} might produce synergistic anti-arrhythmic effects in the chronic AF (cAF) remodelled human atria. The blockade of I_{Kur} and I_{Na} were simulated by modelling effects of Acacetin and optimised I_{Na} blockers using state-dependent blockade models, respectively. Single cell and tissue models were employed to simulate the anti-AF effects of individual and combined blockers.

2. Methods

2.1. Modelling human atrial electrophysiology

A recent mathematical model for human atrial electrophysiology [6] was employed to simulate the human atrial electrical activities under both sinus rhythm (SR) and cAF conditions. The formulation of I_{Kur} was updated based on recently published data [4]. cAF induced electrical remodelling were incorporated based on a previously published model (AF-4 in [6]).

In simulating the effects of sodium blockers on human ventricular AP, the mathematical model published by O'Hara et al. [7] was adopted. The I_{Na} formulation in the ventricular model was replaced with that in [6].

2.2. Modelling I_{Na} blocker

Following previous a study [1] on simulating I_{Na} blockade, a theoretically optimized I_{Na} blocker was used. We employed the guarded receptor model to simulate the binding and unbinding of drug to sodium channel in a drug concentration-dependent manner. It was assumed that the drug predominantly binds to the activated and inactivated states of the sodium channel (denoted as B_A and B_I , respectively). The blockade of I_{Na} is given by:

$$I_{Na} = g_{Na}(1 - B_A - B_I)m^3hj(V - E_{Na}) \quad (1)$$

$$\frac{dB_A}{dt} = K_A[D]m^3hj(1 - B_A - B_I) - L_AB_A \quad (2)$$

$$\frac{dB_I}{dt} = K_I[D](1 - h)(1 - B_A - B_I) - L_IB_I \quad (3)$$

where g_{Na} is the channel conductance of I_{Na} ; B_A and B_I are the fractional activated and inactivated blocks; m is the activation gate, h and j are the inactivation variables of the sodium channel; V is the transmembrane potential; E_{Na} is the reversal potential of Na^+ ; F , R and T are the Faraday's constant, universal gas constant and temperature respectively; K_A , K_I are the binding constants and L_A , L_I are the unbinding constants; $[D]$ is the drug concentration; In the present study, a drug concentration of 60 μM was studied.

2.3. Modelling effects of Acacetin

The effects of Acacetin on human atrial electrophysiology were simulated using the guarded receptor formalism. It was assumed that the drug binds to the open and closed gates of Kv1.5. Therefore, the binding and unbinding kinetics for open and closed states were functions of the binding constants, and given by:

$$I_{Kur} = g_{Kur,v}(1 - B_O - B_C)ai(V - E_K) \quad (4)$$

$$\frac{dB_O}{dt} = K_O \exp\left(Z_{KO} \frac{VF}{RT}\right) [D] ai(1 - B_O - B_C) - L_O \exp\left(-Z_{LO} \frac{VF}{RT}\right) B_O \quad (5)$$

$$\frac{dB_C}{dt} = K_C \exp\left(Z_{KC} \frac{VF}{RT}\right) [D] (1 - a)i(1 - B_O - B_C) - L_C \exp\left(-Z_{LC} \frac{VF}{RT}\right) B_C \quad (6)$$

where $g_{kur,v}$ is the conductance of the Kv1.5 channel; B_O and B_C are the fractional open and closed state block, respectively; a and i are the activation and inactivation gate variables; V is the membrane voltage; E_K is the reversal potential of potassium; F , R and T are the Faraday's constant, universal gas constant and temperature respectively. K_O and K_C are the binding constant, and L_O , L_C are the unbinding constant. Z_{KO} , Z_{LO} , Z_{KC} , Z_{LC} are the drug charge parameters for the

corresponding binding and unbinding processes. $[D]$ is the concentration of acacetin applied. In the present study, a concentration 3.2 μM was considered.

The binding and unbinding parameters were obtained by fitting the model to the experimental data from [3], and are given in Table 1.

Table 1. Print sizes for different parts of the manuscript.

Parameter	Value	Unit
K_O	0.000194	$\mu M^{-1} \cdot ms^{-1}$
L_O	0.000291	ms^{-1}
Z_{KO}	0.257	
Z_{LO}	-0.0128	
K_C	0.00249	$\mu M^{-1} \cdot ms^{-1}$
L_C	0.000286	ms^{-1}
Z_{KC}	0.327	
Z_{LC}	0.837	

2.4. Two-dimensional tissue models

We performed two-dimensional simulations on a 7.5×7.5 cm² sheet of human atrial myocytes with 250 nodes along each direction. To solve the excitation wave propagation problem, the well-known mono-domain equation [6] was used, given by:

$$\frac{\partial V}{\partial t} = \nabla \cdot (D \nabla V) - \frac{I_{ion}}{C_m} \quad (7)$$

where V is the membrane potential, D is the diffusion parameter, I_{ion} is the total ionic current in a single cardiomyocyte and C_m is the membrane capacitance of the cell. The tissue was modelled as homogeneous to conduction. The diffusion coefficient was chosen so that the model gives a conduction velocity of 0.7 m/s measured using plane wave for the tissue with normal human atrial cells. The numerical method used in our previous study [6] was adopted here. The mono-domain equation was integrated at a time step of 0.02 ms.

To initiate re-entrant waves, the cross field (S1-S2) protocol [1] was used. Life span of re-entrant excitations were quantified based on pseudo-ECG signals, which were computed using the equation as described in [8].

3. Results

3.1. Effects of cAF on atrial electrophysiology

Incorporation of cAF induced electrical remodeling to the significantly reduced APD (Figure 1A, 247 ms in SR versus 157 ms in the AF model). The calcium transient (CaT) was also markedly reduced with cAF remodeling (Figure 1B).




Figure 1. Simulated action potentials and calcium transient (CaT) of human atrial cardiomyocytes for sinus rhythm (SR) and cAF (AF). The action potential duration measured at 90% repolarization (APD₉₀) and the amplitude of CaT are quantified and shown in the right panels. The pacing rate was 1Hz.

3.2. Simulated effects of I_{Na} and I_{Kur} blocks on the electrophysiology of human cardiomyocytes

Individual and combined effects of I_{Na} and I_{Kur} blocks on human atrial and ventricular electrophysiology were simulated and the results are shown in Figure 2. For a slow pacing rate (1Hz), blocking I_{Na} alone with the theoretical blocker rendered no significant effects on either AP or peak I_{Na} in both atrial and ventricular cells (Figure 2 A&C). At 1Hz, applying acacetin prolonged APD by 29.3 ms. Combining I_{Na} and I_{Kur} blocks produced negligible change in APD (0.3 ms prolongation) and peak I_{Na} .

At a fast pacing rate (6 Hz), alternans in APD and peak I_{Na} were produced (Figure 2Bi). Applying the I_{Na} blocker exerted pronounced reduction to peak I_{Na} and thus dV/dt_{max} (143.7 versus 172.0 pA/pF for control, 17% reduction); APD was prolonged by 3.5 ms. Applying acacetin alone prolonged APD by 26.1 ms for big AP, and 38.1 ms for the small AP) and marked decrease in peak I_{Na} (by 28.2% for big AP, and 20.1% for small AP). More pronounced effects on AP were produced by combining I_{Kur} and I_{Na} blocks, showing more significant alternans in AP and dV/dt_{max} ; APD was prolonged by 55.8 ms for the big AP, demonstrating a synergistic effect in APD prolongation.




Figure 2. Simulated action potentials and I_{Na} for cAF-remodeled atrial myocytes paced at 1Hz (A) and 6 Hz (B) for control (CTL), I_{Na} channel block (INa-B), I_{Kur} channel block (IKur-B) and combined blockade of I_{Na} and I_{Kur} (INa-B + IKur-B). (C) AP and I_{Na} of a ventricular cardiomyocyte paced at 1Hz with/without I_{Na} block. In the simulations, rate constants for INa-B were: $K_A = 100ms^{-1} \cdot mol^{-1}$, $K_I = 100ms^{-1} \cdot mol^{-1}$, $I_A = 1 ms^{-1}$, $I_I = 0.01 ms^{-1}$.

3.3. Simulated effects of I_{Na} and I_{Kur} blocks on AF termination

Re-entrant excitation waves were initiated in the two-dimensional human atrial tissue. In simulating drug blocks, drugs were applied at 2500 ms following initiation of reentry. Figure 3 shows the simulated pECGs, APs of a single cell in the tissue and the corresponding fractional block of I_{Na} and I_{Kur} of the same myocyte. For the control condition, reentrant waves were sustained within the simulated time of 10 s. For INa-B, the rotors were terminated after 6552 ms following application of the drug. For IKur-B, the excitations diminished after 2082 ms following introducing acacetin.

The application of both drugs exerted most effective termination of AF, with the reentries were sustained for only 1052 ms after applying drugs.




Figure 3. Simulated reentry termination by INa-B and/or IKur-B. (A) Computed pECGs; (B) AP of a single cell from the tissue during the reentrant excitations; (C) Fractional block in I_{Na} ; (D) Fraction of blocked Kv1.5 channels. For the simulation with drugs, the simulation was stopped after the termination of reentry.

4. Discussion and conclusion

Using biophysically detailed models of human cardiac electrophysiology, we show that state-dependent block of sodium channels exerted no effect at slow pacing rate and pronounced effects at fast rates. Combined I_{Na} and I_{Kur} blocks produced much more significant antiarrhythmic effects than an individual application of either blocker. In simulated AF, applying combined blocks was shown to be most effective in terminating rotors.

Combining I_{Na} blockade with I_{Kur} blockade produced synergistic anti-arrhythmic effects in human atria, which

provides a potentially valuable strategy for the treatment of AF.

Acknowledgements

This project was supported by BHF and the University of Manchester.

References

- [1] Aguilar M, Xiong F, Qi XY, Comtois P, Nattel S. Potassium Channel Blockade Enhances Atrial Fibrillation-Selective Antiarrhythmic Effects of Optimized State-Dependent Sodium Channel Blockade. *Circulation* 2015;132(23):2203-11
- [2] Burashnikov A, Diego JMD, Zygmunt AC, Belardinelli L, Antzelevitch C. Atrium-Selective Sodium Channel Block as a Strategy for Suppression of Atrial Fibrillation Differences in Sodium Channel Inactivation Between Atria and Ventricles and the Role of Ranolazine. *Circulation* 2007;116:1449-57.
- [3] Wu H-J, Wu W, Sun H-Y, Qin G-W, Wang H-B, Wang P, et al. Acacetin causes a frequency- and use-dependent blockade of hKv1.5 channels by binding to the S6 domain. *J Mol Cell Cardiol* 2011;51:966-73.
- [4] Christophersen IE, Olesen MS, Liang B, Andersen MN, Larsen AP, Nielsen JB, et al. Genetic variation in KCNA5: impact on the atrial-specific potassium current I_{Kur} in patients with lone atrial fibrillation. *Eur Heart J* 2013;34:1517-25.
- [5] Li G-R, Wang H-B, Qin G-W, Jin M-W, Tang Q, Sun H-Y, et al. Acacetin, a Natural Flavone, Selectively Inhibits Human Atrial Repolarization Potassium Currents and Prevents Atrial Fibrillation in Dogs. *Circulation* 2008;117:2449-57.
- [6] Colman MA, Aslanidi OV, Kharche S, Boyett MR, Garratt C, Hancox JC, et al. Pro-arrhythmogenic effects of atrial fibrillation-induced electrical remodelling: insights from the three-dimensional virtual human atria. *J Physiol* 2013;591:4249-72.
- [7] O'Hara T, Virág L, Varró A, Rudy Y. Simulation of the Undiseased Human Cardiac Ventricular Action Potential: Model Formulation and Experimental Validation. *PLoS Comput Biol* 2011;7:e1002061.
- [8] Baher A, Qu Z, Hayatdavoudi A, Lamp ST, Yang M-J, Xie F, et al. Short-term Cardiac Memory and Mother Rotor Fibrillation. *Am J Physiol - Heart Circ Physiol* 2007;292(1):H180-9.

Address for correspondence.

Haibo Ni.
Room 3.08 Schuster Building, Oxford Road
The University of Manchester
Haibo.ni@manchester.ac.uk