



E-ISSN: 2278-4136  
P-ISSN: 2349-8234  
[www.phytojournal.com](http://www.phytojournal.com)  
JPP 2020; 9(4): 425-427  
Received: 01-04-2020  
Accepted: 03-05-2020

**Y Sudharani**

MS Swaminathan School of  
Agriculture, Centurion  
University of Technology and  
Management, Parlakhemundi,  
Odisha, India

**Priyadarshani P Mohapatra**

College of Agriculture, CAU-I,  
Kyrdemkulai, Meghalaya, India

**Madhuri Pattanaik**

MS Swaminathan School of  
Agriculture, Centurion  
University of Technology and  
Management, Parlakhemundi,  
Odisha, India

**Hansraj Hans**

ICAR Research Complex for  
Eastern Region, Patna, India

**Sagar Maitra**

MS Swaminathan School of  
Agriculture, Centurion  
University of Technology and  
Management, Parlakhemundi,  
Odisha, India

**Corresponding Author:****Hansraj Hans**

ICAR Research Complex for  
Eastern Region, Patna, India

## Effect of Phosphorus on Cowpea (*Vigna unguiculata* L. Walp): A review

**Y Sudharani, Priyadarshani P Mohapatra, Madhuri Pattanaik, Hansraj Hans and Sagar Maitra**

DOI: <https://doi.org/10.22271/phyto.2020.v9.i4e.11721>

**Abstract**

Genotypes and phosphorus levels of other legume crops were also studied and included in the review. It has been seen that plant height of cowpea increased with application of 40 kg P<sub>2</sub>O<sub>5</sub>/ha and 90kg P<sub>2</sub>O<sub>5</sub>/ha. 40 kg P<sub>2</sub>O<sub>5</sub>/ha and 80 kg P<sub>2</sub>O<sub>5</sub>/ha recorded significant increase on number of branches per plant. Number of leaves per plant, dry matter accumulation and days to first flowering is enhanced by 90kg P<sub>2</sub>O<sub>5</sub>/ha, 60 kg P<sub>2</sub>O<sub>5</sub>/ha and 40 kg P<sub>2</sub>O<sub>5</sub>/ha respectively. However cowpea different yield attributes are influenced by 75 kg P<sub>2</sub>O<sub>5</sub>/ha, 60 kg P<sub>2</sub>O<sub>5</sub>/ha, 50 kg P<sub>2</sub>O<sub>5</sub>/ha, 60 kg P<sub>2</sub>O<sub>5</sub>/ha, 70 kg P<sub>2</sub>O<sub>5</sub>/ha, 80 kg P<sub>2</sub>O<sub>5</sub>/ha and 30 kg P<sub>2</sub>O<sub>5</sub>/ha with substantial increase in the growth of different parameters respectively. Hence, application of phosphorus equally influences the growth and yield components of cowpea genotypes.

**Keywords:** Genotypes, effect, phosphorus, application, accumulation

**Introduction**

Cowpea (*Vigna unguiculata* L. Walp) is an annual legume crop which belongs to the family Leguminosae and hence, it is cultivated widely in tropics and subtropics during the warm season. It is commercially grown throughout the India for its seed and green pods which are used as vegetable. The leading states in cowpea production in our country are UP, Bihar, Jharkhand, West Bengal, Odisha and so on. In Odisha, the crop is grown in an area of 1.5 million hectares with a production of 0.49 million t. This clearly imposes the need to identify the reasons for such low productivity in Odisha. Genotypes play an important role in crop production and the potential yield of a genotype within the genetic limit is determined by its growing environment. Hence, combination of genotype and environmental factors can bring about increase in production. Legumes are phosphorus loving plants. They require phosphorus for growth and seed development and most especially in nitrogen fixation which is an energy-driving process (Sanginga *et al.*, 2000) [17]. For sustainable food production to meet the increasing population in developing countries, need for phosphorus fertilizer application is expected to increase yield (Brynes and Bumb, 1998) [5]. Phosphorus is known to cause multiple effects on nutrition, increase in seed yield and nodulation process (Singh *et al.* 2011) [21]. Some researchers also reported that phosphorus application will affect the other nutrients in seed and leaves because of its multiple effects on plant nutrition (Shilpa, 2013) [19]. Earlier researches noted the effect of genotype and phosphorus level on growth and productivity of cowpea in different agro-climatic regions (Singh *et al.*, 2011) [21]. However, there is insufficient research work on impact of phosphorus level on cowpea genotypes under this agro-climatic region. Hence, the present study has been conducted with the following objectives on evaluation and interactive effect of genotypes and phosphorus levels on growth, yield and quality of seed.

**1. Growth of Cowpea as Influenced by Phosphorus Levels****1.1 Effect of Phosphorus Levels on Plant Height of Cowpea**

Baboo and Mishra (2001) [3] stated that increasing phosphorus rate up to 90 kg P<sub>2</sub>O<sub>5</sub>/ha increased the plant height of cowpea. Anil *et al.* (2007) observed that 60 kg P<sub>2</sub>O<sub>5</sub>/ha significantly increased the plant height of cowpea. Jat *et al.* (2013) [8] found that application up to 40 kg P<sub>2</sub>O<sub>5</sub>/ha significantly enhanced the plant height of cowpea compared to control and 20 kg P<sub>2</sub>O<sub>5</sub>/ha.

## 1.2 Effect of Phosphorus Levels on Number of Branches per Plant of Cowpea

Jat *et al.* (2013) <sup>[8]</sup> found that application up to 40 kg P<sub>2</sub>O<sub>5</sub>/ha significantly enhanced number of branches per plant as compared to control and 20 kg P<sub>2</sub>O<sub>5</sub>/ha. Sharma *et al.* (2015) <sup>[18]</sup> investigated the effect of nutrients on productivity of cowpea genotype Pusa Komal and in the experiment application of 80kg P<sub>2</sub>O<sub>5</sub>/ha recorded the maximum number of branches per plant (7.81) at 45 DAS.

## 1.3 Effect of Phosphorus Levels on Number of Leaves per Plant of Cowpea

Baboo and Mishra (2001) <sup>[3]</sup> stated that enhancing phosphorus levels up to 90kg P<sub>2</sub>O<sub>5</sub>/ha increased number of green leaves per plant. Sharma *et al.* (2015) <sup>[18]</sup> noted that application of 80kg P<sub>2</sub>O<sub>5</sub>/ha noted the maximum number of leaves per plant (30.82).

## 1.4 Effect of Phosphorus Levels on Days to First Flowering of Cowpea

Patel and Jadav (2010) <sup>[14]</sup> observed that application of 40kg P<sub>2</sub>O<sub>5</sub>/ha along with rhizobium seed inoculation gave significantly less number of days to first flowering.

## 1.5 Effect of Phosphorus Levels on Dry Matter Accumulation of Cowpea

Bhilare and Patil (2002) <sup>[4]</sup> conducted an experiment at Rahuri to determine the effect of three levels of phosphorus application (0, 30 and 60kg P<sub>2</sub>O<sub>5</sub>/ha) on cowpea. Significantly highest dry matter (48.02 q/ha) were obtained with the application of 60kg P<sub>2</sub>O<sub>5</sub>/ha.

## 2. Yield Attributes and Yields of Cowpea as Influenced by Phosphorus Levels

### 2.1 Effect of phosphorus levels on number of pods per plant of cowpea

Kurdikeri *et al.* (1973) <sup>[11]</sup> recorded more number of pods per plant was produced with the application of 44 kg P<sub>2</sub>O<sub>5</sub>/ha along with 11 kg N/ha. However, Kumar *et al.* (2001) <sup>[10]</sup> reported that application of 50 kg P<sub>2</sub>O<sub>5</sub>/ha to cowpea through di-ammonium phosphate (DAP) exhibited higher number of pods per plant. Khan *et al.* (2002) <sup>[9]</sup> studied on productivity of cowpea as influenced by phosphorus levels on noted that the maximum number of pods per plant was recorded with phosphorus application 75kg P<sub>2</sub>O<sub>5</sub>/ha.

### 2.2 Effect of Phosphorus Levels on Length of Pods of Cowpea

Anil *et al.* (2007) observed that 60 kg P<sub>2</sub>O<sub>5</sub>/ha significantly enhanced the length of pod over control. Prasad *et al.* (2008) concluded that the application of VAM (*Gigaspora calospora*), Rhizobium culture and 80kg P<sub>2</sub>O<sub>5</sub>/ha significantly increased length of pods.

### 2.3 Effect of Phosphorus Levels on 100-seed Weight of Cowpea

Kumar *et al.* (2001) <sup>[10]</sup> found that application of 50 kg P<sub>2</sub>O<sub>5</sub>/ha to cowpea through di-ammonium phosphate (DAP) exhibited highest 100 seed weight.

### 2.4 Effect of phosphorus levels on pod weight of cowpea

Anil *et al.* (2007) observed that 60 kg P<sub>2</sub>O<sub>5</sub>/ha significantly increased the pod weight.

## 2.5 Effect of Phosphorus Levels on Number of Seeds per Pod of Cowpea

Kumar *et al.* (2001) <sup>[10]</sup> recorded application of 50 kg P<sub>2</sub>O<sub>5</sub>/ha in cowpea exhibited higher number of seed per pod.

## 2.6 Effect of Phosphorus Levels on Seed Weight per Pod of Cowpea

Khan *et al.* (2002) <sup>[9]</sup> recorded that maximum seed weight was observed with application of 75 kg P<sub>2</sub>O<sub>5</sub>/ha as compared to higher or lower dose. Tajudeen and Oseni (2009) <sup>[22]</sup> stated that application of phosphorus enhanced seed weight of cowpea at the rate of 40 kg P<sub>2</sub>O<sub>5</sub>/ha.

## 2.7 Effect of Phosphorus Levels on Pod Yield of Cowpea

Arup and Dhananjay (2003) <sup>[2]</sup> reported that phosphorus fertilizers up to 80 kg P<sub>2</sub>O<sub>5</sub>/ha significantly increased the pod yield of cowpea.

## 2.8 Effect of Phosphorus Levels on Seed Yield of Cowpea

Ram and Dixit (2001) <sup>[16]</sup> states that among different levels of phosphorus, 60 kg P<sub>2</sub>O<sub>5</sub>/ha produced significantly more seed yield (10.8 q/ha) than control. Application of 60 kg P<sub>2</sub>O<sub>5</sub>/ha was found better for the production of seed yield. Singh *et al.* (2011) <sup>[21]</sup> noted that there was significant response to applied phosphorus on the seed yield of cowpea. Significantly higher seed yield was recorded in plots applied with 60 kg P<sub>2</sub>O<sub>5</sub>/ha (1353 kg/ha) than control plot (1017 kg/ha). Shilpa *et al.* (2015) <sup>[20]</sup> noted that application of 50 kg P<sub>2</sub>O<sub>5</sub>/ha recorded significantly higher seed yield (1087 kg/ha) of cowpea compared to 25 and 75 kg P<sub>2</sub>O<sub>5</sub>/ha. Mawo *et al.* (2016) <sup>[12]</sup> observed that the application of phosphorus equally influenced the growth and yield components of cowpea at different levels. The results showed the highest seed yield of cowpea was obtained with the application of 30 kg P<sub>2</sub>O<sub>5</sub>/ha.

## 2.9 Effect of Phosphorus Levels on Stover Yield (kg/ha) of Cowpea

Bhilare and Patil (2002) <sup>[4]</sup> conducted experiment at Rahuri to determine the effect of three phosphorus application levels (0, 30 and 60 kg P<sub>2</sub>O<sub>5</sub>/ha) on cowpea. Significantly highest yields of stover yield (244.10 q/ha) were obtained with the application of 60 kg P<sub>2</sub>O<sub>5</sub>/ha. Vikrant *et al.* (2005) at Hisar observed that higher stover yield of cowpea were recorded with 60 kg P<sub>2</sub>O<sub>5</sub>/ha over lower doses of phosphorus.

## 2.10 Effect of Phosphorus Levels on Biological Yield of Cowpea

Choudhary and Yadav (2011) <sup>[6]</sup> at Jobner reported that biological yield of cowpea increased significantly with increasing levels of fertility up to 100% RDF (20 kg N + 40 kg P<sub>2</sub>O<sub>5</sub>/ha).

## 2.11 Effect of Phosphorus Levels on Harvest Index of Cowpea

Ilavarasi *et al.* (2007) <sup>[7]</sup> at Annamalai nagar reported that application of 70 kg P<sub>2</sub>O<sub>5</sub>/ha + 70 kg K<sub>2</sub>O/ha along with basal dose of 20 kg N/ha recorded maximum harvest index in cowpea over lower doses of fertilizers.

## 2.12 Effect of Phosphorus Levels on Growth and Yield of Cowpea

Oka *et al.* (2001) <sup>[13]</sup> worked at Calabar, South Eastern Nigeria four levels of phosphorus fertilizers (10, 20, 40 and 60 kg P<sub>2</sub>O<sub>5</sub>/ha) and they noted that IT 81-D-1228-14 was able to provide higher green pod yields at 20 kg P<sub>2</sub>O<sub>5</sub>/ha. Singh *et*

al. (2011) [21] studied the effect of phosphorus on the growth and yield of cowpea. Results showed significant response to applied phosphorus on pods per plant, seed weight and 100-seed weight with highest response to the application of 60kg P<sub>2</sub>O<sub>5</sub>/ha. Mawo *et al.* (2016) [12] observed that the application of phosphorus equally influences the growth and yield components of cowpea genotypes. The results showed that cowpea gave the highest seed yield with application of 30 kg P<sub>2</sub>O<sub>5</sub>/ha.

### Conclusion

In this review, effect of phosphorus on cowpea, the different yield attributing characters like number of pods per plant, length of pod, pod height, number of seeds, and seed weight per pod were also influenced by the genotypes of cowpea, phosphorus levels and their interaction effects. It has evaluated that application of phosphorus 60kg P<sub>2</sub>O<sub>5</sub>/ha registered the maximum harvest index than other treatment combinations. From the present study, it may be concluded that the crop may be cultivated with 60 kg P<sub>2</sub>O<sub>5</sub>/ha to obtained higher yield and superior quality of cowpea during summer season in clay loam soil under south Odisha conditions.

### References

- Singh AK, Tripathi PIN, Singh R. Effect of Rhizobium inoculation, nitrogen and phosphorus levels on growth, yield and quality of *kharif* cowpea (*Vigna unguiculata* (L.) Walp.). *Crop Research*. 2007; 33(1, 2 & 3):71-73.
- Arup C, Dhananjay D. Response of Vegetable cowpea to phosphorus and Biofertilizers in old alluvial zone of West Bengal. *Legume Research*. 2003; 26(3):196-199.
- Baboo R, Mishra SK. Growth and pod production of cowpea (*Vigna sinensis*) as affected by inoculation, nitrogen and phosphorus. *Annual Agricultural Research*. 2001; 22(1):104-106.
- Bhilare RiL, Patil VS. Response of forage cowpea varieties to phosphorus application. *Forage Research*. 2002; 28(3):179-180.
- Brynes BH, Bumb BL. Population growth, food production and nutrient requirements in nutrient use in crop Production. Ed. Z. Rengel. Food Product Press. New York, London, 1998, 227-265.
- Choudhary GL, Yadav LR. Effect of fertility level of foliar nutrient on cowpea productivity. *Journal of Food Legume*. 2011; 24(1):67-68.
- Iiavarasi K, Anuja S, Shakila A, Angayakanni A. Effect of phosphorus and potassium application on soil available NPK, their uptake and pod yield of cowpea (*Vigna unguiculata* (L.) Walp). *Advances in Plant Science*. 2007; 20(1):205-206.
- Jat SR, Patel, BJ, Shivran AC, Kuri BR, Jat GI. Effect of phosphorus and sulphur levels on growth and yield of cowpea under rainfed conditions. *Annals of Plant and Soil Research*. 2013; 15(2):114-117.
- Khan MA, Aslam M, Sultan T, Mahmood IIA. Response of phosphorus application on growth and yield of inoculated and un-inoculated Mungbean (*Vigna radiata*). *International Journal of Agriculture and Biology*. 2002; 4:523-524.
- Kumar CP, Nagaraju AP, Yogananda SB. Effect of phosphorus sources and zinc level on growth and yield of cowpea (*Vigna unguiculata* L.). *Journal of Ecobiology*. 2001; 13(4):275-278.
- Kurdikeri CIB, Patil RV, Krishnamurthy K. Response of cowpea (*Vigna catjung*) to varying fertilizer levels. *Mysore Journal Agricultural Science*. 1973; 7(2):170-174.
- Mawo YIM, Mohammed IIB, Garko MS. Effect of phosphorus levels on growth, yield and development of cowpea (*Vigna unguiculata* (L.) Walp) varieties. *International Journal of Scientific Engineering and Applied Science*. 2016; 2(5):302-312.
- Oka BFD, Eneji AE, Ereji E, Nwoko C, Shiyam JO. Performance of vegetable cowpea (*Vigna unguiculata* L.) as influenced by P fertilizers in South East Nigeria. *Acta Agronomica Hungarica*. 2001; 49(3):305-309.
- Patel BN, Jadav DIK. Effect of Rhizobium seed inoculation, nitrogen and phosphorus on growth, nodulation, flowering and seed yield of cowpea cv. Pusa Phalguni (*Vigna unguiculata* (L.) Walp). *International Journal of Agricultural Science*. 2010; 6(2):361-364.
- Prasad S, Singh SP, Dwivedi NN, Singh SR, Singh ViK, Singh S, Chandola VIK. Response of mycorrhiza, rhizobium culture and phosphorus levels on yield and its contributing characters of cowpea (*Vigna unguiculata* L.). *Vegetable Science*. 2008; 35(2):210-211.
- Ram SN, Dixit RS. Growth, yield attributing parameters and quality of summer green gram (*Vigna radiata* L. Wilczek) as influenced by dates of sowing and phosphorus. *Indian Journal of Agricultural Research*. 2001; 35:275-277.
- Sanginga N, Lyasse O, Singh BB. Phosphorus use efficiency and nitrogen balance of cowpea breeding lines in a low P soil of the derived savanna zone in West Africa. *Plant and Soil*. 2000; 220:119-128.
- Sharma SIK, Prajapati S, Raghuwanshi O. Effect of organic manures and Inorganic fertilizers on yield and economics of Cowpea production (*Vigna unguiculata* L.). *Indian Research Journal of Genetics & Biotechnology*. 2015; 7(1):152-155.
- Shilpa HD. Performance of cowpea (*Vigna unguiculata* (L.) Walp) genotypes during summer at different phosphorus levels. Thesis, Master of Science in Agronomy. University of Agricultural Sciences, Dharwad, 2013.
- Shilpa HD, Wali SY, Hanamant M, Halli TG. Response of Cowpea (*Vigna unguiculata*) genotypes to different levels of phosphorus during summer under northern dry zone of Karnataka. *International Journal of Technical Research and Applications*. 2015; 3(5):i224-226.
- Singh A, Baoule AL, Ahmed HG, Dikko AIU, Aliyu U, Sokoto MB *et al.* Influence of phosphorus on the performance of cowpea (*Vigna unguiculata* (L.) Walp.) varieties in the Sudan savanna of Nigeria. *Agriculture Sciences*. 2011; 2(3):313-317.
- Tajudeen IO. Growth and Zinc Uptake of Sorghum and Cowpea in Response to Phosphorus and Zinc Fertilization. *World Journal of Agricultural Sciences*. 2009; 5(6):670-674.
- Singh VIH, Malik CVS, Singh BP. Grain yield and protein content of cowpea as influenced by farm yard manure and phosphorus application. *Indian Journal of Pulses Research*. 2005; 18(2):250-251.