

# Analysis on Genetic Diversity of *Radix Astragali* by ISSR Markers

Yaling Liu<sup>1,2</sup>, Pengfei Zhang<sup>1</sup>, Ru Zhang<sup>1</sup>, Meiling Song<sup>1</sup>, Fengbo Liu<sup>2</sup>, Wenquan Wang<sup>2,3\*</sup>, Junling Hou<sup>3\*</sup>

<sup>1</sup>College of Life Science, Shanxi Agricultural University, Taigu, China

<sup>2</sup>School of Chinese Pharmacy, Beijing University of Chinese Medicine, Beijing, China

<sup>3</sup>Institute of Medicinal Plant Development, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing, China

Email: \*wwq57@126.com, \*mshjl@126.com.

**How to cite this paper:** Liu, Y.L., Zhang, P.F., Zhang, R., Song, M.L., Liu, F.B., Wang, W.Q. and Hou, J.L. (2016) Analysis on Genetic Diversity of *Radix Astragali* by ISSR Markers. *Advances in Bioscience and Biotechnology*, 7, 381-391.

<http://dx.doi.org/10.4236/abb.2016.710037>

**Received:** May 3, 2016

**Accepted:** October 16, 2016

**Published:** October 20, 2016

Copyright © 2016 by authors and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

## Abstract

*Radix Astragali* has been an important traditional Chinese herbal medicine for over 2000 years. It is derived from two plant species, namely, *Astragalus mongholicus* [*Astragalus membranaceus* (Fisch.) Bge. var. *mongholicus* (Bge.) Hsiao] and *Astragalus membranaceus* [*Astragalus membranaceus* (Fisch.) Bge.] (*Leguminosae*), according to the Pharmacopoeia of the People's Republic of China. In this study, the genetic diversity and genetic relationships of *Radix Astragali* in China were analyzed by Inter-Simple Sequence Repeat (ISSR) markers. A total of 25 highly polymorphic ISSR primers were selected to amplify 95 *Radix Astragali* samples. Among 273 DNA bands amplified, 213 are polymorphic (percentage of polymorphic bands: 78%). The average value of the amplified bands was 10.9 for each primer, and the number varied from 4 to 20. The genetic diversity of the 95 *Radix Astragali* samples was analyzed by using POPGENE 1.32 software. The Nei's genetic diversity index ( $h$ ) and Shannon's information index ( $I$ ) were 0.3590 and 0.5308, respectively, which indicated the abundant genetic diversity of *Radix Astragali*. The level of genetic diversity in *A. membranaceus* ( $h$ : 0.3109,  $I$ : 0.4657) was slightly lower than that in *A. mongholicus* ( $h$ : 0.3364,  $I$ : 0.4969). Considering the average genetic similarity coefficient by NTSYS analysis to cluster the *A. membranaceus* of nine habitats and *A. mongholicus* of five habitats, *Radix Astragali* samples were clustered into two groups according to place of origin. This clustering is different from traditional clustering, which divides groups according to species. Results obtained from this study will provide a theoretical basis for the molecular study on germplasm resources of *Radix Astragali*.

## Keywords

*Radix Astragali*, ISSR, Genetic Diversity

## 1. Introduction

Radix *Astragali*, which has a long history of medicinal use, is derived from *Astragalus mongholicus* and *Astragalus membranaceus* [1]. *Astragalus* is mainly distributed in North, Northeast, Northwest, and Southwest China [2]. However, the qualitative difference between wild and cultivated *Astragalus* owing to the determination of genuine raw materials results in significant difference in market price, which leads to excavation of wild herbs. Thus, the distribution and population of wild *Astragalus* have rapidly decreased. *Astragalus* has been included in the national list of protected rare and endangered plants [3]. Effective conservation of germplasm resources of Radix *Astragali* is urgently needed. Although the artificial cultivation of *Astragalus* has achieved success, the qualitative and quantitative demands of raw material for medicine have greatly increased with the development of new drugs, medicine, and health products. *Astragalus* cultivation has produced hybrid varieties with unstable yield and low quality. This phenomenon has seriously affected the further development of *Astragalus* industry. Therefore, this preliminary study on *Astragalus* from different habitats was conducted to evaluate its genetic diversity by using the ISSR labeling technique. The findings will provide a theoretical basis for formulating *Astragalus* genetic diversity research, conservation strategies, and cross breeding.

ISSR was conducted by using a synthetic 16 - 18 bp nucleotide repeat sequence as a primer. Genomic DNA was amplified by PCR, which employed a primer anchored at 3 or 5 end of oligonucleotides. The ISSR combines the advantages of RAPD and SSR markers at the same level. Thus, it can produce more polymorphism than RAPD, and the reaction system is more sensitive, more stable, and has good repeatability [4]-[7]. This molecular marker has been widely used in studies on germplasm resource identification, phylogeny of species, plant taxonomy, evolution, and genetic diversity [8]-[10]. *Astragalus* has several variations of morphological features in the actual production, such as stems with red or green color and stems with two colors. Variation and degradation of the *Astragalus* germplasm may occur in the introduction and domestication. Thus, studying the genetic diversity of Radix *Astragali* at the DNA level is necessary. This study provides molecular-level evidence for *Astragalus* germplasm resources by using the ISSR molecular marker.

## 2. Materials and Methods

### 2.1. Plant Materials and DNA Extraction

A total of 95 Radix *Astragali* individuals were sampled representing 10 natural populations, including Shanxi, Jilin, Inner Mongolia, Gansu, Ningxia, and Shaanxi Provinces of China (Table 1). Approximately 1 g of fresh leaf samples from each individual was collected, placed in a plastic bag with silica gel to dry the samples quickly, and brought back to the laboratory for DNA isolation. Total genomic DNA was extracted using a protocol described by Doyle and Doyle [11], and 0.6% agarose gel electrophoresis and a biospec-mini DNA/RNA/protein analyzer were used to detect the quality and yield of DNA. The DNA was later diluted to 50 ng· $\mu\text{L}^{-1}$  working solution and placed in a refri

**Table 1.** Resources of the test samples.

Source	Species	Sample number
China Shanxi Hunyuan	<i>A. mongholicus</i>	33
China Inner Mongolia Hulunbeier	<i>A. mongholicus</i> , <i>A. membranaceus</i>	20
China Ningxia Guyuan	<i>A. membranaceus</i>	3
China Shandong Linyi	<i>A. membranaceus</i>	7
China Shaanxi Baoji	<i>A. mongholicus</i> , <i>A. membranaceus</i>	3
China Gansu Dingxi	<i>A. mongholicus</i> , <i>A. membranaceus</i>	7
China Hebei Anguo	<i>A. mongholicus</i> , <i>A. membranaceus</i>	8
China Heilongjiang Hailin	<i>A. membranaceus</i>	7
China Liaoning Beixi	<i>A. membranaceus</i>	5
China Jilin Tonghua	<i>A. membranaceus</i>	3

generator at 4 °C for future use.

## 2.2. Primer Selection and ISSR-PCR Amplification

To determine the appropriate primers suitable for *Astragali*, we screened 100 ISSR primers purchased from the website of the University of British Columbia in Canada (<https://www.ubc.ca/>) and Synthesis Project. For each primer pair, two samples were amplified, and their amplification products were run on 1% agarose gels. PCR amplifications were performed in a 20 µL reaction that contained 2 µL of 10 × PCR buffer, 2.5 mM dNTPs, 10 µM primer, 5 unit of Taq DNA polymerase (Taka Ra Biotechnology Co., Ltd.), and 50 ng of template DNA. Double distilled water was added to make the volume 20 µL. Amplifications were performed as follows: 95 °C for 5 min, 35 cycles of denaturation at 95 °C for 30 s, annealing at 53 °C - 57 °C for 40 s, extension at 72 °C for 40 s, and a final extension at 72 °C for 10 min. Afterward, the samples were held at 4 °C. Highly polymorphic primers with clear bands and clean background were selected to amplify the genomic DNA of all *Astragalus* species.

## 2.3. Data Analysis

The amplified DNA polymorphic fragments (bands) were scored as present (1) or absent (0), and the data matrix of the ISSR phenotypes was assembled for further analysis. The percentage of polymorphic bands (*PPB*), observed number of alleles (*Na*), effective number of alleles (*Ne*), Nei's genetic diversity index (*h*), and Shannon's information index (*I*) were calculated using POPGENE 1.32 software. NTSYSpc 2.10 software was used to calculate the genetic similarity (Gs) coefficient of the 183 *Astragalus* species. The UPGMA tree [12] was constructed using NTSYS 1.8 software [13] to examine the relationship of the populations.

## 3. Results

### 3.1. DNA Detection

The genomic DNA of the 95 *A. mongholicus* and *A. membranaceus* were extracted and detected by agarose gel electrophoresis. The results showed that the DNA bands were

clear, and no diffusion phenomena were observed. The extracted genomic DNA detected by a nucleic acid protein detector revealed that the ratio of D260 nm/D280 nm was between 1.8 and 2.0, and the yield was between 100 and 1800 ng/ $\mu$ L. These findings indicate that the genomic DNA extracted from the *Astragali* leaves had high quality and high yield.

### 3.2. ISSR Polymorphism

A total of 35 primers were screened from 100 ISSR primers, which were published by the Columbia University, to study the genetic diversity of the two populations of *Radix Astragali*. Among these primers, 25 were polymorphic and 10 were monomorphic. The *PPB* was 71.4%. A total of 273 amplified bands were obtained from the 95 *Radix Astragali* samples using 25 polymorphic ISSR primers (Table 2). Among these bands, 213 were polymorphic, and the *PPB* was 78%. The average value of the amplified bands was 10.9 for each primer, and the number varied from 4 to 20 (Table 3). UBC835 and UBS855 presented the highest number of bands (20/19), whereas UBC811 and UBC808 had the lowest number (4/5). The results showed a significant genetic difference.

**Table 2.** Primer name and sequences used in the ISSR analysis.

Primer	Primer sequence (5' $\rightarrow$ 3')	Tm ( $^{\circ}$ C)
UBC807	(AG) <sub>8</sub> T	52
UBC808	(AG) <sub>8</sub> C	55
UBC810	(GA) <sub>8</sub> T	52
UBC811	(GA) <sub>8</sub> C	55
UBC812	(GA) <sub>8</sub> A	52
UBC817	(CA) <sub>8</sub> A	52
UBC818	(CA) <sub>8</sub> G	55
UBC822	(TC) <sub>8</sub> A	52
UBC825	(AC) <sub>8</sub> T	52
UBC826	(AC) <sub>8</sub> C	55
UBC827	(AC) <sub>8</sub> G	55
UBC830	(TG) <sub>8</sub> G	55
UBC834	(AG) <sub>8</sub> YT	54
UBC835	(AG) <sub>8</sub> YC	56
UBC836	(AG) <sub>8</sub> YA	54
UBC840	(GA) <sub>8</sub> YT	54
UBC842	(GA) <sub>8</sub> YG	56
UBC849	(GT) <sub>8</sub> YA	54
UBC851	(GT) <sub>8</sub> YG	56
UBC855	(AC) <sub>8</sub> YT	54
UBC856	(AC) <sub>8</sub> YA	54
UBC857	(AC) <sub>8</sub> YG	56
UBC864	ATGATGATGATGATGATG	48
UBC899	CATGGTGTGGTCATTGTTCCA	58
UBC900	ACTCCCCACAGGTTAACACA	58

**Table 3.** Statistical results of polymorphic bands of ISSR primers.

Primers	Sequence	Nb	Np	PPB	Primers	Sequence	Nb	Np	PPB
UBC807	(AG) <sub>8</sub> T	11	8	73%	UBC830	(TG) <sub>8</sub> G	9	8	89%
UBC808	(AG) <sub>8</sub> C	5	3	60%	UBC834	(AG) <sub>8</sub> YT	14	13	93%
UBC810	(GA) <sub>8</sub> T	6	5	83%	UBC835	(AG) <sub>8</sub> YC	20	18	90%
UBC811	(GA) <sub>8</sub> C	4	3	75%	UBC836	(AG) <sub>8</sub> YA	15	12	80%
UBC812	(GA) <sub>8</sub> A	13	12	92%	UBC840	(GA) <sub>8</sub> YT	10	6	60%
UBC817	(CA) <sub>8</sub> A	7	4	57%	UBC842	(GA) <sub>8</sub> YG	14	13	93%
UBC818	(CA) <sub>8</sub> G	11	8	73%	UBC849	(GT) <sub>8</sub> YA	6	4	67%
UBC822	(TC) <sub>8</sub> A	10	6	60%	UBC851	(GT) <sub>8</sub> YG	7	5	71%
UBC825	(AC) <sub>8</sub> T	10	9	90%	UBC855	(AC) <sub>8</sub> YT	19	13	68%
UBC826	(AC) <sub>8</sub> C	9	6	67%	UBC856	(AC) <sub>8</sub> YA	12	10	83%
UBC827	(AC) <sub>8</sub> G	13	10	77%	UBC857	(AC) <sub>8</sub> YG	12	10	83%
UBC864	ATGATGATGATGATGATG						11	6	55%
UBC899	CAT GGT GTT GGT CAT TGT TCC A						11	9	82%
UBC900	ACT TCC CCA CAG GTT AAC ACA						14	12	86%
Total							273	213	/
Mean							10.9	8.5	78%

Note: Nb (Number of bands); Np (Number of polymorphic bands); PPB (Percentage of polymorphic bands).

### 3.3. Genetic Diversity of *Radix Astragali*

The binary data matrix was scored as present (1) or absent (0) of homologous bands and loaded into POPGENE 1.32 software to analyze the genetic diversity of *Radix Astragali*. The mean  $N_a$  was 1.9 in the two *Astragalus* species. For *A. membranaceus*, the  $N_e$ ,  $h$ , and  $I$  were 1.5318, 0.3109, and 0.4657, respectively, which were slightly lower than those of *A. mongholicus* ( $N_e$ : 1.5861,  $h$ : 0.3364, and  $I$ : 0.4969). These results indicate that *A. mongholicus* have higher genetic diversity than *A. membranaceus* (Table 4).

### 3.4. Genetic Diversity of Two *Radix Astragali* from Different Regions

*A. membranaceus* was collected from nine provinces including Inner Mongolia, Ningxia, Shandong, Shaanxi, Gansu, Hebei, Heilongjiang, Jilin, and Liaoning (Table 5). The highest number of polymorphic bands ( $N_p$ ) of 190 was observed in Inner Mongolia. The  $PPB$ ,  $h$ , and  $I$  were 89.20%, 0.2538, and 0.3714, respectively. The lowest  $N_p$  was found in Shaanxi with only 83. The  $PPB$ ,  $h$ , and  $I$  were 38.97%, 0.0861, and 0.1260, respectively. These results indicate the different origins and order of *A. membranaceus* genetic diversity from highest to lowest: Inner Mongolia, Heilongjiang, Ningxia, Shandong, Hebei, Gansu, Jilin, Liaoning, and Shaanxi. Meanwhile, *A. mongholicus* was collected from five provinces including Shanxi, Inner Mongolia, Shaanxi, Gansu, and Hebei (Table 5). The highest  $N_p$  of 203 was observed in Shanxi. The  $PPB$ ,  $h$ , and  $I$  were 95.3%, 0.4877, and 0.5212, respectively. The lowest  $N_p$  was observed in Shaanxi with

**Table 4.** Genetic diversity of two species of *radix astragali*.

Primers	<i>Astragalus membranaceus</i> (Fisch.) Bge.var. <i>Mongholicus</i> (Bge.) Hsiao				<i>Astragalus membranaceus</i> (Fisch.) Bge			
	N <sub>a</sub>	N <sub>e</sub>	<i>h</i>	<i>I</i>	N <sub>a</sub>	N <sub>e</sub>	<i>h</i>	<i>I</i>
UBC807	2	1.5279	0.308	0.4662	2	1.4935	0.2956	0.4554
UBC808	2	1.3987	0.2632	0.4241	2	1.5209	0.3165	0.4851
UBC810	2	1.5934	0.3624	0.5454	2	1.5239	0.3087	0.4641
UBC811	2	1.7518	0.4195	0.6075	2	1.3851	0.2692	0.4362
UBC812	2	1.5682	0.3437	0.5203	2	1.4753	0.2825	0.4322
UBC817	2	1.2239	0.1637	0.2867	1.8	1.139	0.1177	0.2221
UBC818	2	1.6939	0.3741	0.5381	2	1.6834	0.391	0.5748
UBC822	1.8	1.3434	0.2232	0.3495	2	1.3814	0.2471	0.398
UBC825	1.6	1.1139	0.0953	0.1768	1.6	1.1974	0.1371	0.2275
UBC826	2	1.7171	0.4152	0.6051	2	1.5242	0.3386	0.5193
UBC827	2	1.7609	0.4288	0.6194	2	1.5266	0.3215	0.4937
UBC830	1.8	1.6351	0.3441	0.4942	1.8	1.6555	0.3521	0.5033
UBC834	1.8	1.5235	0.3072	0.4552	1.8	1.6008	0.3366	0.4875
UBC835	1.6	1.4089	0.2362	0.3478	1.4	1.3159	0.1739	0.25
UBC836	2	1.6018	0.3687	0.553	2	1.7463	0.4165	0.604
UBC840	2	1.8589	0.4606	0.6529	2	1.9524	0.4877	0.6808
UBC842	2	1.7892	0.4287	0.6165	2	1.652	0.3687	0.5444
UBC849	1.8	1.7368	0.3831	0.5374	1.8	1.6119	0.3446	0.4974
UBC851	2	1.7605	0.4239	0.6128	2	1.7846	0.4258	0.6125
UBC855	2	1.7266	0.4081	0.5947	2	1.5569	0.3286	0.4987
UBC856	2	1.6157	0.3703	0.5543	2	1.3582	0.2359	0.3759
UBC857	2	1.8486	0.4558	0.6476	2	1.6908	0.3883	0.5699
UBC864	2	1.7503	0.4172	0.6041	2	1.8313	0.4514	0.6432
UBC899	1.6	1.5046	0.2727	0.3879	1.6	1.3558	0.2222	0.3339
UBC900	1.6	1.1981	0.136	0.2245	1.8	1.3306	0.2136	0.332
Mean	1.9	1.5861	0.3364	0.4969	1.9	1.5318	0.3109	0.4657

Note: N<sub>a</sub> (Number of Aelle), N<sub>e</sub> (Number of effect Aelle); *h* (Nei's gene diversity); *I* (Shannon's information index).

**Table 5.** Genetic diversity of *radix astragali* of different origins.

Origin	N <sub>a</sub>	N <sub>e</sub>	<i>h</i>	<i>I</i>	N <sub>p</sub>	PPB
<i>Astragalus membranaceus</i> (Fisch.) Bge						
Inner Mongolia	1.6452	1.4497	0.2538	0.3714	190	89.20%
Ningxia	1.3306	1.2338	0.1370	0.1999	117	54.93%
Shandong	1.4355	1.2919	0.1660	0.2445	135	63.38%
Shanxi	1.2086	1.1472	0.0861	0.1260	83	38.97%
Gansu	1.3710	1.2194	0.1288	0.1942	121	56.81%
Hebei	1.4274	1.2761	0.1615	0.2400	130	61.03%
Heilongjiang	1.6613	1.4024	0.2321	0.3465	181	84.98%
Jilin	1.3226	1.1913	0.1121	0.1686	111	52.11%
Liaoning	1.2339	1.1467	0.0868	0.1296	92	43.19%
Mean	1.4040	1.2621	0.1516	0.2245	129	60.51%
<i>A. membranaceus</i> (Fisch.) Bge.var. <i>Mongholicus</i> (Bge.) Hsiao						
Shanxi	1.8500	1.6157	0.4877	0.5212	203	95.31%
Inner Mongolia	1.8387	1.5368	0.3109	0.4613	194	91.08%
Shaanxi	1.2097	1.1483	0.0869	0.1268	87	40.85%
Gansu	1.3548	1.2196	0.1306	0.1955	115	53.99%
Hebei	1.3952	1.2528	0.1461	0.2174	127	59.62%
Mean	1.5297	1.3546	0.2324	0.3044	145	68.17%

Note: N<sub>a</sub> (Number of Aelle); N<sub>e</sub> (Number of effect Aelle); *h* (Nei's gene diversity); N<sub>p</sub> (Number of polymorphic bands); *I* (Shannon's information index); PPB (Percentage of polymorphic bands).

only 87. The *PPB*, *h*, and *I* were 40.85%, 0.0869, and 0.126, respectively. These results indicate the different origins and order of *A. mongholicus* genetic diversity from highest to lowest: Shanxi, Heilongjiang, Hebei, Gansu, and Shaanxi. At a general level, the average value (*h*: 0.1516, *I*: 0.2245, *PPB*: 60.51%) of the genetic diversity of *A. membranaceus* from the nine provinces was slightly lower than the value (*h*: 0.2324, *I*: 0.3044, *PPB*: 68.17%) of *A. mongholicus* from the five provinces.

### 3.5. Clustering Analysis of Radix Astragali

The *G<sub>s</sub>* coefficient of nine origins of *A. membranaceus* and five origins of *A. mongholicus* was calculated using NTSYSpc 2.10 software, and UPGMA was used to generate a dendrogram (Figure 1). The 14 different origins of two *Astragali* could be completely separated into two parts based on 0.80 coefficient using the 25 ISSR primers. One group cluster consisted of the northern regions including Inner Mongolia, Shanxi, Shandong, Heilongjiang, Hebei, Jilin, and Liaoning, and the other group cluster consisted of Ningxia, Gansu, and Shaanxi. This first group was clustered according to place of origin and conformed to the geographical characteristic of the traditional Chinese medicinal material geoherbalsism. The large group was also divided into two groups according to production. One group includes Shanxi and Inner Mongolia. Cluster was initially observed with *A. mongolian* of Shanxi and Inner Mongolia at 0.93 and then with *A. membranaceus* of Inner Mongolia at 0.91. The other group includes five districts, namely, Shandong, Heilongjiang, Hebei, Jilin, and Liaoning, which were mostly from *A. membranaceus*. The major second group mainly includes three origins, namely, Ningxia, Gansu, and Shaanxi. The membrane pod of *A. membranaceus* and *A. mongholicus* of Gansu Province was clustered as a group at 0.88. After Shaanxi *A. mongolian* clustering, *A. membranaceus* of Ningxia formed the second group.




Figure 1. Dendrogram of *G<sub>s</sub>* among the radix *astragali* from different habitats.

## 4. Discussion

### 4.1. Genetic Diversity

Genetic diversity is the product of the long-term evolution of a species or a population. It is the basis of existence, development, and evolution of species. Therefore, the study on the genetic diversity of species is fundamental to understand the taxonomy, origin, and evolution of species. Furthermore, such study will provide a theoretical basis for the germplasm resource conservation, development, utilization, and breeding [14]. In the present study, the genetic diversity of 95 *Astragalus* materials was analyzed by using 25 highly repeatable and polymorphic ISSR primers. *A. mongholicus* had higher genetic diversity ( $h$ : 0.3364,  $I$ : 0.4969) than *A. membranaceus* ( $h$ : 0.3109,  $I$ : 0.4657) at the species level (Table 4). The results of this study were consistent with the findings (*A. mongholicus*,  $h$ : 0.822; *A. membranaceus*,  $h$ : 0.689) of the research on genetic diversity of the two species by SSR molecular marker [15]. Compared with a relevant research [16], we speculated that *A. mongholicus* had a higher level of genetic diversity.

*Astragalus* mainly depends on insect for pollination and on humans for seed dispersal; it also possesses distant hybridization properties [17]-[19]. These life history traits are dominated by outcrossing breeding system and considered as an important factor to maintain higher genetic diversity of species [20]-[22]. The wild *Astragalus* resource gradually decreased mainly because of the recent over-excavation and habitat deterioration caused by human activity [23] [24]. However, the interference is likely only the effect of the number and genetic diversity of parts of a local area of *Astragalus*, in addition to the fact that the species are perennial herbaceous plants with high medicinal value. Thus, the large range of distribution, long life cycle, and germplasm (seed) exchanges in the development and propagation can be conducive to the genetic diversity at the species level [21] [25]. Therefore, the genetic diversity of the two species of *Astragalus* is unaffected by severe loss of habitat and human destruction.

### 4.2. Analysis and Diversity of Two Species of *Astragalus* in Different Habitats

The genetic diversity of different habitats of the two species of medicinal *Astragalus* was analyzed. The genetic diversity index of *A. membranaceus* was the highest in Heilongjiang ( $Np$ : 183,  $PPB$ : 86.13%,  $h$ : 0.2321,  $I$ : 0.3465) and the lowest in Shaanxi ( $Np$ : 83,  $PPB$ : 39.1%,  $h$ : 0.0861,  $I$ : 0.1260). Meanwhile, *A. mongholicus* polymorphic loci were the highest in Shanxi ( $Np$ : 203,  $PPB$ : 95.3%,  $h$ : 0.4877,  $I$ : 0.5212) and the lowest in Shaanxi ( $Np$ : 87,  $PPB$ : 40.97%,  $h$ : 0.0869,  $I$ : 0.1268). Thus, the genetic diversity level is basically consistent in *A. membranaceus* and *A. mongholicus*, which have higher genetic diversity in the genuine habitat. These phenomena may be attributed to the abundant germplasm resource and traditional history in development and utilization. At a general level, the genetic diversity of *A. membranaceus* from the nine provinces ( $h$ : 0.1516,  $I$ : 0.2245,  $PPB$ : 60.51%) was slightly lower than that of *A. mongholicus* from the five provinces ( $h$ : 0.2324,  $I$ : 0.3044,  $PPB$ : 68.17%). This result may be due to the fact that *A. mongholicus* has high utilization rate (faster regeneration) and germplasm diversity [15] [26] [27].



### 4.3. Analysis on the Genetic Relationships between Two Species of *Astragalus* in Different Habitats

Results showed that the two medicinal *Astragalus* species from 10 different habitats could be clustered into two groups (Figure 1). In these groups, major clustering was based on the geographical habitats (Shanxi and Inner Mongolia) of *A. membranaceus* and *A. mongholicus*. The morphological characteristics and classification status of the two species showed a very close genetic relationship [2]. In addition, Shanxi and Inner Mongolia are areas where authentic *Astragalus* herbal medicine is real estate areas. Thus, the first clustering was according to place of origin. However, Hebei *A. mongholicus* merged into different geographical positions in the clustering, which may be due to its mixed medicinal material source. Hebei is the largest business center; thus, seed and germplasm mix into the genetic information of the two species. Another group of three habitats of *A. membranaceus* and *A. mongholicus*, including Ningxia, Gansu, and Shaanxi, was clustered into one category from Gansu. The recorded change for the famous region of *Astragalus* in history is from Sichuan to Gansu, Shaanxi, Ningxia, Shanxi, Inner Mongolia, and gradual migration to Northeast Heilongjiang. Main cultivated varieties of *A. membranaceus* also replaced *A. Mongholicus*. Based on the classification of *A. mon-gholicus* from *A. membranaceus* variants, the genetic relationship is very close. The experimental results also presented that these species were divided according to origin and not from species distinction. This clustering also showed that the genetic relationship was close and may be related to the dominant characteristics of ISSR molecular markers.

### 4.4. Protection Strategy of *Astragalus* Germplasm Resources

In summary, two *Astragalus* species currently maintained a middle genetic diversity level. Genetic diversity is the basis of species to adapt to the environment and evolution to avoid extinction and survive. Genetic diversity also provides theoretical basis for further development and utilization of germplasm resources. The current anthropogenic deforesting and picking activities as well as habitat deterioration do not seriously affect the *Astragalus* genetic diversity. However, if the deterioration of the environment and damage induced by human activities are ineffectively prevented, the fragmentation of *Astragalus* habitat and population scale is bound to accelerate. This phenomenon can cause a genetic drift in its population and lead to a substantial loss of the current abundant genetic diversity. Such loss will affect the survival of the species and the availability of excellent germplasm resources. The main reasons of wild *Astragalus* resource loss are deliberate cutting and picking, as well as habitat destruction. We suggest that *Astragalus* protection area should be established as soon as possible in wild *Astragalus* habitat to prohibit unauthorized and wasteful mining. Detailed plan about development and utilization should also be designed. Considering that the current biological research on *Astragalus* is few, we suggest further research on the reproductive biology, conservation genetics, and physiological and ecological aspects to provide basis for the conservation and sustainable utilization of these species.

## Acknowledgements

This study was supported by the National Natural Science Foundation of China (31400285), the Chinese herbal medicine standardization production technology service platform [ministry of consumption (2011) 340], Breed Foundation of Shanxi Agricultural University (No. 2014YZ10).

## References

- [1] China Pharmacopoeia Committee (2015) Pharmacopoeia of People's Republic of China, Vol. 1. Chemical Industry Press, Beijing, 302-303.
- [2] Zhao, Y.Z. (2006) Taxonomy and Floristic Geographical Distribution of the Chinese Medicinal Huangqi. *Bulletin of Botanica Research*, **26**, 532-538.
- [3] Fu, L.G. (1992) In China Plant Red Data Book-Rare and Endangered Plants, Vol. 1. Science Press, Beijing, 372-373.
- [4] Gllbert, J.E., Lewis, R.V., Wilkinson, M.J. and Caligari, P.D.S. (1999) Developing an Appropriate Strategy to Assess Genetic Variability in Plant Germplasm Collections. *Theoretical and Applied Genetics*, **98**, 1125-1131. <http://dx.doi.org/10.1007/s001220051176>
- [5] Prevost, A. and Wilkinson, M.J. (1999) A New System of Comparing PCR Primers Applied to ISSR Fingerprinting of Potato Cultivars. *Theoretical and Applied Genetics*, **98**, 107-112. <http://dx.doi.org/10.1007/s001220051046>
- [6] Farsani, T.M., Etemadi, N., Sayed-Tabatabaei, B.E. and Talebi, M. (2012) Assessment of Genetic Diversity of Bermudagrass (*Cynodon dactylon*) Using ISSR Markers. *International Journal of Molecular Sciences*, **13**, 383-392. <http://dx.doi.org/10.3390/ijms13010383>
- [7] Omondi, E.O., Debener, T., Linde, M., Abukutsa-Onyango, M., Dinssa, F.F. and Winkelmann, T. (2016) Molecular Markers for Genetic Diversity Studies in African Leafy Vegetables. *Advances in Bioscience and Biotechnology*, **7**, 188-197. <http://dx.doi.org/10.4236/abb.2016.73017>
- [8] Ansari, S.A., Narayanan, C., Wali, S.A., Kumar, R., Shukal, N. and Rahangdale, S.K. (2012) ISSR Markers for Analysis of Molecular Diversity and Genetic Structure of Indian Teak (*Tectona grandis* L.f.) Populations. *Annals of Forest Research*, **55**, 11-23.
- [9] Narzary, D., Rana, T.S. and Ranade, S.A. (2012) Genetic Diversity in Inter-Simple Sequence Repeat Profiles across Natural Populations of Indian Pomegranate (*Punica granatum* L.). *Plant Biology*, **12**, 806-813. <http://dx.doi.org/10.1111/j.1438-8677.2009.00273.x>
- [10] Farsani, T.M., Etemadi, N., Sayed-Tabatabaei, B.E. and Talebi, M. (2012) Assessment of Genetic Diversity of Bermudagrass (*Cynodon dactylon*) Using ISSR Markers. *International Journal of Molecular Sciences*, **13**, 383-392. <http://dx.doi.org/10.3390/ijms13010383>
- [11] Doyle, J.J. and Doyle, J.L. (1987) A Rapid DNA Isolation Procedure for Small Quantities of Fresh Leaf Tissue. *Phytochemical Bulletin*, **19**, 11-15.
- [12] Nei, M. (1978) Estimation of Average Heterozygosity and Genetic Distance from a Small Number of Individuals. *Genetics*, **89**, 583-590.
- [13] Rohlf, F.J. (1994) NTSYS-pc, Numerical Taxonomy and Multivariate Analysis System. Ver. 1. 80. Exeter Software, New York.
- [14] Helenurm, K. (2001) High Levels of Genetic Polymorphism in the Insular Endemic Herb *Jepsonia malvifolia*. *Journal Heredity*, **92**, 427-432. <http://dx.doi.org/10.1093/jhered/92.5.427>
- [15] Wang, A. (2013) Genetic Diversity Analysis of *Astragalus mongholicus* and *A. membranaceus*. Minzu University of China, Beijing.

- [16] Jin, Z.F., Piao, X.C., Wu, S.Q. and Wu, R.Z. (2007) Analysis of Genetic Diversity in *Astragalus membranaceus* Germplasms by RAPD Markers. *Hubei Agricultural Sciences*, **46**, 875-876.
- [17] Feng, X.J. (2011) Study on Pollination Characteristic of Mongolia Milkvetch. *Modern Chinese Medicine*, **13**, 24-26.
- [18] Zhang, J., Yang, F.J. and Liu, S. (2010) Study on Pollinating in Sects of *Astragalus membranaceus* (Ficsh) Bunge. *Journal of Anhui Agricultural Sciences*, **38**, 2883-2885.
- [19] Wu, S.Q. and Lian, M.L. (2008) Study on Pollinating on Characteristic of *Radix astragali*. *Journal of Anhui Agricultural Sciences*, **36**, 9184-9303.
- [20] Zhang, J., Yang, F.J., Liu, S., Guo, S. and Ren, Y.Y. (2009) Breeding System Study in *Astragalus membranaceus* (Ficsh) Bunge. *Ginseng Research*, **3**, 17-20.
- [21] Nybom, H. and Bartish, I.V. (2000) Effects of Life History Traits and Sampling Strategies on Genetic Diversity Estimates Obtained with RAPD Markers in Plants. *Perspectives in Plant Ecology, Evolution and Systematics*, **3**, 93-114.  
<http://dx.doi.org/10.1078/1433-8319-00006>
- [22] Nybom, H. (2004) Comparison of Different Nuclear DNA Markers for Estimating Intraspecific Genetic Diversity in Plants. *Molecular Ecology*, **13**, 1143-1155.  
<http://dx.doi.org/10.1111/j.1365-294X.2004.02141.x>
- [23] Zhao, M., Duan, J.A., Huang, W.Z. and Zhou, R.H. (2000) The Analysis of *Astragalus* Linn. Medicinal Plants Resources Present Situation in Chinese. *Chinese Wild Plant Resources*, **19**, 5-9.
- [24] Zhang, L.T., Guo, B.L., Zhu, X.C. and Feng, G.Z. (2006) Survey of *Astragalus* germplasm Resources. *Traditional Chinese Medicinal Materials*, **29**, 771-773.
- [25] Hamrick, J.L. and Godt, M.J.W. (1996) Effects of Life History Traits on Genetic Diversity in Plant Species. *Philosophical Transaction: Biological Sciences*, **351**, 1291-1298  
<http://www.jstor.org/stable/56204> <http://dx.doi.org/10.1098/rstb.1996.0112>
- [26] Feng, X.J., Liu, G.K. and Liang, S.M. (2010) Review of Studies on Germplasm Resources of Mongolia Milkvetch. *Journal of Shanxi Agricultural Sciences*, **38**, 95-98.
- [27] Xie, X.L., Wang, X.S., Zhao, L., Wang, L. and Li, Y. (2004) Morphotype Diversity of *Astragalus Membranaceus* (Fisch.) Bge. Var. *Mongholicus* (Bge.) Hsiao. *Journal of Anhui Agricultural Sciences*, **32**, 1203-1204.



**Submit or recommend next manuscript to SCIRP and we will provide best service for you:**

Accepting pre-submission inquiries through Email, Facebook, LinkedIn, Twitter, etc.

A wide selection of journals (inclusive of 9 subjects, more than 200 journals)

Providing 24-hour high-quality service

User-friendly online submission system

Fair and swift peer-review system

Efficient typesetting and proofreading procedure

Display of the result of downloads and visits, as well as the number of cited articles

Maximum dissemination of your research work

Submit your manuscript at: <http://papersubmission.scirp.org/>

Or contact [abb@scirp.org](mailto:abb@scirp.org)