

Tracking Microorganisms in Production and Sale Operations of Spiced Geese

Huadong Xie¹, Lijun Bu¹, Zhengze Zhong¹, Yifan Zhang¹, Jun Lin², Zhixi Li^{3*}

¹Chongqing Academy of Animal Science, Key Laboratory of Pig Industry Sciences, Ministry of Agriculture, Chongqing Key Laboratory of Pig Industry Sciences, Chongqing, China; ²Chongqing Municipal Agricultural School, Chongqing, China; ³Department of Food Science and Engineering, Northwest A & F University, Xianyang, China.

Email: *lzx580721@yahoo.com.cn

Received June 14th, 2013; revised July 14th, 2013; accepted July 21st, 2013

Copyright © 2013 Huadong Xie *et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Cooling, transportation and sale processes of spiced geese were studied, eight spiced geese meat samples with different sampling time, Airborne microorganism samples of three different workplaces and five different environmental contact substance samples were test, measures of special mediums, biochemical identification and DNA sequencing were carried out, then *Escherichia coli*, *Yeast*, *Mildew*, *Lactic acid bacteria*, *Staphylococcus aureus* and *Janthinobacterium* were detected. For spiced geese meat samples, microorganisms were significant ($p < 0.05$) increased with the prolong of sampling time. *Lactic acid bacteria*, *Staphylococcus aureus* and *Janthinobacterium* were detected in each processing operation and the total aerobic counts of each sample was increased or significant ($p < 0.05$) increased with the prolong of sampling time; *Escherichia coli*, *Yeast* and *Mildew* were detected on samples entered into the retail outlet mainly and the total aerobic counts of each sample was increased or significant ($p < 0.05$) increased also. In the household workshop, *Mildew* and *Janthinobacterium* were the superior microorganisms. In the transport vehicle, *Staphylococcus aureus* and *Janthinobacterium* were the superior microorganisms; *Staphylococcus aureus* was the superior microorganism in the retail outlet. For environmental contact substances, Cooling platform, pallet, chopping block were the most serious contaminated environmental contact substances and the total bacteria counts were significant ($p < 0.05$) more than stainless steel barrel and chopper; *Janthinobacterium* was the superior microorganism on pallet, stainless steel barrel and chopper; *Lactic acid bacteria* was the superior microorganism on chopping block and stainless steel barrel; *Staphylococcus aureus* was the superior microorganism on cooling platform. Findings indicate that *Escherichia coli*, *Yeast*, *Mildew*, *Lactic acid bacteria*, *Staphylococcus aureus*, *Janthinobacterium* were the main microorganisms; Household workshop and the retail outlet were the main place microorganisms contaminated; Pallet, stainless steel barrel and chopper were the main environmental contact substances.

Keywords: Spiced Geese Meat; Microbial Contamination; Production Operations; Sale Operations

1. Introduction

China is the largest country with geese producing and consuming [1]. 2010 Statistical data (FAO) shown that, about 321.90 millions of geese amount of livestock on hand in China, 89.92% of the world's total amount of livestock on hand (about 357.98 millions); 601.65 millions geese were Slaughtered in China, 94.32% of the world's total amount of slaughtering (637.89 millions); 240.72 ten thousand tons geese meat were produced in China, 95.47% of the world's total amount of geese meat (252.14 ten thousand tons).

Market for geese is growing, its production and con-

sumption have increased 65.53 ten thousand tons since 2000 (FAO). Clearly, the continued growth and prosperity of the geese industry will depend, to a large degree, on its ability to supply the consumer with wholesome and safe products [2].

Spiced goose meat was one of the favorite spiced meat to Chinese because of the delicious taste and abundant nutrition [3].

In general case, traditional spiced geese meat were made in household workshop, sold in the retail stalls directly and without any microbial prevention or control measures during the whole process, such as packaging and sterilization, so microbiological safety can't be safeguard [4].

*Corresponding author.

Even though the nutritional value of geese meat is well documented, very little information is available worldwide on the microbiological aspects, especially of the spiced geese meat and the production process. The purpose of the present study was to examine the number and types of microorganisms associated with spiced geese meat and the production process. These findings may provide information that will be useful in controlling contamination during processing and extending the shelf life of spiced geese meat [5].

2. Materials and Methods

2.1. Samples

All spiced geese meat samples, airborne microorganisms samples and environmental contact substances samples were collected from a famous household workshop in Rongchang county Chongqing (China) and one of the retail outlets.

2.2. Spiced Geese Meat Sampling

According to the production process and the HACCP analysis, cooling, transportation and sale process should be the critical points of microbial contamination. So time points when products at the 5th and the 30th minute in cooling process, when products in transport process, when products at the 0th h, 1st h, 2nd h, 3rd h, 4th h in the retail stores were selected as the sampling time, corresponding samples were named C5 min, C30 min, T, S0 h, S1 h, S2 h, S3 h, S4 h.

Three parallel samples were collected at each sampling time, each sample was placed in a separate sterile plastic bag and transported to the laboratory immediately for analysis.

A 1g sample of each obtained spiced geese meat was taken aseptically by scalpel excision and placed in a sterile conical flask with 9 ml sterile saline as bacterial suspension. Decimal dilutions were carried out using the same diluents with pure water [6].

2.3. Airborne Microorganisms Sampling

Airborne microorganisms in the household workshop, transport vehicle and retail outlet were collected.

Natural sedimentation method has been used for the collection of air microorganism samples according to the national standard of the People's Republic of China GB/T18204.1-2000. In order to make air flow minimized, points where keep more than 1 meters distance away from wall and far away from the vents were selected. Then culture dishes (diameter 9 centimeters) with different special medium were exposed 5 minutes on the sample points. Three parallel samples were collected on each sampling point, then transported to the laboratory immediately [7].

Plates were incubated at 37°C for 48 hours, colony forming units (CFU) on the plates were counted and total aerobic counts per 1 m³ was determined according the following formula.

$$\text{Total aerobic counts (CFU/m}^3\text{)} = 5000 \frac{N}{A} \times T$$

A: represent area of the culture dish (cm²);

T: represent culture dish exposure time (min);

N: represent mean colony content (CFU).

2.4. Environmental Contact Substances Sampling

Microbial samples on cooling platform, stainless steel barrel, pallet, chopping block, chopper were collected.

Sampling plane with 100 cm² was selected randomly on surface of each contact substances. Four sterile cotton balls were used to wipe the sampling plane, then the cotton balls were put into a sterile conical flask with 20 ml sterile saline as bacterial suspension. Three parallel samples were collected of each sampling plane, then transported to the laboratory immediately after collection.

A 200 µl sample of each 20 ml bacterial suspension was taken aseptically, decimal dilutions were carried out using the same diluents with pure water [8].

2.5. Microbiological Identification and Enumeration

Bacteria were enumerated on six different media. Total aerobic counts were determined using Nutrient agar (Hangzhou Microbial Reagent CO., Ltd.) spread plates incubated at 37°C for 48 hours [2]; *Escherichia coli* (*E. coli*) and *Salmonella* were determined using Maconkey agar (Beijing Aobox Biotechnology Co., Ltd.) spread plates incubated at 37°C for 48 hours (AduGyamfil, Torgby-Tetteh&Appiah, 2012); *Fungus* was determined by the spread plate method using improved Martin medium (Hangzhou Microbial Reagent Co., Ltd.); *Lactic acid bacteria* was determined using MRS agar (Beijing Aobox Biotechnology Co., Ltd.) spread plates incubated at 37°C for 48 hours; *Staphylococcus aureus* (*S. aureus*) was determined by the spread plate method using Baird-Parker agar base. The plates were incubated at 37°C for 48 h [9]. In order to determine *S. aureus* counts, random isolates from suitable plates were picked, purified and tested for electron microscopy, gram stain [10], Mannitol fermentation and catalase activity [11]; *Janthinobacterium* was determined using medium with peptone 20 g, potassium dihydrogenphosphate 1.5 g, magnesium sulfate 1.5 g, agar medium 15 g, distilled water 1000 ml, pH: 6.9 - 7.1, spread plates incubated at 37°C for 48 hours. In order to determine *Janthinobacterium* counts, random isolates from suitable plates were picked, purified and tested for gram stain [10], electron microscopy, urea de-

composition, starch hydrolysis, hydrogen sulfide production, lactose hydrolysis, sucrose hydrolysis and indole production. Furthermore, PCR/RFLP marker of *Janthinobacterium* was cloned and 16S rDNA was sequenced.

2.6. Statistical Analysis

For spiced geese meat sample, total aerobic counts were transformed to \log_{10} CFU/g. For Airborne microorganisms counts, the data were transformed to \log_{10} CFU/m³. For environmental contact substance counts, the data were also transformed to \log_{10} CFU/cm² so as to enable a true comparison of the different counts reported by other authors with that determined in this study [12].

One-Way Analysis of Variance (ANOVA) was performed, and if the ANOVA detected significant differences in group means, the Duncan Multiple Comparisons Test was used to determine which treatment groups differed significantly. All significant differences were determined at $p < 0.05$.

Analysis was carried out using the "SAS 8. 2" software package (SAS Institute Inc., Cary, NC, USA) for Windows XP.

3. Results and Discussion

3.1. Microorganism Identification

Special mediums were used to identify the microorganisms isolated from different samples, furthermore methods of gram stain and electron microscopy were also used, results were shown in **Figure 1**. The preliminary identification results indicated that *E. coli*, *Yeast*, *Mildew*, *Lactic acid bacteria*, *S. aureus*, *Janthinobacterium* were the contamination microorganisms.

In order to make sure the results correct, biochemical identification was carried out on *S. aureus* and *Janthinobacterium*, the results (**Table 1**) indicated that, biochemical identification results of *S. aureus* and *Janthinobacterium* were agreed with the reference answers. In order to verify the results, method of DNA sequencing was carried out, the PCR fragments of *Janthinobacterium* were shown in **Figure 2**. DNA sequence was obtained and compared to the Basic Local Alignment Search Tool (BLAST), the result shown that the obtained DNA sequence has 99% similarity with *Janthinobacterium*.

3.2. Microbial Quality of Geese Meat Samples

Microbial loads of spiced geese meat samples were given in **Table 2**, the total aerobic counts for C5 min, C30 min, T, S0 h, S1 h, S2 h, S3 h, S4 h were 2.68, 3.04, 3.16, 3.54, 4.07, 3.95, 4.21 and 4.86 \log_{10} CFU/g respectively; The mean total *E. coli*, *Yeast*, *Mildew*, *Lactic acid bacteria*, *S.*




Figure 1. Gram stain and electron microscopy results of *S. aureus* and *Janthinobacterium*. (a) *Staphylococcus aureus*; (b) *Janthinobacterium*.




Figure 2. 16S rDNA fragments of agarose gel electrophoresis of *Janthinobacterium*.

Table 1. Biochemical identification results of *S. aureus* and *Janthinobacterium*.

Microbial Category	Test Projects	Results	Reference Results
<i>S. aureus</i>	Electron microscopy	Spherical, without Spores and capsular	Spherical., without Spores and Capsular
	Gram stain	+	+
	Mannitol fermentation	+	+
	Catalase activity	-	-
	Thrombin activity	+	+
	Electron microscopy	Double spherical	Double spherical
	Gram stain	-	-
	Ureade composition	+	+
<i>Janthinobacterium</i>	Starchhydrolysis	-	-
	Hydrogensulfide production	-	-
	Lactose hydrolysis	-	-
	Sucrose hydrolysis	-	-
	Indole production	+	+

Table 2. The results of contaminated microorganism on spiced goose meat with different sampling time. Unit: log₁₀CFU/g.

	C _{5min}	C _{30min}	T	S _{0h}	S _{1h}	S _{2h}	S _{3h}	S _{4h}
Total aerobic counts	2.68 ± 0.014 ^H ^a	3.04 ± 0.021 ^G ^a	3.16 ± 0.0065 ^F ^b	3.54 ± 0.024 ^E ^b	4.07 ± 0.043 ^C ^b	3.95 ± 0.017 ^D ^b	4.21 ± 0.016 ^B ^a	4.86 ± 0.014 ^A ^a
<i>E. coli</i>	2.00 ± 0.066 ^D ^b	0.00 ± 0.00 ^E ^d	0.00 ± 0.00 ^E ^d	0.00 ± 0.00 ^E ^c	3.54 ± 0.024 ^C ^c	3.54 ± 0.029 ^C ^d	3.83 ± 0.011 ^B ^c	4.48 ± 0.018 ^A ^c
<i>Salmonella</i>	0.00 ± 0.00 ^A ^d	0.00 ± 0.00 ^A ^d	0.00 ± 0.00 ^A ^d	0.00 ± 0.00 ^A ^c	0.00 ± 0.00 ^A ^f	0.00 ± 0.00 ^A ^g	0.00 ± 0.00 ^A ^e	0.00 ± 0.00 ^A ^g
<i>Yeast</i>	1.69 ± 0.088 ^E ^c	0.00 ± 0.00 ^F ^d	0.00 ± 0.00 ^F ^d	0.00 ± 0.00 ^F ^e	2.90 ± 0.016 ^D ^c	3.30 ± 0.025 ^C ^f	3.61 ± 0.028 ^B ^d	4.05 ± 0.052 ^A ^e
<i>Mildew</i>	0.00 ± 0.00 ^B ^d	0.00 ± 0.00 ^B ^d	0.00 ± 0.00 ^B ^d	0.00 ± 0.00 ^B ^c	0.00 ± 0.00 ^B ^f	0.00 ± 0.00 ^B ^g	0.00 ± 0.00 ^B ^e	3.09 ± 0.043 ^F ^f
<i>Lactic acid bacteria</i>	0.00 ± 0.00 ^H ^d	1.70 ± 0.056 ^G ^c	2.00 ± 0.035 ^F ^c	2.98 ± 0.016 ^E ^d	3.39 ± 0.016 ^D ^d	3.48 ± 0.015 ^C ^e	4.20 ± 0.015 ^B ^a	4.63 ± 0.037 ^A ^b
<i>S. aureus</i>	2.65 ± 0.019 ^E ^b	3.00 ± 0.020 ^F ^d	3.23 ± 0.014 ^D ^b	3.90 ± 0.011 ^C ^c	4.12 ± 0.0063 ^C ^d	4.18 ± 0.021 ^B ^c	4.16 ± 0.012 ^A ^b	4.89 ± 0.021 ^A ^d
<i>Janthinobacterium</i>	1.97 ± 0.19 ^G ^a	0.00 ± 0.00 ^F ^b	3.18 ± 0.021 ^E ^a	3.32 ± 0.027 ^D ^a	3.41 ± 0.019 ^C ^a	3.90 ± 0.011 ^B ^a	4.15 ± 0.013 ^B ^b	4.18 ± 0.032 ^A ^a

Values are log₁₀ (mean CFU/g) ± SD; Different subscript uppercase letters in the same raw indicated significantly ($p < 0.05$) different; Different superscript lower case letters in the same column indicated significantly ($p < 0.05$) different.

aureus, *Janthinobacterium* count for C5 min, C30 min, T, S0 h, S1 h, S2 h, S3 h, S4 h were 2.00, 0, 0, 0, 3.54, 3.54, 3.83 and 4.48 log₁₀CFU/g; 1.69, 0, 0, 0, 2.90, 3.30, 3.61 and 4.05 log₁₀CFU/g; 1.69, 0, 0, 0, 2.90, 3.30, 3.61 and 4.05 log₁₀CFU/g; 0, 1.70, 2.00, 2.98, 3.39, 3.48, 4.20 and 4.63 log₁₀CFU/g; 1.97, 0, 3.18, 3.32, 3.41, 3.90, 4.15 and 4.18 log₁₀CFU/g; 2.65, 3.00, 3.23, 3.90, 4.12, 4.18, 4.16 and 4.89 log₁₀CFU/g respectively; *Salmonella* not detected on each samples.

Table 2 indicated that total aerobic counts was significant ($p < 0.05$) increased of each spiced geese meat sample. *Lactic acid bacteria*, *S. aureus* and *Janthinobacterium* were detected in each processing operations and the total aerobic counts of each was increased or signifi-

cant ($p < 0.05$) increased; *E. coli*, *Yeast* and *Mildew* were detected on samples entered into the retail outlet mainly and the total aerobic counts of each was increased or significant ($p < 0.05$) increased also.

3.3. Airborne Microorganisms Quality

Results of airborne microorganisms in the household workshop, transport vehicle, retail outlet were shown in **Table 3**. The results indicated that, total aerobic counts for household workshop, transport vehicle, retail outlet were 3.96, 3.04 and 3.54 log₁₀CFU/g respectively; The mean total *E. coli*, *Yeast*, *Mildew*, *Lactic acid bacteria*, *S. aureus*, *Janthinobacterium* count for household workshop, transport vehicle, retail outlet were 2.03, 0 and 2.56

\log_{10} CFU/g, 2.62, 2.00 and 0 \log_{10} CFU/g, 3.16, 2.88 and 2.50 \log_{10} CFU/g, 2.98, 2.20 and 2.75 \log_{10} CFU/g, 3.46, 3.05 and 2.86 \log_{10} CFU/g, 2.87, 3.11 and 3.18 \log_{10} CFU/g respectively;

In the household workshop, *Janthinobacterium* was the most serious contamination microorganisms, while there were no significant differences ($p > 0.05$) on contamination of *Yeast*, *Mildew*, *Lactic acid bacteria* and *S. aureus*; In the transport vehicle, *Mildew*, *S. aureus*, *Janthino bacterium* were the most serious contamination microorganisms; In the retail outlet, *S. aureus* was the most serious contamination microorganism, the total aerobic counts of *S. aureus* was significant ($p < 0.05$) superior to other microorganisms.

Compared to air in transport vehicle and retail outlet, air in household workshop contaminated more microorganisms.

3.4. Microbial Quality of Environmental Contact Substances

Different environmental contact substances such as cooling platform, pallet, stainless steel barrel, chopper and chopping block were studied, the results were shown in **Table 4**. **Table 4** shown that, total aerobic counts for cooling platform, pallet, stainless steel barrel, chopper and chopping block were 3.70, 3.65, 3.52, 1.94, 3.72 \log_{10} CFU/cm² respectively; The mean total *E. coli*, *Yeast*, *Mildew*, *Lactic acid bacteria*, *S. aureus*, *Janthinobacterium* count for cooling platform, pallet, stainless steel barrel, chopper and chopping block were 3.60, 3.48, 3.25, 1.36, 3.68 \log_{10} CFU/cm², 0, 0, 1.10, 0, 0 \log_{10} CFU/cm², 3.54, 3.08, 3.54, 1.10, 4.00 \log_{10} CFU/cm², 3.60, 3.60, 3.54, 2.33, 3.64 \log_{10} CFU/cm², 3.65, 3.17, 3.48, 1.00, 3.56 \log_{10} CFU/cm² respectively.

Table 3. The results of contaminated microorganism in different workplace. Unit: \log_{10} CFU/m³.

	Household workshop	Transport vehicle	Retail outlet
Total aerobic counts	3.96 ± 0.14 _A ^a	3.04 ± 0.45 _B ^a	3.54 ± 0.18 _{AB} ^a
<i>E. coli</i>	2.03 ± 0.96 _A ^d	0.00 ± 0.00 _B ^c	2.56 ± 0.10 _A ^d
<i>Salmonella</i>	0.00 ± 0.00 _A ^e	0.00 ± 0.00 _A ^c	0.00 ± 0.00 _A ^e
<i>Yeast</i>	2.62 ± 0.10 _A ^{cd}	2.20 ± 0.00 _B ^b	0.00 ± 0.00 _C ^e
<i>Mildew</i>	3.16 ± 0.16 _A ^{bc}	2.88 ± 0.14 _B ^a	2.50 ± 0.00 _C ^d
<i>Lactic acid bacteria</i>	2.98 ± 0.15 _A ^{bc}	2.20 ± 0.00 _B ^b	2.75 ± 0.13 _B ^c
<i>S. aureus</i>	2.87 ± 0.34 _A ^{bc}	3.11 ± 0.19 _{AB} ^a	3.18 ± 0.09 _B ^b
<i>Janthinobacterium</i>	3.46 ± 0.28 _A ^{ab}	3.05 ± 0.22 _A ^a	2.86 ± 0.06 _A ^c

Values are \log_{10} (mean CFU/g) ± SD; Different subscript uppercase letters in the same raw indicated significantly ($p < 0.05$) different; Different superscript lower case letters in the same column indicated significantly ($p < 0.05$) different.

Table 4. The results of contaminated microorganism on different environmental contact substances. Unit: \log_{10} CFU/cm².

	Cooling platform	Pallet	stainless steel barrel	chopper	chopping block
Total aerobic counts	3.70 ± 0.01 7 _A ^a	3.65 ± 0.0048 _A ^a	3.52 ± 0.026 _B ^a	1.94 ± 0.12 _C ^b	3.72 ± 0.008 _A ^b
<i>E. coli</i>	3.60 ± 0.011 _A ^c	3.48 ± 0.017 _B ^b	3.25 ± 0.022 _C ^b	1.36 ± 0.10 _D ^c	3.68 ± 0.014 _A ^c
<i>Salmonella</i>	0.00 ± 0.00 _A ^f	0.00 ± 0.00 _A ^f	0.00 ± 0.00 _A ^c	0.00 ± 0.00 _A ^f	0.00 ± 0.00 _A ^f
<i>Yeast</i>	2.82 ± 0.015 _B ^e	2.75 ± 0.042 _B ^e	2.90 ± 0.016 _B ^c	1.26 ± 0.24 _C ^{cd}	3.62 ± 0.010 _A ^d
<i>Mildew</i>	0.00 ± 0.00 _B ^f	0.00 ± 0.00 _B ^f	1.10 ± 0.017 _A ^d	0.00 ± 0.00 _B ^f	0.00 ± 0.00 _B ^f
<i>Lactic acid bacteria</i>	3.54 ± 0.019 _B ^d	3.08 ± 0.025 _C ^d	3.54 ± 0.006 _B ^a	1.10 ± 0.17 _D ^{de}	4.00 ± 0.017 _A ^a
<i>S. aureus</i>	3.65 ± 0.024 _A ^b	3.17 ± 0.073 _D ^c	3.48 ± 0.029 _C ^a	1.00 ± 0.00 _E ^e	3.56 ± 0.006 _B ^c
<i>Janthinobacterium</i>	3.60 ± 0.011 _A ^c	3.60 ± 0.022 _A ^a	3.54 ± 0.019 _A ^a	2.33 ± 0.14 _B ^a	3.64 ± 0.027 _A ^d

Values are \log_{10} (mean CFU/g) ± SD; Different subscript uppercase letters in the same raw indicated significantly ($p < 0.05$) different; Different superscript lower case letters in the same column indicated significantly ($p < 0.05$) different.

Cooling platform, pallet, chopping block were the most serious contaminated environmental contact substances and total aerobic counts were significant ($p < 0.05$) more than stainless steel barrel and chopper.

Janthinobacterium was the superior microorganism on pallet, stainless steel barrel and chopper; *Lactic acid bacteria* was the superior microorganism on chopping block and stainless steel barrel; *S. aureus* was the superior microorganism on cooling platform.

4. Acknowledgements

We are grateful to the earmarked fund for Modern Agro-industry Technology Research System (CARS-43-15) and Chongqing Fundamental Research Project (11616, 11622, 12619 and 13419).

REFERENCES

- [1] Y. Liu and J. W. Ren, "Primary Approach to Promote Goose Farming in Shanxi Province," *Journal of Shanxi Agricultural Sciences*, Vol. 39, No. 5, 2011, pp. 474-476.
- [2] A. C. Carlos, M. F. Beatriz, P. Miguel and R. Capita, "Microbiological Quality of Vacuum-Packed Retail Ostrich Meat in Spain," *Food Microbiology*, Vol. 21, No. 2, 2004, pp. 241-246. [doi:10.1016/S0740-0020\(03\)00060-1](https://doi.org/10.1016/S0740-0020(03)00060-1)
- [3] J. W. Ren, Y. Liu, R. Q. Fan and S. Y. Yuan, "How to Utilize Genetic Resources of Goose in China to Progress Goose Industry in Shanxi Province," *Journal of Shanxi Agricultural Sciences*, Vol. 39, No. 12, 2011, pp. 1337-1340.
- [4] L. J. Bu, T. Xiong, J. Lin, B. Z. Lin and H. D. Xie, "Research on Major Pollution Microorganisms and Growth Law in Spiced Goose during the Production and Sales Process," *Academic Periodical of Farm products Processing*, Vol. 9, 2013, in press.
- [5] A. Hinton Jr., J. A. Cason and D. I. Kimberly, "Tracking Spoilage Bacteria in Commercial Poultry Processing and Refrigerated Storage of Poultry Carcasses," *International Journal of Food Microbiology*, Vol. 91, No. 2, 2004, pp. 155-165. [doi:10.1016/S0168-1605\(03\)00377-5](https://doi.org/10.1016/S0168-1605(03)00377-5)
- [6] M. Alvarez-Astorga, R. Capita, C. Alonso-Calleja, B. Moreno and M. C. García-Fernández, "Microbiological Quality of Retail Chicken By-Products in Spain," *Meat Science*, Vol. 62, No. 1, 2002, pp. 45-50. [doi:10.1016/S0309-1740\(01\)00225-X](https://doi.org/10.1016/S0309-1740(01)00225-X)
- [7] R. Fries and C. Graw, "Water and Air in Two Poultry Processing Plants' Chilling Facilities—A Bacteriological Survey," *British Poultry Science*, Vol. 40, No. 1, 1999, pp. 52-58. [doi:10.1080/00071669987836](https://doi.org/10.1080/00071669987836)
- [8] C. J. Thomas and T. A. McMeekin, "Contamination of Broiler Carcass Skin during Commercial Processing Procedures: An Electron Microscopic Study," *Applied and Environmental Microbiology*, Vol. 40, No. 1, 1980, pp. 133-144.
- [9] G. A. Lancette and S. R. Tatini, "*Staphylococcus Aureus*," In: C. Vanderzant and D. F. Splittstoesser, Eds., *Compendium of Methods for the Microbiological Examination of Foods*, APHA, Washington DC, 1992, p. 533.
- [10] W. F. Harrigan and M. E. McCance, "Métodos de Laboratorio en Microbiología de Alimentos y Productos Lácteos," Academia, León, 1976, p. 9.
- [11] S. T. Cowan, "Cowan and Steel's Manual for the Identification of Medical Bacteria," Cambridge University Press, Cambridge, 1974, p. 163.
- [12] A. Adu-Gyamfi, E. Torgby-Tetteh and V. Appiah, "Microbiological Quality of Chicken Sold in Accra and Determination of D₁₀-Value of *E. coli*," *Food and Nutrition Sciences*, Vol. 3, No. 5, 2012, pp. 693-698. [doi:10.4236/fns.2012.35094](https://doi.org/10.4236/fns.2012.35094)