

Activation of Various Downstream Signaling Molecules by IGFBP-3

Hanief Mohammad Shahjee^{1,2*}, Nisan Bhattacharyya^{1,3}

¹Diabetes Branch, NIDDK, National Institutes of Health, Bethesda, USA

²Present Address: Department of Biochemistry and Biophysics, University of Rochester Medical Center, Rochester, USA

³Present Address: Skeletal Clinical Studies Unit, CSDB, National Institute of Dental and Craniofacial Research, National Institutes of Health, Bethesda, USA

Email: hanief_shahjee@urmc.rochester.edu

Received 1 June 2014; revised 1 July 2014; accepted 28 July 2014

Copyright © 2014 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Insulin-like growth factor binding protein-3 (IGFBP-3), a secretory protein, is the most abundant IGF binding protein present in human serum among all IGF binding proteins. IGFBP-3 shows decreased level of expression in cancerous cells but has been known to be present in significant amounts in normal or non-cancerous cells. IGFBP-3 can induce apoptosis in prostate cancer cells either in an IGF-dependent manner or independently of IGF binding. Although putative cell death specific Insulin-like growth factor binding protein-3 (IGFBP-3R) receptor(s) has recently been identified by which IGFBP-3 may induce its anti-tumor effects, IGFBP-3 has also been known to activate various downstream intracellular signaling molecules via a different mechanistic pathway. Stat-1 has been known to be one of the candidate molecules activated by IGFBP-3. IGFBP-3 can also inhibit Akt/IGF-1 survival pathway in MCF-7 breast cancer cells which ultimately leads to the induction of apoptosis in these cells. All these studies clearly demonstrate that IGFBP-3 regulates cell proliferation and promotes its pro-apoptotic effects in cancer cells in two different pathways: 1) sequester IGF-I to bind to IGF-I receptor to inhibit cell proliferation and induce apoptosis, 2) independent of IGF-I pathway, IGFBP-3 binds to some putative receptor and activate various downstream pro-apoptotic molecules involved in cell death.

Keywords

Apoptosis, IGFBP-3, Stat-1, IGF-I, TGF- β

*Corresponding author.

1. IGFBP-3 Expression and Function

Insulin-like growth factor binding protein-3 (IGFBP-3) is one of the six known IGF binding proteins present in human serum and is comprised of 264-amino acid mature protein, with a 27-amino acid signal peptide [1]. IGFBP-3 is expressed in a wide variety of tissues. Epidemiological studies have clearly suggested that there is an inverse relationship between IGFBP-3 levels and occurrence of cancers [2]. Increased expression levels of serum IGFBP-3 result in decreased prevalence of prostate [3] [4] and colorectal [5] [6] cancers. It has also been shown that IGFBP-3 over expression results in decreased tumor formation in xenografts of non-small lung cancer cells [7] and M12 human prostate cancer cells [8]. Patient samples with hepatocellular and non-small cell lung carcinoma have been shown to have decreased expression of IGFBP-3 [9] [10]. Besides, it has also been reported that there is an increased level of IGFBP-3 in senescence [11]. In addition, IGFBP-3 expression level was decreased in cells immortalized with the human papillomavirus type 16 oncoprotein E7 due to proteasomal degradation [12]. Although IGFBP-3 is a secretory protein, but active protein could also be found in nucleus and cytoplasm [13]. Besides its full length form, IGFBP-3 could also be found in an N-terminal truncated form [14] [15].

IGFBP-3 has been known to play an important role in cell proliferation by inducing its anti-proliferative and pro-apoptotic effects in breast and prostate cancer cells [16]-[18]. It has also been shown that IGFBP-3 not only triggers growth inhibitory effects by inducing apoptosis, but also can function in G1 cell cycle arrest in human breast, kidney and lung cancer cells [19] [20]. IGFBP-3 may also play an important role in mediating inhibitory effects of transforming growth factor (TGF)- β (Figure 1), retinoic acid, tumor necrosis factor (TNF)- α and p53 [16] [21]-[26], but modulation in IGFBP-3 synthesis or action could regulate these anti-proliferative effects.

2. IGF-Dependent and Independent Effect of IGFBP-3

IGF dependent studies revealed that Insulin-like growth factor binding protein (IGFBP)-3 could induce apoptosis by binding to IGF-I and form a binary complex with IGF-I and prevent it to activate (IGF-IR) IGF-1 receptor (Figure 2) to stimulate cell proliferation and survival [27] [28]. In addition, Insulin-like growth factor binding protein (IGFBP)-3 has been shown to potently inhibit cell proliferation and induce apoptosis in an insulin-like growth factor (IGF)-independent manner [13] [18] [29]. IGFBP-3 in the media may exert its pro-apoptotic effects by binding to some cell surface receptors resulting in the activation of various signal transduction pathways.




Figure 1. Key candidate molecules activated by IGFBP-3. IGFBP-3 can activate various downstream signaling molecules either by binding to its putative IGFBP-3R (receptor) to monitor its signal and induce apoptosis or bind to TGF- β receptor resulting in Smad activation which leads to apoptosis. IGFBP-3 can also activate Stat-1 or bind to RXR- α to induce its anti-proliferative and pro-apoptotic effects.




Figure 2. IGF dependent and independent effects of IGFBP-3. IGF-1 can bind to IGF-I receptor and stimulate cell proliferation. IGFBP-3 blocks IGF-I to bind to its receptor resulting in the induction of apoptosis. IGFBP-3 can also induce apoptosis on its own by IGF-I independent mechanism.




Figure 3. IGFBP-3 entry into the cell by different pathways. IGFBP-3 in the media seemingly binds with plasma membrane IGFBP-3 receptor to activate signal transduction pathway or enters into the cell by endocytosis to induce apoptosis. IGFBP-3 may either directly activate signal transduction pathway by binding to some receptor to induce apoptosis or in the form of vesicular IGFBP-3 which enters through endoplasmic reticulum membrane into the cytosol and possibly gets translocated into the nucleus.

Alternatively, IGFBP-3 may possibly enter into the cell by endocytosis (**Figure 3**). It has also been reported in some other studies that IGFBP-3 binds to RXR α (**Figure 1**) and induce its pro-apoptotic effects in PC-3 human prostate cancer cells [30]. Surprisingly, recent studies have shown that IGFBP-3 fail to gain entry or be internalized upon binding to IGF-1 in the extracellular media to induce its pro-apoptotic effects in non-transformed mammary epithelial cells [31].

3. Pathways Involved in IGFBP-3 Signaling

IGFBP-3 action results in the activation of various signal transduction pathways [32] and Stat-1 (signal transducer and activator of transcription 1) has been known to have a functional role in IGFBP-3-induced apoptosis

(Figure 1) in rat chondroprogenitor cells [33], although our previous studies [34] showed a protective role of Stat-1 on IGFBP-3 induced apoptosis in PC-3 human prostate cancer cells, implying the fact that role of Stat-1 may be cell type dependent.

Some other downstream signaling molecules of IGFBP-3 have also been reported in various other studies [16] [35] [36]. It has been shown from these studies that IGFBP-3 can bind to transforming growth factor- β (TGF- β) cell surface receptors and one of the studies have shown direct interaction of IGFBP-3 with TGF- β -receptor typeV (TGF- β -RV) in mink lung epithelial cells [37].

IGFBP-3 has also been known to bind and activate intracellular signaling by forming a hetero-meric complex with other TGF- β receptors (TGF- β RII and TGF- β RI) in T47D breast cancer cells [35] [36]. Besides, addition of IGFBP-3 resulted in the activation of Smad 2 phosphorylation and cell growth inhibition in these cells. It has also been shown that IGFBP-3 mediates its pro-apoptotic effects via TGF- β in PC-3 human prostate cancer cells [16]. Results from our previous studies indicated the inhibition of TGF- β signaling in presence of IGFBP-3 in PC-3 human prostate cancer cells [34], although we did not study the Smad activation in these cells.

It has also been reported that IGFBP-3 down-regulates Akt activity in human epidermal growth factor receptor-2 (HER-2) over expressed MCF-7 breast cancer cells [38], in this regard recent studies have shown that IGFBP-3 induces apoptosis in MCF-7 breast cancer cells by inhibiting IGF-I/Akt survival pathway [39]. Previous studies also indicated that IGFBP-3 can bind to a new cell death receptor (IGFBP-3R), a single-span membrane protein which specifically binds to IGFBP-3 but doesn't bind to other IGFBP's. *In vivo* studies using prostate and breast cancer xenografts in athymic nude mice, showed anti-tumor effects of IGFBP-3R [40]. It was shown from the *in vitro* studies that IGFBP-3R triggers IGFBP-3 induced apoptosis in various cancer cells via a caspase-8 dependent pathway. IGFBP-3R directly interacts and activates caspase-8 in inducing apoptosis and knockdown of caspase-8 expression or activity can lead to inhibition of IGFBP-3/IGFBP-3R induced apoptosis. All these studies clearly indicate that IGFBP-3 induces its anti-proliferative and pro-apoptotic effects either by sequestering IGF-1 to prevent it to bind to Insulin-like growth factor-I receptor (IGF-IR) in an IGF dependent manner or by activating several candidate molecules via signal transduction pathway independent of IGF binding. Although insulin-like growth factor binding protein-3 receptor (IGFBP-3R) has been known to be directly involved in IGFBP-3 action and may prove to be an important target molecule for the treatment of cancer, further studies are still needed to clarify the role of this receptor or other IGFBP-3 cell surface binding protein(s) in IGFBP-3 mediated cell signaling events.

4. Conclusion

IGFBP-3 is known to activate various downstream signaling molecules. Some of these signaling events could be initiated after binding to its receptor independent of IGF-I binding, to induce anti-proliferative and pro-apoptotic effects in a certain type of cancer cells. On the other hand, IGFBP-3 can also bind to IGF-1 to modulate its ability to bind to IGF-1 receptor to inhibit cell proliferation and cell survival, resulting in cell death.

Funding

This work was supported by an intramural grant from NIDDK, NIH.

References

- [1] Wood, W.I., Cachianes, G., Henzel, W.J., Winslow, G.A., Spencer, S.A., Hellmiss, R., Martin, J.L. and Baxter, R.C. (1988) Cloning and Expression of the Growth Hormone-Dependent Insulin-Like Growth Factor-Binding Protein. *Molecular Endocrinology*, **2**, 1176-1185. <http://dx.doi.org/10.1210/mend-2-12-1176>
- [2] Mehta, H.H., Gao, Q., Galet, C., Paharkova, V., Wan, J., Said, J., Sohn, J.J., Lawson, G., Cohen, P., Cobb, L.J. and Lee, K.W. (2011) IGFBP-3 Is a Metastasis Suppression Gene in Prostate Cancer. *Cancer Research*, **71**, 5154-5163. <http://dx.doi.org/10.1158/0008-5472.CAN-10-4513>
- [3] Chan, J.M., Stampfer, M.J., Giovannucci, E., Gann, P.H., Ma, J., Wilkinson, P., Hennekens, C.H. and Pollak, M. (1998) Plasma Insulin-Like Growth Factor-I and Prostate Cancer Risk: A Prospective Study. *Science*, **279**, 563-566. <http://dx.doi.org/10.1126/science.279.5350.563>
- [4] Chan, J.M., Stampfer, M.J., Ma, J., Gann, P., Gaziano, J.M., Pollak, M. and Giovannucci, E. (2002) Insulin-Like Growth Factor-I (IGF-I) and IGF Binding Protein-3 as Predictors of Advanced-Stage Prostate Cancer. *Journal of the National Cancer Institute*, **94**, 1099-1106. <http://dx.doi.org/10.1093/jnci/94.14.1099>

- [5] Ma, J., Pollak, M.N., Giovannucci, E., Chan, J.M., Tao, Y., Hennekens, C.H. and Stampfer, M.J. (1999) Prospective Study of Colorectal Cancer Risk in Men and Plasma Levels of Insulin-Like Growth Factor (IGF)-I and IGF-Binding Protein-3. *Journal of the National Cancer Institute*, **91**, 620-625. <http://dx.doi.org/10.1093/jnci/91.7.620>
- [6] Giovannucci, E., Pollak, M.N., Platz, E.A., Willett, W.C., Stampfer, M.J., Majeed, N., Colditz, G.A., Speizer, F.E. and Hankinson, S.E. (2000) A Prospective Study of Plasma Insulin-Like Growth Factor-1 and Binding Protein-3 and Risk of Colorectal Neoplasia in Women. *Cancer Epidemiological Biomarkers Preview*, **9**, 345-349.
- [7] Hochscheid, R., Jaques, G. and Wegmann, B. (2000) Transfection of Human Insulin-Like Growth Factor-Binding Protein 3 Gene Inhibits Cell Growth and Tumorigenicity: A Cell Culture Model for Lung Cancer. *Journal of Endocrinology*, **166**, 553-563. <http://dx.doi.org/10.1677/joe.0.1660553>
- [8] Devi, G.R., Sprenger, C.C., Plymate, S.R. and Rosenfeld, R.G. (2002) Insulin-Like Growth Factor Binding Protein-3 Induces Early Apoptosis in Malignant Prostate Cancer Cells and Inhibits Tumor Formation *in Vivo*. *Prostate*, **51**, 141-152. <http://dx.doi.org/10.1002/pros.10068>
- [9] Hanafusa, T., Yumoto, Y., Nouse, K., Nakatsukasa, H., Onishi, T., Fujikawa, T., Taniyama, M., Nakamura, S., Uemura, M., Takuma, Y., Yumoto, E., Higashi, T. and Tsuji, T. (2002) Reduced Expression of Insulin-Like Growth Factor Binding Protein-3 and Its Promoter Hypermethylation in Human Hepatocellular Carcinoma. *Cancer Letters*, **176**, 149-158. [http://dx.doi.org/10.1016/S0304-3835\(01\)00736-4](http://dx.doi.org/10.1016/S0304-3835(01)00736-4)
- [10] Chang, Y.S., Kong, G., Sun, S., Liu, D., El-Naggar, A.K., Khuri, F.R., Hong, W.K. and Lee, H.Y. (2002) Clinical Significance of Insulin-Like Growth Factor-Binding Protein-3 Expression in Stage I Non-Small Cell Lung Cancer. *Clinical Cancer Research*, **8**, 3796-3802.
- [11] Lu, X.F., Jiang, X.G., Lu, Y.B., Bai, J.H. and Mao, Z.B. (2005) Characterization of a Novel Positive Transcription Regulatory Element That Differentially Regulates the Insulin-Like Growth Factor Binding Protein-3 (IGFBP-3) Gene in Senescent Cells. *Journal of Biological Chemistry*, **280**, 22606-22615. <http://dx.doi.org/10.1074/jbc.M412073200>
- [12] Mannhardt, B., Weinzimer, S.A., Wagner, M., Fiedler, M., Cohen, P., Jansen-Durr, P. and Zwerschke, W. (2000) Human Papillomavirus Type 16 E7 Oncoprotein Binds and Inactivates Growth-Inhibitory Insulin-Like Growth Factor Binding Protein 3. *Molecular and Cellular Biology*, **20**, 6483-6495. <http://dx.doi.org/10.1128/MCB.20.17.6483-6495.2000>
- [13] Bhattacharyya, N., Pechhold, K., Shahjee, H., Zappala, G., Elbi, C., Raaka, B., Wiench, M., Hong, J. and Rechler, M.M. (2006) Nonsecreted Insulin-Like Growth Factor Binding Protein-3 (IGFBP-3) Can Induce Apoptosis in Human Prostate Cancer Cells by IGF-Independent Mechanisms without Being Concentrated in the Nucleus. *Journal of Biological Chemistry*, **281**, 24588-24601. <http://dx.doi.org/10.1074/jbc.M509463200>
- [14] Lalou, C., Lassarre, C. and Binoux, M. (1996) A Proteolytic Fragment of Insulin-Like Growth Factor (IGF) Binding Protein-3 That Fails to Bind IGFs Inhibits the Mitogenic Effects of IGF-I and Insulin. *Endocrinology*, **137**, 3206-3212.
- [15] Shahjee, H., Bhattacharyya, N., Zappala, G., Wiench, M., Prakash, S. and Rechler, M.M. (2008) An N-Terminal Fragment of Insulin-Like Growth Factor Binding Protein-3 (IGFBP-3) Induces Apoptosis in Human Prostate Cancer Cells in an IGF-Independent Manner. *Growth Hormone & IGF Research*, **18**, 188-197. <http://dx.doi.org/10.1016/j.ghir.2007.08.006>
- [16] Rajah, R., Valentinis, B. and Cohen, P. (1997) Insulin-Like Growth Factor IGF Binding Protein-3 Induces Apoptosis and Mediates the Effects of Transforming Growth Factor- β 1 on Programmed Cell Death through a p53- and IGF-Independent Mechanism. *Journal of Biological Chemistry*, **272**, 12181-12188. <http://dx.doi.org/10.1074/jbc.272.18.12181>
- [17] Butt, A.J., Firth, S.M., King, M.A. and Baxter, R.C. (2000) Insulin-Like Growth Factor-Binding Protein-3 Modulates Expression of Bax and Bcl-2 and Potentiates p53-Independent Radiation-Induced Apoptosis in Human Breast Cancer Cells. *Journal of Biological Chemistry*, **275**, 39174-39181. <http://dx.doi.org/10.1074/jbc.M908888199>
- [18] Hong, J., Zhang, G., Dong, F. and Rechler, M.M. (2002) Insulin-Like Growth Factor (IGF)-Binding Protein-3 Mutants That Do Not Bind IGF-I or IGF-II Stimulate Apoptosis in Human Prostate Cancer Cells. *Journal of Biological Chemistry*, **277**, 10489-10497. <http://dx.doi.org/10.1074/jbc.M109604200>
- [19] Kim, H.S., Lee, W.J., Lee, S.W., Chae, H.W., Kim, D.H. and Oh, Y. (2010) Insulin-Like Growth Factor Binding Protein-3 Induces G1 Cell Cycle Arrest with Inhibition of Cyclin-Dependent Kinase 2 and 4 in MCF-7 Human Breast Cancer Cells. *Hormone and Metabolic Research*, **42**, 165-172.
- [20] Wu, C., Liu, X.B., Wang, Y.Y., Tian, H.N., Xie, Y.Y., Li, Q.J., Zhang, X.K. and Liu, F.M. (2013) Insulin-Like Factor Binding Protein-3 Promotes the G1 Cell Cycle Arrest in Several Cancer Cell Lines. *Gene*, **512**, 127-133. <http://dx.doi.org/10.1016/j.gene.2012.09.080>
- [21] Gucev, Z.S., Oh, Y., Kelley, K.M. and Rosenfeld, R.G. (1996) Insulin-Like Growth Factor Binding Protein-3 Mediates Retinoic Acid- and Transforming Growth Factor Beta2-Induced Growth Inhibition in Human Breast Cancer Cells. *Cancer Research*, **56**, 1545-1550.
- [22] Rajah, R., Lee, K.W. and Cohen, P. (2002) Insulin-Like Growth Factor Binding Protein-3 Mediates Tumor Necrosis

- Factor-Alpha-Induced Apoptosis: Role of Bcl-2 Phosphorylation. *Cell Growth & Differentiation*, **13**, 163-171.
- [23] Vasylyeva, T.L., Chen, X. and Ferry Jr., R.J. (2005) Insulin-Like Growth Factor Binding Protein-3 Mediates Cytokine-Induced Mesangial Cell Apoptosis. *Growth Hormone & IGF Research*, **15**, 207-214. <http://dx.doi.org/10.1016/j.ghir.2005.02.008>
- [24] Ryan, K.M. and Vousden, K.H. (1998) Characterization of Structural p53 Mutants Which Show Selective Defects in Apoptosis but Not Cell Cycle Arrest. *Molecular and Cellular Biology*, **18**, 3692-3698.
- [25] Grimberg, A., Liu, B., Bannerman, P., El-Deiry, W.S. and Cohen, P. (2002) IGFBP-3 Mediates p53-Induced Apoptosis during Serum Starvation. *International Journal of Oncology*, **21**, 327-335.
- [26] Harms, K.L. and Chen, X. (2005) The C Terminus of p53 Family Proteins Is a Cell Fate Determinant. *Molecular and Cellular Biology*, **25**, 2014-2030. <http://dx.doi.org/10.1128/MCB.25.5.2014-2030.2005>
- [27] Rechler, M.M. and Clemmons, D.R. (1998) Regulatory Actions of Insulin-Like Growth Factor-Binding Proteins. *Trends in Endocrinology and Metabolism*, **9**, 176-183. [http://dx.doi.org/10.1016/S1043-2760\(98\)00047-2](http://dx.doi.org/10.1016/S1043-2760(98)00047-2)
- [28] Firth, S.M. and Baxter, R.C. (2002) Cellular Actions of the Insulin-Like Growth Factor Binding Proteins. *Endocrine Reviews*, **23**, 824-854. <http://dx.doi.org/10.1210/er.2001-0033>
- [29] Han, J., Jogie-Brahmin, S., Harada, A. and Oh, Y. (2011) Insulin-Like Growth Factor-Binding Protein-3 Suppresses tumor Growth via Activation of Caspase-Dependent Apoptosis and Cross-Talk with NF- κ B Signaling. *Cancer Letters*, **307**, 200-210. <http://dx.doi.org/10.1016/j.canlet.2011.04.004>
- [30] Zappala, G., Elbi, C., Edwards, J., Gorenstein, J., Rechler, M.M. and Bhattacharyya, N. (2008) Induction of Apoptosis in Human Prostate Cancer Cells by Insulin-Like Growth Factor Binding Protein-3 Does Not Require Binding to Retinoid X Receptor- α . *Endocrinology*, **149**, 1802-1812. <http://dx.doi.org/10.1210/en.2007-1315>
- [31] Leibowitz, B.J., Agostini-Dreyer, A., Jetzt, A.E., Krumm, C.S. and Cohick, W.S. (2013) IGF Binding Protein-3 Mediates Stress-Induced Apoptosis in Non-Transformed Mammary Epithelial Cells. *Journal of Cellular Physiology*, **228**, 734-742. <http://dx.doi.org/10.1002/jcp.24220>
- [32] Ricort, J.M. (2004) Insulin-Like Growth Factor Binding Protein (IGFBP) Signalling. *Growth Hormone & IGF Research*, **14**, 277-286. <http://dx.doi.org/10.1016/j.ghir.2004.02.002>
- [33] Spagnoli, A., Torello, M., Nagalla, S.R., Horton, W.A., Pattee, P., Chiarelli, F., Roberts Jr., C.T. and Rosenfeld, R.G. (2002) Identification of STAT-1 as a Molecular Target of IGFBP-3 in the Process of Chondrogenesis. *Journal of Biological Chemistry*, **277**, 18860-18867. <http://dx.doi.org/10.1074/jbc.M200218200>
- [34] Shahjee, H.M., Kefas, B., Bhattacharyya, N. and Radwan, M.K. (2013) Signal Transduction Pathways Mediated by Secreted and Non-Secreted Forms of Intact Insulin-Like Growth Factor Binding Protein-3 and Its 1-97 N-Terminal Fragment in PC-3 Human Prostate Cancer Cells. *Journal of Cancer Therapy*, **4**, 1290-1297. <http://dx.doi.org/10.4236/jct.2013.48152>
- [35] Fanayan, S., Firth, S.M., Butt, A.J. and Baxter, R.C. (2000) Growth Inhibition by Insulin-Like Growth Factor-Binding Protein-3 in T47D Breast Cancer Cells Requires Transforming Growth Factor- β (TGF- β) and the Type II TGF- β Receptor. *Journal of Biological Chemistry*, **275**, 39146-39151. <http://dx.doi.org/10.1074/jbc.M006964200>
- [36] Fanayan, S., Firth, S.M. and Baxter, R.C. (2002) Signalling through the Smad Pathway by Insulin-Like Growth Factor-Binding Protein-3 in Breast Cancer Cells. Relationship to Transforming Growth Factor- β 1 Signaling. *Journal of Biological Chemistry*, **277**, 7255-7261. <http://dx.doi.org/10.1074/jbc.M108038200>
- [37] Leal, S.M., Huang, S.S. and Huang, J.S. (1999) Interactions of High Affinity Insulin-Like Growth Factor-Binding Proteins with the Type V Transforming Growth Factor- β Receptor in Mink Lung Epithelial Cells. *Journal of Biological Chemistry*, **274**, 6711-6717. <http://dx.doi.org/10.1074/jbc.274.10.6711>
- [38] Jerome, L., Alami, N., Belanger, S., Page, V., Yu, Q., Paterson, J., Shiry, L., Pegram, M. and Leyland-Jones, B. (2006) Recombinant Human Insulin-Like Growth Factor Binding Protein-3 Inhibits Growth of Human Epidermal Growth Factor Receptor-2 Overexpressing Breast Tumors and Potentiates Herceptin Activity *in Vivo*. *Cancer Research*, **66**, 7245-7252. <http://dx.doi.org/10.1158/0008-5472.CAN-05-3555>
- [39] Brosseau, C., Pirianov, G. and Colston, K.W. (2013) Role of Insulin-Like Growth Factor Binding Protein-3 in 1,25-Dihydroxyvitamin-D₃-Induced Breast Cancer Cell Apoptosis. *International Journal of Cell Biology*, **2013**, Article ID: 960378. <http://dx.doi.org/10.1155/2013/960378>
- [40] Ingermann, A.R., Yang, Y.F., Han, J., Mikami, A., Garza, A.E., Mohanraj, L., Fan, L., Idowu, M., Ware, J.L., Kim, H.S., Lee, D.Y. and Oh, Y. (2010) Identification of a Novel Cell Death Receptor Mediating IGFBP-3-Induced Anti-Tumor Effects in Breast and Prostate Cancer. *Journal of Biological Chemistry*, **285**, 30233-30246. <http://dx.doi.org/10.1074/jbc.M110.122226>

Scientific Research Publishing (SCIRP) is one of the largest Open Access journal publishers. It is currently publishing more than 200 open access, online, peer-reviewed journals covering a wide range of academic disciplines. SCIRP serves the worldwide academic communities and contributes to the progress and application of science with its publication.

Other selected journals from SCIRP are listed as below. Submit your manuscript to us via either submit@scirp.org or [Online Submission Portal](#).

