

# Interacting generalized chaplygin gas model in bianchi type-I universe

Raghavendra Chaubey

Applied Mathematics, DST-Centre for Interdisciplinary Mathematical Sciences, Faculty of Science, Banaras Hindu University, Varanasi, India; \*Corresponding Author: [rchaubey@bhu.ac.in](mailto:rchaubey@bhu.ac.in)

Received 15 February 2011; revised 3 April 2011; accepted 18 April 2011.

## ABSTRACT

In this paper, we have studied the generalized chaplygin gas of interacting dark energy to obtain the equation of state for the generalized chaplygin gas energy density in anisotropic Bianchi type-I cosmological model. For negative value of  $B$  in equation of state of generalized chaplygin gas, we see that  $\gamma_{\Lambda}^{eff} < -1$ , that corresponds to a universe dominated by phantom dark energy.

**Keywords:** Cosmological Models; Chaplygin Gas; Cosmological Parameters

## 1. INTRODUCTION

One of the most important problems of cosmology, is the problem of so-called dark energy (DE). The type Ia supernova observations suggests that the universe is dominated by dark energy with negative pressure which provides the dynamical mechanism of the accelerating expansion of the universe [1-3]. The strength of this acceleration is presently matter of debate, mainly because it depends on the theoretical model implied when interpreting the data. Most of these models are based on dynamics of a scalar or multi-scalar fields. Primary scalar field candidate for dark energy was quintessence scenario [4,5], a fluid with the parameter of the equation of state lying in the range,  $-1 < \gamma < -1/3$ .

In a very interesting paper Kamenshchik, Moschella and Pasquier [6] have studied a homogeneous model based on a single fluid obeying the Chaplygin gas equation of state

$$p = -\frac{A}{\rho} \quad (1.1)$$

where  $p$  and  $\rho$  are respectively pressure and energy density in comoving reference frame, with  $\rho > 0$ ;  $A$  is a positive constant. This equation of state has raised a certain interest [7] because of its many interesting and, in some sense, intriguingly unique features. Some possi-

ble motivations for this model from the field theory points of view are investigated in [8]. The Chaplygin gas emerges as an effective fluid associated with d-branes [9] and can also be obtained from the Born-infield action [10].

Inserting the equation of state (1.1) into the relativistic energy conservation equation, leads to a density evolving as

$$\rho_{\Lambda} = \sqrt{A + B/V^2} \quad (1.2)$$

where  $B$  is an integration constant.

There exist a wide class of anisotropic cosmological models, which also often studying in cosmology [11]. There are theoretical arguments that sustain the existence of an anisotropic phase that approaches an isotropic case [12]. Also, anisotropic cosmological models are found a suitable candidate to avoid the assumption of specific initial conditions in FRW models. The early universe could also characterized by irregular expansion mechanism. Therefore, it would be useful to explore cosmological models in which anisotropic, existing at early stage of expansion, are damped out in the course of evolution. Interest in such models have been received much attention since 1978 [13].

Setare [14] has obtained the equation of state for the generalized Chaplygin gas energy density in non-flat universe. Chaubey [15] has obtained the role of modified chaplygin gas in Bianchi type - I universe. In the present paper, using the generalized Chaplygin gas model of dark energy, we obtain equation of state for interacting Chaplygin gas energy density in anisotropic Bianchi type-I cosmological model. For negative value of  $B$  in equation of state of generalized chaplygin gas, we see that  $\gamma_{\Lambda}^{eff} < -1$ , that corresponds to a universe dominated by phantom dark energy.

## 2. INTERACTING GENERALIZED CHAPLYGIN GAS

In this section we obtain the equation of state for the generalized Chaplygin gas when there is an interaction

between generalized Chaplygin gas energy density  $\rho_\Lambda$  and a Cold Dark Matter (CDM) with  $\gamma_m = 0$ .

The continuity equations for dark energy and CDM are

$$\dot{\rho}_\Lambda + 3H(1 + \gamma_\Lambda)\rho_\Lambda = -Q \tag{2.1}$$

$$\dot{\rho}_m + 3H\rho_m = Q. \tag{2.2}$$

The interaction is given by the quality  $Q = \Gamma\rho_\Lambda$ . This is a decaying of the generalized Chaplygin gas component into CDM with the decay rate  $\Gamma$ . Taking a ratio of two energy densities as  $r = \rho_m/\rho_\Lambda$ , the above equations lead to

$$\dot{r} = 3Hr \left[ \gamma_\Lambda + \frac{1+r}{r} \frac{\Gamma}{3H} \right] \tag{2.3}$$

Following [3], if we define

$$\gamma_\Lambda^{eff} = \gamma_\Lambda + \frac{\Gamma}{3H}, \quad \gamma_m^{eff} = -\frac{1}{r} \frac{\Gamma}{3H} \tag{2.4}$$

Then, the continuity equations can be written in their standard form

$$\dot{\rho}_\Lambda + 3H(1 + \gamma_\Lambda^{eff})\rho_\Lambda = 0 \tag{2.5}$$

$$\dot{\rho}_m + 3H(1 + \gamma_m^{eff})\rho_m = 0 \tag{2.6}$$

We consider the homogeneous anisotropic Bianchi type-I cosmological model with line element

$$ds^2 = dt^2 - a_1^2 dx^2 - a_2^2 dy^2 - a_3^2 dz^2 \tag{2.7}$$

where  $a_1, a_2$  and  $a_3$  are function of  $t$  only.

The Einstein field equations for the metric (2.1) are written in the form

$$\frac{\ddot{a}_2}{a_2} + \frac{\ddot{a}_3}{a_3} + \frac{\dot{a}_2 \dot{a}_3}{a_2 a_3} = \kappa[\rho_\Lambda + \rho_m]. \tag{2.8}$$

$$\frac{\ddot{a}_1}{a_1} + \frac{\ddot{a}_3}{a_3} + \frac{\dot{a}_1 \dot{a}_3}{a_1 a_3} = -\kappa[\rho_\Lambda]. \tag{2.9}$$

$$\frac{\ddot{a}_1}{a_1} + \frac{\ddot{a}_2}{a_2} + \frac{\dot{a}_1 \dot{a}_2}{a_1 a_2} = -\kappa[\rho_\Lambda]. \tag{2.10}$$

$$\frac{\dot{a}_1 \dot{a}_2}{a_1 a_2} + \frac{\dot{a}_2 \dot{a}_3}{a_2 a_3} + \frac{\dot{a}_3 \dot{a}_1}{a_3 a_1} = -\kappa[\rho_\Lambda] \tag{2.11}$$

where  $\kappa \equiv 8\pi G/c^4$  is constant.

We define

$$V = a_1 a_2 a_3 \tag{2.12}$$

By using the method of Singh *et al.* [16-19], we obtain

$$a_1(t) = D_1 V^{1/3} \exp\left(X_1 \int \frac{dt}{V(t)}\right) \tag{2.13}$$

$$a_2(t) = D_2 V^{1/3} \exp\left(X_2 \int \frac{dt}{V(t)}\right) \tag{2.14}$$

$$a_3(t) = D_3 V^{1/3} \exp\left(X_3 \int \frac{dt}{V(t)}\right) \tag{2.15}$$

where  $D_i$  ( $i = 1, 2, 3$ ) and  $X_i$  ( $i = 1, 2, 3$ ) satisfy the relation  $D_1 D_2 D_3 = 1$  and  $X_1 + X_2 + X_3 = 0$ .

Now, adding **Eqs.2.9, 2.10** and **2.11** and three times **Eq.2.8**, we get

$$\begin{aligned} & \left( \frac{\ddot{a}_1}{a_1} + \frac{\ddot{a}_2}{a_2} + \frac{\ddot{a}_3}{a_3} \right) + 2 \left( \frac{\dot{a}_1 \dot{a}_2}{a_1 a_2} + \frac{\dot{a}_2 \dot{a}_3}{a_2 a_3} + \frac{\dot{a}_3 \dot{a}_1}{a_3 a_1} \right) \\ & = \frac{3}{2} \kappa [(\rho_\Lambda + \rho_m) - p_\Lambda]. \end{aligned} \tag{2.16}$$

From **Eqs.2.12** and **2.16**, we have

$$\frac{\ddot{V}}{V} = \frac{3}{2} \kappa [(\rho_\Lambda + \rho_m) - p_\Lambda]. \tag{2.17}$$

Define as usual

$$\Omega_m = \frac{\rho_m}{\rho_{cr}} = \frac{3\kappa\rho_m V^2}{\dot{V}^2}; \quad \Omega_\Lambda = \frac{\rho_\Lambda}{\rho_{cr}} = \frac{3\kappa\rho_\Lambda V^2}{\dot{V}^2} \tag{2.18}$$

From above, we obtain following relation for ratio of energy densities  $r$  as

$$r = \frac{\Omega_m}{\Omega_\Lambda} \tag{2.19}$$

In the generalized Chaplygin gas approach [10], the equation of state to (1.1) is generalized to

$$p_\Lambda = -\frac{A}{\rho_\Lambda^\alpha} \tag{2.20}$$

The above equation of state leads to a density evolution as

$$\rho_\Lambda = \left[ A + \frac{B}{V^{(1+\alpha)}} \right]^{\frac{1}{1+\alpha}} \tag{2.21}$$

Taking derivatives in both sides of above equation with respect to cosmic time, we obtain

$$\dot{\rho}_\Lambda = -B \left( \frac{\dot{V}}{V} \right) V^{-(1+\alpha)} \left[ A + B V^{-(1+\alpha)} \right]^{-\frac{\alpha}{1+\alpha}} \tag{2.22}$$

Substituting this relation into **Eq.2.1** and using definition  $Q = \Gamma\rho_\Lambda$ , we obtain

$$\gamma_\Lambda = \frac{B}{V^{(1+\alpha)} [A + B V^{-(1+\alpha)}]} - \frac{\Gamma}{(\dot{V}/V)} - 1 \tag{2.23}$$

Here as in Ref. [20], we choose the following relation for decay rate

$$\Gamma = b^2 (1+r) \frac{\dot{V}}{V} \tag{2.24}$$

with the coupling constant  $b^2$ . Using **Eq.2.14**, the

above decay rate take following form

$$\Gamma = b^2 \left( \frac{\dot{V}}{V} \right) \left( \frac{\Omega_\Lambda + \Omega_m}{\Omega_\Lambda} \right) \quad (2.25)$$

Substituting this relation into **Eq.2.23**, one finds the generalized Chaplygin gas energy equation of state

$$\gamma_\Lambda = \frac{B}{V^{1+\alpha} [A + BV^{-(1+\alpha)}]} - b^2 \left( \frac{\Omega_\Lambda + \Omega_m}{\Omega_\Lambda} \right) - 1. \quad (2.26)$$

Now using the definition generalized Chaplygin gas energy  $\rho_\Lambda$ , and using  $\Omega_\Lambda$ , we can rewrite the above equation as

$$\gamma_\Lambda = \frac{3B}{\left\{ \frac{1}{\kappa} \left( \frac{\dot{V}^2}{V} \right) \Omega_\Lambda \right\}^{(1+\alpha)}} - b^2 \left( \frac{\Omega_\Lambda + \Omega_m}{\Omega_\Lambda} \right) - 1 \quad (2.27)$$

From **Eqs.2.4, 2.25** and **2.27**, we have the effective equation of state as

$$\gamma_\Lambda^{eff} = \frac{3B}{\left[ \frac{1}{\kappa} \left( \frac{\dot{V}^2}{V} \right) \Omega_\Lambda \right]^{(1+\alpha)}} - 1 \quad (2.28)$$

By choosing a negative value for  $B$  we see that  $\gamma_\Lambda^{eff} < -1$ , that corresponds to a universe dominated by phantom dark energy, **Eq.2.28**, for  $\alpha = 1$ , is the effective parameter of state for Chaplygin gas. In this case, in the expression for energy density (1.2), term under square root should be positive, *i.e.*  $V^2 > -B/A$ , then the minimal value of the volume factor is given by

$$V_{min} = \left( -\frac{B}{A} \right)^{\frac{1}{2}} \quad (2.29)$$

Now, from **Eqs.2.13-2.15** and **2.29**, we have find the minimal value of the scale factors are given by

$$a_{1min} = D_1 \left( -\frac{B}{A} \right)^{\frac{1}{6}} \exp \left[ X_1 \left( -\frac{B}{A} \right)^{\frac{1}{2}} t \right] \quad (2.30)$$

$$a_{2min} = D_2 \left( -\frac{B}{A} \right)^{\frac{1}{6}} \exp \left[ X_2 \left( -\frac{B}{A} \right)^{\frac{1}{2}} t \right] \quad (2.31)$$

$$a_{3min} = D_3 \left( -\frac{B}{A} \right)^{\frac{1}{6}} \exp \left[ X_3 \left( -\frac{B}{A} \right)^{\frac{1}{2}} t \right] \quad (2.32)$$

According to this model we have a bouncing universe. Generally for this model  $A > 0, B < 0$  and  $1 + \alpha > 0$ . From **Eq.2.21**, we can realize that the cosmic scalar fac-

tors take values in the interval  $a_{i min} < a_i < \infty$  (for  $i = 1, 2, 3$ ) which corresponds to  $0 < \rho < (2A)^{\frac{1}{1+\alpha}}$ , where

$$V_{min} = \left( -\frac{B}{A} \right)^{\frac{1}{1+\alpha}} \quad (2.33)$$

and

$$a_{i min} = D_i \left( -\frac{B}{A} \right)^{\frac{1}{3(1+\alpha)}} \exp \left[ X_i \left( -\frac{B}{A} \right)^{\frac{1}{(1+\alpha)}} t \right], i = 1, 2, 3. \quad (2.34)$$

Using **Eq.1.2**, one can see that the Chaplygin gas interpolates between dust at small  $a_i$  and a cosmological constant at large  $a_i$ , but choosing a negative value of  $B$ , this quartessence idea lose. Following [6] if we consider a homogeneous scalar field  $\phi(t)$  and a potential  $V(\phi)$  to describe the Chaplygin cosmology, we find

$$\dot{\phi}^2 = \frac{B}{V^2 \sqrt{A + B/V^2}} \quad (2.35)$$

Now, by choosing a negative value for  $B$  we see that  $\dot{\phi}^2 < 0$ , then we can write

$$\phi = i\psi \quad (2.36)$$

In this case the lagrangian of scalar field  $\phi(t)$  can rewritten as

$$L = -\frac{1}{2} \dot{\phi}^2 - V(\phi) = -\frac{1}{2} \dot{\psi}^2 - V(i\psi) \quad (2.37)$$

The energy density and the pressure corresponding to the scalar field  $\psi$  are as respectively

$$\rho_\psi = -\frac{1}{2} \dot{\psi}^2 + V(i\psi) \quad (2.38)$$

$$p_\psi = -\frac{1}{2} \dot{\psi}^2 - V(i\psi) \quad (2.38)$$

Therefore, the scalar field  $\psi$  is a phantom field. This implies that one can generate phantom-like equation of state from an interacting generalized Chaplygin gas dark energy model in anisotropic universe.

### 3. CONCLUSIONS

We have studied the generalized chaplygin gas of interacting dark energy to obtain the equation of state for the generalized chaplygin gas energy density in anisotropic Bianchi type-I cosmological model. By choosing a negative value for  $B$  we see that  $\gamma_\Lambda^{eff} < -1$ , that cor-

responds to a universe dominated by phantom dark energy.

## REFERENCES

- [1] Perlmutter, S., *et al.* (1999) Measurements of  $\Omega$  and  $\Lambda$  from 42 high-redshift supernovae. *The Astrophysical Journal*, **517**, 565-586. [doi:10.1086/307221](https://doi.org/10.1086/307221)
- [2] Garnavich, P.M., *et al.* (1998) Constraints on cosmological models from *Hubble Space Telescope* observations of high- $z$  supernovae. *The Astrophysical Journal*, **493**, L53. [doi:10.1086/311140](https://doi.org/10.1086/311140)
- [3] Riess, A.G., *et al.* (1998) Observational evidence from supernovae for an accelerating universe and cosmological constant. *The Astrophysical Journal*, **116**, 1009-1038. [doi:10.1086/300499](https://doi.org/10.1086/300499)
- [4] Ratra, B. and Peebles, P.J.E. (1988) Cosmological consequences of a rolling homogeneous scalar field. *Physical Review D*, **37**, 3406-3427.
- [5] Zlatev, I., Wang, L. and Steinhardt, P.J. (1999) Quintessence, cosmic coincidence, and the cosmological constant. *Physics Review Letters*, **82**, 896-899. [doi:10.1103/PhysRevD.59.123504](https://doi.org/10.1103/PhysRevD.59.123504)
- [6] Kamenshchik, A.Y., Moschella, U. and Pasquier, V. (2001) An alternative to quintessence. *Physics Letters B*, **511**, 265-268. [doi:10.1016/S0370-2693\(01\)00571-8](https://doi.org/10.1016/S0370-2693(01)00571-8)
- [7] Bazeia, D. and Jackiw, R. (1998) Nonlinear realization of a dynamical Poincaré symmetry by a field-dependent diffeomorphism. *Annals of Physics*, **270**, 246-259. [doi:10.1103/PhysRevD.69.023506](https://doi.org/10.1103/PhysRevD.69.023506)
- [8] Bilic, N., Tupper, G.B. and Viollier, R.D. (2002) Unification of dark matter and dark energy: the inhomogeneous Chaplygin gas. *Physics Letters B*, **535**, 17-21.
- [9] Bordemann, M. and Hoppe, J. (1993) The dynamics of relativistic membranes. Reduction to 2-dimensional fluid dynamics. *Physics Letters B*, **317**, 315-320. [doi:10.1023/A:1015266421750](https://doi.org/10.1023/A:1015266421750)
- [10] Bento, M.C., Bertolami, O. and Sen, A.A. (2003) WMAP constraints on the generalized Chaplygin gas model. *Physics Letters B*, **575**, 172-180. [doi:10.1103/PhysRevD.72.063511](https://doi.org/10.1103/PhysRevD.72.063511)
- [11] Misner, C.W., Thorne, K.S. and Wheeler, J.A. (1973) *Gravitation*. W.H. Freeman, New York.
- [12] Misner, C.W. (1968) The isotropy of the universe. *Astrophysics Journal*, **151**, 431-457.
- [13] Hu, B.L. and Parker, L. (1978) Anisotropy damping through quantum effects in the early universe. *Physics Review D*, **17**, 933-945. [doi:10.1103/PhysRevD.17.933](https://doi.org/10.1103/PhysRevD.17.933)
- [14] Setare, M.R. (2007) Interacting generalized Chaplygin gas model in non-flat universe. *European Physical Journal*, **C52**, 689-692. [doi:10.1140/epjc/s10052-007-0405-5](https://doi.org/10.1140/epjc/s10052-007-0405-5)
- [15] Chaubey, R. (2009) Role of modified Chaplygin gas in Bianchi type-I universe. *International Journal of Theoretical Physics*, **48**, 952-960.
- [16] Singh, T. and Chaubey, R. (2009) Bianchi type-I, III, V, VI<sub>0</sub> and Kantowski-Sachs models in scalar-tensor theories with dynamic cosmological constant. *Astrophysics and Space Science*, **318**, 231-236. [doi:10.1007/s10509-008-9917-1](https://doi.org/10.1007/s10509-008-9917-1)
- [17] Singh, T. and Chaubey, R. (2008) Bianchi type-I universe with wet dark fluid. *Pramana—Journal of Physics*, **71**, 447-458.
- [18] Singh, T. and Chaubey, R. (2009) Bianchi type-I, III, V, VI<sub>0</sub> and Kantowski-Sachs universes in creation-field cosmology. *Astrophysics and Space Science*, **321**, 5-18. [doi:10.1007/s10509-009-9989-6](https://doi.org/10.1007/s10509-009-9989-6)
- [19] Singh, T. and Chaubey, R. (2009) Bianchi type-III, V, VI<sub>0</sub> and Kantowski-Sachs universes with varying  $\Lambda$ ,  $G$  and  $\sigma^2$  simultaneously. *Proceedings of the National Academy of Sciences, India Section A*, **79**, 337-354.
- [20] Wang, B., Gong, Y. and Abdalla, E. (2005) Transition of the dark energy equation of state in an interacting holographic dark energy model. *Physics Letters*, **B624**, 141-146. [doi:10.1016/j.physletb.2005.08.008](https://doi.org/10.1016/j.physletb.2005.08.008)