

Periodate Oxidation of a Ternary Complex of Nitrilotriacetatochromium(III) Involving β -Alanine as Co-Ligand

Hassan A. Ewais^{1*}, Ahmed H. Abdel-Salam², Amal S. Basaleh¹, Mohamed A. Habib³

¹Chemistry Department, Faculty of Science, King Abdulaziz University, Jeddah, KSA

²Chemistry Department, Faculty of Science, University of Jeddah, Jeddah, KSA

³Chemistry Department Faculty of Science, Al-Tahadi University, Serit, Lybia

Email: *ahmedhassan179@yahoo.com

How to cite this paper: Ewais, H.A., Abdel-Salam, A.H., Basaleh, A.S. and Habib, M.A. (2018) Periodate Oxidation of a Ternary Complex of Nitrilotriacetatochromium(III) Involving β -Alanine as Co-Ligand. *Open Journal of Inorganic Chemistry*, 8, 91-104. <https://doi.org/10.4236/ojic.2018.84008>

Received: September 5, 2018

Accepted: October 12, 2018

Published: October 15, 2018

Copyright © 2018 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

The kinetics of the periodate oxidation of chromium(III)-complex, $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]$ - (NTA = Nitrilotriacetate and Ala = β -alanine) to Cr(VI) have been carried out for the temperature range 15°C - 35°C under pseudo-first order conditions, $[\text{IO}_4^-] \gg [\text{complex}]$. Reaction obeyed first order dependence with respect to $[\text{IO}_4^-]$ and $[\text{Cr}(\text{III})]$, and the rate of reaction increases with increasing of pH for the range 3.40 - 4.45. Experimentally, the mechanism of this reaction is found to be consistent with the rate law in which the hydroxo species, $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{OH})]^{2-}$ is considerably much more reactive than their conjugate acid. ΔH^\ddagger and ΔS^\ddagger have been calculated. It is proposed that electron transfer occurs through an inner-sphere mechanism *via* coordination of IO_4^- to chromium(III).

Keywords

Nitrilotriacetatochromium(III), Ternary Complex, Periodate Oxidation, Inner-Sphere Mechanism, Thermodynamic Activation Parameters

1. Introduction

The Ternary complexes of oxygen-donor ligands and heteroaromatic N-bases such as nitrilotriacetic acid (NTA) and iminodiacetic acid (IDA) with transition metals have attracted much interest, as they can display exceptionally high stability and may be biologically relevant [1] [2]. The use of transition metal complexes of NTA is gaining increasing use in biotechnology, particularly in the

protein purification technique known as immobilized metal-ion chromatography [3]. The chromium(III) complexes of α amino acids are biologically available, depending on the complexing ability of the ligands for chromium against OH^- . Chromium can also aid in the transportation of amino acids through the cell membrane [4]. The biological oxidation of chromium from the trivalent to the hexavalent state is an important environmental process because of the high mobility and toxicity of chromium(VI) [5]. Recently, Cr(III) oxidation to Cr(V) and/or Cr(VI) in biological systems came into consideration as a possible reason for the anti-diabetic activities of some Cr(III) complexes, as well as the long-term toxicities of such complexes [6]. The specific interactions of chromium ions with cellular insulin receptors [7] are a consequence of intra- or extracellular oxidations of Cr(III) to Cr(V) and/or Cr(VI) compounds, which act as protein tyrosine phosphatase (PTP) inhibitors. Periodate oxidations have been reported to play an important role in biological processes [8] [9] [10].

Studies of the kinetics of periodate oxidations on a series of dextran oligomers, polymers and some dimeric carbohydrates [8] revealed a dependence of the kinetic rates on the molecular weight. The oxidation of caffeic acid (3,4-dihydroxy cinnamic acid) by sodium periodate was found to mimic the mechanism of polyphenol oxidase. The antioxidant product 2-s-cysteinyl caffeic acid exhibits slightly improved antiradical activity compared to the parent molecule (caffeic acid) [9]. The imidazol-modified M-salophen/ NaIO_4 system can be applied to oxidize a large number of primary aromatic amines in good yield at short times and room temperature [10].

An inner-sphere mechanism for oxidation of chromium(III) complexes of some amino acids [11] [12] [13] and nucleosides [14] [15] [16] by periodate has been proposed with the hydroxo group acting as bridging ligand, or through the substitution of coordinated H_2O by $[\text{IO}_4]^-$. Oxidation of ternary nitrilotriacetatocobalt(II) complexes involving succinate, malonate, tartrate, maleate and benzoate as secondary ligands by periodate has been investigated [17] [18] [19]. In all cases, initial cobalt(III) products were formed, and these changed slowly to the final cobalt(III) products. It is proposed that the reaction follows an inner-sphere mechanism, involving a ring closure step that is faster than the oxidation step. The IV^{VI} in the initial product is probably substitutional by water at a very slow rate due to the substituted inertness of Co(III) and also the Co(II)- OIO_3 bond being stronger than Co- H_2O bond. The oxidation of cobalt(II) complexes of propylenediaminetetraacetate (PDTA) [20], 1,3-diamino-2-hydroxypropanetetraacetate (HPDTA) [20], diethylenetriamine-pentaacetate (DPTA) [21], trimethylenediaminetetraacetate (TMDTA) [22] and ethyleneglycol,bis(2-aminoethyl)ether,N,N,N',N'-tetraacetate (EGTA) [22] by periodate gave only the final product. Periodate oxidations of the chromium(III) complexes of NTA [23], 2-aminopyridine [24] and IDA [25] were studied. In all cases, the electron transfer proceeds through an inner-sphere mechanism *via* coordination of IO_4^- to chromium(III).

In this paper, we report on the kinetics and mechanism of the periodate oxidation of ternary complexes of chromium(III) involving NTA as primary ligand and β -alanine as a secondary ligand, in order to study the effect of secondary ligand on the stability of $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$ [23] toward oxidation.

2. Experimental

2.1. Materials and Methods

The ternary complexes of chromium(III) involving nitrilotriacetato and β -alanine was prepared according to the report method [26]. All chemicals used in this study were of analar grade (BDH, Aldrich and Sigma). Buffer solutions were prepared from CH_3COONa (Sigma 99%) and CH_3COOH (BHD 99.9%) of known concentration. NaNO_3 (Aldrich 99.99%) was used to adjust ionic strength in the different buffered solutions. Doubly distilled H_2O was used in all kinetic runs. A stock solution of NaIO_4 (Aldrich 99.9%) was prepared by accurate weighing and wrapped in aluminum foil to avoid photochemical decomposition [27].

2.2. Instrumentation

UV-vis spectrophotometer model JASCO UV-530 was used to record the electronic spectra of the investigated complexes. The oxidation of complex, $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$ by IO_4^- were followed spectrophotometrically. The absorption measurements for the oxidation of reaction products are maximum at the reaction pH. Automatic circulation thermostat was used to regulate the temperature of solution. The average stabilizing accuracy was $\pm 0.1^\circ\text{C}$. Large excess of IO_4^- (>10-fold) was used in all measurements to get the pseudo-first order situation. NaNO_3 solution was utilized to make a constant ionic strength. It is noticed that during the course of the reaction the pH of the reaction is constant.

2.3. Kinetic Measurements


The UV-Visible absorption spectra of the products of oxidation of the complex $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$ by IO_4^- was followed spectrophotometrically for a measured period of time using a JASCO UV-530 spectrophotometer. All reactants were thermally equilibrated for ca 15 min in an automatic circulation thermostat, then mixed thoroughly and quickly transferred to an absorption cell. The oxidation rates were measured by monitoring the absorbance of Cr(VI) at 350 nm, on a Jenway 3600 spectrophotometer, where the absorption of the oxidation products is maximal at the reaction pH. The pH of the reaction mixture was measured using a G-C825 pH-meter. Pseudo-first-order conditions were maintained in all runs by maintaining a large excess (>10-fold) of IO_4^- over complexes. The ionic strength was kept constant by the addition of NaNO_3 solution. The pH of the reaction mixture was found to be constant during the reaction runs. Potentiometric measurements were performed with a Metrohm 702

SM titrino, using Irving and Rossotti techniques [28].

3. Results and Discussion

The UV-Visible spectra of the oxidation product of the complex, $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$ by periodate were recorded over time on a JASCO UV-530 spectrophotometer (Figure 1). The spectrum gives a maxima at 564 and 410 nm for $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$ complex which disappeared and replaced by a single peak at 350 nm due to the formation of chromium (VI). The presence of one isosbestic point at 501 nm in the absorption spectra (Figure 1) indicates the presence of two absorbing species in equilibrium. To measure the stoichiometry, a known excess of Cr(III) complex was added to IO_4^- solution and the absorbance of Cr(VI) produced was measured at 350 nm after 24 h. The quantity of Cr(III) consumed was calculated using the molar absorptivity of Cr(VI) at the utilized pH.

The oxidation of $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$ -complex by periodate was carried out in the pH range 3.40 - 4.45, 0.2 M ionic strength, $[\text{IO}_4^-]$ range $(0.5 - 5.0) \times 10^{-2}$ M and with temperature range $15^\circ\text{C} - 35^\circ\text{C}$ ($\pm 0.1^\circ\text{C}$). The stoichiometry of the reaction can be represented by Equation (1):



where Cr(III) and I(VII) represent total chromium(III)-complex and periodate, respectively. The concentration ratio of IO_4^- initially present to Cr(VI)




Figure 1. Absorbance spectra of reaction mixtures at interval times; $[\text{complex}] = 2.5 \times 10^{-4} \text{ mol}\cdot\text{dm}^{-3}$, $[\text{IO}_4^-] = 0.02 \text{ mol}\cdot\text{dm}^{-3}$, $I = 0.2 \text{ mol}\cdot\text{dm}^{-3}$, $\text{pH} = 4.05$ and $T = 30^\circ\text{C}$. Curve (.....) spectrum of the complex ($2.5 \times 10^{-4} \text{ mol}\cdot\text{dm}^{-3}$) at the same pH.

produced was found to be 3:2. The stoichiometry is also consistent with the observation that IO_3^- does not oxidize the Cr(III)-complex over the studied pH range. **Table 1** shows pseudo-first order rate constants, k_{obs} . Data obtained exhibits that k_{obs} does not have any effect, when we change the concentration of $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$ complex with constant IO_4^- concentration of $2.0 \times 10^{-2} \text{ mol}\cdot\text{dm}^{-3}$, pH = 4.05, ionic strength $0.20 \text{ mol}\cdot\text{dm}^{-3}$, temperature 25°C and at different concentrations of complex over the range $(1.25 - 6.25) \times 10^{-4} \text{ mol}\cdot\text{dm}^{-3}$, confirming that this reaction is first order and related to the concentration of Cr(III) complex, $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$. This behavior is represented by Equation (2).

$$\text{Rate} = k_{\text{obs}} [\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^- \quad (2)$$

The effect of periodate on the rate of the reaction of $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$ was studied over the temperature range ($15^\circ\text{C} - 35^\circ\text{C}$). The variation of rate constant, k_{obs} , with different concentrations of $[\text{IO}_4^-]$ at different temperatures are summarized in **Table 1**. Plotting k_{obs} against $[\text{IO}_4^-]$, was found to be linear without intercept as shown in **Figure 2**. The dependence of k_{obs} on $[\text{IO}_4^-]$ is thus described by Equation (3):

$$k_{\text{obs}} = k_1 [\text{IO}_4^-] \quad (3)$$

The dependence of the reaction rate on pH was investigated over the 3.40 - 4.45 pH range at constant $[\text{IO}_4^-] = 2.0 \times 10^{-2} \text{ mol}\cdot\text{dm}^{-3}$, $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^- = 2.5 \times 10^{-4} \text{ mol}\cdot\text{dm}^{-3}$, $I = 0.20 \text{ mol}\cdot\text{dm}^{-3}$ and $T = 25^\circ\text{C}$. The kinetic data are graphically represented in **Figure 3**. Variation of the k_{obs} with pH is summarized in (**Table 2**), which indicates that the reaction rate increases with increasing pH values. Plot of k_{obs} against $[\text{IO}_4^-]$ at different pH values are given in **Figure 3**. From **Figure 3**, it was found that, the slopes are dependent on pH (**Table 3**). Plot of these slopes (k_1) versus $1/\text{H}^+$ are linear with slope (k_3) and an intercept (k_2) according to Equation (4).

$$k_1 = k_2 + k_3 / [\text{H}^+] \quad (4)$$

Table 1. Dependence of k_{obs} on $[\text{IO}_4^-]$ at pH = 4.05, $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^- = 2.5 \times 10^{-4} \text{ mol}\cdot\text{dm}^{-3}$, and $I = 0.2 \text{ mol}\cdot\text{dm}^{-3}$ at different temperatures.


$10^2 [\text{IO}_4^-]$ (mol·dm ⁻³)	$10^4 k_{\text{obs}}$ (s ⁻¹)	15°C	20°C	25°C	30°C	35°C
0.5	0.500	0.88	1.20	1.51	2.31	
1.0	1.25	1.95	2.96	3.48	6.23	
1.5	2.51	3.25	4.15	4.81	8.50	
2.0	2.98	4.5	5.65	6.33	12.00	
3.0	4.46	6.68	8.31	10.15	13.50	
4.0	-	8.80	9.86	12.25	-	
5.0	7.25	10.20	12.75	-	20.05	

Table 2. Effect of pH on k_{obs} at $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^- = 2.5 \times 10^{-4} \text{ mol}\cdot\text{dm}^{-3}$, $I = 0.2 \text{ mol}\cdot\text{dm}^{-3}$, and $T = 25^\circ\text{C}$.

$10^2 [\text{IO}_4^-]$ ($\text{mol}\cdot\text{dm}^{-3}$)	$10^4 k_{\text{obs}}$ (s^{-1}) pH = 3.40	pH = 3.72	pH = 4.05	pH = 4.27	pH = 4.45
0.5	0.66	0.83	1.20	1.66	3.60
1.0	0.88	1.51	2.96	5.51	-
1.5	1.20	2.28	4.14	6.4	8.45
2.0	1.58	2.66	5.65	10.51	12.16
3.0	2.95	4.01	8.31	15.5	19.01
4.0	4.16	5.66	9.86	19.03	22.56
5.0	5.00	7.58	12.75	23.68	31.20

Table 3. Values of (k_1) at different temperatures.

T ($^\circ\text{C}$)	$10^3/T$ (K^{-1})	$10^2 k_1$ ($\text{mol}^{-1}\cdot\text{dm}^3\cdot\text{s}^{-1}$)	$-\ln k_1/T$ ($\text{mol}^{-1}\cdot\text{dm}^3\cdot\text{s}^{-1}\cdot\text{K}^{-1}$)
15	3.47	1.42	9.92
20	3.41	2.13	9.53
25	3.35	2.64	9.33
30	3.30	3.21	9.15
35	3.25	4.05	8.93

**Figure 2.** Plot of k_{obs} versus $[\text{IO}_4^-]$ at different temperatures.




Figure 3. Plot of k_{obs} versus $[\text{IO}_4^-]$ at different pH.

The values of k_2 and k_3 were obtained from the intercept and slope as $4.28 \times 10^{-3} \text{ mol}^{-1} \cdot \text{dm}^3 \cdot \text{s}^{-1}$ and $2.09 \times 10^{-6} \text{ s}^{-1}$ respectively at $T = 25^\circ \text{C}$.


From Equations (2), (3) and (4), the rate law for the oxidation of $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$ by periodate is given by Equation (5):

$$d[\text{Cr}^{\text{VI}}]/dt = [\text{IO}_4^-][\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^- (k_2 + k_3/[\text{H}^+]) \quad (5)$$

and




$$k_{\text{obs}} = (k_2 + k_3/[\text{H}^+])[\text{IO}_4^-] \quad (6)$$

Table 3 shows the values of k_1 which obtained from the slopes of **Figure 2** at different temperatures. From these results, thermodynamic activation parameters ΔH^\ddagger and ΔS^\ddagger associated with constant (k_1) in Equation (3) were calculated using Eyring approximation. ΔH^\ddagger and ΔS^\ddagger are equal to $35.75 \text{ kJ} \cdot \text{mol}^{-1}$ and $-155.3 \text{ J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$ respectively. According to the data reported, The effect of hydrogen ion concentration was investigated over the pH range 3.40 - 4.45, we noticed that in acidic aqueous medium the chromium(III) complex may be involved in the equilibrium shown in Equation (7).








The value of K_1 can be determined potentiometrically and has the value 1.70×10^{-5} at 25°C . From the pH range and K_1 value, it may be suggested that the involvement of the deprotonated form of the chromium(III)-complex in the rate-determining step. There are possibilities for the coordination of IO_4^- due

to the following reasons. Firstly, the H₂O ligand in [Cr^{III}(NTA)(Ala)(H₂O)]⁻ may be labile and hence substitution by IO₄⁻ is likely [29] [30] [31]. Secondly, periodate ion is capable of acting as a ligand, as evidenced from its coordination to copper(III) [32] and nickel(IV) [33]. Also there is a direct relationship between the reaction rate and ionic strength, where the values of 10⁴k_{obs} obtained at I = 0.30, 0.40, 0.50 and 0.60 mol·dm⁻³, pH = 4.05, [IO₄⁻] = 0.02 mol·dm⁻³ and T = 25°C are 5.83, 6.05, 6.27 and 6.57, respectively which is attributed to the reaction between similar charged species. It may be concluded that from the reported equilibrium constants of aqueous periodate solutions over the pH range used that, the periodate species likely to be present are IO₄⁻, H₄IO₆⁻ and H₃IO₆²⁻ [34], according to the equilibria, Equations (8)-(10):



From K₄ value, H₃IO₆²⁻ is not the predominant species (IO₄⁻ will be used to represent H₄IO₆⁻).

The mechanistic pathway for the oxidation of nitrilotriacetatetrisodium salt chromium(III) complex by periodate over the studied pH range may be represented by Equations (11)-(23):



From the above mechanism, the rate of the reaction is given by:

$$\begin{aligned} \frac{d[\text{Cr}^{\text{VI}}]}{dt} = & k_4 [\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{OIO}_3)]^{2-} \\ & + k_5 [\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{OH})(\text{OIO}_3)]^{3-} \end{aligned} \quad (16)$$

Since

$$[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{OIO}_3)]^{2-} = K_5 [\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^- [\text{IO}_4^-] \quad (17)$$

and

$$[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{OH})(\text{OIO}_3)]^{3-} = K_6 [\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{OH})]^{2-} [\text{IO}_4^-] \quad (18)$$

Substitution in Equations (17) and (18) in Equation (16) leads to:

$$\begin{aligned} d[\text{Cr}^{\text{VI}}]/dt = & K_2 k_4 [\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^- [\text{IO}_4^-] \\ & + k_5 K_6 [\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{OH})]^{2-} [\text{IO}_4^-] \end{aligned} \quad (19)$$

Since

$$[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{OH})]^{2-} = K_1 [\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^- / [\text{H}^+] \quad (20)$$

Substitution Equation (20) in Equation (19) we obtained:

$$\begin{aligned} d[\text{Cr}^{\text{VI}}]/dt = & K_5 k_4 [\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^- [\text{IO}_4^-] \\ & + (k_5 K_6 K_1 / [\text{H}^+]) [\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^- [\text{IO}_4^-] \end{aligned} \quad (21)$$

On rearrangement:

$$d[\text{Cr}^{\text{VI}}]/dt = (k_4 K_5 + k_5 K_6 K_1 / [\text{H}^+]) [\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^- [\text{IO}_4^-] \quad (22)$$

Hence,

$$k_{\text{obs}} = [\text{IO}_4^-] \left\{ k_4 K_5 + (k_5 K_1 K_6 / [\text{H}^+]) \right\} \quad (23)$$


From a comparison of Equations (6) and (23) one obtains $k_2 = k_4 K_5$ and $k_3 = k_5 K_1 K_6$. Equation (23) contains two terms, first term represents path independent of $[\text{H}^+]$ and the second term represents path dependent on $[\text{H}^+]$. In comparison with the oxidation of $[\text{Cr}(\text{NTA})(\text{H}_2\text{O})_2]$ [23] under the same conditions, the deprotonated complexes are significantly found to be more reactive than their conjugate acids. The rate of oxidation of this $[\text{Cr}(\text{NTA})(\text{H}_2\text{O})_2]$ is more than $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$. This means that the stability of the ternary complex, $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$, is more than the binary one, $[\text{Cr}(\text{NTA})(\text{H}_2\text{O})_2]$, toward oxidation. This may be due to the presence of the amino acid as a secondary ligand in the ternary complex, increase the stability of chromium(III) towards oxidation than binary complex, $[\text{Cr}^{\text{III}}(\text{NTA})(\text{H}_2\text{O})_2]$.

The small ΔH^\ddagger values and large negative activation entropies reasonably could reflect some nonadibatically in the electron transfer process [35]. Both ΔH^\ddagger and ΔS^\ddagger then may be expected to systematically increases as the orientation of the oxidant in the precursor complex is alter so as to enhance overlap between donor and acceptor redox orbitals and consequently the probability of adiabatic electron transfer [35]. The relatively low value of ΔH^\ddagger for $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$ is due to its composite value including formation which may be exothermic and intramolecular electron transfer which may be endothermic.

Enthalpies and entropies of activation for the oxidation of chromium(III) complexes by periodate are collected in **Table 4**. ΔH^\ddagger and ΔS^\ddagger for the oxidation of these complexes were calculated related to intramolecular electron transfer steps except for $[\text{Cr}^{\text{III}}(\text{HIDA})_2(\text{H}_2\text{O})]$, and $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Hist})(\text{H}_2\text{O})]^-$, ΔH^\ddagger and ΔS^\ddagger are composite values including the enthalpy of formation of the precursor complexes and the intramolecular electron transfer steps. A plot of ΔH^\ddagger versus ΔS^\ddagger for these complexes is shown in **Figure 4**, and an excellent linear relationship

Table 4. Enthalpies and entropies of activation for the oxidation of chromium(III) complexes by periodate.

Complex	$10^3 k^{\text{et}}$ (s^{-1})	ΔH^* (KJ/mol)	$-\Delta S^*$ (J/Kmol)	Ref.	Figure 4 key
$[\text{Cr}^{\text{III}}(\text{TOH})(\text{H}_2\text{O})]$	2.95	76	38.7	29	1
$[\text{Cr}^{\text{III}}(\text{NTA})(\text{Asp})(\text{H}_2\text{O})]^-$	3.93	64.6	76	13	2
$[\text{Cr}^{\text{III}}(\text{Ud})(\text{Asp})(\text{H}_2\text{O})_3]^{2+}$	0.70	59.5	106.8	16	3
$[\text{Cr}^{\text{III}}(\text{NTA})(\text{Hist})(\text{H}_2\text{O})]^-$	32.00	36.5	148	13	4
$[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$	26.40	35.75	155.3	This work	5
$[\text{Cr}^{\text{III}}(\text{Arg})_2(\text{H}_2\text{O})_2]^+$	3.46	30	192	13	6
$[\text{Cr}^{\text{III}}(\text{NTA})(\text{H}_2\text{O})_2]$	62.00	14	220	23	7
$[\text{Cr}^{\text{III}}(\text{HIDA})_2(\text{H}_2\text{O})]$	10.90	12.3	240.7	25	8

**Figure 4.** Enthalpies and entropies of activation of some chromium(III) complexes.

was obtained. Similar linear plots were found for a large number of redox reactions [36] [37] and for each reaction series a common rate-determining step is proposed. The isokinetic relation lends support a common mechanism for the oxidation of chromium(III) complexes, reported here, by periodate.

This consists of a periodate ion coordination to the chromium(III) complexes in step preceding the rate-determining intramolecular electron transfer within the precursor complex. Isokinetic compensation between ΔH^* and ΔS^* in a series of related reactions usually implies that one interaction between the reactants varies within the series, the remainder of the mechanism being invariant [32]. The electron transfer reactivities of these complexes with periodate are comparable, as the coordination of periodate with these complexes are identical. All of

this suggests that the excellent correlation often observed between ΔS^\ddagger and ΔH^\ddagger mainly reflects the fact that both thermodynamic parameters are in reality two measures of the same thing, and that measuring a compensation temperature is just a rather indirect way of measuring the average temperature at which the experiments were carried out. As this temperature will often be in a range that the experimenter expects to have some biological significance, it is not surprising if the compensation temperature turns out to have a biologically suggestive value [38] [39] [40].

4. Conclusion

Oxidation of $[\text{Cr}^{\text{III}}(\text{NTA})(\text{Ala})(\text{H}_2\text{O})]^-$ by periodate proceeds via an inner-sphere mechanism. Rate of oxidation increases with increasing pH. These reactions proceed through two-electron transfer process leading to the formation of chromium(VI). A common mechanism for the oxidation of ternary chromium(III) complex by periodate is proposed, and is supported by the excellent isokinetic relationship between ΔH^\ddagger and ΔS^\ddagger values for these reactions.

Conflicts of Interest

The authors declare no conflicts of interest regarding the publication of this paper.

References

- [1] Sigel, H. (1980) Ternary Complexes in Solution. 34. Discriminating and Stability increasing Properties of the Imidazole Moiety in Mixed-Ligand Complexes. *Inorganic Chemistry*, **19**, 1411. <https://doi.org/10.1021/ic50207a069>
- [2] Sigel, H., Operschall, B.P., Massoud, S.S., Song, B. and Griesser, R. (2006) Evidence for Intramolecular Aromatic-Ring Stacking in the Physiological pH Range of the Monodeprotonated Xanthine Residue in Mixed-Ligand Complexes Containing Xanthosinate 5'-monophosphate (XMP). *Dalton Transactions*, **46**, 5521. <https://pubs.rsc.org/en/content/articlelanding/2006/dt/b610082a#!divAbstract> <https://doi.org/10.1039/B610082A>
- [3] Arnold, H.F. (1991) Metal-Affinity Separations: A New Dimension in Protein Processing. *Bio/Technology*, **9**, 151. <https://www.nature.com/articles/nbt0291-151>
- [4] Roginski, E.E. and Mertz, W. (1969) Effects of Chromium(III) Supplementation on Glucose and Amino Acid Metabolism in Rats Fed a Low Protein Diet. *The Journal of Nutrition*, **97**, 525. <https://academic.oup.com/jn/article-abstract/97/4/525/4776397?redirectedFrom=fulltext> <https://doi.org/10.1093/jn/97.4.525>
- [5] Levina, A. and Lay, P. (2008) Chemical Properties and Toxicity of Chromium(III). *Chemical Research in Toxicology*, **21**, 563-571. <https://doi.org/10.1021/tx700385t>
- [6] Levina, A. and Lay, P.A. (2005) Mechanistic Studies of Relevance to the Biological Activities of Chromium. *Coordination Chemistry Reviews*, **249**, 281. <https://www.sciencedirect.com/science/article/pii/S0010854504000438> <https://doi.org/10.1016/j.ccr.2004.02.017>
- [7] Lilian, J., Yanjie, S., Jason, H., Weiwei, G., Stephen, P.C., Michael, W., Gary, A. L.,

- John, B.V. and Jean-Marc, L. (2003) Characterization of Chromodulin by X-Ray Absorption and Electron Paramagnetic Resonance Spectroscopies and Magnetic Susceptibility Measurements. *Journal of the American Chemical Society*, **125**, 774.
<https://doi.org/10.1021/ja0202661>
- [8] Tiziani, S., Sussich, F. and Cesaro, A. (2003) The Kinetics of Periodate Oxidation of Carbohydrates 2, Polymeric Substrates. *Carbohydrate Research*, **338**, 1083.
<https://www.sciencedirect.com/science/article/pii/S000862150300082X>
[https://doi.org/10.1016/S0008-6215\(03\)00082-X](https://doi.org/10.1016/S0008-6215(03)00082-X)
- [9] Bassil, D., Makris, D.P. and Kefalas, P. (2005) Oxidation of Caffeic Acid in the Presence of L-Cysteine: Isolation of 2-S-Cysteinylcaffeic Acid and Evaluation of Its Antioxidant Properties. *Food Research International*, **38**, 395.
<https://www.sciencedirect.com/science/article/pii/S0963996904002297>
<https://doi.org/10.1016/j.foodres.2004.10.009>
- [10] Mirkhani, V., Tangestaninejad, S., Moghadom, M. and Moghbel, M. (2004) Cytochrome P-450 Dependent Monooxygenases Model System: Rapid and Efficient Oxidation of Primary Aromatic Amines to Azo Derivatives with Sodium Periodate Catalyzed by Manganese(III) Schiff Base Complexes. *Bioorganic & Medicinal Chemistry*, **12**, 4673.
<https://www.sciencedirect.com/science/article/pii/S0968089604004742>
<https://doi.org/10.1016/j.bmc.2004.06.029>
- [11] Ewais, H.A. and Ismail, M.I. (2013) Mechanism of Electron Transfer Reaction of Ternary Dipicolinatochromium(III) Complex Involving Oxalate as Secondary Ligand. *Journal of Chemical Sciences*, **125**, 1152-1159.
<https://doi.org/10.1007/s12039-013-0491-x>
- [12] Abdel-Khalek, A.A., Sayyah, S.M. and Ewais, H.A. (1997) Kinetics and Mechanism of Oxidation of the chromium(III)-dl-valine Complex/Periodate Reaction. Evidence for Iron(II) Catalysis. *Transition Metal Chemistry*, **22**, 375.
<https://link.springer.com/article/10.1023/A%3A1018518019925#citeas>
<https://doi.org/10.1023/A:1018518019925>
- [13] Ewais, H.A., Habib, M.A. and Elroby, S.A.K. (2010) Kinetics and Mechanism of Periodate Oxidation of Two Ternary Nitrilotriacetatochromium(III) Complexes Involving Histidine and Aspartate Co-Ligands. *Transition Metal Chemistry*, **35**, 73.
<https://link.springer.com/article/10.1007/s11243-009-9297-6>
<https://doi.org/10.1007/s11243-009-9297-6>
- [14] Ewais, H.A., Ahmed, S.A. and Abdel-Khalek, A.A. (2004) Kinetics and Mechanism of Oxidation of Chromium(III) Complex of Guanosine by Periodate. *Inorganic Reaction Mechanism*, **5**, 125.
- [15] Abdel-Khalek, A.A., Ewais, H.A. and Ahmed, S.A. (2004) Kinetics and Mechanism of Oxidation of Chromium(III)-Guanosine 5-Monophosphate Complex by Periodate. *Journal of the Chinese Chemical Society*, **51**, 713.
<https://onlinelibrary.wiley.com/doi/abs/10.1002/jccs.200400108>
<https://doi.org/10.1002/jccs.200400108>
- [16] Khaled, S.E. (2006) Inner-Sphere Oxidation of Binary and Ternary Uridine Chromium(III) Complexes Involving Aspartate by Periodate. *Inorganic Reaction Mechanisms*, **6**, 247-256.
- [17] Ewais, H.A. (2008) Mechanism of Electron Transfer Reaction of Ternary Nitrilotriacetatocobalt(II) Complexes Involving Succinate and Malonate as Secondary Ligands. *International Journal of Chemical Kinetics*, **40**, 103.
<https://onlinelibrary.wiley.com/doi/abs/10.1002/kin.20289>
<https://doi.org/10.1002/kin.20289>

- [18] Abdel-Khalek, A.A., Khaled, E.S.H. and Mohamed, R.A. (2007) Kinetics Mechanism of Electron Transfer Reactions of Ternary Nitrilotriacetatocobaltate(II) Complexes Involving Maleate and Tartarate by Periodate. *Journal of Coordination Chemistry*, **61**, 152-161. <https://doi.org/10.1080/00958970701310902>
- [19] Ewais, H.A. (2009) Kinetics and Mechanism of the Oxidation of a Ternary Complex of Cobalt(II) Involving Nitrilotriacetate and Benzoate by Periodate in Acetate Medium. Evidence for Manganese(II) as a Catalyst. *Journal of Coordination Chemistry*, **62**, 152. <https://www.tandfonline.com/doi/full/10.1080/00958970802381190> <https://doi.org/10.1080/00958970802381190>
- [20] Naik, R.M., Sarker, J., Chaturvedi, D.D., Verma, A. and Singh, S.K. (2003) Kinetics of Oxidation of Cobalt(II) Complexes of Propylenediaminetetraacetate and 1,3-diamino-2-hydroxyprpanetetra Acetate by Periodate. *Indian Journal of Chemistry Section A*, **42**, 1639-1647. <http://hdl.handle.net/123456789/20691>
- [21] Abu-Elenien, M.H., Al-Shatti, N.I., Hussein, M.A. and Sulfab, Y. (1990) Kinetics and Mechanism of the Oxidation of Diethylenetriaminepentaacetatocobaltate(II) by Periodate. *Polyhedron*, **9**, 99-105. <https://www.sciencedirect.com/science/article/pii/S0277538700842527>
- [22] Naik, R.M., Srivastava, A., Tiwari, A.K., Yadav, S.B.S. and Verma, A.K. (2007) Kinetic and Mechanistic Studies of Oxidation of Amine-N-Polycarboxylates Complexes of Cobalt(II) by Periodate Ions in Aqueous Medium. *Journal of the Iranian Chemical Society*, **4**, 63-71. <https://link.springer.com/article/10.1007/BF03245804>
- [23] Abdel-Khalek, A.A. and Elsemongy, M.M. (1989) Kinetics of the Oxidation of Diaqua(nitrilotriacetato)-Chromium(III) by Periodate in Aqueous Solutions. *Transition Metal Chemistry*, **14**, 206-208. <https://doi.org/10.1007/BF01043796> <https://link.springer.com/article/10.1007/BF01043796>
- [24] Abdel-Hady, A.M. (2000) Kinetics of Oxidation of a 2-Aminomethyl Pyridine Chromium (III) Complex by Periodate. *Transition Metal Chemistry*, **25**, 437-442. <https://link.springer.com/article/10.1023/A%3A1007019528319> <https://doi.org/10.1023/A:1007019528319>
- [25] Ewais, H.A., Al-Otaibi, F.D. and Abdel-Khalek, A.A. (2006) Kinetics and Mechanism of Oxidation of Iminodiacetatochromium(III) by Periodate. *Inorganic Reaction Mechanisms*, **6**, 39-47.
- [26] Sharma, C.L., Jain, P.K. and De, T.K. (1980) Preparation and Characterization of Some mixed Ligand Complexes of Chromium Nitrilotriacetate with Some Amino Acids. *Inorganic Nuclear Chemistry*, **42**, 1681-1687. <https://ac.els-cdn.com/0022190280801394/1-s2.0-0022190280801394-main.pdf>
- [27] Symons, M.C.R. (1955) Photochemical Decomposition of Periodate. *Journal of the Chemical Society*, 2794-2799. <https://doi.org/10.1039/jr9550002794>
- [28] Irving, H.M. and Rossotti, H.S. (1953) The Calculation of Formation Curves of Metal Complexes from pH Titration Curves in Mixed Solvents. *Journal of the Chemical Society*, 2904-2910. <https://doi.org/10.1039/jr9530003397> <https://pubs.rsc.org/en/content/articlelanding/1954/jr/jr9540002904#!divAbstract>
- [29] Abdel Khalek, A.A., Sayyah, S.M. and Abdel-Hameed, F.F. (1994) Kinetics and Mechanism of Oxidation of Chromium(III)-Tetraoxalylurea Complex by Periodate. *Transition Metal Chemistry*, **19**, 108-110. <https://doi.org/10.1007/BF00166281> <https://link.springer.com/article/10.1007/BF00166281>
- [30] Amal, S., Basaleh, A. and Abdel-Salam, H. (2016) Inner-Sphere Oxidation of a Ternary Dipicolinatochromium(III) Complex Involving a Malonic Acid Co-Ligand. *Transition Metal Chemistry*, **41**, 713-719.

- <https://doi.org/10.1007/s11243-016-0073-0>
<https://link.springer.com/article/10.1007%2Fs11243-016-0073-0>
- [31] Hassan, A., Ewais, R., Al-Orabi, O. and Obaid, A.Y. (2016) Manganese(II) Catalyzed Periodate Oxidation of a Ternary Dipicolinatochromium(III) Complex with Iminodiacetate as Co-Ligand: Mechanistic and Kinetic Study. *Transition Metal Chemistry*, **41**, 427-434. <https://doi.org/10.1007/s11243-016-0038-3>
<https://link.springer.com/article/10.1007/s11243-016-0038-3>
- [32] Hadince, I., Jenosovsky, L., Linek, A. and Syncecek, V. (1960) Über die Struktur der komplexen Perkuprate. *Naturwiss*, **47**, 377.
<https://link.springer.com/article/10.1007/BF00631384>
- [33] Ray, P. (1957) Sodium and Potassium Nickel (IV) Paraperiodates. *Inorganic Synthesis*, **5**, 201-208.
- [34] Crouthamel, C.E., Meek, H.V., Martin, D.S. and Banks, C.V. (1949) Spectrophotometric Studies of Dilute Aqueous Periodate Solutions. *Journal of the American Chemical Society*, **71**, 3031-3035. <https://pubs.acs.org/doi/10.1021/ja01177a026>
<https://doi.org/10.1021/ja01177a026>
- [35] Mcardle, J.V., Coyle, C.L., Gray, H.B., Yoneda, C.S. and Howerda, R.A. (1977) Kinetics Studies of the Oxidation of Blue Copper Proteins by Tris(1,10-phenanthroline)-Cobalt(III) Ions. *Journal of the American Chemical Society*, **99**, 2483-2489. <https://pubs.acs.org/doi/10.1021/ja00450a014>
<https://doi.org/10.1021/ja00450a014>
- [36] Holwerda, R.A. and Clemmer, J.D. (1979) Isokinetic Relationship in the Oxidation of Cuprous Stellacyanin by Cobalt(III) Complexes. *Bioinorganic Chemistry*, **11**, 7-15. <https://www.sciencedirect.com/science/article/pii/S0162013400800483>
- [37] Wherland, S. and Gray, H.B. (1977) Biological Aspects of Inorganic Chemistry. Wiley, New York, 189.
- [38] Bowden, A.C. (2002) Enthalpy-Entropy Compensation: A Phantom Phenomenon. *Journal of Biosciences*, **27**, 121-126. <https://doi.org/10.1007/BF02703768>
<https://link.springer.com/article/10.1007/BF02703768>
- [39] Abu-Gharib, E.A., EL-Khatib, R.M., Nassr, L.A.E. and Abu-Dief, A.M. (2011) Kinetics of Base Hydrolysis of Some Chromen-2-One Indicator Dyes in Different Solvents at Different Temperatures. *Journal of the Korean Chemical Society*, **55**, 346-353. <https://doi.org/10.5012/jkcs.2011.55.3.346>
- [40] Abu-Gharib, E.A., EL-Khatib, R.M., Nassr, L.A.E. and Abu-Dief, A.M. (2017) Kinetics, Reactivity, Initial-Transition State Analysis and Thermodynamic Parameters of Base-Catalyzed Hydrolysis of Coumalic Acid in Solvents with Different Polarities. *Arabian Journal of Chemistry*, **10**, S988-S995.
<https://www.sciencedirect.com/science/article/pii/S1878535213000117>