

# Endothelin-2 Differential Expression in Normal and Early-Stages of Colon Cancer Development

Mariana Bianchi<sup>1</sup>, Javier Adur<sup>1</sup>, Maria F. Izaguirre<sup>1</sup>, Silvia Viale<sup>1</sup>, Carlos L. Cesar<sup>2</sup>, Víctor H. Casco<sup>1\*</sup>

<sup>1</sup>Microscopy Laboratory Applied to Cellular and Molecular Studies, Bioengineering and Bioinformatic School, National University of Entre Ríos, Concepción del Uruguay, Argentina; <sup>2</sup>INFABiC-National Institute of Science and Technology on Photonics Applied to Cell Biology, Campinas, Brazil.

Email: \*vcasco@bioingenieria.edu.ar

Received May 23<sup>rd</sup>, 2013; revised June 22<sup>nd</sup>, 2013; accepted June 30<sup>th</sup>, 2013

Copyright © 2013 Mariana Bianchi *et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

The endothelin family has been related with several pathological diseases including cardiovascular disorders, hypertension and cancer. However, little is known about endothelin system in early stages of colorectal cancer and there are no studies evaluating differences in proximal and distal colon segments. To deepen in this issue, we have studied the endothelin's family gene and protein expression in normal mice and early stage of a mice model of Azoxymethane (AOM) and Dextran Sodium Sulphate (DSS) induced colorectal cancer in proximal and distal segments. Additionally, using nonlinear microscopy (NLM) techniques, we have characterized collagen changes in early stages of cancer disease development. In the present study, we have found significant differential gene expression and protein localization between these colon regions, which allow us to hypothesize a new role for the ET-2 as an early marker of colon cancer development.

**Keywords:** Colon Cancer; Endothelin-2; Azoxymethane; Dextran Sodium Sulphate

## 1. Introduction

Endothelin family includes three cyclic peptides (ET-1, ET-2 and ET-3) of 21-amino acid with two disulfide bridges established between cysteine residues located at positions 1 - 15 and 3 - 11, and two general types of receptors (ETA and ETB). ET-2 and ET-3 differ from ET-1 in two and six amino acid residues respectively. In mice, a homolog of ET-2 that only diverges in one amino acid residue has been called vasoactive intestinal contractor (VIC) [1]. It is widely accepted that endothelin axis is closely related with several pathologies including cardiovascular disorders, pulmonary disease, renal diseases, cancer and many others [2]. Endothelin axis also takes part in the physiology [3-6] and pathology of gastrointestinal tract [2] although their function is not clear yet.

Alterations in the expression of ET-1 ligand and the receptors ETA and ETB have been shown in human colon cancer. While in premalignant adenomas and malignant colon carcinomas, exhibit increased levels of pre-proET-1, endothelin-converting enzymes, and ET-1 [7,8],

there is no consensus about how the ETA and ETB receptors behave [9,10]. On the other hand, it has been recently proved that ET-2 and ET-3 are silenced in colon cancer and their induced overexpressions inhibit cell migration and invasion [11]. In that work, it was hypothesized that ET-2 and ET-3 might be silenced in early stages of cancer avoiding "mixed messages" in the ET axis. Both the disappearance of ET-2 and ET-3 and the overexpression of ET-1 could lead to the cancer progress. In the present report, we prove that this relation is not linear for ET-2, since its gene expression goes down only in distal colon region.

## 2. Materials and Methods

### 2.1. Animal Proceedings

Adult male BALB/c mice ( $n = 10$ ) 20 - 30 g body weight, was purchased from Center of Biological Experimentations and Biotery of National University of Litoral, Santa Fe, Argentina. Mice were killed by cervical dislocation and colon segments (proximal and distal) were removed. Our experimental procedures were in accordance with the Guidelines on Handling of Laboratory Animals for

\*Corresponding author.

our institution.

## 2.2. Murine Model of Chemical-Induced Colon Cancer (AOM/DSS)

Adult male BALB/c mice were intraperitoneal injected with Azoxymethane (AOM) 10 mg/Kg body weight and a week later, Dextran Sodium Sulphate (DSS) was given in the drinking water for seven days following the procedure reviewed by Rosenberg *et al.* (2009) [12], Tanaka (2009) [13] and De Robertis *et al.*, (2011) [14]. Control animal were injected with saline solution and were not given DSS. Samples were taken at day zero and at week four after injection. Proximal and distal segments were destined to histological procedure, immunofluorescence analysis and semi-quantitative RT-PCR for their study.

## 2.3. Histological Procedure

Colon segments were Carnoy's fixed (2 h, room temperature) dehydrated in ethanol series and paraffin embedded. Samples were cut at 4  $\mu$ m thick in microtome, dewaxed, rehydrated in a decreasing ethanolic series and Eosin and Hematoxylin stained.

## 2.4. Immunofluorescence

The immunoreactions were made on 4  $\mu$ m-thick microtome (Reichert, Austria) sections. The sections were permeated with Triton X-100 0.3% in PBS. To avoid non-specific reactions, sections were blocked with heat-inactivated normal goat serum and heat-inactivated normal donkey serum, as required. The antibodies used were rabbit polyclonal IgG (Immuno-Biological Laboratories Co., Ltd., Gunma, Japan) against ET-1, diluted 1:50, goat polyclonal IgG against ET-2 (Santa Cruz, sc-21627), diluted 1:50, rabbit polyclonal IgG against ETA receptor (Santa Cruz, sc-33535), diluted 1:50 and rabbit polyclonal IgG (Immuno-Biological Laboratories Co., Ltd., Gunma, Japan) against ETB receptor, diluted 1:50. The secondary antibodies were goat anti-rabbit IgG-Cy3 (Chemicon, AP187C) and donkey anti-goat IgG-Cy3 (Chemicon, AP180C). Control staining was carried out jumping the primary antibody to the dilution buffer. Af-

ter reaction, nuclei were stained with Hoechst 33342 (Sigma B-2261). Finally the sections were mounted using Vectashield, fluorescence mounting medium (Vector Laboratories) and observed and recorded using an Olympus BX-50 epifluorescence microscope equipped with an Apogee AM4 CCD camera.

## 2.5. Semi-Quantitative PCR

Samples were RNA-extracted with TRIzol reagent (Invitrogen) and retro-transcribed with Revert Aid First Strand cDNA Synthesis Kit (Fermentas) using oligo (dT) primers to synthesize cDNA only from mRNA. The cDNAs obtained were PCR amplified and solved in an agarose gel. The primer sequences for each amplified gene are listed in **Table 1**. The housekeeping gene GAPDH was used as internal control and each analyzed gene was normalized using this control by gel-analysis software, developed by ourselves [15].

## 2.6. Non-Linear Optical Microscopy

Two-photon excitation fluorescence (TPEF) and second-harmonic generation (SHG) images were acquired with an inverted microscope Axio Observer Z.1 (Carl Zeiss AG, Germany) equipped with a Zeiss LSM 780-NLO confocal scan head (Carl Zeiss AG, Germany), as was recently described in detail [16]. TPEF and SHG images were excited with 20 mW (at the sample) of a 940 nm fs beam, generating a SHG at 470 nm and a TPEF > 490 nm.

## 2.7. Statistical Analysis

The results are presented as means  $\pm$  SD and analyzed by unpaired t-test with Welch correction to determine significant differences among data groups. P values lower than 0.05 were considered statistically significant and lower than 0.01 were very significant.

## 3. Results

### 3.1. Cancer Induction Model

We run this experience in order to study the behavior of

**Table 1. Optimal primers for real-time PCR of murine endothelin system and GAPDH.**

Gene	Sense	Antisense
ET-2/VIC	5'-CTGCGTTTTTCGTCGATGCT-3'	5'-TGCAGCTCATGGTGTATCTCTTC-3'
ET-1	5'-TTCCCGTGATCTTCTCTCTGCT-3'	5'-CCTGCTTGGCAGAAATTCCA-3'
ETA	5'-GCTGGTTCCTCTTCACTTAAGC-3'	5'-TCATGGTTGCCAGGTTAATGC-3'
ETB	5'-TGTGCTCTAAGTATTGACAGATATCGAG-3'	5'-GGCTGTCTTGTA AAACTGCATGA-3'
GAPDH	5'-CTTACCACCATGGAGAAGGC-3'	5'-GGCATGGACTGTGGTCATGAG-3'

endothelin axis in the initial step of cancer development. At second week of the experience, while animals were drinking DSS, they suffer diarrhea and anal bleeding probably caused by the DSS-generated inflammation. At week three all of them started getting better and they looked completely recovered at fourth week. Comparing with normal colon (**Figure 1(A)**), Eosin and Hematoxylin histology showed signs of inflammation and aberrant crypt foci and occasionally little microadenomas (**Figure 1(B)**) as was previously informed by De Robertis *et al.*, (2011), especially in distal colon.


### 3.2. Immunofluorescence

In normal conditions, ET-2 was mainly immunolocalized at the base of distal colon crypts (**Figure 1(C)**), in contrast with proximal colon where the localization is much more changeful, including middle and upper part of the crypts. Furthermore the quantification of fluorescence in the immunoreaction gave significant differences between the segments, having proximal colon higher expression than distal one (not showed). Contrary to ET-2, ET-1 was homogeneously expressed both in proximal and distal region. ETA receptor was localized weakly in distal colon, mainly in the basal membrane (**Figure 1(E)**), while its immunolocalization is significant higher in proximal colon (not showed). On the other hand, ETB receptor was present in the nuclei of epithelial and some muscle cells, according with the observations of Takizawa *et al.*, (2004) [6], Bianchi *et al.*, (2012) [4] and several reports that have proved that the density of ETB receptor is higher in the nucleus than in the cytosol of all human cell types tested [17-20].

We studied distal colon in early stages of cancer induction, where the major rate of tumors is reached in this mice model [12]. The results of this preliminary study have demonstrated that ET-2 peptide was not localized in all the analyzed crypts, but the peptide seems to be reduced in epithelial cells while increase in immune cells like lymphocytes (**Figure 1(D)**). ET-1 peptide and ETB receptor did not show significant differences in normal and induced animals, ETA receptor seem to reach a higher expression in tissue-infiltrating lymphocytes (**Figure 1(F)**).

### 3.3. Gene Expression

In normal mice, the semi-quantitative gene expression analysis showed that the mean of ET-2 transcripts in distal colon was significantly lower than in proximal one (**Figure 2(a)**), these results seems to be in line with the quantification values of immunofluorescence. Despite having a tendency, the other analyzed genes gave no significant differences (**Figure 2(a)**). In early stages of induced cancer mice, ET-2 was not detected in distal colon.



**Figure 1.** Histological sections of E&H-stained slices from normal (A) and AOM/DSS induced animals (B). Immunofluorescence of ET-2 in normal (C) and AOM/DSS induced animals (D), and immunofluorescence of ETA receptor in normal (E) and AOM/DSS induced animals (F),  $n = 5$ .


Although the other analyzed genes in the induced model did not exhibit significant differences between proximal and distal colon, they did display the same tendency that was observed in normal mice colon (**Figure 2(b)**).

There were no significant differences between ET-2 gene expression in ascending colon between normal and pathological mice. By the contrary, in descending colon, the peptide in the induced cancer model diminished its expression level and it could not be detected by the used approach (**Figure 2(c)**).

Comparing ETA expression in normal and pathological conditions, it was verified that the gene was overexpressed both in proximal and distal colon, being the differences very significant in distal and significant in proximal colon (**Figure 2(c)**).

### 3.4. Non-Linear Optical Microscopy

Crypts were identified with TPEF and the connective tis-



**Figure 2. Semi-quantification of gene expression relative to GAPDH of ET-2, ET1, ETA and ETB receptors in proximal and distal portions of control animals (a). Semi-quantification of gene expression relative to GAPDH of ET-2, ET1, ETA and ETB receptors in proximal and distal portions of AOM/DSS induced animals (b). Comparison of the gene expression both in proximal (PC) and distal colon (DC) and in control and AOM/DSS treated animals (c). n=5, ND: not detectable, \* ( $p < 0.05$ ), \*\* ( $p < 0.001$ ).**

sue between crypts was detected via SHG of collagen (**Figure 3**). In the early stage of tumor tissue, SHG images showed changes in the fibrous stroma. Additionally to collagen fibers morphology, collagen density changes were detected (**Figure 3**).


#### 4. Discussion

Since most of the current studies are focused on well-established colon cancer model behavior, the main objective of the present work was to analyze the endothelin system in early stages of colon cancer development. In normal mice, endothelins are highly expressed in the gastrointestinal tract [3,21,22]. The previously unreported-differential distribution of ET-2 (both mRNA and peptide) along colon must be focused in order to understand how this peptide acts. This particular distribution is more interesting yet, when we verify that within the crypt axis this peptide is not uniformly expressed, being mainly

immunolocalized at the base of distal colon crypts, whereas in proximal colon it was localized also at the middle and upper part of the crypts. This expression pattern suggests a pleiotropic activity, which seems to be more extent in proximal colon than in distal one.

This particular distribution is deeply modified in early stages of the mice AOM/SDD colon cancer model showing a significant decrease of the mRNA-ET-2 expression on distal colon segment, where it became no detectable. The immunofluorescence pattern is coincident with these results, since most of the crypts are negatives for that peptide and only a weak reaction can be seen. Coincident with our results, recent studies on colon cancer induction models, have reported an epigenetic silencing of the ET-2 gene several weeks before the onset of frank tumors [11]. As was previously mentioned, distal colon exhibit the major tumors rate in early stages of cancer induction [12], therefore ET-2 down regulation





**Figure 3.** Eosin and hematoxylin, SHG and TPEF images obtained from colon of normal and AOM/DSS week 4 treated mice. Barr: 200  $\mu$ m.

may be an early prognostic marker of the disease. We hypothesize that at this stage silencing could start in distal portion, and lengthen to the rest of the colon progressively, while malignancy increases. Although it was reported that significantly elevated ET-1 expression occurs in 80% of primary human colon cancers [23,24], we found no significant differences comparing ET-1 gene expression in normal and early stage colon cancer conditions, thus this peptide might become overexpressed just in later stages.

The ETA receptor plays an important role in tumor cell migration, metastasis, and proliferation. The ETB receptor plays a critical role in angiogenesis and the inhibition of anti-tumor immune cell recruitment [25]. In this work, ETA receptor underwent an overexpression both in proximal and distal segments; this behavior was reported several times in many kinds of tumors [26-32], including colorectal cancer [9]. Recent evidence showed that ETA receptor is expressed on CD133+ cancer stem cells in both cell lines and primary human tumor cells [25] and that CD133 expression exhibit enhanced tumorigenicity over CD133-negative (CD133-) cells, thus the former would be a more malignant phenotype than the latter [33]. Increased expression of ETA receptor in distal colon in early stage of cancer could be related to its increasing presence in cancer stem cells. Furthermore, ETA receptor inhibition prevents the induction of

ICAM1, necessary for immune cell recruitment [25]. Since we observed ETA immunoreactivity in lymphocytes, we could hypothesize that its increased gene expression would respond to the increased number of infiltrated lymphocytes. In contrast, ETB receptor, kept on in the same level expression, despite being reported it is down-regulated in colorectal cancer [9] this does not seem to occur in the early stages of the disease.

Collagen forms the structural network of the extracellular matrix (ECM) in tissue and plays an important role during epithelial tumor progression. SHG has already been shown to have potential applicability for cancer diagnosis by revealing changes in the ECM in tumors relative to normal tissues [34]. An advantage of SHG compared with other visualization techniques, such as bright field H&E images, is the SHG selectivity, which is blind to the background populated with too many other things and only sees the collagen fibril's pattern. This allows an automatic digital imaging processing of collagen network only. The results obtained in this work from SHG are in line with previous studies [35-37]. Jain and coworkers demonstrated that an increase of collagen concentration (*i.e.*, desmoplasia) was associated with implanted melanoma-cell tumor [38]. On the other hand, in a complete analysis of human gastrointestinal tract, Nathanson and coworkers find out small amount of collagen in benign tissues [39]. Furthermore, in a recent

study was demonstrated that ET-1 stimulates processes involved in tumor stroma production by colonic fibroblasts [40]. Because the collagen deposition in stroma is a dynamic process which regulates epithelial proliferation/tumor formation cycle, future studies must be performed. However, the present results suggest that collagen concentration could be an important factor in tumoral progression and can be used also as biomarker.

## 5. Conclusion

Our findings emphasize the idea that the colon must not be considered as a homogeneously functional structure, at least in the endothelin system behavior, according to our study, it allows us dividing it in at least proximal and distal portions. Furthermore, our work found that at early stages of colon cancer, the first modifications in the endothelin axis are the decrease of ET-2 in distal portion and the overexpression of ETA receptor in whole colon.

Since ET-1 acts as a strong stimulus for tumor stroma secretion, collagen generation in the tumor might be increased as a result of ET-1 overexpression. As the disappearance of ET-2 occurs before that, we support that ET-2 decrease might serve as an early biomarker of colon cancer risk, even more than stroma or ET-1 increment.

## 6. Acknowledgements

This paper was supported by the PID-UNER 6116. The authors are grateful to CEPOF (Optics and Photonics Research Center, FAPESP) and National Institute of Photonics Applied to Cell Biology (INFABIC), UNICAMP, Campinas, Brazil.

## REFERENCES

- [1] K. Saida, Y. Mitsui and N. Ishida, "A Novel Peptide, vasoactive Intestinal Contractor, of a New (Endothelin) Peptide Family. Molecular Cloning, Expression, and Biological Activity," *Journal of Biological Chemistry*, Vol. 264, No. 25, 1989, pp. 14613-14616.
- [2] A. K. Khimji and D. C. Rokey, "Endothelin—Biology and Disease," *Cellular Signalling*, Vol. 22, No. 11, 2010, pp. 1615-1625. [doi:10.1016/j.cellsig.2010.05.002](https://doi.org/10.1016/j.cellsig.2010.05.002)
- [3] I. Chang, A. N. Bramall, A. G. Baynash, A. Rattner, D. Rakheja, M. Post, S. Joza, C. McKerlie, D. J. Stewart, R. R. McInnes and M. Yanagisawa, "Endothelin-2 Deficiency Causes Growth Retardation, Hypothermia, and Emphysema in Mice," *The Journal of Clinical Investigation*, 2013. [doi:10.1172/JCI66735](https://doi.org/10.1172/JCI66735)
- [4] M. Bianchi, J. F. Adur, S. Takizawa, K. Saida and V. H. Casco, "Endothelin System in Intestinal Villi: A Possible Role of Endothelin-2/Vasoactive Intestinal Contractor in the Maintenance of Intestinal Architecture," *Biochemical and Biophysical Research Communications*, Vol. 417, No. 4, 2012, pp. 1113-1118. [doi:10.1016/j.bbrc.2011.12.053](https://doi.org/10.1016/j.bbrc.2011.12.053)
- [5] S. Takizawa, T. Uchide, J. Adur, T. Kozakai, E. Kotake-Nara, J. Quan and K. Saida, "Differential Expression of Endothelin-2 along the Mouse Intestinal Tract," *Journal of Molecular Endocrinology*, Vol. 35, No. 2, 2005, pp. 201-209. [doi:10.1677/jme.1.01787](https://doi.org/10.1677/jme.1.01787)
- [6] S. Takizawa, T. Uchide, T. Kozakai, J. Adur, J. Quan and K. Saida, "Immunolocalization of Endothelin-B Receptor in Mouse Intestinal Tract," *Journal of Cardiovascular Pharmacology*, Vol. 44, 2004, pp. S329-S331. [doi:10.1097/01.fjc.0000166285.85785.7a](https://doi.org/10.1097/01.fjc.0000166285.85785.7a)
- [7] H. Ali, M. Dashwood, K. Dawas, M. Loizidou, F. Savage and I. Taylor, "Endothelin Receptor Expression in Colorectal Cancer," *Journal of Cardiovascular Pharmacology*, Vol. 36, Suppl. 1, 2000, pp. S69-S71.
- [8] G. Egidy, L. Juillerat-Jeanneret, J. F. Jeannin, P. Korth, F. T. Bosman and F. Pinet, "Modulation of Human Colon Tumor-Stromal Interactions by the Endothelin System," *The American Journal of Pathology*, Vol. 157, No. 6, 2000, pp. 1863-1874. [doi:10.1016/S000](https://doi.org/10.1016/S000)
- [9] M. M. Hoosein, M. R. Dashwood, K. Dawas, H. M. Ali, K. Grant, F. Savage, I. Taylor and M. Loizidou, "Altered Endothelin Receptor Subtypes in Colorectal Cancer," *European Journal of Gastroenterology & Hepatology*, Vol. 19, No. 9, 2007, pp. 775-782. [doi:10.1097/MEG.0b013e3282c563de](https://doi.org/10.1097/MEG.0b013e3282c563de)
- [10] L. P. Eberl, O. Valdenaire, V. Saintgiorgio, J. F. Jeannin and L. Juillerat-Jeanneret, "Endothelin Receptor Blockade Potentiates FasL-Induced Apoptosis in Rat Colon Carcinoma Cells," *International Journal of Cancer*, Vol. 86, No. 2, 2000, pp. 182-187. [doi:10.1002/\(SICI\)1097-0215\(20000415\)86:2<182::AID-IJC6>3.0.CO;2-G](https://doi.org/10.1002/(SICI)1097-0215(20000415)86:2<182::AID-IJC6>3.0.CO;2-G)
- [11] R. Wang, C. V. Löhr, K. Fischer, W. M. Dashwood, J. A. Greenwood, E. Ho, D. E. Williams, H. Ashktorab, M. R. Dashwood and R. H. Dashwood, "Epigenetic Inactivation of Endothelin-2 and Endothelin-3 in Colon Cancer," *International Journal of Cancer*, Vol. 132, No. 5, 2013, pp. 1004-1012.
- [12] D. W. Rosenberg, C. Giardina and T. Tanaka, "Mouse Models for the Study of Colon Carcinogenesis," *Carcinogenesis*, Vol. 30, No. 2, 2009, pp. 183-196. [doi:10.1093/carcin/bgn267](https://doi.org/10.1093/carcin/bgn267)
- [13] T. Tanaka, "Colorectal Carcinogenesis: Review of Human and Experimental Animal Studies," *Journal of Carcinogenesis*, Vol. 8, No. 1, 2009, p. 5. [doi:10.4103/1477-3163.49014](https://doi.org/10.4103/1477-3163.49014)
- [14] M. D. Robertis, E. Massi, M. L. Poeta, S. Carotti, S. Morini, L. Cecchetelli, E. Signori and V. M. Fazio, "The AOM/DSS Murine Model for the Study of Colon Carcinogenesis: From Pathways to Diagnosis and Therapy Studies," *Journal of Carcinogenesis*, Vol. 10, No. 1, 2011, p. 9. [doi:10.4103/1477-3163.78279](https://doi.org/10.4103/1477-3163.78279)
- [15] L. Bugnon, "Diseño y Desarrollo de un Sistema para Fotodocumentación y análisis Cuantitativo de Geles de Electroforesis," Degree Bioengineering Thesis, Facultad de Ingeniería, Universidad Nacional de Entre Ríos, Concepción del Uruguay, 2013.
- [16] J. Adur, V. B. Pelegati, A. A. de Thomaz, M. O. Baratti, D. B. Almeida, L. A. Andrade, F. Bottcher-Luiz, H. F.

- Carvalho, C. L. Cesar, "Optical Biomarkers of Serous and Mucinous Human Ovarian Tumor Assessed with Nonlinear Optics Microscopies," *Plos One*, Vol. 7, No. 10, 2012, Article ID: e47007. [doi:10.1371/journal.pone.0047007](https://doi.org/10.1371/journal.pone.0047007)
- [17] G. Bkaily, L. Avedanian, J. Al-Khoury, C. Provost, M. Nader, P. D'Orléans-Juste and D. Jacques, "Nuclear Membrane Receptors for ET-1 in Cardiovascular Function," *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, Vol. 300, No. 2, 2011, pp. R251-R263. [doi:10.1152/ajpregu.00736.2009](https://doi.org/10.1152/ajpregu.00736.2009)
- [18] G. Bkaily, L. Avedanian, J. Al-Khoury, L. Ahmarani, C. Perreault and D. Jacques, "Receptors and Ionic Transporters in Nuclear Membranes: New Targets for Therapeutic Pharmacological Interventions," *Canadian Journal of Physiology and Pharmacology*, Vol. 90, No. 8, 2012, pp. 953-965. [doi:10.1139/y2012-077](https://doi.org/10.1139/y2012-077)
- [19] J. C. Honore, E. Carrier, M. H. Fecteau, C. R. Tirapelli, G. Bkaily and P. D'Orléans-Juste, "Nonselective ETA/ETB-Receptor Blockade Increases Systemic Blood Pressure of Bio 14. 6 Cardiomyopathic Hamsters," This Article Is One of a Selection of Papers Published in the Special Issue (Part 1 of 2) on Frontiers in Endothelin, *Canadian Journal of Physiology and Pharmacology*, Vol. 86, No. 6, 2008, pp. 394-401. [doi:10.1139/Y08-041](https://doi.org/10.1139/Y08-041)
- [20] L. Avedanian, D. Jacques and G. Bkaily, "Presence of Tubular and Reticular Structures in the Nucleus of Human Vascular Smooth Muscle Cells," *Journal of Molecular and Cellular Cardiology*, Vol. 50, No. 1, 2011, pp. 175-186. [doi:10.1016/j.yjmcc.2010.10.005](https://doi.org/10.1016/j.yjmcc.2010.10.005)
- [21] J. Adur, T. Uchide, S. Takizawa, J. Quan and K. Saida, "Real-Time Polymerase Chain Reaction Quantification of Gene Expression Levels of Murine Endothelin-A and Endothelin-B Receptors: Gene Expression Profiles by the Standard Curve Method," *Journal of Cardiovascular Pharmacology*, Vol. 44, Suppl. 1, 2004, pp. S321-S328. [doi:10.1097/01.fjc.0000166270.01926.00](https://doi.org/10.1097/01.fjc.0000166270.01926.00)
- [22] J. F. Adur and V. H. Casco, "Endotelinas Murinas," Editorial Académica Española, Saarbrücken, 2011.
- [23] T. H. Kim, H. Xiong, Z. Zhang and B. Ren, " $\beta$ -Catenin Activates the Growth Factor Endothelin-1 in Colon Cancer Cells," *Oncogene*, Vol. 24, No. 4, 2004, pp. 597-604. [doi:10.1038/sj.onc.1208237](https://doi.org/10.1038/sj.onc.1208237)
- [24] E. Asham, A. Shankar, M. Loizidou, S. Fredericks, K. Miller, P. B.oulos, G. Burnstock and I. Taylor, "Increased Endothelin-1 in Colorectal Cancer and Reduction of Tumour Growth by ET A Receptor Antagonism," *British Journal of Cancer*, Vol. 85, No. 11, 2001, p. 1759. [doi:10.1054/bjoc.2001.2193](https://doi.org/10.1054/bjoc.2001.2193)
- [25] L. Coffman, C. Mooney, J. Lim, S. Bai, I. Silva, Y. Gong, K. Yang and R. J. Buckanovich, "Endothelin Receptor-A Is Required for the Recruitment of Antitumor T Cells and Modulates Chemotherapy Induction of Cancer Stem Cells," *Cancer Biology & Therapy*, Vol. 14, No. 2, 2013, pp. 184-192. [doi:10.4161/cbt.22959](https://doi.org/10.4161/cbt.22959)
- [26] A. Bagnato, F. Spinella and L. Rosanò, "Emerging Role of the Endothelin Axis in Ovarian Tumor Progression," *Endocrine-related Cancer*, Vol. 12, No. 4, 2005, pp. 761-772. [doi:10.1677/erc.1.01077](https://doi.org/10.1677/erc.1.01077)
- [27] L. Rosanò, V. Di Castro, F. Spinella, G. Tortora, M. R. Nicotra, P. G. Natali and A. Bagnato, "Combined Targeting of Endothelin A Receptor and Epidermal Growth Factor Receptor in Ovarian Cancer Shows Enhanced Antitumor Activity," *Cancer Research*, Vol. 67, No. 13, 2007, pp. 6351-6359. [doi:10.1158/0008-5472.CAN-07-0883](https://doi.org/10.1158/0008-5472.CAN-07-0883)
- [28] L. Rosanò, V. Di Castro, F. Spinella, M. R. Nicotra, P. G. Natali and A. Bagnato, "ZD4054, a Specific Antagonist of the Endothelin A Receptor, Inhibits Tumor Growth and Enhances Paclitaxel Activity in Human Ovarian Carcinoma *in Vitro* and *in Vivo*," *Molecular Cancer Therapeutics*, Vol. 6, No. 7, 2007, pp. 2003-2011. [doi:10.1158/1535-7163.MCT-07-0151](https://doi.org/10.1158/1535-7163.MCT-07-0151)
- [29] M. A. Carducci and A. Jimeno, "Targeting Bone Metastasis in Prostate Cancer with Endothelin Receptor Antagonists," *Clinical Cancer Research*, Vol. 12, No. 20, 2006, pp. 6296s-6300s. [doi:10.1158/1078-0432.CCR-06-0929](https://doi.org/10.1158/1078-0432.CCR-06-0929)
- [30] H. Q. Mai, Z. Y. Zeng, K. T. Feng, Y. L. Ye, C. Q. Zhang, W. J. Liang, X. Guo, H. Y. Mo and M. H. Hong, "Therapeutic Targeting of the Endothelin A Receptor in Human Nasopharyngeal Carcinoma," *Cancer Science*, Vol. 97, No. 12, 2006, pp. 1388-1395. [doi:10.1111/j.1349-7006.2006.00333.x](https://doi.org/10.1111/j.1349-7006.2006.00333.x)
- [31] L. Boldrini, S. Gisfredi, S. Ursino, P. Faviana, M. Lucchi, F. Melfi, A. Mussi, F. Basolo and G. Fontanini, "Expression of Endothelin-1 Is Related to Poor Prognosis in Non-Small Cell Lung Carcinoma," *European Journal of Cancer*, Vol. 41, No. 18, 2005, pp. 2828-2835. [doi:10.1016/j.ejca.2005.08.030](https://doi.org/10.1016/j.ejca.2005.08.030)
- [32] M. Felx, M. Guyot, M. Isler, R. Turcotte, J. Doyon, A. M. Khatib, S. Leclerc A. Moreau and F. Moldovan, "Endothelin-1 (ET-1) Promotes MMP-2 and MMP-9 Induction Involving the Transcription Factor NF-kappaB in Human Osteosarcoma," *Clinical Science*, Vol. 110, 2006, pp. 645-654. [doi:10.1042/CS20050286](https://doi.org/10.1042/CS20050286)
- [33] C. Chao, J. R. Carmical, K. L. Ives, T. G. Wood, J. F. Aronson, G. A. Gomez, C. D. Djukom and M. R. Helmich, "CD133+ Colon Cancer Cells Are More Interactive with the Tumor Microenvironment than CD133- Cells," *Laboratory Investigation* Vol. 92, No. 3, 2012, pp. 420-436. [doi:10.1038/labinvest.2011.185](https://doi.org/10.1038/labinvest.2011.185)
- [34] S. Zhuo, J. Chen, W. G. Wu, S. Xie, L. Zheng, X. Jiang and X. Zhu, "Quantitatively Linking Collagen Alteration and Epithelial Tumor Progression by Second Harmonic Generation Microscopy," *Applied Physics Letters*, Vol. 96, No. 21, 2010, pp. 1-3. [doi:10.1063/1.3441337](https://doi.org/10.1063/1.3441337)
- [35] P. P. Provenzano, C. T. Rueden, S. M. Trier, L. Yan, S. M. Ponik, D. R. Inman, P. J. Keely and K. W. Eliceiri, "Nonlinear Optical Imaging and Spectral-Lifetime Computational Analysis of Endogenous and Exogenous Fluorophores in Breast Cancer," *Journal of Biomedical Optics*, Vol. 13, No. 3, 2008, Article ID: 031220. [doi:10.1117/1.2940365](https://doi.org/10.1117/1.2940365)
- [36] P. P. Provenzano, K. W. Eliceiri, L. Yan, A. Ada-Nguema, M. W. Conklin, D. R. Inman and P. J. Keely, "Nonlinear Optical Imaging of Cellular Processes in Breast Cancer," *Microscopy and Microanalysis*, Vol. 14, No. 6, 2008, pp. 532-548. [doi:10.1017/S1431927608080884](https://doi.org/10.1017/S1431927608080884)

- [37] P. P. Provenzano, D. R. Inman, K. W. Eliceiri, S. M. Trier and P. J. Keely, "Contact Guidance Mediated Three-Dimensional Cell Migration Is Regulated by Rho/Rock-Dependent Matrix Reorganization," *Biophysical Journal*, Vol. 95, No. 11, 2008, pp. 5374-5384. [doi:10.1529/biophysj.108.133116](https://doi.org/10.1529/biophysj.108.133116)
- [38] E. Brown, T. McKee, E. diTomaso, A. Pluen, B. Seed, Y. Boucher and R. K. Jain, "Dynamic Imaging of Collagen and Its Modulation in Tumors *In Vivo* Using Second-Harmonic Generation," *Nature Medicine*, Vol. 9, No. 6, 2003, pp. 796-800. [doi:10.1038/nm879](https://doi.org/10.1038/nm879)
- [39] J. N. Rogart, J. Nagata, C. S. Loeser, R. D. Roorda, H. Aslanian, M. E. Robert, W. R. Zipfel and M. H. Nathanson, "Multiphoton Imaging Can Be Used for Microscopic Examination of Intact Human Gastrointestinal Mucosa *ex Vivo*," *Clinical Gastroenterology and Hepatology*, Vol. 6, No. 1, 2008, pp. 95-101. [doi:10.1016/j.cgh.2007.10.008](https://doi.org/10.1016/j.cgh.2007.10.008)
- [40] J. P. Knowles, X. Shi-Wen, S. U. Haque, A. Bhalla, M. R. Dashwood, S. Yang, I. Taylor, M. C. Winslet, D. J. Abraham and M. Loizidou, "Endothelin-1 Stimulates Colon Cancer Adjacent Fibroblasts," *International Journal of Cancer*, Vol. 130, No. 6, 2012, pp. 1264-1272. [doi:10.1002/ijc.26090](https://doi.org/10.1002/ijc.26090)