

The Role for AVE0991 (MAS-Receptor Angiotensin II (1-7) Agonist) in Reducing Cisplatin-Induced Acute Kidney Injury on C57BL/6 Mice

Chris Mathew^{1,2}

¹Faculty of Science, St Mary's Coptic Orthodox College, Coolaroo, Australia

²Institute of Health and Sports (IHES), Victoria University, Melbourne, Australia

Email: c.mathew@stmarys.vic.edu.au, chris.pittappillil@live.vu.edu.au

How to cite this paper: Mathew, C. (2023)

The Role for AVE0991 (MAS-Receptor Angiotensin II (1-7) Agonist) in Reducing Cisplatin-Induced Acute Kidney Injury on C57BL/6 Mice. *Journal of Biosciences and Medicines*, 11, 195-214.

<https://doi.org/10.4236/jbm.2023.111019>

Received: December 11, 2022

Accepted: January 28, 2023

Published: January 31, 2023

Copyright © 2023 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Acute Kidney Injury (AKI) is a condition that causes nephrotoxicity in kidney tissues due to cisplatin-induced cancer treatments. Hence, it is proposed in this review that AVE0991 (a MAS-receptor Angiotensin II (1-7) agonist) may reduce cisplatin-induced acute kidney injury by promoting nitric oxide production.

Keywords

Cisplatin, Acute Kidney Injury, AKI, Cisplatin-Induced Acute Kidney Injury, Nephrotoxicity, Renal Renin Angiotensin System, RAS, AVE0991, MAS-Receptor Angiotensin II (1-7) Agonist

1. Introduction

Kidney injury is normally classified as functional disruption [1], as the decrease in Glomerular Filtration Rate (GFR) is evident [1] [2]. The adequacy of renal blood flow is important, but understanding the kidney injury model on a microvascular and macrovascular scale is essential [1]. For instance, insufficient renal perfusion occurs pre-renally on a macrovascular level during the state of shock and intra-renally on a microvascular level, during ischemic reperfusion injury [1]. Renal blood flow may or may not correlate with glomerular perfusion, because changes in glomerular perfusion are evident in periods of preserved blood pressure via differential effects on afferent and efferent arterioles [1].

In normal human physiology, the afferent arteriolar tone is controlled via tubule-glomerular feedback from the Juxtaglomerular Apparatus (JGA) mediated

by angiotensin II, thromboxane, catecholamines, Nitric Oxide (NO) and adenosine [1]. The efferent tone is controlled via angiotensin II in response to Renin Angiotensin Aldosterone System (RAAS) [1] [3]. Studies over the past 2 decades have reported changes to afferent and efferent microvascular tone altered in the diseased state, therefore, predicting with the increase in glomerular perfusion pressure leads to kidney injury [1]. Renal perfusion is important, however, there are many factors contributing to kidney injury such as the absence of hypotension and glomerular hypoperfusion may cause tubular damage and also associate with oxidative stress and inflammation [1].

Figure 1 illustrates the vast potentiality of toxic or ischemic injury to cause tubular damage resulting in cell death via complex pathways mediated by microvascular dysfunction due to increased Reactive Oxygen Species (ROS), causing oxidative stress, inflammation and immune dysregulation ultimately resulting in Acute Kidney Injury (AKI) [1] [4]. Tumour Necrosis Factor- α (TNF- α), Interleukin (IL) 1, 6 and 8 [5], along with Transforming Growth Factor- β (TGF- β) and Toll-Like Receptors (TLRs) causes AKI [1] [2]. Pathophysiology of AKI is very complex and nephrotoxicity is one of the major side-effects of toxic-induced renal injury. It is reported in 25% of cases of severe AKI, toxic drugs affect critically ill patients [2] [6] and 19% of failures in phase 3 clinical trial is due to kidney toxicity [2] [7]. Serum Creatinine (sCr) and Blood Urea and Nitrogen (BUN) levels are increased during AKI, causing kidneys to dysregulate electrolyte balance, hence failing to excrete fluids and waste products [8] [9]. Current therapies involving fluid strategies tentatively increase intra-abdominal pressure, thereby resulting in ischemic oedema and congestion in the renal vasculature, which increases the chances of AKI [1]. Macrophages and lymphocytes are upregulated to repair the epithelial cells from injury; however, the depletion of these inflammatory responses causes prolonged kidney injury [10]. Studies have reported regulatory T cells (T-reg) play a crucial role in limiting tissue injury via RAG-1 strain [10]. When T-reg is inhibited, then worse conditions are evident that leads to AKI caused by ischemic-reperfusion injury [10].

Toxic drugs pass through the tubular organic ion transporters (CTR1 and OCT2) across the luminal membranes of the tubule [11] and are reabsorbed again into the lumen of the tubule, generating a greater risk of injury on renal epithelial cells [11] [12]. Many cancers are treated with the aid of platinum drugs, but unfortunately cause an increase in oxidative stress, which affects renal proximal tubules resulting in AKI [4] [13] [14]. Platinum drugs such as cisplatin are identified to accumulate in the proximal tubule of the nephron, where toxic effects take place [15] [16] [17] [18]. Cell death associated with AKI may also be facilitated via genetic factors such as Bcl-2 X protein (BAX) causing the cell-cycle arrest, thereby preventing cell division as the DNA could be damaged [1] [15]. Even though studies involving animals indicated the use of B cells slowing tissue repair post-injury; in humans, the adaptive immune system and immunological memory play a crucial role in the genesis of AKI and Chronic Kidney Disease (CKD) [10].




Figure 1. Toxic or ischemic injury causes tubular damage resulting in cell death via various mechanisms [1].

This review will examine the effects of cisplatin on kidney tissue and also the role of AVE0991, MAS receptor angiotensin (1-7) agonist, which may reduce cisplatin-induced AKI by promoting NO production.

2. Renal Renin Angiotensin System (RAS)

The Renin Angiotensin System (RAS) plays a crucial role in regulating blood pressure and electrolyte metabolism [19]. Several clinical studies indicate diverse involvement of RAS in various physiological and pathological processes such as cellular proliferation, inflammation and tissue fibrosis via several receptors [19]. Renin is released from the JGA cells into the blood, but some portions are filtered through glomeruli and reabsorbed by proximal tubules [3]. However synthesis of renin occurs in collecting ducts, and the release of renin is altered as regulatory production of cyclic Adenosine Monophosphate (cAMP) and cyclic Guanosine Monophosphate (cGMP) changes intracellular calcium and sodium levels [3]. Stimulation of cAMP via β -adrenergic activation releases renin from JGA cells triggering Sympathetic Nervous System (SNS), enabling prostanoids to stimulate secretion of renin via Prostaglandin E2 (EP2), Prostaglandin E4 (EP4) and Prostacyclin (IP) receptors [3]. NO and atrial natriuretic peptides are also involved in the release of renin via cGMP dependent pathway [3].

Endothelial NOS regulates vascular function and the production of NO can be enhanced by several stimuli such as [20] [21] shear stress, Acetylcholine (ACh), bradykinin and histamine via specific receptors, increases intracellular Ca^{2+} concentration, which binds to Calmodulin (CaM) activating eNOS, that facilitates an electron flux from the reductase to the oxygenase domain of the enzyme to produce NO [20] [21]. Phosphorylation of eNOS via independent Ca^{2+} pathway is equally crucial because it binds to Ca^{2+} concentration and facilitates active electron flux causing reductase to transform into oxygenase domain to produce NO [21]. Endothelial derived NO is a vasodilator as it stimulates soluble Guany-

late Cyclase (sGC) in vascular smooth muscle cells, inducing cGMP, which primarily activates protein kinase G promoting reuptake of cytosolic Ca^{2+} into the Sarcoplasm Reticulum (SR) [21]. As the Ca^{2+} exits the cell, opening of Ca^{2+} -activated K^+ channels is identified [21]. Since cGMP is induced, decreased intracellular Ca^{2+} concentration inhibits Myosin Light Chain Kinase (MLCK) phosphorylation of myosin, resulting in smooth muscle cell relaxation as illustrated in **Figure 2** [21].

NO can also affect cellular activity independently via sGC, Stimulating sarcoplasmic/Endoplasmic Reticulum Ca^{2+} ATPase (SERCA) resulting in relaxation of smooth muscle cell [21]. Dysfunction in endothelium is not only due to decreased NO production by eNOS but a mix combination of variables, that decreases the availability of L-arginine and enzyme dysfunction resulting in increased degradation. Hence, there is a need to consider various signalling pathways and changes in bioavailability while examining the role of NO in vascular wall [21].

Release of renin is inhibited by an increase in intraglomerular hydrostatic pressure, solely depended on angiotensin converting enzyme-Ang-II Type 1 Receptors (AT_1R) [3]. Several clinical studies have indicated that a high sodium intake inhibits renin secretion and angiotensin-II inhibits renin release directly via short negative feedback loop AT_1R and indirectly via the suppression of prostanoids synthesis by Cyclo-Oxygenase 2 (COX2) [3] [10].

Angiotensinogen, when broken down by renin form Ang-I and ACE, consequently converting Ang-I into Ang-II [23], as illustrated in **Figure 3** [22] and cisplatin entry through ACE via AT_1R promotes sodium and water retention, oxidative stress, vasoconstriction, cell proliferation, inflammation and fibrosis [23] [24]. Activated AT_1R is phosphorylated and binds to arrestins, which mediate rapid receptor desensitisation and internalisation [24] [25]. AT_1R couples to various soluble and receptor tyrosine kinases such as p38-Mitogen Activated Protein Kinases (MAPK), Extracellular Regulated Kinases (ERK1, ERK2) and Jun N-terminal Kinases (JNK) functioned through jak-STAT pathway, which increases ROS [24] [26] [27]. The mechanism linking AT_1R and tyrosine kinase pathway may involve de novo synthesis of cytokine and Epidermal Growth Factor (EGF) ligands to initiate signalling [24] [26]. There are many isomers for EGF receptor, however, metalloproteinase-dependent shedding of EGF ligands are crucial in Ang-II- AT_1R mediated growth, hypertrophy and proliferation of cardiac, vascular and renal cells [3] [24] [26].

RAS has demonstrated to play a crucial role in the molecular mechanism of AKI, as elevated renin activity results in ischemia, thereby increasing Ang-II via AT_1R , which decreases the vasculature of the tubular region [19] leading to endothelial dysfunction causing vasoconstriction by decreased O_2 delivery [10]. Further promoting inflammatory neutrophils and monocytes causing tubular dysfunction and affecting GFR via tubulo-glomerular feedback [10]. These imbalances indicate the relationship between arterial pressure and vascular resistance, where outer medulla of kidney is usually affected [10]. Thus, lack of NO




Figure 2. Increased intracellular Ca^{2+} via eNOS promotes relaxation in smooth muscle cells by eliciting NO production [20].




Figure 3. In kidney, angiotensinogen is broken down by ACE and renin forming ACE2 and Ang-II. An increase in Ang-II results in vasoconstriction whereas reduction of Ang-II elicits vasodilation [22].

production from blood vessels increasing vascular permeability, which causes tissue swelling [10].

The synthesis of inactive Ang-(1-9) from Ang-I and the catabolism of Ang-II to form Ang-(1-7), binding to MAS receptor (specific membrane receptor) as illustrated in **Figure 2** [22] [23], counteracts the effects of Ang-II, which is also the main functions of ACE2 [22] [23]. Ang-I is also a substrate of Neprilysin, cleaving to produce Ang-(1-7) [23]. ACE2 non classical RAS Ang-(1-7) counteracts with effects of the ACE-Ang-II-AT₁ axis [23] as specific functions include

natriuresis, reduced oxidative stress, vasodilation, anti-proliferative activity and diuresis by upregulating the concentrations of NO and prostaglandins [23] [28] [29] [30]. These processes aids in protecting the kidney from damage [23]. Ang-(1-7) is also metabolized by ACE, furthermore, accumulating evidences indicates that, ACE/ACE2 ratio regulates the production and accumulation of Ang-II and that ACE2 deficiencies leads to higher Ang-II concentrations [23] [31]. Balance between the effects of these two molecules affects the RAS and hence, the ACE/ACE2 ratio might represent the key parameter that is driving the regulation of RAS [23]. In healthy conditions, ACE2 activity increases along with ACE, whereas imbalances can develop under diseased conditions [23] [32] [33] [34]. In human kidneys, biopsies have indicated that hypertensive patients have higher ACE/ACE2 mRNA ratios [23] [34]. Battle and colleagues suggested that ACE and ACE2 are regulated via different mechanisms and that ACE/ACE2 ratio could be misleading [35], however, Pohl and colleagues indicated ACE2 is expressed along the entire renal tubular segment, while ACE is only expressed in the brush membrane border of proximal tubule and that surface expression of ACE and ACE2 differed as a function of endocytosis [36]. Thereby indicating an increase in ACE/ACE2 ratio induced via the ACE-Ang-II-AT₁ axis leads to kidney damage, whereas renal ACE/ACE2 ratio are regulated via independent mechanisms [23].

3. Cisplatin-Induced Oxidative Stress

Cisplatin enters renal cells via passive and or facilitated mechanisms [37], either via CTR1 or Organic Cation Transporter 2 (OCT2), which are cell death promoting via MAPK, ROS, p53 or cytochrome p21 pathways [37] [38]. Generation of TNF- α productions in tubular cells, stimulates inflammatory response leading to tubular cell injury or cell death [37] [39]. Mitochondrial dysfunction, inhibition of lysosomal hydrolyse, along with phospholipid damage and increase in intracellular Ca²⁺ leads to toxic build up in proximal tubular cells, influenced directly by ROS and oxidative stress [11] [13] [40] [41]. By doing so cisplatin induces Acute Tubular Necrosis (ATN), which evidently causes oxidative stress and these conditions promotes ROS formations via two mechanisms [11] [39].

Firstly, cisplatin in its highly reactive form reacts with thiol-containing molecules including Glutathione (GSH), a well-recognised cellular antioxidant [11] [13] [39]. Inactivation or depletion of GSH and related antioxidants, generates endogenous ROS accumulation within the cells [11] [13] by activating MAPK, p53 and possibly p21, leading to renal tubular cell death [11] [13] [15] [42]. Consequently, ROS stimulates fibrotic process either directly or indirectly via enhanced inflammation [11] [13] [42].

Fibrosis and inflammation might further increase ROS formation or stimulate the production of cytokines and growth factors [11] [12] [13] [40].

Secondly, cisplatin may induce mitochondrial dysfunction and increase ROS production via its disrupted respiratory chain, reducing cellular respiration and

Adenosine Triphosphate (ATP) [11] [13] [43] [44]. ROS targets lipid components of membrane, causing peroxidation, denaturing of proteins and enzyme inactivation [11] [13] [40]. A variety of cellular enzyme systems such as NADPH oxidase, xanthine oxidase, uncoupled endothelial Nitric Oxide Synthase (e-NOS) and arachidonic acid metabolizing enzymes including cytochrome P450, lipoxigenase and COX generates ROS [16] [40] [45] [46].

Uncontrolled production of mitochondrial ROS activates the opening of the mitochondrial Permeability Transition Pore (PTP) as illustrated in **Figure 4** induces necrosis in microsomes via cytochrome P450 enzymes [11] [45] [47] [48], which activates Glucose-6-Phosphate Dehydrogenase (G6PD) and hexokinase causing an increase in free radicals and depletion of antioxidants such as Superoxide Dismutase (SOD), glutathione peroxidase and catalase [40] [47]. G6PD stimulates an increase in intracellular Ca^{2+} levels, activating NADPH oxidase, stimulating ROS production by damaged mitochondria [40] [47] [49] [50] [51] [52].

Increase in Reactive Nitrogen Species (RNS), decreases cellular defences by oxidising thiol pools, by inducing structure and function of proteins and lipid peroxidation [40].

Absorption of various plasma proteins and molecules by tubular cells may cause secretion of chemotactic and inflammatory mediators in the interstitium [11] [53]. Nuclear Factor-kappa B (NF-kB) regulates DNA transcriptase and upregulates inflammatory mediators such as Damage-Associated Molecular Patterns (DAMPs) through TLR4 [11] [53]. Manganese Superoxide Dismutase (Mn-SOD), primary mitochondrial antioxidant enzyme, essential for maintaining normal cell functions [45] decreases post-cisplatin administration, which causes cell injury, indicating poor antioxidant response for cell survival [40] [45] [54].

TNF- α is dependent on ROS and activates NF-kB via p38-MAPK pathway [40] [55]. It is mediated by tumour necrosis factor receptors TNFR1 and TNFR2 via p55 and p75 respectively [40] [55]. Cisplatin administration also increases inflammatory cytokines and chemokines such as IL-1 β , IL-18, IL-6 and CX3CL1 mediated via TNFR2 [40] [49] [55]-[62]. Moreover, studies reports that cisplatin induced apoptosis in p53 deficient renal cells is via the intrinsic mitochondrial pathway resulting in cell death as p53 is observed to cause nephrotoxicity through the administration of cisplatin by inducing tubular cell apoptosis as a dependent mechanism. However, independent mechanisms may also induce apoptosis by activating BAX and releasing cytochrome c, resulting in mitochondrial damage [45] [59] [61] [63] [64].

4. Endoplasmic Reticular Stress (ERS)

Endoplasmic Reticulum (ER) performs several important functions including post-translation modification, folding, and the assembly of newly synthesised secretory and cell membrane proteins and its proper function is essential for cell survival [65] [66]. Cells tentatively functions by responding to increased ERS via




Figure 4. Decreased NO production via eNOS increases ROS resulting in oxidative stress causes mitochondrial damage [47].

Unfolded Protein Response (UPR), which aims to generate more homeostatic environment, however, can also promote cell death, if ERS is prolonged or severe [67] [68]. When ER pathway is interrupted due to an increase in oxidative stress, decreases ATP, thereby implicating inflammation in its pathogenesis [69]. ERS is a form of cellular stress caused by excessive protein accumulation at the ER [69]. Three major ERS pathways are by the accumulation of Pancreatic ER Kinase (PERK), eukaryotic initiation factor 2 α (eIF2 α); or Inositol-Requirement Enzyme 1 (IRE1), X-box binding protein 1 (XBP1); or by Activating Transcription Factor (ATF6) such as C/EBP Homologous Protein (CHOP), resulting in growth arrest via the activation of c-jun-NH₂-terminal kinase signalling causing increased production of Transforming Growth Factor- β 1 (TGF- β 1), which enhances proliferation, apoptosis and DNA damage [44] [65] [66].

CHOP induces ER stress resulting in apoptosis [65] [67] [70]. Therefore, unattended ERS could result in cell death, inflammation and excessive oxidant production [69] [71]. Even though, ERS could be a result of atherosclerosis, diabetes and neurodegenerative diseases; AKI generated via hypoxia, free radical generation and decreased amounts of glucose and amino acids are few of the causes of ERS [69].

Cisplatin induces caspase 3 activation, involving Ca²⁺ and Ca²⁺ dependent calpain protease [73], which inactivates calpain-dependent ER-specific caspase 12 and upregulates Heat Shock Proteins (HSPs) subfamily; composed of four isomers HSC70-inducible isoform, HSP72, mHSP75 and GRP78 enables to restore cellular homeostasis [21] [74]. Glucose Regulated Protein 78 (GRP78) upregulation are due to the administration of platinum drugs [73] [75]. GRP78 acts in accordance with Ca²⁺ dependent pathway, which acts as the main regulator of ER function [67] [73]. Cisplatin can also induce apoptosis in enucleated mouse kidney proximal tubular cells as indicated by Yu *et al.* 2007 causing cell death via cytoplasmic signalling, which acts independently to nucleus but regulated by cyclin-dependent kinase 2 (cdk2)-E2F1 pathway as the cytoplasmic lo-

cations are in ER and Golgi complex as illustrated in **Figure 5** [72] [73] [75] [76]. Evidently, upregulation of GRP78 and CHOP via caspase 4, found in damaged tubular cells post toxic kidney injury indicates that cisplatin induces apoptosis through ERS pathway [73] [77] [78]. HSP72 and GRP78 are expressed in response to cellular stress and its induction can be as great as 15% of the total protein [74]. Studies on rat showed proteinuria in early stages, and further examinations showed accumulation of transgene substances in the ER inducing ERS and subsequent kidney injury [67]. Given that HSP72, GRP78 and CHOP are induced in renal tubules during AKI and that proximal detachment is projected to the urinary space; levels of these proteins can serve as a possible early biomarker to detect cisplatin-induced AKI and will be studied in this thesis [65] [66] [70] [74] [79] [80].

5. Cisplatin-Induced Nephrotoxicity

Cisplatin induces direct tubular epithelial cell toxicity as well as reduction in renal blood flow as a result of endothelial dysfunction and vasoconstriction [24] [27] [81]. Many cancer patients show a decrease in appetite, lowering blood pressure, consequently deteriorates and form cisplatin-induced nephrotoxicity despite routine administration of Intravenous (IV) fluids [50] [81] [82]. Patients with low blood pressure are expected to have delayed urinary excretion of cisplatin due to decrease in GFR [8] [81] [83]. Komaki reported that, patients are at high risk of AKI during intercurrent illness such as volume depletion because, the contraction of efferent arterioles is inhibited due to oxidative stress [81] [84]. RAS can also be activated by volume depletion from sodium wasting or polyuria and vascular contractions post-cisplatin administration [81]. Increase in plasma renin activity and plasma aldosterone concentrations were also identified with cisplatin administration [81] [85] [86]. Malignant transformations are associated with at least six acquired, functional capabilities: sustained angiogenesis, evasion of apoptosis, self-sufficiency in growth signals, insensitivity to anti-growth signals, tissue invasion and metastasis and limitless replicative potential [24] [37]. RAS regulates all of these capabilities; however, most prominent effects are on angiogenesis, invasion, pro-survival and proliferation [24]. Much of the increased metastasis and invasion that is associated with an activated RAS is likely to be a direct consequence of angiogenesis [24].

6. Current Therapeutic Strategies

Hydration, electrolyte replacement with saline and avoiding other nephrotoxic drugs has been the most supportive measures in reducing severe AKI [39]. GFR must be assessed routinely as well as hydration, which should have commenced prior to the treatment and continued for at least 3 days post treatment [39]. Adequacy for hydration solely depends on urine output, which needs to be maintained 3 - 4 L/day [39]. Magnesium (Mg^{2+}) also has been reported to decrease in cisplatin administration and several patients are advised to routinely




Figure 5. ER stress induces UPR resulting in apoptosis through various mechanisms [72].

assess their serum Mg^{2+} and needs to be replaced adequately [39]. However, therapeutic strategies are required to eliminate AKI completely and several researchers have tried to implement many experimental studies on animals mostly for the prevention of cisplatin-induced AKI [39] because cisplatin based chemotherapy induces nephrotoxicity in approximately 25% - 30% of patients treated for solid cancers like small cell lung carcinoma and prostate cancer [87]. Accumulative research studies indicate that autophagy from cytoplasm, organelles or membrane engulfed by double-membrane structure targeted for destruction in lysosomes, could act as a protective measure against cisplatin-induced cell death; and activation of mTOR signalling may regulate autophagy that affects tubular cell death via various mechanisms [87].

In a recent study, metformin was found as a potential protective agent by targeting cisplatin-induced tubular cell death in cultured NRK-52E cells and AKI in mice [87]. It functions by stimulating Adenosine Monophosphate Kinase alpha ($AMPK\alpha$) and thus causing induction of autophagy [87]. However, this study was interesting, there was also a down regulation of $AMPK\alpha$ phosphorylation that led to diminished metformin promoted cell survival [87] and this study was done on cultured cell whereas it didn't express the effects on in-vivo mice model [87].

ACE2 upregulation as a treatment was based on reduced levels of ACE2 in patients with renal disease and in experimental models [28] [88] [89]. Studies in mice showed significant improvement in kidney function by lowering urinary albumin excretion rate and improved creatinine clearance [27] [28]. This improvement was associated with less severe renal fibrosis, which attributes to inhibition of excess canonical and non-canonical TGF- β signaling by ACE2 [28]. Pro-inflammatory cytokine expression and macrophage infiltration were attenuated by renal ACE2 associated with a decrease in MAPK signaling [28]. RAS activation has an important role in the pathological processes that leads to kidney injury by regulating body fluid balance and blood pressure, and inhibition of

this pathway improves renal outcomes [28] [90]. Expression of ACE2 is altered in human kidney disease [29]. Neo-expression of ACE2 is found in glomerular and peritubular capillary endothelium in various renal disorders and in renal transplants [29]. Further studies should elucidate the pathophysiological significance of these changes in ACE2 expression and explore its role as a possible protective mechanism [29]. The significance of reduction in renal ACE coupled with an increase in ACE2 proteins needs to be clarified, but it seems logical to propose such combination could attenuate Ang-II and could exert protective effect against necrosis at early stages of AKI and the increase in ACE2 should further prevent Ang-II accumulation by favoring conversion of Ang-I to Ang-(1-9) and Ang-II to Ang-(1-7) [30] [31] [91] [92].

There were several studies that linked many mechanisms that could reduce inflammation and tumour cell death that results in necrosis; however, there was not enough studies conducted on endothelial cell pathway, which could potentially increase NO production, which could cause vasodilation via cGMP pathway. Therefore, the study proposal of using AVE0991, of Ang-(1-7) mimetic could prove as a protective agent against cisplatin-induced AKI.

7. Relationship between AVE0991 and RAS

AVE0991 (5-formyl-4-methoxy-2-phenyl-1[[4-[2-ethylaminocarbonylsulfonamido]-5-isobutyl-3-thienyl]-phenyl]-methyl]-imidazole [93] [94], is a derivative of imidazole [93] and is soluble in alkaline water solutions [93]. AVE0991 is an orally active and physiologically well tolerated compound, which mimics the action of Ang-(1-7) in several tissues [93] [95]. It is also reported that, AVE0991 has a longer biological half-life and evidently more stable compared to Ang-(1-7) [96]. AVE0991 helps to release Nitric Oxide (NO) and to a lesser extent superoxide (O_2^-) in endothelial cells [93] and it is also capable of releasing 5 times as much NO as Ang-(1-7) [93] [94]. Generation of O_2^- and consequent fast reactions with O_2^- and NO forms peroxy-nitrite and other cytotoxic radicals, while the release of NO associated with low concomitant production of O_2^- might contribute to the preservation of the vasculature [93] [97] [98].

Studies on AVE0991 have shown beneficial effects in atherosclerosis and hypertension [21]. Prolonged treatment using Ang-(1-7) mimetic has led to an increase in glucose uptake and insulin resistance in rats [21].

Pinheiro reported that AVE0991 increases water reabsorption, along with a decrease in urinary volume in control mice with a rise in urine osmolarity [101]. Ang-(1-7) increases osmotic water permeability in distal nephron [101]. It is also reported of inducing concentration-dependent vasodilator effects on aortic rings and this mechanism was evidently present in intact endothelium [93] by stimulating endothelial function facilitated by release of NO [93]. Another study reported that AVE0991 attenuated inflammatory effects on ACE2-Ang-(1-7) MAS axis by attenuating an elevation of sCr, decreasing neutrophil influx in both kidney and lungs [102]. Decrease in neutrophil accumulation reduced chemokine production as identified with low circulating levels of chemokine in AVE

treated animals [102]. Administration of AVE0991 improves renal damage by significantly decreasing index of renal injury on glomerular and tubular regions [102] by attenuating glomerular sclerosis in experimental glomerulonephritis and over-expression of ACE2 in diabetic nephropathy may also be due to an increase in Ang-(1-7) levels [102]. In correlation with MAS receptor, mRNA levels are increased whereas AVE0991 may exhibit beneficial effects through various mechanisms by interacting with ACE2 by inhibiting renal activities of this enzyme [102]. An increase in MAS receptor, could represent a compensatory response to renal damage as compared to endogenous regulatory mechanisms [102]. Many studies have indicated the potentiality of MAS receptor and putative drug AVE0991 is found to stimulate ACE2-Ang-(1-7) MAS axis by promoting anti-inflammatory responses as illustrated in **Figure 6** [102].

Yuedong and colleagues indicated the potentiality of AVE0991 to reduce oxidative stress by demonstrating it on aortic banding mice [103]. Their results indicated that, vehicle treated mice had a higher expression of NADPH Oxidase 2 (NOX2) and NOX4 proteins, compared with sham-operated group [103]. Elevation in NADPH oxidase generates endothelial ROS that leads to proliferation [104]. When, AVE0991 was administered, it was found to suppress the increase of NOX2 and NOX4 mRNA and protein levels indicating AVE0991 reduces oxidative stress in mice [103]. By reducing oxidative stress, indicates fewer ROS in endothelial cell, which may reduce ERS and fewer inflammatory responses that could preserve the vascular tone of the kidney tissues, as illustrated in **Figure 6**.

8. Research Rationale

Treatment with AVE0991 is found to reduce inflammation, and glomerular and tubulointerstitial damage in ischemic/reperfusion AKI-induced animals (via occlusion of the renal pedicle), thereby reducing acute and chronic kidney injury,




Figure 6. AVE0991 stimulates the MASR to elicit the production of NO resulting in vasodilation [99] [100], whereas cisplatin enters renal epithelial cells via the OCT2 or CTR1, induces epithelial cell death by causing DNA damage, hence, increasing the production of ROS, ER stress and TNF- α resulting in vasoconstriction [46].

hence, it is proposed that AVE0991 via stimulation of the ACE2-Ang-(1-7) MAS axis on kidney tissues can counteract the effects of cisplatin-induced AKI (as illustrated in **Figure 6**) such as anti-oxidative stress, decreased production of ROS and decreased inflammatory responses and may preserve the vasculature of kidney tissues, which may restore reno-protective mechanisms [46] [93] [96] [105].

9. Conclusion

Cisplatin is the most widely used drug in treating cancers, but can also cause many side effects such as AKI. Although this review consolidates many research studies and current therapies by illustrating the mechanisms, it showcases the pathway to reduce cisplatin-induced AKI by understanding the role of AVE0991 in decreasing ROS and inflammatory responses in preserving the kidney tissues, which may counteract the effects and provide better reno-protective mechanisms.

Conflicts of Interest

The author declares no conflicts of interest regarding the publication of this paper.

References

- [1] Chen, H. and Busse, L.W. (2017) Novel Therapies for Acute Kidney Injury. *Kidney International Reports*, **2**, 785-799. <https://doi.org/10.1016/j.ekir.2017.06.020>
- [2] Qu, Y., et al. (2018) A Nephron Model for Study of Drug-Induced Acute Kidney Injury and Assessment of Drug-Induced Nephrotoxicity. *Biomaterials*, **155**, 41-53. <https://doi.org/10.1016/j.biomaterials.2017.11.010>
- [3] Szczepanska-Sadowska, E., Czarzasta, K. and Cudnoch-Jedrzejewska, A. (2018) Dysregulation of the Renin-Angiotensin System and the Vasopressinergic System Interactions in Cardiovascular Disorders. *Current Hypertension Reports*, **20**, Article No. 19. <https://doi.org/10.1007/s11906-018-0823-9>
- [4] Chen, Y., et al. (2013) Assessment of Cisplatin-Induced Kidney Injury Using an Integrated Rodent Platform. *Toxicology and Applied Pharmacology*, **268**, 352-361. <https://doi.org/10.1016/j.taap.2013.01.032>
- [5] Mitazaki, S., et al. (2013) Interleukin-6 Modulates Oxidative Stress Produced during the Development of Cisplatin Nephrotoxicity. *Life Sciences*, **92**, 694-700. <https://doi.org/10.1016/j.lfs.2013.01.026>
- [6] Rameshkumar, R., et al. (2017) Histopathological Changes in Septic Acute Kidney Injury in Critically Ill Children: A Cohort of Post-Mortem Renal Biopsies. *Clinical and Experimental Nephrology*, **21**, 1075-1082. <https://doi.org/10.1007/s10157-016-1343-z>
- [7] Naughton, C. (2008) Drug-Induced Nephrotoxicity. *American Academy of Family Physicians*, **78**, 743-750.
- [8] Santos, N.A., et al. (2007) Cisplatin-Induced Nephrotoxicity Is Associated with Oxidative Stress, Redox State Unbalance, Impairment of Energetic Metabolism and Apoptosis in Rat Kidney Mitochondria. *Archives of Toxicology*, **81**, 495-504. <https://doi.org/10.1007/s00204-006-0173-2>
- [9] Sung, M.J., et al. (2008) Genistein Protects the Kidney from Cisplatin-Induced In-

- jury. *Kidney International*, **74**, 1538-1547. <https://doi.org/10.1038/ki.2008.409>
- [10] Ferenbach, D.A. and Bonventre, J.V. (2016) Acute Kidney Injury and Chronic Kidney Disease: From the Laboratory to the Clinic. *Néphrologie & Thérapeutique*, **12**, S41-S48. <https://doi.org/10.1016/j.nephro.2016.02.005>
- [11] Hosohata, K. (2016) Role of Oxidative Stress in Drug-Induced Kidney Injury. *International Journal of Molecular Sciences*, **17**, Article 1826. <https://doi.org/10.3390/ijms17111826>
- [12] Hosohata, K., *et al.* (2016) Early Prediction of Cisplatin-Induced Nephrotoxicity by Urinary Vanin-1 in Patients with Urothelial Carcinoma. *Toxicology*, **359-360**, 71-75. <https://doi.org/10.1016/j.tox.2016.06.011>
- [13] Dasari, S. and Tchounwou, P.B. (2014) Cisplatin in Cancer Therapy: Molecular Mechanisms of Action. *European Journal of Pharmacology*, **740**, 364-378. <https://doi.org/10.1016/j.ejphar.2014.07.025>
- [14] Dehghani, A., Saberi, S. and Nematbakhsh, M. (2016) Cisplatin-Induced Nephrotoxicity Alters Blood Pressure Response to Angiotensin II Administration in Rats. *Advanced Biomedical Research*, **5**, Article No. 53. <https://doi.org/10.4103/2277-9175.178797>
- [15] Xu, Y., *et al.* (2015) A Role for Tubular Necroptosis in Cisplatin-Induced AKI. *Journal of the American Society of Nephrology*, **26**, 2647-2658. <https://doi.org/10.1681/ASN.2014080741>
- [16] Meng, X.M., *et al.* (2018) NADPH Oxidase 4 Promotes Cisplatin-Induced Acute Kidney Injury via ROS-Mediated Programmed Cell Death and Inflammation. *Laboratory Investigation*, **98**, 63-78. <https://doi.org/10.1038/labinvest.2017.120>
- [17] Portilla, D., *et al.* (2006) Metabolomic Study of Cisplatin-Induced Nephrotoxicity. *Kidney International*, **69**, 2194-2204. <https://doi.org/10.1038/sj.ki.5000433>
- [18] Stojanovska, V., *et al.* (2017) Neurotoxicity Associated with Platinum-Based Anti-Cancer Agents: What Are the Implications of Copper Transporters? *Current Medicinal Chemistry*, **24**, 1520-1536. <https://doi.org/10.2174/0929867324666170112095428>
- [19] Ono, M., *et al.* (2015) Role of Intrarenal (Pro)renin Receptor in Ischemic Acute Kidney Injury in Rats. *Clinical and Experimental Nephrology*, **19**, 185-196. <https://doi.org/10.1007/s10157-014-0979-9>
- [20] Zhao, Y., Vanhoutte, P.M. and Leung, S.W. (2015) Vascular Nitric Oxide: Beyond eNOS. *Journal of Pharmacological Sciences*, **129**, 83-94. <https://doi.org/10.1016/j.jphs.2015.09.002>
- [21] Karpe, P.A. and Tikoo, K. (2014) Heat Shock Prevents Insulin Resistance-Induced Vascular Complications by Augmenting Angiotensin-(1-7) Signaling. *Diabetes*, **63**, 1124-1139. <https://doi.org/10.2337/db13-1267>
- [22] Patel, V.B., *et al.* (2016) Role of the ACE2/Angiotensin 1-7 Axis of the Renin-Angiotensin System in Heart Failure. *Circulation Research*, **118**, 1313-1326. <https://doi.org/10.1161/CIRCRESAHA.116.307708>
- [23] Mizuiri, S. and Ohashi, Y. (2015) ACE and ACE2 in Kidney Disease. *World Journal of Nephrology*, **4**, 74-82. <https://doi.org/10.5527/wjn.v4.i1.74>
- [24] George, A.J., Thomas, W.G. and Hannan, R.D. (2010) The Renin-Angiotensin System and Cancer: Old Dog, New Tricks. *Nature Reviews Cancer*, **10**, 745-759. <https://doi.org/10.1038/nrc2945>
- [25] Kaschina, E., Namsolleck, P. and Unger, T. (2017) AT2 Receptors in Cardiovascular and Renal Diseases. *Pharmacological Research*, **125**, 39-47.

- <https://doi.org/10.1016/j.phrs.2017.07.008>
- [26] Dugbartey, G.J., Peppone, L.J. and de Graaf, I.A. (2016) An Integrative View of Cisplatin-Induced Renal and Cardiac Toxicities: Molecular Mechanisms, Current Treatment Challenges and Potential Protective Measures. *Toxicology*, **371**, 58-66. <https://doi.org/10.1016/j.tox.2016.10.001>
- [27] Chang, C., et al. (2017) Pharmacogenomic Variants May Influence the Urinary Excretion of Novel Kidney Injury Biomarkers in Patients Receiving Cisplatin. *International Journal of Molecular Sciences*, **18**, Article 1333. <https://doi.org/10.3390/ijms18071333>
- [28] Lieben, L. (2017) Alport Syndrome: ACE2 Administration Slows Kidney Damage. *Nature Reviews Nephrology*, **13**, Article No. 261. <https://doi.org/10.1038/nrneph.2017.36>
- [29] Lely, A.T., et al. (2004) Renal ACE2 Expression in Human Kidney Disease. *The Journal of Pathology*, **204**, 587-593. <https://doi.org/10.1002/path.1670>
- [30] Xia, H. and Lazartigues, E. (2010) Angiotensin-Converting Enzyme 2: Central Regulator for Cardiovascular Function. *Current Hypertension Reports*, **12**, 170-175. <https://doi.org/10.1007/s11906-010-0105-7>
- [31] Ye, M., et al. (2004) Increased ACE 2 and Decreased ACE Protein in Renal Tubules from Diabetic Mice: A Renoprotective Combination? *Hypertension*, **43**, 1120-1125. <https://doi.org/10.1161/01.HYP.0000126192.27644.76>
- [32] Iwai, M. and Horiuchi, M. (2009) Devil and Angel in the Renin-Angiotensin System: ACE-Angiotensin II-AT1 Receptor Axis vs. ACE2-Angiotensin-(1-7)-Mas Receptor Axis. *Hypertension Research*, **32**, 533-536. <https://doi.org/10.1038/hr.2009.74>
- [33] Ma, C., et al. (2014) Relationship between Renal Injury and the Antagonistic Roles of Angiotensin-Converting Enzyme (ACE) and ACE2. *Genetics and Molecular Research*, **13**, 2333-2342. <https://doi.org/10.4238/2014.April.3.5>
- [34] Wakahara, S., et al. (2007) Synergistic Expression of Angiotensin-Converting Enzyme (ACE) and ACE2 in Human Renal Tissue and Confounding Effects of Hypertension on the ACE to ACE2 Ratio. *Endocrinology*, **148**, 2453-2457. <https://doi.org/10.1210/en.2006-1287>
- [35] Batlle, D., et al. (2012) Angiotensin-Converting Enzyme 2: Enhancing the Degradation of Angiotensin II as a Potential Therapy for Diabetic Nephropathy. *Kidney International*, **81**, 520-528. <https://doi.org/10.1038/ki.2011.381>
- [36] Pohl, M., et al. (2010) Intrarenal Renin Angiotensin System Revisited: Role of Megalin-Dependent Endocytosis along the Proximal Nephron. *Journal of Biological Chemistry*, **285**, 41935-41946. <https://doi.org/10.1074/jbc.M110.150284>
- [37] Pabla, N. and Dong, Z. (2008) Cisplatin Nephrotoxicity: Mechanisms and Renoprotective Strategies. *Kidney International*, **73**, 994-1007. <https://doi.org/10.1038/sj.ki.5002786>
- [38] Harrach, S. and Ciarimboli, G. (2015) Role of Transporters in the Distribution of Platinum-Based Drugs. *Frontiers in Pharmacology*, **6**, Article 85. <https://doi.org/10.3389/fphar.2015.00085>
- [39] Ozkok, A. and Edelstein, C.L. (2014) Pathophysiology of Cisplatin-Induced Acute Kidney Injury. *BioMed Research International*, **2014**, Article ID: 967826. <https://doi.org/10.1155/2014/967826>
- [40] Oh, G.S., et al. (2014) Cisplatin-Induced Kidney Dysfunction and Perspectives on Improving Treatment Strategies. *Electrolytes & Blood Pressure*, **12**, 55-65.

- <https://doi.org/10.5049/EBP.2014.12.2.55>
- [41] Han, W.K., *et al.* (2002) Kidney Injury Molecule-1 (KIM-1): A Novel Biomarker for Human Renal Proximal Tubule Injury. *Kidney International*, **62**, 237-244. <https://doi.org/10.1046/j.1523-1755.2002.00433.x>
- [42] Tian, L., *et al.* (2017) Kidney Injury Molecule-1 Is Elevated in Nephropathy and Mediates Macrophage Activation via the Mapk Signalling Pathway. *Cellular Physiology and Biochemistry*, **41**, 769-783. <https://doi.org/10.1159/000458737>
- [43] Kruidering, M., *et al.* (1997) Cisplatin-Induced Nephrotoxicity in Porcine Proximal Tubular Cells: Mitochondrial Dysfunction by Inhibition of Complexes I to IV of the Respiratory Chain. *Journal of Pharmacology and Experimental Therapeutics*, **280**, 638-649.
- [44] Chou, Y.H., Huang, T.M. and Chu, T.S. (2017) Novel Insights into Acute Kidney Injury-Chronic Kidney Disease Continuum and the Role of Renin-Angiotensin System. *Journal of the Formosan Medical Association*, **116**, 652-659. <https://doi.org/10.1016/j.jfma.2017.04.026>
- [45] Yuan, Y., *et al.* (2015) P53 Contributes to Cisplatin Induced Renal Oxidative Damage via Regulating P66shc and MnSOD. *Cellular Physiology and Biochemistry*, **37**, 1240-1256. <https://doi.org/10.1159/000430247>
- [46] Miller, R.P., *et al.* (2010) Mechanisms of Cisplatin Nephrotoxicity. *Toxins (Basel)*, **2**, 2490-2518. <https://doi.org/10.3390/toxins2112490>
- [47] Chang, J.C., *et al.* (2010) Regulatory Role of Mitochondria in Oxidative Stress and Atherosclerosis. *World Journal of Cardiology*, **2**, 150-159. <https://doi.org/10.4330/wjc.v2.i6.150>
- [48] Qian, W., *et al.* (2005) Mitochondrial Density Determines the Cellular Sensitivity to Cisplatin-Induced Cell Death. *American Journal of Physiology-Cell Physiology*, **289**, C1466-C1475. <https://doi.org/10.1152/ajpcell.00265.2005>
- [49] Bhat, Z.Y., *et al.* (2015) Understanding the Risk Factors and Long-Term Consequences of Cisplatin-Associated Acute Kidney Injury: An Observational Cohort Study. *PLOS ONE*, **10**, e0142225. <https://doi.org/10.1371/journal.pone.0142225>
- [50] Hanigan, M.H. and Devarajan, P. (2008) Cisplatin Nephrotoxicity: Molecular Mechanisms. National Institute of Health, Bethesda, 47-61.
- [51] Mohamed, F., *et al.* (2017) Nephrotoxicity-Induced Proteinuria Increases Biomarker Diagnostic Thresholds in Acute Kidney Injury. *BMC Nephrology*, **18**, Article No. 122. <https://doi.org/10.1186/s12882-017-0532-7>
- [52] Husi, H. and Human, C. (2015) Molecular Determinants of Acute Kidney Injury. *Injury and Violence*, **75**, 75-86.
- [53] Ozkok, A., *et al.* (2016) NF-kappaB Transcriptional Inhibition Ameliorates Cisplatin-Induced Acute Kidney Injury (AKI). *Toxicology Letters*, **240**, 105-113. <https://doi.org/10.1016/j.toxlet.2015.10.028>
- [54] Rybalka, E., *et al.* (2017) Chemotherapeutic Agents Induce Mitochondrial Superoxide Production and Toxicity but Do Not Alter Respiration in Skeletal Muscle *in Vitro*. *Mitochondrion*, **42**, 33-49. <https://doi.org/10.1016/j.mito.2017.10.010>
- [55] Zhang, B., *et al.* (2007) Cisplatin-Induced Nephrotoxicity Is Mediated by Tumor Necrosis Factor-Alpha Produced by Renal Parenchymal Cells. *Kidney International*, **72**, 37-44. <https://doi.org/10.1038/sj.ki.5002242>
- [56] Faubel, S., *et al.* (2007) Cisplatin-Induced Acute Renal Failure Is Associated with an Increase in the Cytokines Interleukin (IL)-1beta, IL-18, IL-6, and Neutrophil Infiltration in the Kidney. *Journal of Pharmacology and Experimental Therapeutics*, **322**,

- 8-15. <https://doi.org/10.1124/jpet.107.119792>
- [57] Lu, L.H., *et al.* (2008) Increased Macrophage Infiltration and Fractalkine Expression in Cisplatin-Induced Acute Renal Failure in Mice. *Journal of Pharmacology and Experimental Therapeutics*, **324**, 111-117. <https://doi.org/10.1124/jpet.107.130161>
- [58] Nozaki, Y., *et al.* (2012) Signaling through the Interleukin-18 Receptor Alpha Attenuates Inflammation in Cisplatin-Induced Acute Kidney Injury. *Kidney International*, **82**, 892-902. <https://doi.org/10.1038/ki.2012.226>
- [59] Tsuruya, K., *et al.* (2003) Direct Involvement of the Receptor-Mediated Apoptotic Pathways in Cisplatin-Induced Renal Tubular Cell Death. *Kidney International*, **63**, 72-82. <https://doi.org/10.1046/j.1523-1755.2003.00709.x>
- [60] Liang, H., *et al.* (2016) CXCL16 Regulates Cisplatin-Induced Acute Kidney Injury. *Oncotarget*, **7**, 31652-31662. <https://doi.org/10.18632/oncotarget.9386>
- [61] Bonavia, A. and Singbartl, K. (2017) A Review of the Role of Immune Cells in Acute Kidney Injury. *Pediatric Nephrology*, **33**, 1629-1639. <https://doi.org/10.1007/s00467-017-3774-5>
- [62] Bunel, V., *et al.* (2017) Early Detection of Acute Cisplatin Nephrotoxicity: Interest of Urinary Monitoring of Proximal Tubular Biomarkers. *Clinical Kidney Journal*, **10**, 639-647. <https://doi.org/10.1093/ckj/sfx007>
- [63] Wang, H.H. (2015) Oxidative Stress and Potential Renal Damage in Neonates. *Pediatrics & Neonatology*, **56**, 209-210. <https://doi.org/10.1016/j.pedneo.2015.03.003>
- [64] Jiang, M., *et al.* (2009) Cisplatin-Induced Apoptosis in p53-Deficient Renal Cells via the Intrinsic Mitochondrial Pathway. *American Journal of Physiology-Renal Physiology*, **296**, F983-F993. <https://doi.org/10.1152/ajprenal.90579.2008>
- [65] Endo, M., *et al.* (2006) C/EBP Homologous Protein (CHOP) Is Crucial for the Induction of Caspase-11 and the Pathogenesis of Lipopolysaccharide-Induced Inflammation. *The Journal of Immunology*, **176**, 6245-6253. <https://doi.org/10.4049/jimmunol.176.10.6245>
- [66] Chikka, M.R., *et al.* (2013) C/EBP Homologous Protein (CHOP) Contributes to Suppression of Metabolic Genes during Endoplasmic Reticulum Stress in the Liver. *Journal of Biological Chemistry*, **288**, 4405-4415. <https://doi.org/10.1074/jbc.M112.432344>
- [67] Inagi, R. (2009) Endoplasmic Reticulum Stress in the Kidney as a Novel Mediator of Kidney Injury. *Nephron Experimental Nephrology*, **112**, e1-e9. <https://doi.org/10.1159/000210573>
- [68] Rabik, C.A., *et al.* (2008) Enhancement of Cisplatin [*cis*-diammine dichloroplatinum (II)] Cytotoxicity by *O*-Benzylguanine Involves Endoplasmic Reticulum Stress. *Journal of Pharmacology and Experimental Therapeutics*, **327**, 442-452. <https://doi.org/10.1124/jpet.108.141291>
- [69] Huang, L., *et al.* (2011) Increased Susceptibility to Acute Kidney Injury Due to Endoplasmic Reticulum Stress in Mice Lacking Tumor Necrosis Factor-Alpha and Its Receptor 1. *Kidney International*, **79**, 613-623. <https://doi.org/10.1038/ki.2010.469>
- [70] Oyadomari, S. and Mori, M. (2004) Roles of CHOP/GADD153 in Endoplasmic Reticulum Stress. *Cell Death and Differentiation*, **11**, 381-389. <https://doi.org/10.1038/sj.cdd.4401373>
- [71] Xu, Y., *et al.* (2016) Endoplasmic Reticulum Stress and Its Effects on Renal Tubular Cells Apoptosis in Ischemic Acute Kidney Injury. *Renal Failure*, **38**, 831-837. <https://doi.org/10.3109/0886022X.2016.1160724>
- [72] Vandewynckel, Y.-P., *et al.* (2013) The Paradox of the Unfolded Protein Response

- in Cancer. *Anticancer Research*, **33**, 4683-4694.
- [73] Xu, Y., Wang, C. and Li, Z. (2014) A New Strategy of Promoting Cisplatin Chemotherapeutic Efficiency by Targeting Endoplasmic Reticulum Stress. *Molecular and Clinical Oncology*, **2**, 3-7. <https://doi.org/10.3892/mco.2013.202>
- [74] Barrera-Chimal, J., *et al.* (2011) Hsp72 Is an Early and Sensitive Biomarker to Detect Acute Kidney Injury. *EMBO Molecular Medicine*, **3**, 5-20. <https://doi.org/10.1002/emmm.201000105>
- [75] Yu, F., *et al.* (2007) Involvement of the CDK2-E2F1 Pathway in Cisplatin Cytotoxicity *in Vitro* and *in Vivo*. *American Journal of Physiology-Renal Physiology*, **293**, F52-F59. <https://doi.org/10.1152/ajprenal.00119.2007>
- [76] Mandic, A., *et al.* (2003) Cisplatin Induces Endoplasmic Reticulum Stress and Nucleus-Independent Apoptotic Signaling. *Journal of Biological Chemistry*, **278**, 9100-9106. <https://doi.org/10.1074/jbc.M210284200>
- [77] Yu, H., *et al.* (2011) p62/SQSTM1 Involved in Cisplatin Resistance in Human Ovarian Cancer Cells by Clearing Ubiquitinated Proteins. *European Journal of Cancer*, **47**, 1585-1594. <https://doi.org/10.1016/j.ejca.2011.01.019>
- [78] Xu, Y., *et al.* (2012) Inhibition of Autophagy Enhances Cisplatin Cytotoxicity through Endoplasmic Reticulum Stress in Human Cervical Cancer Cells. *Cancer Letters*, **314**, 232-243. <https://doi.org/10.1016/j.canlet.2011.09.034>
- [79] Niu, Z., *et al.* (2015) Elevated GRP78 Expression Is Associated with Poor Prognosis in Patients with Pancreatic Cancer. *Scientific Reports*, **5**, Article No. 16067. <https://doi.org/10.1038/srep16067>
- [80] Zahedi, K., *et al.* (2017) Activation of Endoplasmic Reticulum Stress Response by Enhanced Polyamine Catabolism Is Important in the Mediation of Cisplatin-Induced Acute Kidney Injury. *PLOS ONE*, **12**, e0184570. <https://doi.org/10.1371/journal.pone.0184570>
- [81] Komaki, K., *et al.* (2017) Lower Blood Pressure and Risk of Cisplatin Nephrotoxicity: A Retrospective Cohort Study. *BMC Cancer*, **17**, Article No. 144. <https://doi.org/10.1186/s12885-017-3135-6>
- [82] Glezerman, I.G. and Jaimes, E.A. (2016) Chemotherapy and Kidney Injury. *American Society of Nephrology*, Washington DC, 1-10.
- [83] Peres, L.A. and Da Cunha Jr., A.D. (2013) Acute Nephrotoxicity of Cisplatin: Molecular Mechanisms. *Journal Brasileiro de Nefrologia*, **35**, 332-340. <https://doi.org/10.5935/0101-2800.20130052>
- [84] Astashkina, A.I., *et al.* (2012) A 3-D Organoid Kidney Culture Model Engineered for High-Throughput Nephrotoxicity Assays. *Biomaterials*, **33**, 4700-4711. <https://doi.org/10.1016/j.biomaterials.2012.02.063>
- [85] Kanki, M., *et al.* (2014) Identification of Urinary miRNA Biomarkers for Detecting Cisplatin-Induced Proximal Tubular Injury in Rats. *Toxicology*, **324**, 158-168. <https://doi.org/10.1016/j.tox.2014.05.004>
- [86] Kidera, Y., *et al.* (2014) Risk Factors for Cisplatin-Induced Nephrotoxicity and Potential of Magnesium Supplementation for Renal Protection. *PLOS ONE*, **9**, e101902. <https://doi.org/10.1371/journal.pone.0101902>
- [87] Li, J., *et al.* (2016) Metformin Protects against Cisplatin-Induced Tubular Cell Apoptosis and Acute Kidney Injury via AMPK α -Regulated Autophagy Induction. *Scientific Reports*, **6**, Article No. 23975. <https://doi.org/10.1038/srep23975>
- [88] Cheng, Q., *et al.* (2016) ACE2 Overexpression Inhibits Acquired Platinum Resistance-Induced Tumor Angiogenesis in NSCLC. *Oncology Reports*, **36**, 1403-1410.

- <https://doi.org/10.3892/or.2016.4967>
- [89] Cheng, T.C., *et al.* (2018) Nephroprotective Effect of Electrolyzed Reduced Water against Cisplatin-Induced Kidney Toxicity and Oxidative Damage in Mice. *Journal of the Chinese Medical Association*, **81**, 119-126. <https://doi.org/10.1016/j.jcma.2017.08.014>
- [90] Ferrao, F.M., Lara, L.S. and Lowe, J. (2014) Renin-Angiotensin System in the Kidney: What Is New? *World Journal of Nephrology*, **3**, 64-76. <https://doi.org/10.5527/wjn.v3.i3.64>
- [91] Habiyakare, B., *et al.* (2014) Reduction of Angiotensin A and Alamandine Vasoactivity in the Rabbit Model of Atherogenesis: Differential Effects of Alamandine and Ang(1-7). *International Journal of Experimental Pathology*, **95**, 290-295. <https://doi.org/10.1111/iep.12087>
- [92] Hayati, F., *et al.* (2015) Prevention of Cisplatin Nephrotoxicity. *Journal of Nephro pharmacology*, **5**, 57-60.
- [93] Santos, R.A.S. and Ferreira, A.J. (2006) Pharmacological Effects of AVE 0991, a Nonpeptide Angiotensin-(1-7) Receptor Agonist. *Cardiovascular Drug Reviews*, **24**, 239-246. <https://doi.org/10.1111/j.1527-3466.2006.00239.x>
- [94] Cunha, T.M., *et al.* (2013) The Nonpeptide ANG-(1-7) Mimic AVE 0991 Attenuates Cardiac Remodeling and Improves Baroreflex Sensitivity in Renovascular Hypertensive Rats. *Life Sciences*, **92**, 266-275. <https://doi.org/10.1016/j.lfs.2012.12.008>
- [95] Burns, W.C., *et al.* (2010) Angiotensin II Mediates Epithelial-to-Mesenchymal Transformation in Tubular Cells by ANG 1-7/MAS-1-Dependent Pathways. *American Journal of Physiology-Renal Physiology*, **299**, F585-F593. <https://doi.org/10.1152/ajprenal.00538.2009>
- [96] Singh, K., *et al.* (2012) Possible Mechanism of the Cardio-Renal Protective Effects of AVE-0991, a Non-Peptide Mas-Receptor Agonist, in Diabetic Rats. *Journal of the Renin-Angiotensin-Aldosterone System*, **13**, 334-340. <https://doi.org/10.1177/1470320311435534>
- [97] Qaradakhi, T., *et al.* (2017) Alamandine Reverses Hyperhomocysteinemia-Induced Vascular Dysfunction via PKA-Dependent Mechanisms. *Cardiovascular Therapeutics*, **35**, e12306. <https://doi.org/10.1111/1755-5922.12306>
- [98] El-Hawli, A., *et al.* (2017) IRAP Inhibition Using HFI419 Prevents Moderate to Severe Acetylcholine Mediated Vasoconstriction in a Rabbit Model. *Biomedicine & Pharmacotherapy*, **86**, 23-26. <https://doi.org/10.1016/j.biopha.2016.11.142>
- [99] Klein, S., *et al.* (2015) Hemodynamic Effects of the Non-Peptidic Angiotensin-(1-7) Agonist AVE0991 in Liver Cirrhosis. *PLOS ONE*, **10**, e0138732. <https://doi.org/10.1371/journal.pone.0138732>
- [100] Harrison, C.B., *et al.* (2010) Evidence That Nitric Oxide Inhibits Vascular Inflammation and Superoxide Production via a p47phox-Dependent Mechanism in Mice. *Clinical and Experimental Pharmacology and Physiology*, **37**, 429-434. <https://doi.org/10.1111/j.1440-1681.2009.05317.x>
- [101] Pinheiro, S.V., *et al.* (2004) Nonpeptide AVE 0991 Is an Angiotensin-(1-7) Receptor Mas Agonist in the Mouse Kidney. *Hypertension*, **44**, 490-496. <https://doi.org/10.1161/01.HYP.0000141438.64887.42>
- [102] Barroso, L.C., *et al.* (2012) Renoprotective Effects of AVE0991, a Nonpeptide Mas Receptor Agonist, in Experimental Acute Renal Injury. *International Journal of Hypertension*, **2012**, Article ID: 808726. <https://doi.org/10.1155/2012/808726>
- [103] Ma, Y., *et al.* (2016) AVE 0991 Attenuates Cardiac Hypertrophy through Reducing Oxidative Stress. *Biochemical and Biophysical Research Communications*, **474**,

- 621-625. <https://doi.org/10.1016/j.bbrc.2015.09.050>
- [104] Al-Harbi, N.O., *et al.* (2018) Short Chain Fatty Acid, Acetate Ameliorates Sepsis-Induced Acute Kidney Injury by Inhibition of NADPH Oxidase Signaling in T Cells. *International Immunopharmacology*, **58**, 24-31. <https://doi.org/10.1016/j.intimp.2018.02.023>
- [105] Romero, C.A., Orias, M. and Weir, M.R. (2015) Novel RAAS Agonists and Antagonists: Clinical Applications and Controversies. *Nature Reviews Endocrinology*, **11**, 242-252. <https://doi.org/10.1038/nrendo.2015.6>