

Serum Protein Level Is Related to ADL in Patients with Parkinson's Disease

Lihe Huang^{1*}, Xuezhen Deng¹, Jieyi Yang², Xuemei He¹, Yanchun Zhu¹, Lijian Chen¹

¹Neurology Department, Affiliated Brain Hospital of Guangzhou Medical University, Guangzhou, China

²Guangzhou Medical University, Guangzhou, China

Email: *yangjizhong@126.com

How to cite this paper: Huang, L.H., Deng, X.Z., Yang, J.Y., He, X.M., Zhu, Y.C. and Chen, L.J. (2022) Serum Protein Level Is Related to ADL in Patients with Parkinson's Disease. *Open Journal of Nursing*, 12, 738-744.

<https://doi.org/10.4236/ojn.2022.1211051>

Received: September 24, 2022

Accepted: October 28, 2022

Published: October 31, 2022

Copyright © 2022 by author(s) and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Objectives: To establish an ADL prediction model for Parkinson's inpatients as an auxiliary evaluation scheme. **Methods:** The data of Parkinson's patients hospitalized in the Department of Neurology of Affiliated Brain Hospital of Guangzhou Medical University from 2019 to 2022, which suited the criteria were collected, and a multiple linear regression model was established with serum total protein, serum albumin, age, BMI and education level as independent variables and BI scores as dependent variables. **Results:** A total of 95 PD patients were included (mean 70.05 ± 10.87 years): 53 males and 42 females. The correlation analysis showed that the serum total protein ($r = 0.398$, $P < 0.001$), serum albumin ($r = 0.217$, $P = 0.035$), age ($r = -0.434$, $P < 0.001$), BMI ($r = 0.269$, $P = 0.008$) were respectively linearly correlated with BI total score, and education level is negatively correlated with BI scores (Kendall's tau-b = -0.2 , $P = 0.011$). A multiple linear regression model which established with serum total protein, serum albumin, age, BMI and education level as independent variables and BI scores as dependent variables, is statistically significant ($F = 9.041$, $P < 0.001$, adjust $R^2 = 0.3$). **Conclusion:** The ADL multiple linear regression model can be used as an important means to evaluate the ADL ability of PD patients in hospital.

Keywords

Parkinson's Disease, Nursing, Serum Total Protein, Serum Albumin, Activities of Daily Living

1. Introduction

Parkinson's disease (PD) is a neurodegenerative disease characterized by rest tremor, bradykinesia, rigidity and abnormal posture and pace [1]. Globally, the prevalence of Parkinson's disease is expected to double in the next two decades

as the population ages [2]. As the symptoms of Parkinson's disease will gradually worsen, its activities of daily living (ADL) will also be significantly affected [3]. Nurses often use Barthel Index (BI) to evaluate the activities of daily living of inpatients such as Parkinson's disease [3]. The Barthel Index (BI), with 10 items for mobility and ADL areas, is considered the most convenient measure of competency in basic ADL in neurological and musculoskeletal disorders [4]. However, the BI is easy to use, but often with a certain degree of subjectivity.

Some studies have found that there is a significant correlation between serum protein level and ADL ability in patients with Parkinson's disease, and with the decrease of albumin level, the BI score increased [5]. Therefore, we speculate that serum protein levels can predict the ADL ability of patients with PD to some extent. The purpose of this study is to construct a prediction model of ADL ability of PD patients by analyzing serum total protein, serum albumin and some basic physiological indexes, in order to help judge the ADL ability of Parkinson's patients.

2. Patients and Methods

2.1. Patients

Patients with Parkinson's disease hospitalized in the Department of Neurology of the Affiliated Brain Hospital of Guangzhou Medical University from 2019 to 2022. Inclusion criteria: 1) Meet the diagnostic criteria of PD of the International Parkinson and Movement Disorder Society [6]. 2) No severe cognitive impairment. 3) No mental illness. 4) At present, the symptoms of Parkinson's disease are stable and the medication pattern is regular. Exclusion criteria: 1) Secondary Parkinsonism such as drug-induced, vascular and traumatic Parkinson's syndrome. 2) Parkinsonism-plus syndrome such as multiple system atrophy and progressive supranuclear palsy. 3) Severe respiratory and cardiovascular diseases. 4) Severe cognitive, mental illness.

2.2. Methods

The Parkinson's patients hospitalized in the Department of Neurology of our hospital from 2019 to 2022 were reviewed by professionals, and the patients who met the admission criteria were further selected. The general data such as sex, age, BMI, serum total protein, serum albumin and the scores of BI were collected.

2.3. Related Clinical Data

The serum protein level is a good index to reflect the recent nutritional status of patients. Serum total protein is a complex mixture of various proteins, which can be separated by different methods. Plasma albumin, α 1-globulin, α 2-globulin, β -globulin, fibrinogen, prothrombin and other coagulation factors are synthesized from the liver [7]. Albumin is the most abundant serum protein, produced by hepatocytes, up to 9 - 12 g/days, accounting for about 50% of the total human protein, and the normal serum range is 3.5 - 5 g/L. It is generally considered that

albumin lower than 35 g/L is considered to be hypoalbuminemia [8] [9].

The BI was routinely used to evaluate the ADL of inpatients in the department of neurology of our hospital, which consists of 10 items (bowels, bladder, grooming, toilet use, feeding, transfer, mobility, dressing, stairs, and bathing) [10]. Scores are categorized into 5 levels: 100 (no dependency), 61 - 99 (mild dependency), 41 - 60 (moderate dependency), 20 - 40 (severe dependency), 0 - 19 (complete dependency).

2.4. Statistical Analysis

Data were analyzed using SPSS26.0 (SPSS Inc., Chicago, IL, USA). The enumeration data are expressed as numbers and percentages, and the measurement data are expressed as mean \pm standard deviation ($\bar{x} \pm s$). Correlation analysis between the measurement data (serum total protein, serum albumin, age, BMI) and the BI scores was conducted using Pearson correlation analysis. The correlation between gender and the total score of BI was examined with independent sample t-test. Kendall's tau-b correlation analysis was used to analyze the correlation between education level and BI total score. Finally, the statistically significant variables are obtained as independent variables and incorporated into the multivariate linear regression equation with BI as the dependent variable for statistical processing. A value of $P < 0.05$ was considered significant.

3. Results

3.1. General Data

A total of 95 PD patients were included (mean 70.05 ± 10.87 years): 53 males and 42 females. Other information is shown in **Table 1**.

3.2. Univariate Correlation

As showed in **Table 1**. The serum total protein, serum albumin, age, BMI and BI total scores showed normal distribution as indicated by the P-P plot, and the Pearson correlation analysis showed that the serum total protein ($r = 0.398$, $P <$

Table 1. Univariate correlation.

Variables	$\bar{x} \pm s/n$	r/t/Kendall's tau-b	P
TP	64.84 ± 5.95	0.398	0.001
ALB	38.19 ± 5.35	0.217	0.035
Age	70.05 ± 10.87	-0.434	0.001
BMI	20.46 ± 2.55	0.269	0.008
Education level (illiterate/primary/junior high/senior high school/college and above)	8/26/24/19/18	-0.2	0.011

Abbreviations: TP: serum total protein; ALB: serum albumin; BMI: body mass index.

0.001), serum albumin ($r = 0.217$, $P = 0.035$), age ($r = -0.434$, $P < 0.001$), BMI ($r = 0.269$, $P = 0.008$) were respectively linearly correlated with BI total score. The results of independent sample t-test indicated that there was no statistical difference in BI scores between men and women, which means there was no correlation between gender and BI scores. The correlation between education level and BI scores was conducted by Kendall's tau-b correlation analysis, and it is concluded that education level is negatively correlated with BI scores (Kendall's tau-b = -0.2 , $P = 0.011$).

3.3. Multiple Linear Regressions

A multiple linear regression model was established with serum total protein, serum albumin, age, BMI and education level as independent variables and BI scores as dependent variables. The target sample size for this study was calculated based on the recommendation of five to ten outcome events per predictor variable (EPV) [11]. In this study, there are 5 variables of regression, so the number of people included in regression ($n = 95 > 50$) is sufficient. By drawing the partial regression scatter plot and the scatter plot of biochemical residual and predicted value, we can judge that there is a linear relationship between independent variable and dependent variable. It has been verified that the variables are independent of each other, and the Durbin-Watson test value is 1.5%. By drawing the scatter diagram between the biochemical residual and the unstandardized predicted value, it is proved that the variance of the residual is homogeneous. The variance inflation factors (VIF) of the regression are all less than 10, suggesting that there is no multicollinearity. And the standardized residual P-P plot of regression suggests that the residual is close to normal distribution. In summary, the regression model is statistically significant ($F = 9.041$, $P < 0.001$, $\text{adjust } R^2 = 0.3$) (showed in **Table 2**). And the effects of the five independent variables included in the model on BI scores were statistically significant ($P < 0.05$).

4. Discussion

In this study, the univariate correlation analysis revealed that total protein, albumin, and BMI were positively correlated with BI scores. It may be that these indicators often represent the nutritional status of PD patients. Such as in patients with poor nutritional status, their exercise ability will be further impaired, and their ADL ability will also decrease accordingly, which is consistent with the conclusions of some recent studies [12] [13] [14]. In addition, we found that the educational level of PD patients was negatively correlated with BI. The lower the educational level, the higher their ADL ability. This finding is not consistent with Palacios's study [15]. However, some studies have shown that there is no significant correlation between education level and ADL ability [16] [17]. So far, no additional research has been found to support our findings, thus we speculate that AD patients with low education level may be more involved in physical activity in the past, so that their motor ability and ADL ability are better preserved.

Furthermore, our multiple linear regression model has been proved to be

Table 2. Multiple linear regression.

Variables	β	SE	B'	t	P
Constant	45.931	411.383		1.110	0.027
TP	-0.115	0.526	-0.023	-0.218	0.028
ALB	1.313	0.624	0.238	2.104	0.038
Age	-0.913	0.253	-0.336	-3.610	0.001
BMI	-5.008	2.038	-0.212	-2.457	0.016
Education level	2.247	1.024	0.194	2.196	0.031

Abbreviations: TP: serum total protein; ALB: serum albumin; BMI: body mass index.

statistically significant, which proves that we can predict the ADL ability of PD patients by their serum protein levels and simple physiological indexes. In the past, the BI was used to evaluate the ADL ability of PD patients when they were hospitalized, which is often subjective [14]. Also, different nurses evaluate them in different periods of time during hospitalization, and the results may be inconsistent. Accordingly, our prediction model is often more convenient for ADL assessment of hospitalized PD patients, and the results are more consistent. Aside from that, plasma protein level is often an item that must be checked when PD patients are hospitalized. While evaluating the nutritional status of PD patients, they can also evaluate their ADL ability, which makes nursing work more convenient and more suitable to implement nursing programs and ADL tracking of PD patients.

In this study, we still have some shortcomings that should be improved in the future. For instance, the model is mainly applied to inpatients, but not to home patients.

5. Conclusion

The ADL multiple linear regression model based on serum total protein, serum albumin, age, BMI and education level can be used as an important means to evaluate the ADL ability of PD patients in hospital, and can provide reference for nurses' evaluation plan.

Fund

This work was supported by the Guangzhou Medical Health Science and Technology Project (Grant NO. 20201A011044).

Conflicts of Interest

The authors declare no conflicts of interest regarding the publication of this paper.

References

- [1] Bloem, B.R., Okun M.S. and Klein, C. (2021) Parkinson's Disease. *The Lancet*, **397**,

- 2284-2303. [https://doi.org/10.1016/S0140-6736\(21\)00218-X](https://doi.org/10.1016/S0140-6736(21)00218-X)
- [2] Dorsey, E.R., Sherer, T., Okun, M.S. and Bloem, B.R. (2018) The Emerging Evidence of the Parkinson Pandemic. *Journal of Parkinson's Disease*, **8**, S3-S8. <https://doi.org/10.3233/JPD-181474>
- [3] Clarke, C.E., et al. (2016) Physiotherapy and Occupational Therapy vs No Therapy in Mild to Moderate Parkinson Disease: A Randomized Clinical Trial. *JAMA Neurology*, **73**, 291-299. <https://doi.org/10.1001/jamaneuro.2015.4452>
- [4] Taghizadeh, G., et al. (2020) Barthel Index and Modified Rankin Scale: Psychometric Properties during Medication Phases in Idiopathic Parkinson Disease. *Annals of Physical and Rehabilitation Medicine*, **63**, 500-504. <https://doi.org/10.1016/j.rehab.2019.08.006>
- [5] Li, S., Zhang, Q.X., et al. (2021) Serum Folate, Vitamin B12 Levels, and Systemic Immune-Inflammation Index Correlate with Motor Performance in Parkinson's Disease: A Cross-Sectional Study. *Frontiers in Neurology*, **12**, Article No. 665075. <https://doi.org/10.3389/fneur.2021.665075>
- [6] Postuma, R.B., et al. (2015) MDS Clinical Diagnostic Criteria for Parkinson's Disease. *Movement Disorders*, **30**, 1591-1601. <https://doi.org/10.1002/mds.26424>
- [7] Tian, C.R., Qian, L., Shen, X.Z., Li, J.J. and Wen, J.T. (2014) Distribution of Serum Total Protein in Elderly Chinese. *PLOS ONE*, **29**, e101242. <https://doi.org/10.1371/journal.pone.0101242>
- [8] Cabrerizo, S., Cuadras, D., Gomez-Busto, F., Artaza-Artabe, I., Marín-Ciancas, F. and Malafarina, V. (2015) Serum Albumin and Health in Older People: Review and Meta Analysis. *Maturitas*, **81**, 17-27. <https://doi.org/10.1016/j.maturitas.2015.02.009>
- [9] Bernardi, M., Maggioli, C. and Zaccherini, G. (2012) Human Albumin in the Management of Complications of Liver Cirrhosis. *Critical Care*, **16**, Article No. 211. <https://doi.org/10.1186/cc11218>
- [10] Mahoney, F.I. and Barthel, D.W. (1965) Functional Evaluation: The Barthel Index. *Maryland State Medical Journal*, **14**, 61-65.
- [11] Vittinghoff, E. and McCulloch, C.E. (2007) Relaxing the Rule of Ten Events per Variable in Logistic and Cox Regression. *American Journal of Epidemiology*, **165**, 710-718. <https://doi.org/10.1093/aje/kwk052>
- [12] Wei, K., et al. (2018) Association of Frailty and Malnutrition with Long-Term Functional and Mortality Outcomes among Community-Dwelling Older Adults: Results from the Singapore Longitudinal Aging Study 1. *JAMA Netw Open*, **1**, e180650. <https://doi.org/10.1001/jamanetworkopen.2018.0650>
- [13] Stange, I., et al. (2013) Effects of a Low-Volume, Nutrient- and Energy-Dense Oral Nutritional Supplement on Nutritional and Functional Status: A Randomized, Controlled Trial in Nursing Home Residents. *The Journal of Post-Acute and Long-Term Care Medicine*, **14**, 628.E1-628.E8. <https://doi.org/10.1016/j.jamda.2013.05.011>
- [14] Rolland, Y., et al. (2007) Exercise Program for Nursing Home Residents with Alzheimer's Disease: A 1-Year Randomized, Controlled Trial. *Journal of the American Geriatrics Society*, **55**, 158-165. <https://doi.org/10.1111/j.1532-5415.2007.01035.x>
- [15] Palacios-Cena, D., et al. (2012) Has the Prevalence of Disability Increased over the Past Decade (2000-2007) in Elderly People? A Spanish Population-Based Survey. *The Journal of Post-Acute and Long-Term Care Medicine*, **13**, 136-142. <https://doi.org/10.1016/j.jamda.2010.05.007>
- [16] Yang, L., Konttinen, H., Martikainen, P. and Silventoinen, K. (2018) Socioeconomic

Status and Physical Functioning: A Longitudinal Study of Older Chinese People. *The Journals of Gerontology. Series B*, **73**, 1315-1329.

<https://doi.org/10.1093/geronb/gbx010>

- [17] Xu, X., Liang, J., Bennett, J.M., Botosaneanu, A., and Allore, H.G. (2015) Socioeconomic Stratification and Multidimensional Health Trajectories: Evidence of Convergence in Later Old Age. *The Journals of Gerontology. Series B*, **70**, 661-671.

<https://doi.org/10.1093/geronb/gbu095>