

EFFECT OF SYNTHETIC ANTIMICROBIAL PEPTIDES ON *NAEGLERIA FOWLERI* TROPHOZOITES

Supathra Tiewcharoen¹, Watchara Phurttikul^{1,2}, Jundee Rabablert³, Prasert Auewarakul², Sittiruk Roytrakul⁴, Pruksawan Chetanachan⁵, Thassanant Atitthep⁶ and Virach Junnu¹

¹Department of Parasitology, ²Department of Microbiology, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok; ³Department of Biology, Faculty of Science, Silpakorn University, Nakhon Pathom; ⁴Genome Institute, National Center for Genetic Engineering and Biotechnology, Pathum Thani; ⁵National Institute of Health, Department of Medical Sciences, Nonthaburi; ⁶Center of Nanoimaging, Faculty of Science, Mahidol University, Bangkok, Thailand

Abstract. We evaluated the effect of tritripticin, lactoferrin, killer decapeptide and scrambled peptide *in vitro* against *Naegleria fowleri* trophozoites compared with amphotericin B. Tritripticin (100 µg/ml) caused apoptosis of *N. fowleri* trophozoites (2×10^5 cells/ml), while lactoferrin, killer decapeptide and scrambled peptide did not. On Gormori trichrome staining, tritripticin affected the elasticity of the surface membrane and reduced the size of the nuclei of *N. fowleri* trophozoites. The ultrastructure surface membrane and food cup formation of the trophozoites were 100% inhibited. These results are consistent with inhibition of the *nfa1*, *Mp2CL5* of the treated trophozoite, which plays a role in food cup formation. Tritripticin 100 µg/ml was not toxic against SK-N-MC cells. Our findings suggest tritripticin has activity against the surface membrane and *nfa1* and *Mp2CL5* of *N. fowleri* trophozoites and could be developed as a potential therapeutic agent.

Keywords: *Naegleria fowleri*, antiamebic peptide, tritripticin

INTRODUCTION

The free-living ameba *Naegleria fowleri* is a causative agent of primary amebic meningoencephalitis (PAM) in humans and animals (Madarova *et al*, 2010). PAM may occur, when an otherwise healthy person is exposed to contaminated water in the nose (Yoder *et al*, 2010). Clinical manifestations begin a few days after exposure

(Visvesvara, 2010). Sources of contaminated water include environmental pools, inadequately chlorinated swimming pool water and heated and contaminated tap water (Yoder *et al*, 2012). Infection initially occurs when the *N. fowleri* ameba penetrates the mucous membranes of the nasal cavity and travels to the brain through the olfactory nerves (Visvesvara *et al*, 2005). The *nfa1* and *Mp2CL5* genes are found only in *N. fowleri* (Tiewcharoen *et al*, 2011). Nfa1 protein is expressed from the *nfa1* gene and is located in the pseudopodia and around food vacuoles (Kang *et al*, 2005). Nfa1 protein is localized

Correspondence: Jundee Rabablert, Department of Biology, Faculty of Science, Silpakorn University, Nakhon Pathom 73000, Thailand. Tel: +66 (0) 34 243429; Fax: +66 (0) 34 273046 E-mail: jundee04@gmail.com; jundee@su.ac.th

in the food cups which are involved in phagocytic activity (Marciano-Cabral and Cabral, 2007; Tiewcharoen *et al*, 2011). The *Naegleria pore B* gene encodes Naegleria pore A and B proteins that display pore-forming activities and kill prokaryotic and eukaryotic target cells and the *nf actin* is a housekeeping gene (Tiewcharoen *et al*, 2012). The *ITS* is located on the 5.8S rRNA gene and species-specific chromosomal DNA *pB2.5* genes are used to identify pathogenic *N. fowleri* at the molecular level (Rabablert *et al*, 2011).

Patients with *N. fowleri* infection often have initial symptoms of high grade fever (38°-40°C), sore throat, stuffy nose, severe headache and then rapidly progress to meningitis, encephalitis, ataxia, mental confusion and coma a few days prior to death (Yoder *et al*, 2012). Treatment includes amphotericin B (AMB) and multiple other drugs: rifampicin, fluconazole, dexamethasone and ceftriaxone (Vargas-Zepeda *et al*, 2005). Most patients with *N. fowleri* infection die; the rare cases who survive may have neurological sequel (Visvesvara *et al*, 2010).

Because of the poor treatment results, several medications such as AMB, miconazole, fluconazole, ketoconazole and rifampin had been used in combination (CDC, 2013) with the new drug, Miltefosine (Kim *et al*, 2008). However, the mortality of PAM is still high (Yoder *et al*, 2010). Treatment problems include limited availability of AMB and side effects of drugs (Kim *et al*, 2008). AMB resistance by *N. fowleri* has been reported (Donadio *et al*, 2010). Several studies have reported antifungal resistance to the azole group can be spread via resistant *Candida albicans* (Sterling and Merz, 1998; Barker and Rogers, 2006). *N. fowleri* resistance to AMB has been attributed to a virulent gene (Tiewcharoen *et al*, 2011).

Antimicrobial peptides (AMPs) are recognized as an important component of the nonspecific host defense system against invading pathogens (Hancock and Chapple, 1999; Wilcox, 2004; McGregor, 2008). The characteristic of AMPs include small molecular size and cationic affinity (Dürr *et al*, 2006), it is usually non-immunogenic and has a short half-life. Their activity includes targeting the membrane, disrupting protein-protein interaction and the ability to penetrate tissues (McGregor, 2008). The activity of AMPs can be triggered by binding to negatively charged sites on the surface of brain capillary endothelial cells (Mahurkar *et al*, 2014). AMPs have a wide spectrum of activity, acting against gram-positive and gram-negative bacteria, protozoa, fungi, viruses, and mammalian cells (Brogden, 2005; Cironi *et al*, 2006; Bagheri *et al*, 2011). The aim of the current study was to test for the first time the effectiveness of AMPs against *N. fowleri* trophozoites and to evaluate cell damage and alternations in luminescence by scanning electron microscopy.

MATERIALS AND METHODS

Culture of *Naegleria fowleri*

N. fowleri (Siriraj strain) was isolated in 1986 from a PAM patient at Siriraj Hospital, Bangkok, Thailand. The trophozoites were cultured in T 75-cm² flasks (Corning, Corning, NY) containing Nelson's medium supplemented with 5% fetal calf serum (FCS) without antibiotics at 37°C. The trophozoites were incubated at 4°C for 10 minutes, scraped and then centrifuged at 5,000 rpm for 2 minutes. The pellet was then dissolved in 2 ml of Nelson's medium. The number of cells was counted using the Trypan blue exclusion method: 10 µl of medium containing trophozoites was mixed with 90 µl of 0.4%

Table 1
List of antimicrobial peptide names and sequences.

Peptides	Amino acid sequences	Molecular weight
Tritrpticin	VRRFPWWPFLRR	1902.30
Lactoferrin	RRWQWRMKKLG	1544.90
Killer decapeptide	AKVTMTCSAS	998.19
Scrambled peptide	MSTAVSKCAT	998.19

Trypan blue stain and then the cells were counted under a light microscope. The procedure was conducted in duplicate.

Human neuroblastoma cultivation

Human neuroblastoma SK-N-MC cells isolated from a female Caucasian patient with Askin's tumor were purchased from Cell Line Service (Eppelheim, Germany). The cells were maintained in Dulbecco's Modified Eagle Medium and HAM's F-12 (DMEM: HAM'S F-12) medium with 10% fetal bovine serum, 4 mM L-glutamine, 100 µl/ml penicillin and 100 µg/ml streptomycin and were grown in monolayer cultures at 37°C in 5% CO₂ (Tiewcharoen *et al*, 2008).

Antimicrobial peptides

Tritrpticin (Trp; Infante *et al*, 2011), lactoferrin (LF; León-Sicairos *et al*, 2006), killer decapeptide (Kp) and scrambled peptide (Sp) (Fiori *et al*, 2006) are AMPs with properties against protozoa (Arrighi *et al*, 2002). The AMPs were obtained from China Peptides (Shanghai, China) (Table 1). The AMPs, which came as a powder, were dissolved in distilled water at a concentration of 1 mg/ml and then stored at -80°C, until used.

Reagents

Caspase-Glo[®] 3/7 Substrate and Caspase-Glo[®] 3/7 Buffer obtained from Promega (Madison, WI), were mixed by inversion until the substrate was dis-

solved and then stored at -20°C until used. Five milligrams of MTT (Invitrogen/ Molecular probes, Eugene, OR) was dissolved in 1 ml of phosphate buffer (PBS pH 7.4) and used immediately.

Caspase-Glo[®] 3/7 assay

N. fowleri trophozoites at a concentration of 1x10⁶ cells/ml, were dissolved in Nelson's medium. One hundred microliters of each studied peptide was then added to the trophozoites and incubated at 37°C for 0.5, 1, 3, 6 or 12 hours. As a negative control, trophozoites in Nelson's medium were studied alone. The untreated and treated trophozoites were then harvested and centrifuged at 5,000g for 2 minutes. A pellet each of untreated and treated trophozoites were dissolved in PBS 7.4 (250 µl) and then 50 µl of Caspase-Glo[®] 3/7 reagent was added and incubated at room temperature (RT) for 30 minutes in darkness. Caspase activity was measured at a wavelength of 485 nm with an emission wavelength of 527 nm by a luminometer (luminescence-Octa AB-2270, Tokyo, Japan) (Renault *et al*, 2010).

Gormori trichrome staining

Untreated and treated trophozoites were fixed with Schaudinn's fixative at RT for 24 hours, stained with 2% iodine for 2 minutes, washed with 70% ethanol for 2 minutes, counterstained with Gormori trichrome for 10-15 minutes, washed with acid alcohol for 5 seconds, dehydrated

Table 2
Primers used for RT-PCR.

Genes	Primer sequences	bp	PCR product (bp)
<i>nfa1</i>	Forward: 5'ATGGCACTACTATTCCATCACCA 3'	23	360
	Reverse : 5'TTAAAGCACTCCCTTGTACTTCAT 3'	24	
<i>Mp2CL5</i>	Forward: 5'TCTAGAGATCCAACCAATGG 3'	20	166
	Reverse : 5'ATTCTATTCACTCCACAATCC 3'	21	
<i>ITS</i>	Forward: 5'GAACCTGCGTAGGGATCATT 3'	21	450
	Reverse : 5'TTTCTTTTCCTCCCCTTATTA 3'	21	
<i>pB2.5</i>	Forward: 5'GTGAAAACCTTTTTTCCATTTACA 3'	24	310
	Reverse : 5'AAATAAAAAGATTGACCATTGAAA 3'	24	
<i>Naegleria pore B</i>	Forward: 5'TTGATGTCAATGCTGTCAAGC 3'	21	165
	Reverse : 5'CTTGGGCAGACATCAACG 3'	19	
<i>nf actin</i>	Forward: 5'ACTCTGGTGATGGTGTCTCTCACAC 3'	25	170
	Reverse : 5'CTCTGACAATTCTCTCTCAGTGG 3'	24	

bp, base pair.

using 90% and then 95% ethanol for 1 minute each xylene for 1 minute and then mounted on a glass slide and observed under a light microscope.

Scanning electron microscope

Untreated and treated trophozoites were pre-warmed in 2.5% glutaraldehyde and 0.1 M PBS at a pH of 7.3 at 37°C for 30 minutes and then cooled to 4°C for 24 hours. The trophozoites were then fixed in 1% osmium tetroxide (OsO₄) and 0.1 M PBS at RT for 90 minutes, rinsed with 0.1 M PBS, sequentially dehydrated in serial dilutions of ethanol, critical point dried, and then coated with gold-palladium (Au-Pd). Finally, the trophozoites were examined and photographed under a scanning electron microscope (SEM) (Hitachi S-51, Tokyo, Japan) at an accelerating voltage of 25 kV.

Total RNA extraction

Untreated and treated trophozoite pellets were extracted with a Favorprep™ Tissue total RNA mini kit (Favorgen, Ping-Tung, Taiwan) at 1, 3, 6, and 12 hours post-

incubation. Briefly, approximately 1 x 10⁶ trophozoites were added to 350 µl of FARB buffer and 3.5 µl β-ME and vortexed vigorously to lyse the cells. The sample was transferred to a filter column placed on a collection tube and centrifuged at 10,000g for 2 minutes. The supernatant from the collection tube was then transferred to a microcentrifuge tube. Ethanol (70%) was added to the lysate and mixed well by vortexing. The ethanol-treated sample was transferred to a FARB Mini column, centrifuged at 10,000g for 1 minute. Wash buffer 1 (500 µl) was added to wash the FARB Mini column and centrifuged at 10,000g for 1 minute. Wash buffer 2 (750 µl) was added to wash the FARB Mini column and centrifuged at 10,000g for 1 minute, twice. The FARB Mini column was centrifuged at 10,000g for 3 minutes to dry the column. RNase-free ddH₂O (45 µl) was added to the membrane center of the FARB Mini column for 1 minute and centrifuged at 10,000g for 2 minutes to elute the RNA. The RNA was stored at -80°C until used.




Fig 1—Apoptosis of *N. fowleri* trophozoites (2×10^5 cell/ml) treated with Trp, LF, Kp, Sp (100 $\mu\text{g/ml}$) and AMB (10 $\mu\text{g/ml}$) evaluated with the Caspase-Glo[®] 3/7 Assay at 0.5, 1, 3, 6 and 12 hours post-incubation. Data is expressed as relative light units (RLU). These experiments were performed in triplicate. Error bars represent standard deviations ($p < 0.05$). hpi, hours post incubation.

cDNA synthesis

Untreated and treated trophozoite pellets were extracted with the Favorprep[™] Tissue total RNA mini kit (Favorgen, Ping-Tung, Taiwan). A total of 20 μl of RNA was used as a template to synthesize the first strand of the cDNA using the Maxime RT PreMix Kit (iNtRON Biotechnology, Gyeonggi-do, Korea), following the manufacturer's instructions. Briefly, template RNA and RNase-free water were added to the MaximeRT PreMix tubes (Random primer) to give a total volume of 20 μl ; this was done in duplicate. The cDNA reaction was carried out at 45°C for 60 minutes and then 95°C for 5 minutes and then stored at -20°C until used for PCR amplification.

Polymerase chain reaction

The PCR reaction was carried out in a volume of 20 μl , in the presence of 10 mM Tris-HCl, 2 mM MgCl₂, 0.2 mM

dNTP, 0.2 μM of specific primers (Table 2), 2.5 μmol of *Taq* polymerase and 2 μl of single stranded cDNA. The cDNA templates were amplified at an initial incubation at 94°C for 5 minutes followed by 35 cycles at 94°C for 30 seconds, 55°C for 30 seconds and 72°C for 45 seconds in a Gene Amp PCR 2400 thermal cycle system (Perkin-Elmer, Cetus Crop, Waltham, MA). The PCR product was incubated at 72°C for 10 minutes to ensure complete extension of all amplified molecules. Finally, the PCR products were subjected to 2% agarose Tris-borate-EDTA gel electrophoresis at 100 V

for 30 minutes. The gel was stained with ethidium bromide and then visualized under ultraviolet light.

The effect of tritrpticin on morphology of SK-N-MC co-culture with *N. fowleri*

SK-N-MC cells (1×10^6 cells/ml) were cultured in Dulbecco's Modified Eagle Medium HAM's F-12 for 24 hours and then *N. fowleri* trophozoites were added to the SK-N-MC cells and incubated with or without Trp (100 $\mu\text{g/ml}$) at 37°C for 0.5, 1, 2, or 3 hours. As a negative control, the SK-N-MC cells were treated with medium alone. The morphologies of the cultures were observed with a SEM. The preparation of SEM was described above.

Statistical analysis

The results are expressed as means \pm standard deviations for the three experiments carried out in triplicate. A Student's *t*-test was used for analysis; a *p*-value < 0.05 was considered significant.




Fig 2—Photomicrograph of *N. fowleri* trophozoites in the absence or presence of Trp (100 µg/ml) in Gormori's trichrome stain at 1 and 3 hours post-exposure: (a) control trophozoites exposed to medium alone showing a vesicular nucleus and small granules in the cytoplasm; (b) 1 hour post-exposure showing a small nucleolus; (c) 3 hours post-exposure appears the same as control (x 400).

RESULTS

Effect of antimicrobial peptides on apoptosis of *N. fowleri* trophozoites

As shown in Fig 1, Trp induced apoptosis in *N. fowleri* trophozoites after a 30-minute exposure period at a concentration of 100 µg/ml. The activity of Trp is also comparable to AMB (10 µg/ml). In contrast, LF, Kp and Sp did not induce apoptosis. Trp (100 µg/ml) did not damage human neuroblastoma SK-N-MC cells at indicated times (data not shown).

Effect of tritrypticin on the morphology of *N. fowleri* trophozoites

The morphological characteristics of *N. fowleri* trophozoites exposed to Trp (100 µg/ml) and stained with Gormori trichrome at 0.5, 1, 2 and 3 hours post-exposure were compared to untreated trophozoites. Untreated trophozoites had homogenous cytoplasm with a normal elastic membrane and a large nucleolus in the central nucleus surrounded by a complete nuclear membrane (Fig 2a). Trp-treated trophozoites had an elongated shape with a loose elastic membrane and

a small nucleolus 1 hour post-exposure (Fig 2b). However, treated trophozoites had a similar morphology to untreated trophozoites 3 hours after exposure (Fig 2c). A SEM micrograph of *N. fowleri* trophozoites showed a sucker like apparatus and wrinkled membrane (Fig 3a). Trp-treated trophozoites were small in size and few in number. In addition, the ultrastructural surface membrane and food cup formation were 100% inhibited at 1 hour post-exposure (Fig 3b). However, the abnormalities of the surface membrane recovered by 3 hours post-exposure (Fig 3c).

Effect of tritrypticin on *N. fowleri* trophozoites at the gene level

Trp inhibited the *nfa1* and *Mp2CL5* genes of *N. fowleri* trophozoites 1 hour post-incubation, but did not inhibit the *nfa1* or *Mp2CL5* genes at 3, 6 and 12 hours post-incubation. This indicates Trp had time-dependent activity against *N. fowleri* trophozoites. Trp did not inhibit the *ITS*, *pB2.5*, *Naegleria pore B* or *nfactin* genes of *Naegleria* trophozoites at 1, 3, 6 or 12 hours post-incubation (Fig 4).




Fig 3—Scanning electron micrograph of *Naegleria* trophozoites; (a) control trophozoites exposed to medium alone showing a sucker like apparatus and wrinkled membrane; (b) Trp (100 µg/ml) treated amebae are small in size and few in number; (c) 3 hours post-exposure is similar to control. Bars represent 10, 20 µm. Sa, sucker apparatus.

The effect of tritrypticin on the morphology of the SK-N-MC/*N. fowleri* co-culture

A SEM micrograph of the SK-N-MC cells not exposed to *N. fowleri* trophozoites showed an elongated shape with dendrites and axons. The SK-N-MC cells cultured with *N. fowleri* trophozoites had trophozoites attached to the surface membrane of the SK-N-MC cells 1 hour post-incubation. Pre-incubation, the tritrypticin treated SK-N-MC/*N. fowleri* co-culture had a reduction in the size and number of trophozoites. This suggests Trp may prevent phagocytosis of *N. fowleri* by SK-N-MC cells.

DISCUSSION

Free living *N. fowleri* amoeba can cause acute, fulminant, necrotizing, hemorrhagic PAM leading to death (Budge *et al*, 2013). AMB is the only medication with proven clinical efficacy in treating PAM (Brunton *et al*, 2006). However, AMB is not always successful in treating PAM and is associated with severe adverse effects (Soltow and Brenner, 2007).

AMPs are important components of the nonspecific host defense system against invading pathogens (Brogden, 2005). Typically, these peptides are relatively short, positively charged, and amphiphilic (Reddy *et al*, 2004). Previous publications have reported AMP activity against bacteria, fungi, viruses, and protozoa (Cirioni *et al*, 2006; Bagheri *et al*, 2011). It has been believed AMPs act at the membrane level by permeabilization of the cytoplasmic membrane of the microorganism (Franco *et al*, 2006; Jaeyong *et al*, 2012).

In our study, Trp at 100 µg/ml reduced *N. fowleri* trophozoite viability (Fig 1). Our findings are consistent with other reports in which Trp at 100 µg/ml reduced




Fig 4—Expression of *nfa1*, *Mp2CL5*, *ITS*, *pB2.5*, *Naegleria pore B* and *nf actin* genes in untreated and Trp-treated *N. fowleri* trophozoites at 1, 3, 6 and 12 hours post-exposure using RT-PCR.

viability of *Trichomonas vaginalis* (Infante *et al*, 2011). Trp has also been shown to lyse *Bacillus subtilis* and *Escherichia coli* membranes (Bagheri *et al*, 2011). Trp caused *N. fowleri* trophozoites to have abnormal membranes and damaged food cups in our study (Figs 2 and 3). AMB and chlorpromazine damage *N. fowleri* trophozoites, causing bleb formation and disappearance of suckers and pseudopodia (Tiewcharoen *et al*, 2011). Trp activity is comparable to AMB and chlorpromazine, but it appears to be ameba cestatic.

Previous studies have found *nfa1* and *Mp2CL5* genes of *N. fowleri* trophozoites to be pathogenic (Kang *et al*, 2005). Nfa1 protein, expressed by the *nfa1* gene, is located in the pseudopodia and around food vacuoles (Shin *et al*, 2001). Nfa1 protein is specifically localized to food cups, which are involved in phagocytic activity (Kang *et al*, 2005). In our study, Trp inhibited the *nfa1* gene which can cause the amebae to lose their phagocytic activity. Réveiller *et al* (2002) found expression of *Mp2CL5* protein in *N. fowleri* during the growth phase to be regulated. *Mp2CL5* protein is increased in expression during

the stationary phase of growth, when the cells are experiencing nutrients. In our study, Trp inhibited the *Mp2CL5* gene, suggesting the activity of Trp may cause a loss of cell recognition, sensing the environment and growth of the ameba (Fig 4).

Tritrpticin (>150 µg/ml) retains most of its antimicrobial activity, but has enhanced hemolytic and membrane-disruption activity (Yang *et al*, 2002). Trp (>150 µg/ml)

has an inhibitory effect against human MDA-MB-361 and A549 cells (Yang *et al*, 2009). In our study, Trp (100 µg/ml) did not damage human neuroblastoma SK-N-MC cells. In contrast, AMP (10 µg/ml) decreased cell viability by 40% 12 hours post-incubation (data not shown). Trp (100 µg/ml) had an effect on *N. fowleri* trophozoites at both the cellular and molecular levels (Figs 1 and 4). We believe Trp may be a good candidate for development as a potential drug against *N. fowleri* trophozoites.

ACKNOWLEDGEMENTS

This work was partially supported by the Chalermprakiat Foundation, and a Siriraj Graduate Thesis Scholarship from the Faculty of Medicine, Siriraj Hospital, Mahidol University and by a grant number RPG 2554/11 from the Department of Biology, Faculty of Science, Silpakorn University at Sanamchan Palace, Nakhon Pathom, Thailand. We thank Dr Pathom Awanpanyalert, Director of the National Institute of Health, Department of Medical Sciences for providing the electron

microscopy facilities.

REFERENCES

- Arrighi RBG, Nakamura C, Miyake J, Hurd H, Burgess JG. Design and activity of antimicrobial peptides against sporogonic-stage parasites causing murine malaras. *Antimicrob Agents Chemother* 2002; 46: 2104.
- Bagheri M, Beyermann M, Dathe M. Mode of action of cationic antimicrobial peptides defines the tethering position and the efficacy of biocidal surfaces. *Bioconjug Chem* 2011; 23: 66-74.
- Barker KS, Rogers PD. Recent insights into the mechanisms of antifungal resistance. *Curr Infect Dis Rep* 2006; 8: 449-6.
- Brogden KA. Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria? *Nat Rev Microbiol* 2005; 3: 238-50.
- Brunton LL, Lazo JS, Parker KL. Goodman & Gilman's the pharmacological basis of therapeutics. 11th ed. New York: McGraw-Hill, 2006: 1984 pp.
- Budge PJ, Lazensky B, Van Zile KW, et al. Primary amebic meningoencephalitis in Florida: a case report and epidemiological review of Florida cases. *J Environ Health* 2013; 75: 26-31.
- Centers for Diseases Control and Prevention (CDC). Primary amebic meningoencephalitis associated with ritual nasal rinsing – St Thomas, US Virgin Islands, 2012. *MMWR Morb Mortal Wkly Rep* 2013; 62: 903.
- Cirioni O, Giacometti A, Silvestri C, et al. In vitro activities of tritrypticin alone and in combination with other antimicrobial agents against *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2006; 50: 3923-5.
- Donadio S, Maffioli S, Monciardini P, Sosio M, Jabes D. Antibiotic discovery in the twenty-first century: current trends and future perspective. *J Antibiotics* 2010; 63: 423-30.
- Dürr UHN, Sudheendra US, Ramamoorthy A. LL-37, the only human member of the cathelicidin family of antimicrobial peptides. *Biochim Biophys Acta Biomemb* 2006; 1758: 1408-25.
- Fiori PL, Mattana A, Dessi D, Conti S, Magliani W, Polonelli L. *In vitro* acanthamoebicidal activity of a killer monoclonal antibody and a synthetic peptide. *J Antimicrob Chemother* 2006; 57: 891-8.
- Franco R, Bortner CD, Cidlowski JA. Potential roles of electrogenic ion transport and plasma membrane depolarization in apoptosis. *J Membr Biol* 2006; 209: 43-58.
- Hancock REW, Chapple DS. Peptide antibiotics. *Antimicrob Agents Chemother* 1999; 43: 1317-23.
- Infante VV, Miranda-Olvera AD, De Leon-Rodriguez LM, Anaya-Velazquez F, Rodriguez MC, Avila EE. Effect of the antimicrobial peptide tritrypticin on the *in vitro* viability and growth of *Trichomonas vaginalis*. *Curr Microbiol* 2011; 62: 301-6.
- Jaeyong C, Hwang I-s, Choi H, Hwang JH, Hwang JS, Lee DG. The novel biological action of antimicrobial peptides via apoptosis induction. *J Microbiol Biotechnol* 2012; 22: 1457-66.
- Kang SY, Song KJ, Jeong SR, et al. Role of the Nfa1 protein in pathogenic *Naegleria fowleri* cocultured with CHO target cells. *Clin Diagn Lab Immunol* 2005; 12: 873-6.
- Kim JH, Jung SY, Lee YJ, et al. Effect of therapeutic chemical agents *in vitro* and on experimental meningoencephalitis due to *Naegleria fowleri*. *Antimicrob Agents Chemother* 2008; 52: 4010-6.
- León-Sicairos N, Reyes-López M, Ordaz-Pichardo C, De la Garza M. Microbicidal action of lactoferrin and lactoferricin and their synergistic effect with metronidazole in *Entamoeba histolytica*. *Biochem Cell Biol* 2006; 84: 327-36.
- Madarova L, Trnkov K, Feiková S, Klement C, Obernauerov M. A real-time PCR diagnostic method for detection of *Naegleria fowleri*. *Exp Parasitol* 2010; 126: 37-41.
- Mahurkar S, Suppiah V, O'Doherty C. Pharmacogenomics of interferon beta and glatiramer acetate response: a review of the literature. *Autoimmun Rev* 2014; 13: 178-86.

- Marciano-Cabral F, Cabral G. The immune response to *Naegleria fowleri* amebae and pathogenesis of infection. *FEMS Immunol Med Microbiol* 2007; 51: 243-59.
- McGregor DP. Discovering and improving novel peptide therapeutics. *Curr Opin Pharmacol* 2008; 8: 616-9.
- Rabablert J, Tiewcharoen S, Junnu V. ITS and pB2.5 gene expression of *Naegleria fowleri* in drug resistance. *Health* 2011; 3: 529-33.
- Reddy KVR, Yedery RD, Aranha C. Antimicrobial peptides: premises and promises. *Int J Antimicrob Agents* 2004; 24: 536-47.
- Renault CA, Traore A, Machekano RN, Israel-ski DM. Validation of microcapillary flow cytometry for community- based CD4+ T lymphocyte enumeration in remote Burkina Faso. *Open AIDS J* 2010; 4: 171-5.
- Réveiller FL, Cabanes PA, Marciano-Cabral F. Development of a nested PCR assay to detect the pathogenic free living amoeba *Naegleria fowleri*. *Parasitol Res* 2002; 88: 443-50.
- Shin HJ, Cho MS, Jung SU, Kim HI, Park S, Kim HJ. Molecular cloning and characterization of a gene encoding a 13.1 kDa antigenic protein of *Naegleria fowleri*. *J Eukaryot Microbiol* 2001; 48: 713-7.
- Soltow SM, Brenner GM. Synergistic activities of azithromycin and amphotericin B against *Naegleria fowleri* in vitro and in a mouse model of primary amebic meningoencephalitis. *Antimicrob Agents Chemother* 2007; 51: 23-7.
- Stering TR, Merz WG. Resistance to amphotericin B: emerging clinical and microbiological patterns. *Drug Resist Updat* 1998; 1: 161-5.
- Tiewcharoen S, Malainual N, Junnu V, Chetanachan P, Rabablert J. Cytopathogenesis of *Naegleria fowleri* Thai strains for cultured human neuroblastoma cells. *Parasitol Res* 2008; 102: 997-1000.
- Tiewcharoen S, Rabablert J, Worawirunwong D, Pratumrikajorn T, Limsangurai S, Junnu V. Activity of chlorpromazine on *nfa1* and *Mp2CL5* genes of *Naegleria fowleri* trophozoites. *Health* 2011; 3: 166-71.
- Tiewcharoen S, Rabablert J, Roytrakul S, Wiyawuth W, Junnu V, Kumchol C. Differentially expressed genes of *Naegleria fowleri* during exposure to human neuroblastoma cells. *Asian Biomed* 2012; 6: 909-15.
- Vargas-Zepeda J, Gómez-Alcalá AV, Vásquez-Morales JA, Licea-Amaya L, De Jonckheere JF, Lares-Villa F. Successful treatment of *Naegleria fowleri* meningoencephalitis by using intravenous amphotericin B, fluconazole and rifampicin. *Arch Med Res* 2005; 36: 83-6.
- Visvesvara GS. Free-living amebae as opportunistic agents of human disease. *J Neuroparasitol* 2010;1. doi: 10.403/jnp/N100802.
- Visvesvara GS, De Jonckheere JF, Sriram R, Daft B. Isolation and molecular typing of *Naegleria fowleri* from the brain of a cow that died of primary amebic meningoencephalitis. *J Clin Microbiol* 2005; 43: 4203-4.
- Wilcox S. The new antimicrobials: cationic peptides. *Bio Teach J* 2004; 2: 88-91.
- Yang ST, Yub Shin SY, et al. Conformation-dependent antibiotic activity of tritrypticin, a cathelicidin-derived antimicrobial peptide. *Biochem Biophys Res Commun* 2002; 296: 1044-50.
- Yang ST, Kim JI, Shin SY. Effect of dimerization of a beta-turn antimicrobial peptide, PST13-RK, on antimicrobial activity and mammalian cell toxicity. *Biotechnol Lett* 2009; 31: 233-7.
- Yoder JS, Eddy BA, Visvesvara GS, Capewell L, Beach MJ. The epidemiology of primary amoebic meningoencephalitis in the USA, 1962-2008. *Epidemiol Infect* 2010; 138: 968-75.
- Yoder JS, Straif-Bourgeois S, Roy SL, et al. Primary amebic meningoencephalitis deaths associated with sinus irrigation using contaminated tap water. *Clin Infect Dis* 2012; 55: 79-85.