



*Supplement of*

**Measurement report: Assessing the impacts of emission uncertainty on aerosol optical properties and radiative forcing from biomass burning in peninsular Southeast Asia**

**Yinbao Jin et al.**

*Correspondence to:* Yiming Liu (liuym88@mail.sysu.edu.cn) and Qi Fan (eesfq@mail.sysu.edu.cn)

The copyright of individual parts of the supplement might differ from the article licence.

# Supplement




Figure S1. WRF-Chem uses land use classification data from 24 USGS classifications.




Figure S2. Time series of the observed (green lines) and simulated (red lines) daily average of  $PM_{2.5}$  concentrations from WRF-Chem with FINN 1.5 in the 23 cities in Table S6 during March 2019.





Figure S3. Satellite-AERONET linear regression, where (a)-(h) denote MODIS combined, MODIS DB, MODIS DT, MYDIS combined, MYDIS DB, MYDIS DT, VIIRS DB, VIIRS DT different satellite inversion AOD (550 nm) products compared with AERONET, (i) is Himawari-8 AOD (500 nm) vs. AERONET. The number of samples ( $N$ ),  $R^2$  correlation coefficient squared, and linear regression function and RMSE are also labeled in the upper left corner of each figure, and the lower right corner characterizes the expected values.



**Figure. S4. (a) The frequency distribution of six aerosol types within an 8 km altitude over the PSEA region in March 2019 (the different colors represent the six aerosol types). (b) 48-hour mean backward trajectory air mass (5500 m) at the Laos site (20.93,101.41) in March 2019, where the red boundary is the PSEA region.**

**Table S1. Emission factors (g kg<sup>-1</sup>) for species emitted from different types of biomass burning in GFED.**

Species	Sava	Borf	Temf	Defo	Peat	Agri
CO	63	127	88	93	210	102
NO <sub>x</sub>	3.9	0.9	1.92	2.55	1	3.11
SO <sub>2</sub>	0.48	1.1	1.1	0.4	0.4	0.4
BIGALK	0.055	0.349	0.225	0.072	0.072	0.34
BIGENE	0.133	0.385	0.369	0.267	0.267	0.333
C <sub>2</sub> H <sub>4</sub>	0.82	1.42	1.17	1.06	2.57	1.46
C <sub>2</sub> H <sub>5</sub> OH	0.024	0.055	0.1	0.037	0.037	0.035
C <sub>2</sub> H <sub>6</sub>	0.66	1.79	0.63	0.71	0.71	0.91
C <sub>3</sub> H <sub>8</sub>	0.1	0.44	0.22	0.126	0.126	0.28
C <sub>3</sub> H <sub>6</sub>	0.79	1.13	0.61	0.64	3.05	0.68
CH <sub>2</sub> O	0.73	1.86	2.09	1.73	1.4	2.08
CH <sub>3</sub> CHO	0.84	0.81	1.21	2.26	1.16	1.8
CH <sub>3</sub> COCH <sub>3</sub> (C <sub>3</sub> H <sub>6</sub> O)	0.47	1.59	0.76	0.63	0.91	0.71
CH <sub>3</sub> OH	1.18	2.82	1.74	2.43	8.46	3.29
MEK	0.181	0.22	0.13	0.5	0.5	0.9
TOLUENE	0.27	1.626	0.54	0.697	4.36	0.451
NH <sub>3</sub>	0.52	2.72	0.84	1.33	1.33	2.17
NO <sub>2</sub>	2.92	0.67	1.44	1.91	0.75	2.33
Open/BIGALD(C <sub>5</sub> H <sub>6</sub> O <sub>2</sub> )	0.02	0.01	0.01	0.01	0	0.01
C <sub>10</sub> H <sub>16</sub>	0.081	2.003	2.003	0.15	0.15	0.005
CH <sub>3</sub> COOH	3.55	4.41	2.13	3.05	8.97	5.59
CRESOL	0.44	0.85	0.07	0.17	0	0.6
GLYALD(HOCH <sub>2</sub> CHO)	0.25	0.86	0.86	0.74	0.74	0.71
Mgly/CH <sub>3</sub> COCHO	0.73	0.73	0.73	0.73	0.73	0.73
GLY	0.33	0.59	0.54	0.5	1.3	0.24
ACETOL/HYAC	1.01	0.77	8.03	0.55	0.64	0
ISOP(C <sub>5</sub> H <sub>8</sub> )	0.039	0.15	0.099	0.13	1.38	0.38
MACR	0	0	0	0.08	0	0
MVK	0	0	0	0.2	0	0
OC	2.62	9.6	9.6	4.71	6.02	2.3
BC	0.37	0.5	0.5	0.52	0.04	0.75
PM <sub>10</sub>	7.2	18.4	16.97	18.5	0	7.02
PM <sub>2.5</sub>	7.17	15.3	12.9	9.1	9.1	6.26

Note: Sava: Savanna, grassland, and shrubland fires; Borf: Boreal forest fires; Temf: Temperate forest fires; Defo: Tropical deforestation & degradation; Peat: Peat fires; Agri: Agricultural waste burning. Compared to the FINNs scheme, the missing compounds and aerosols were added based on the methodology of Akagi et al. (2011) and Heil A. (2020).

**Table S2. Emission factors (g kg<sup>-1</sup>) for species emitted from different types of biomass burning in FINN1.5.**

Species	Savanna	Tropical	Temperate	Agriculture	Boreal	Shrublands
CO	59	92	102	111	118	68
NO	0.38	0.74	0.26	0.09	0.70	0.74
SO <sub>2</sub>	0.48	0.45	1	0.4	1	0.68
BIGALK	0.02	0.13	0.11	0.09	0.16	0.42
BIGENE	0.45	0.52	0.22	0.37	0.35	0.63
C <sub>2</sub> H <sub>4</sub>	2.27	1.38	1.11	1.08	1.62	2.30
C <sub>2</sub> H <sub>5</sub> OH	0.02	0.01	0.01	0.01	0.01	0.02
C <sub>2</sub> H <sub>6</sub>	0.82	0.82	0.29	0.43	1.63	1.01
C <sub>3</sub> H <sub>8</sub>	0.18	0.10	0.10	0.08	0.13	0.37
C <sub>3</sub> H <sub>6</sub>	0.43	0.56	0.26	0.38	0.76	0.77
CH <sub>2</sub> O	2.12	2.08	1.33	1.84	1.46	2.23
CH <sub>3</sub> CHO	1.03	1.27	0.38	3.05	0.67	0.96
CH <sub>3</sub> COCH <sub>3</sub>	0.22	0.39	0.20	0.83	0.20	0.71
CH <sub>3</sub> OH	1.92	2.60	1.51	2.11	2.50	2.49
MEK	1.31	0.85	0.41	0.79	1.64	1.16
TOLUENE	1.16	2.06	0.61	1.07	1.30	1.30
NH <sub>3</sub>	0.49	0.76	1.5	2.3	3.5	1.2
NO <sub>2</sub>	3.2	3.6	2.7	3.9	3	1.4
Open/BIGALD	0.02	0.01	0.01	0.01	0.01	0.02
C <sub>10</sub> H <sub>16</sub>	0.01	0.04	0.03	0.00	0.04	0.01
CH <sub>3</sub> COOH	2.08	1.87	0.53	2.19	1.80	1.24
CRESOL	0.44	0.17	0.07	0.60	0.85	0.00
GLYALD	0.5	0.79	0.28	1.68	0.25	1.39
Mgly/CH <sub>3</sub> COCHO	0.81	0.37	0.17	0.19	0.28	0.86
GLY	0.81	0.37	0.17	0.19	0.28	0.86
ACETOL/HYAC	1.01	0.55	8.03	0.00	0.77	0.00
ISOP	0.05	0.07	0.03	0.60	0.14	0.03
MACR	0.0	0.08	0.0	0.0	0.0	0.0
MVK	0.0	0.20	0.0	0.0	0.0	0.0
OC	2.6	4.7	9.2	3.3	7.8	6.6
BC	0.37	0.52	0.56	0.69	0.2	0.5
PM <sub>2.5</sub>	5.4	9.7	13	5.8	13	9.3

**Table S3. Emission factors (g kg<sup>-1</sup>) for species emitted from different types of biomass burning in FINN2.5.**

Species	Savanna	Tropical	Temperate	Agriculture	Boreal	Shrublands	
CO	63	93	122	91	111	67	
NO	2.16	0.9	0.95	1.18	0.83	0.77	
SO <sub>2</sub>	0.9	0.4	1.1	0.4	1	0.68	
BIGALK	0.156	0.219	0.415	0.246	1.821	0.644	
BIGENE	1.467	0.662	1.393	0.674	0.627	1.274	
C <sub>2</sub> H <sub>4</sub>	1.218	1.505	1.930	1.412	1.407	2.886	
C <sub>2</sub> H <sub>5</sub> OH	0.00	0.00	0.066	0.0	0.023	0.055	
C <sub>2</sub> H <sub>6</sub>	0.859	0.939	0.611	0.673	1.168	0.641	
C <sub>3</sub> H <sub>8</sub>	0.09	0.114	0.149	0.142	0.194	0.561	
C <sub>3</sub> H <sub>6</sub>	0.647	0.603	0.487	0.457	0.499	0.557	
CH <sub>2</sub> O	1.532	2.299	2.181	1.716	1.361	2.285	
CH <sub>3</sub> CHO	1.037	1.404	0.758	0.929	0.416	0.792	
CH <sub>3</sub> COCH <sub>3</sub>	0.201	0.433	0.297	0.162	0.242	0.242	
CH <sub>3</sub> OH	1.451	3.031	1.744	2.328	1.608	1.650	
MEK	0.37	0.666	0.274	0.387	0.104	0.286	
TOLUENE	0.457	0.769	0.605	0.375	1.327	0.531	
NH <sub>3</sub>	0.56	1.3	2.47	2.12	1.8	1.2	
NO <sub>2</sub>	3.22	3.6	2.34	2.99	0.63	2.58	
Open/BAZLD	0.791	0.12	0.298	0.325	0.166	0.272	
C <sub>10</sub> H <sub>16</sub>	APIN	0.009	0.0	0.261	0.01	0.259	0.053
	BPIN	0.0	0.0	0.008	0.0	0.209	0.004
CH <sub>3</sub> COOH	2.371	2.029	1.292	2.349	1.36	1.353	
CRESOL	0.059	0.0	0.059	0.074	0.04	0.058	
GLYALD	0.390	1.886	0.210	0.800	0.233	0.128	
MGLY(CH <sub>3</sub> COCHO)	0.347	0.0	0.135	0.171	0.09	0.094	
GLY	0.347	0.0	0.135	0.171	0.09	0.094	
ACETOL/ HYAC	0.309	0.609	0.223	1.548	0.149	0.118	
ISOP	0.069	0.029	0.129	0.062	0.085	0.138	
MACR	0.0	0.222	0.113	0.0	0.024	0.147	
MVK	0.317	0.222	0.247	0.193	0.087	0.301	
OC	2.6	4.7	7.6	2.66	7.8	3.7	
BC	0.37	0.52	0.56	0.51	0.2	1.31	
PM <sub>10</sub>	7.2	18.5	16.97	7.02	18.4	11.4	
PM <sub>2.5</sub>	7.17	9.9	15	6.43	18.4	7.1	



**Table S4. Emission factors (g kg<sup>-1</sup>) for species emitted from different types of biomass burning in GFAS.**

Species	Savanna	Tropical	Temperate	Agriculture
CO	61	101	106	92
NO <sub>x</sub>	2.1	2.3	3.4	2.3
SO <sub>2</sub>	0.37	0.71	1.0	0.37
BIGALK	0.13	0.17	0.29	0.41
BIGENE	0.32	0.51	0.47	0.28
C <sub>2</sub> H <sub>4</sub>	0.84	1.5	1.2	1.3
C <sub>2</sub> H <sub>5</sub> OH	0.018	0.018	0.018	0.018
C <sub>2</sub> H <sub>6</sub>	0.32	1.1	0.72	1.2
C <sub>3</sub> H <sub>8</sub>	0.087	0.54	0.27	0.16
C <sub>3</sub> H <sub>6</sub>	0.34	0.76	0.57	0.57
CH <sub>2</sub> O	1.06	2.2	2.2	2.1
CH <sub>3</sub> CHO(C <sub>2</sub> H <sub>4</sub> O)	0.5	2.3	0.98	2.8
CH <sub>3</sub> COCH <sub>3</sub> (C <sub>3</sub> H <sub>6</sub> O)	0.48	0.63	0.67	1.1
CH <sub>3</sub> OH	1.5	3.0	1.9	3.7
MEK	0.37	0.666	0.274	0.387
TOLUENE(C <sub>7</sub> H <sub>8</sub> )	0.18	0.24	0.40	0.18
NH <sub>3</sub>	0.90	0.93	1.6	1.6
NO <sub>2</sub>	1.575	1.725	2.55	1.725
Open/BAZLD	0.791	0.12	0.298	0.325
C <sub>10</sub> H <sub>16</sub>	0.009	0.0	0.269	0.01
CH <sub>3</sub> COOH	2.371	2.029	1.292	2.349
CRESOL	0.059	0.0	0.059	0.074
GLYALD	0.390	1.886	0.210	0.800
MGLY(CH <sub>3</sub> COCHO)	0.347	0.0	0.135	0.171
GLY(CH <sub>3</sub> COCHO)	0.347	0.0	0.135	0.171
ACETOL/ HYAC	0.309	0.609	0.223	1.548
ISOP(C <sub>5</sub> H <sub>8</sub> )	0.026	0.22	0.11	0.40
MACR	0.0	0.222	0.113	0.0
MVK	0.317	0.222	0.247	0.193
OC	3.2	4.3	9.1	4.2
BC	0.46	0.57	0.56	0.42
PM <sub>10</sub>	7.2	18.5	16.97	7.02
PM <sub>2.5</sub>	4.9	9.1	13.8	8.3

Note: Compared to the FINNs scheme, the missing compounds and aerosols were added based on the methodology of Andreae and Merlet (2001;2019).

**Table S5. Missing data supplementation of FEER, QFED and IS4FIRES emission inventories based on the proportionality approach when WRF-Chem uses the MOZART-MOSAIC chemical mechanism.**

Species	FEER	QFED	IS4FIRES
CO	NM	NM	9.18
NO <sub>x</sub>	NM	NM	0.29
SO <sub>2</sub>	NM	NM	0.05
BIGALK	0.001	0.001	0.001
BIGENE	0.0041	0.0041	0.0041
C2H4	0.0109	0.0109	0.093
C2H5OH	0.0001	0.0001	0.0001
C2H6	NM	NM	0.2520
C3H8	NM	NM	0.0270
C3H6	NM	NM	0.059
CH2O	NM	NM	0.18
CH3CHO	0.01	NM	0.14
CH3COCH3	NM	NM	0.07
CH3OH	NM	0.0206	0.27
MEK	NM	NM	0.0067
TOLUENE	0.0163	0.0163	0.0163
NH <sub>3</sub>	NM	NM	0.1
NO <sub>2</sub>	0.75*NO <sub>x</sub>	NM	0.08
Open	0.0001	0.0001	0.0001
C10H16	0.0003	0.0003	0.0003
CH3COOH	0.0148	0.0148	0.0148
CRESOL	0.0013	0.0013	0.0013
GLYALD	0.0062	0.0062	0.0062
MGLY	0.0029	0.0029	0.0029
GLY	0.0029	0.0029	0.0029
ACETOL	0.0044	0.0044	0.0044
ISOP	0.0006	0.0006	0.03
MACR	0.0006	0.0006	0.0006
MVK	0.0016	0.0016	0.0016
OC	NM	NM	0.373
BC	NM	NM	0.054
PM <sub>10</sub>	0.3*PM <sub>2.5</sub>	NM	0.3
PM <sub>2.5</sub>	NM	NM	NM

Notes: FEER, QFED emission inventories compared to the MOZART-MOSAIC mechanism missing data were supplemented through a combination of methodology of Jose et al. (2017), Andreae and Merlet (2001;2019) and the actual FINN data (March 2019 over PESA), where CO was used as a scaling factor. The IS4FIRES was supplemented with other substances through a combination of methodology of Jose et al. (2017), Andreae and Merlet (2001;2019), Baró et al. (2021), and Wiedinmyer et al. (2011) using PM<sub>2.5</sub> and CO as scaling factors. NO<sub>2</sub> using 0.75\*NO<sub>x</sub>. PM<sub>10</sub> using 0.3\*PM<sub>2.5</sub>. NM: No missing data.

**Table S6. Comparison statistics of meteorological variables simulated by WRF-Chem with FINN1.5 scheme and observation stations**

Stations	Latitude(°N)	Longitude(°E)	variables	MB	RMSE	IOA	R
56964	22.47	100.58	T(°C)	-0.85	2.56	0.91	0.91
			RH (%)	5.17	17.67	0.83	0.7
			WS(m/s)	1.76	2.35	0.64	0.51
56969	21.28	101.35	T(°C)	-0.95	2.33	0.91	0.95
			RH (%)	-13.58	20.69	0.82	0.82
			WS(m/s)	2.69	3.08	0.61	0.52
59431	22.38	108.13	T(°C)	0.97	3.71	0.79	0.65
			RH (%)	0.78	9.51	0.8	0.62
			WS(m/s)	1.85	2.71	0.66	0.31
59644	21.27	109.08	T(°C)	1.29	3.87	0.71	0.54
			RH (%)	7.62	11.2	0.72	0.4
			WS(m/s)	2.91	3.78	0.6	0.30
59758	20	110.15	T(°C)	-2.47	3.55	0.72	0.65
			RH (%)	11.33	15.09	0.74	0.66
			WS(m/s)	1.72	2.59	0.65	0.33
59287	23.13	113.29	T(°C)	1.0	2.75	0.83	0.79
			RH (%)	-8.01	15.03	0.77	0.65
			WS(m/s)	0.74	1.69	0.69	0.32
59663	21.86	111.96	T(°C)	-0.43	2.33	0.82	0.73
			RH (%)	-0.18	8.16	0.81	0.65
			WS(m/s)	0.55	1.82	0.71	0.33
45011	22.15	113.59	T(°C)	0.35	2.13	0.78	0.65
			RH (%)	0.82	7.64	0.79	0.65
			WS(m/s)	1.36	2.27	0.67	0.31
45007	22.3	113.91	T(°C)	-1.37	2.63	0.78	0.68
			RH (%)	7.66	12.62	0.74	0.66
			WS(m/s)	-2.09	3.16	0.71	0.40
48855	16.04	108.19	T(°C)	-2.76	3.35	0.72	0.83
			RH (%)	2.4	9.67	0.81	0.74
			WS(m/s)	1.62	2.12	0.66	0.62
48930	19.87	102.16	T(°C)	-3.42	4.61	0.82	0.85
			RH (%)	-7.51	18.09	0.82	0.72
			WS(m/s)	2.43	3.01	0.60	0.30
48327	18.76	98.96	T(°C)	-3.45	4.29	0.83	0.91
			RH (%)	-2.45	13.63	0.82	0.74
			WS(m/s)	1.86	2.54	0.6	0.30
58968	25.03	121.51	T(°C)	-1.71	2.56	0.82	0.85
			RH (%)	8.06	13.95	0.76	0.61
			WS(m/s)	1.61	2.48	0.71	0.45

**Table S7. Comparison statistics of PM<sub>2.5</sub> simulated by WRF-Chem with FINN1.5 and air quality stations**

Cites	latitude(°N)	longitude(°E)	MB	RMSE	IOA	R
Jing Hong	22	100.79	-11.2	38.44	0.82	0.75
Pu ER	22.76	100.98	11.92	26.32	0.8	0.84
Wen Shan	23.35	104.25	-2.77	24.97	0.7	0.41
Hong He Zhou	23.36	103.37	-13.24	23.04	0.77	0.62
Mao Ming	21.46	111.02	-6.69	12.35	0.72	0.51
Yang Jiang	21.85	111.95	-12.09	18.47	0.71	0.40
Guang Zhou	23.55	113.58	9.65	16.06	0.67	0.50
Hai Kou	20.05	110.32	6.53	11.34	0.66	0.45
San Ya	18.24	109.5	-5.55	10.62	0.67	0.69
Zhu Hai	22.42	113.62	13.6	24.72	0.69	0.51
Hongkong	22.32	114.25	82.21	88.43	0.54	0.43
Macao	22.15	113.56	-17.77	25.36	0.7	0.48
TaiWan	25.013	121.511	-30.37	35.23	0.66	0.38
Saraburi	14.68	100.87	38.57	44.28	0.6	0.44
Ratchaburi	13.52	99.81	40.18	52.03	0.54	0.57
Chiang Mai	18.79	98.99	9.54	34.37	0.71	0.45
Lampang	18.25	99.76	24.0	37.95	0.63	0.35
Nakhon Sawan	15.68	100.11	-2.5	26.31	0.73	0.45
Chiang Rai Mueang	19.9	99.82	-33.81	56.82	0.74	0.64
Sa Kaeo	13.69	102.51	-43.66	56.3	0.68	0.36
Kanchanaburi	14.02	99.53	26.98	43.48	0.58	0.47
Nan	18.78	100.77	30.32	46.74	0.64	0.36
Hochi	10.78	106.7	-15.6	21.47	0.75	0.68

**Table S8. Statistical metrics for observation-model comparisons**

Mathematical formulas	Number
$MB = \frac{1}{N} \sum_{i=1}^N (P_i - O_i)$	(1)
$RMSE = \left[ \frac{1}{N} \sum_{i=1}^N (P_i - O_i)^2 \right]^{\frac{1}{2}}$	(2)
$IOA = 1 - \frac{\sum_{i=1}^N (P_i - O_i)^2}{\sum_{i=1}^N ( P_i - \bar{P}  +  O_i - \bar{O} )^2}$	(3)

Note: Where  $P_i$  and  $O_i$  are the predicted and observed T<sub>2</sub>, RH<sub>2</sub>, WS<sub>10</sub>, and PM<sub>2.5</sub>, respectively. N is the total number of the predictions used for comparisons, and  $\bar{P}$  and  $\bar{O}$  represents the average of the prediction and observation, respectively.

**Table S9. WRF-Chem AAOD at 500 nm vs. AERONET in HAAOD (97-110°E, 15-22.5°N) during the wildfire period, where HAAOD includes Laos, Doi Ang Khang, Fang, Nong Khai (all statistical days in 4 stations are greater than 10 days).**

Stations	Variables	BB emission inventories							
		GFED	FINN1.5	FINN2.5MOS	FINN2.5 MOSVIS	GFAS	FEER	QFED	IS4FIRES
Laos	R	0.77	0.64	0.59	0.65	0.58	0.69	0.65	0.71
	IOA	0.63	0.67	0.71	0.71	0.63	0.65	0.63	0.64
Doi Ang Khang	R	0.07	0.37	0.31	0.64	NA	0.24	NA	0.14
	IOA	0.64	0.7	0.69	0.75	0.64	0.65	0.64	0.63
Fang	R	0.59	0.77	0.77	0.79	0.55	0.7	0.64	0.67
	IOA	0.66	0.7	0.76	0.74	0.65	0.66	0.66	0.67
Nong Khai	R	0.35	0.37	0.52	0.74	0.31	0.35	0.34	0.43
	IOA	0.7	0.72	0.76	0.78	0.68	0.73	0.7	0.73

Notes: The blank data in the table are not counted because the statistical correlation coefficient is negative and the value is particularly small, which we consider an error.

NA: Not available.

## Reference

- Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmos. Chem. Phys.*, 19, 8523-8546, 10.5194/acp-19-8523-2019, 2019.
- Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, 15, 955-966, <https://doi.org/10.1029/2000GB001382>, 2001.
- Baró, R., Maurer, C., Brioude, J., Arnold, D., and Hirtl, M.: The Environmental Effects of the April 2020 Wildfires and the Cs-137 Re-Suspension in the Chernobyl Exclusion Zone: A Multi-Hazard Threat, 12, 467, 2021.
- Heil A., B. I.: ESA CCI ECV Fire Disturbance: D5.1 Product Validation and Intercomparison Report, version 2.1, 2020.
- Jose, R. S., Pérez, J. L., González, R. M., Pecci, J., and Palacios, M.: Improving air quality modelling systems by using on-line wild land fire forecasting tools coupled into WRF/Chem simulations over Europe, *Urban Climate*, 22, 2-18, <https://doi.org/10.1016/j.uclim.2016.09.001>, 2017.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625-641, 10.5194/gmd-4-625-2011, 2011.