

1 **Supplementary Material:**

2 **Tropospheric ozone changes, radiative forcing and**
3 **attribution to emissions in the Atmospheric Chemistry and**
4 **Climate Model Inter-comparison Project (ACCMIP)**


5

6 **D.S. Stevenson¹, P.J. Young^{2,3}, V. Naik⁴, J.-F. Lamarque⁵, D.T. Shindell⁶, A.**
7 **Voulgarakis⁷, R. Skeie⁸, S. Dalsoren⁸, G. Myhre⁸, T. Berntsen⁸, G.A. Folberth⁹,**
8 **S.T. Rumbold⁹, W.J. Collins⁹, I.A. MacKenzie¹, R.M. Doherty¹, G. Zeng¹⁰, T. van**
9 **Noije¹¹, A. Strunk¹¹, D. Bergmann¹², P. Cameron-Smith¹², D.A. Plummer¹³, S.A.**
10 **Strode¹⁴, L. Horowitz¹⁵, Y.H. Lee⁶, S. Szopa¹⁶, K. Sudo¹⁷, T. Nagashima¹⁸, B.**
11 **Josse¹⁹, I. Cionni²⁰, M. Righi²¹, V. Eyring²¹, A. Conley⁵, K.W. Bowman²², O.**
12 **Wild²³**

13


14

1




2

3 **Figure S1:** Annual zonal mean ozone (ppbv) for the 1850s and 2000s. For scale see Figure 1.




1

2 Figure S1 (continued)




1

2 **Figure S2:** Annual mean tropospheric column ozone (DU), for the 1850s and 2000s, using
 3 the MASKZMT. For scale see Figure 2.




1

2 Figure S2 (continued)




1
 2 **Figure S3:** Changes (2000s-1850s) in annual zonal mean ozone (ppbv) and tropospheric
 3 column ozone (both masked using MASKZMT). See Figure 3 for scales.



1


2 Figure S3 (continued)




1

2 **Figure S4a:** Total ozone RFs for all models, masked using MASKZMT.


IGAC/SPARC database 2000s-1850s O3t RF (315) mW/m²



IGAC/SPARC database 2000s-1850s O3t SW RF (60) mW/m²



IGAC/SPARC database 2000s-1850s O3t LW RF (255) mW/m²




1

2 **Figure S4b:** Annual mean tropospheric ozone total, SW, and LW RFs for the IGAC/SPARC
3 ozone dataset, as used in Cionni et al (2011). Compare to the multi-model mean in Figure 4
4 and also compare to Figure 15a of Cionni et al. (2011), which shows the total RF calculated
5 with an earlier version of the E-S radiation code, and finds a total RF 27% lower.


6

1



2


3 **Figure S5:** Normalised RFs for MASKZMT. See Figure 5 for scale.



1

2 **Figure S6:** An example of one model's results (model B) for the attribution experiments. Left
 3 hand side shows contributions to changes in zonal annual mean ozone, right hand side shows
 4 contributions to change in annual mean tropospheric ozone column. Referring to the
 5 experiment numbers in Table 3, rows from the top show: experiment #1-#0 (all components);
 6 #1-#2 (CH₄); #1-#3 (NO_x); #1-#4 (CO); #1-#5 (NMVOC).

7




1

2 **Figure S7:** Radiative forcings for model B, from the attribution experiments. Left-hand plot
 3 shows total 2000s-1850s (#1-#0); middle shows the CH₄ component (#1-#2); and the right-
 4 hand plot shows the NO_x component (#1-#3). The CO and NMVOC components are
 5 significantly less (38 and 37 mWm⁻², respectively) and are not shown.


6

1



2

3 **Figure S8:** As Figure 6, but showing the impact of climate change (scenario RCP8.5) on
4 ozone from 1850s to the (a) 2030s; (b) 2100s.



1

2 Figure S8 continued


3

1 **1 Radiative forcing of increased CO₂ due to emissions of CH₄, CO and**
2 **NMVOCs from fossil sources**

3

4 Emissions of CH₄, CO and NMVOCs from fossil sources leads to increased levels of CO₂ in
5 the atmosphere. Assessing the emission-based impact on radiative forcing these CO₂
6 contributions need be attributed to the emissions of the source gases. Here we have used the
7 anthropogenic emissions from Lamarque et al. (2010), covering 1850 onwards for CO and
8 NMVOCs, while methane emissions were taken from the emission data recommended for
9 CMIP5 use (<http://www.iiasa.ac.at/web-apps/tnt/RcpDb/dsd?Action=htmlpage&page=download>) . Emissions from the power plants,
10 energy conversion, extraction and distribution sectors are regarded as fossil, while other
11 sectors are assumed to be non-fossil.
12

13



14

15 **Figure S9:** Historic emissions of CH₄, CO and NMVOCs (all in Tg/yr) from
16 fossil/anthropogenic sources.

17

18 The oxidation of CH₄, CO and NMVOCs in the atmosphere leads to an atmospheric source of
19 CO₂. For methane the lifetime in the atmosphere is long enough to allow for a non-negligible
20 fraction of the historical emissions to be left in the atmosphere. Using a lifetime for methane
21 of 9.6 years (IPCC, 2001) we find that 12% of the methane has not yet been oxidised to CO₂.

22

1 From the emissions and the atmospheric lifetime of CH₄ we calculate the corresponding
 2 atmospheric source of CO₂ as a function of time. For the NMVOCs we have assumed an
 3 average carbon content of 80% by mass. We then calculate the resulting development in the
 4 CO₂ concentrations using the impulse response function for CO₂ (IRF¹) given in IPCC (2007).
 5 The change in the mixing ratio of CO₂ (X_{CO2}(t)) is given by

$$X_{CO_2}(t) = \int_0^t Em_{CO_2}(t') \cdot IRF(t - t') dt'$$

6
 7
 8
 9 The contribution to atmospheric CO₂ is given in figure 2a.

10
 11 The radiative forcing due to the CO₂ from the fossil/anthropogenic emissions of methane, CO
 12 and NMVOCs are calculated by the simple parameterization given in IPCC (2001)


$$RF = 5.35 \cdot \ln \left[\frac{X}{X_0} \right]$$

13
 14
 15
 16 The calculations are done with X₀ = 278 ppm. The radiative forcings are shown in figure 2b.
 17 In 2010 the RFs are 18, 87 and 33 mW m⁻² for emissions of methane, CO and NMVOC
 18 respectively.

19

20

¹ $IRF(t) = \alpha_0 + \sum_{i=1}^3 \alpha_i \cdot e^{-t/\tau_i}$



1 **Figure S10:** Contribution from fossil/anthropogenic emissions of CH₄, CO and NMVOCs to
 2 atmospheric CO₂ and radiative forcing.
 3