

1 **Supplemental material to:**

2 **Chemistry of new particle growth in mixed urban and biogenic**
3 **emissions - Insights from CARES**

4
5 A. Setyan^{1,#}, C. Song², M. Merkel³, W. B. Knighton⁴, T. B. Onasch⁵, M. R. Canagaratna⁵, D.
6 R. Worsnop^{5,6}, A. Wiedensohler³, J. E. Shilling², Q. Zhang^{1,*},

7
8 ¹ Department of Environmental Toxicology, University of California, Davis, CA 95616,
9 United States

10 ² Atmospheric Sciences and Global Change Division, Pacific Northwest National Laboratory,
11 Richmond, WA 99352, United States

12 ³ Leibniz Institute for Tropospheric Research, 04318 Leipzig, Germany

13 ⁴ Montana State University, Bozeman, MT 59717, United States

14 ⁵ Aerodyne Research Inc., Billerica, MA 01821, United States

15 ⁶ Department of Physics, University of Helsinki, FI-00014 Helsinki, Finland

16 [#] Now at: Département Chimie & Environnement, Ecole Nationale Supérieure des Mines de
17 Douai, 59508 Douai Cedex, France


18 ^{*} Corresponding author, Department of Environmental Toxicology, University of California,
19 Davis, CA 95616, United States, dkwzhang@ucdavis.edu; 530-752-5779

20

21 **Size distributions of ammonium and sulfate using high resolution data**


22 PToF data in AMS is usually used in unit mass resolution to determine size distributions of
23 species. However, during the present study, PToF data of ammonium was too noisy and not
24 usable to determine size distributions for short selected periods. Therefore, in the present case,
25 PToF data has been used in high resolution, in which ammonium fragments had satisfactory
26 signal to noise (S/N) ratios. First, 86 size bins recorded in the PToF mode and covering 40-
27 1400 nm (in D_{va}) have been grouped into 7 different size ranges in order to increase the S/N
28 ratio. Given that the PToF data processed in the PIKA software is without DC markers
29 applied, an eighth size range between 1400-2200 nm has been used as a background signal to
30 subtract the signals of the other size ranges (Fig. S1). Then, for each size range, average high
31 resolution mass spectra have been plotted, and the signals of the ammonium and sulfate
32 fragments having the best S/N ratios have been integrated. For that purpose, we chose NH_3^+
33 (m/z 17) for ammonium and SO^+ (m/z 48) for sulfate (Fig. S1). The scatterplots of NH_3^+ vs.
34 total ammonium and SO^+ vs. total sulfate (Fig. S2) have then been used to reconstruct the
35 concentrations of ammonium and sulfate for each of the 7 size ranges. Finally, these results
36 have been used to reconstruct size distributions of the two species in Hz, and converted to
37 $\mu\text{g}/\text{m}^3$ by scaling the size distributions to the concentrations of these species in MS mode.

38 **Figure S1.** Average high resolution mass spectra between 14:00-15:00 during NPE days for
 39 particles in the range 250-400 nm (a, b), 1400-2200 nm (c, d), and top MS minus bottom MS
 40 (e, f).



41
 42


43 **Figure S2.** Scatterplots of (a) NH_3^+ vs. total ammonium, and (b) SO^+ vs. total sulfate. The
44 data fitting was performed using the orthogonal distance regression (ODR).



45

46


47 **Figure S3.** Diurnal patterns of (a) temperature, (b) relative humidity, and (c) broadband solar
48 radiation during new particle event (NPE) days and non-NPE days.



49

50


51 **Figure S4.** Diurnal patterns of (a) isoprene, (b) terpenes, (c) sum of methacrolein (MACR)
 52 and methyl vinyl ketone (MVK), (d) formaldehyde, (e) acetic acid, (f) acetaldehyde, (g)
 53 benzene, (h) toluene, (i) black carbon, (j) CO, (k) O₃, and (l) NO_x, and (m) during NPE and
 54 non-NPE days.



55

56

57 **Figure S5.** Time series of (a) broadband solar radiation, O₃ and NO_x, (b) Org, SO₄²⁻, NO₃⁻,
 58 NH₄⁺ and Cl⁻, (c) OA components, and (d) particle number concentration and condensation
 59 sink. Periods of T0 to T1 transport and northwesterly wind periods are shaded in pink and
 60 green, respectively.



61
 62
 63