

1 **Table S1** The input of organic compounds for MCM model (pmol mol⁻¹)^a

Species	PM _{2.5} < 75 µg m ⁻³	PM _{2.5} ≥ 75 µg m ⁻³
Methane	1900	1900
Formaldehyde	3.49	7.65
Acetaldehyde	2.36	5.89
Ethane	3.675	10.34
Propane	2.165	6.93
isobutane	0.815	2.52
n-Butane	1.135	3.58
isopentane	0.65	2.26
n-Pentane	0.465	1.51
2-methylpentane	0.235	0.77
n-Hexane	0.265	0.84
Ethylene	2.245	6.64
Propene	0.505	1.19
1-Butene	0.14	0.36
isoprene	0.085	0.12
Acetylene	1.945	7.82
Benzene	0.63	1.95
Toluene	0.925	2.84
Ethylbenzene	0.295	1.08
m/p-Xylene	0.605	2.03
o-Xylene	0.175	0.63
Acrolein	0.1	0.25
Propanal	0.31	0.66
Acetone	2.41	6
MTBE	0.2	0.71
Methylvinylketone	0.095	0.32
Methylethylketone	0.435	1.41
n-Hexanal	0.18	0.48
Chloromethane	0.32	0.91
Dichloromethane	1.895	4.4
Chloroform	0.395	1.31
1,2-Dichloroethane	0.5	1.91
1,2-Dichloropropane	0.35	1.46
1,1,2,2-Tetrachloroethane	0.315	0.32

2 ^a The mixing ratio of methane is from Wang et al. (2001). The mixing ratios of formaldehyde and
 3 acetaldehyde are from Rao et al. (2016). The mixing ratios of other organic compounds are from Wu et
 4 al. (2016)

5 **Reference**

6 Rao, Z., Chen, Z., Liang, H., Huang, L., and Huang, D.: Carbonyl compounds over urban Beijing:
 7 Concentrations on haze and non-haze days and effects on radical chemistry, *Atmos. Environ.*, 124,

- 8 207-216, 2016.
- 9 Wang, Y., Zhou, L., Wang, M., and Zheng, X.: Trends of atmospheric methane in Beijing,
10 Chemosphere, 3, 65-71, 2001.
- 11 Wu, R., Li, J., Hao, Y., Li, Y., Zeng, L., and Xie, S.: Evolution process and sources of ambient volatile
12 organic compounds during a severe haze event in Beijing, China, Sci. Total Environ., 560, 62-72,
13 2016.