

Interactive comment on “Effects of NO_x and seed aerosol on highly oxygenated organic molecules (HOM) from cyclohexene ozonolysis” by Meri Rätty et al.

Anonymous Referee #1

Received and published: 15 October 2020

This manuscript presents gas-phase measurements of highly oxygenated organic molecules (HOMs) derived from cyclohexene ozonolysis, made using a nitrate chemical ionization mass spectrometer (NO₃-CIMS), as a function of initial NO_x mixing ratio and ammonium bisulfate (ABS) seed concentration. Comparison of mass spectra for experiments with and without NO addition reveals a general decrease in the abundance of C₁₁–12H₁₆–22O₇–16 dimers, varied responses for C₃–6H₄–12O₃–10 monomers, and increases in both N-containing monomers (C₅–6H₇–11NO₄–10) and dimers (C₁₁–12H₁₇–20NO₉–14). Based on the observed uptake of compounds to the ABS seed, the authors propose that condensation is primarily controlled by the number of oxygen atoms present in each molecule. The measured gas-phase losses of

Printer-friendly version

Discussion paper



monomers, dimers, and N-containing compounds due to condensation are also compared to those predicted by two models designed to estimate volatilities of atmospheric oxidation products.

In many respects, this work is a replication of the recent study by Peräkylä et al. (2020), focusing on HOMs derived from ozonolysis of cyclohexene rather than α -pinene. Atmospheric interest in cyclohexene ozonolysis is as a surrogate for α -pinene, given the potential to derive generalizable mechanistic insight from its comparatively simpler symmetric structure. Although this manuscript has the potential to complement and extend the findings of Peräkylä et al. (2020), in its current form it presents few novel results and the discussion is lacking. In particular, molecular-level interpretation of the observed effects of NO_x on the abundance and distribution of cyclohexene-derived HOMs is limited and often overlooks key findings from recent papers by co-author Matti Rissanen on HOMs formation in the cyclohexene ozonolysis system. For these reasons, I recommend that publication be considered only after the comments detailed below are addressed.

Specific Comments

1. Line 207: Dependence of C₆H₈O₇ and C₆H₈O₉ on NO_x. Given that Rissanen et al. (2014) proposes structures and formation mechanisms for these HOMs based on quantum chemical calculations and experiments with isotopic labeling, “speculating the reason for this difference is difficult. . .” is a rather unsatisfying interpretation. A discussion that specifically addresses whether the observed NO_x trends are consistent with the Rissanen et al. (2014) structures/mechanisms would strengthen this section considerably. Notably, Rissanen et al. (2014) states that “it seems more likely that C₆H₈O₇ product is formed through bimolecular reactions of the intermediate peroxy radicals (i.e., by RO₂ + RO₂ reactions),” which is consistent with the observed decrease in the C₆H₈O₇ signal with increasing NO. Additionally, Rissanen et al. (2018) states that “at a low addition level, NO aids the C₆H₈O₈ HOM product formation by reactive alkoxy radical (RO) formation and illustrates the oxidation enhancing influence of gen-

[Printer-friendly version](#)[Discussion paper](#)

erating highly reactive RO radicals.” A similar role of RO reactions in the formation of the C₆H₈O₉ HOM may account for the observed increase in signal with increasing NO.

2. Line 184: Role of NO₂. Despite the statement that “only NO mainly has an impact on the radical chemistry,” NO₂ should also exert a considerable influence in this system via reaction with acylperoxy radicals [RC(O)O₂]. Rissanen et al. (2014) proposes that acylperoxy radicals (C₆H₉O₆ and C₆H₉O₈) are key intermediates in the formation of the major C₆H₈O₇ and C₆H₈O₉ HOMs, while Rissanen et al. (2018) implicates acylperoxy radicals in the production of C₁₂ dimers: “these observations imply the special importance of acylperoxy radicals in directing autoxidation phenomena.” Given the reported NO/NO_x ratio of ~3%, the role of NO₂ should be accounted for.

3. Kinetic Modeling. Interpretation of the observed NO_x dependences would greatly benefit from a kinetic box model, parameterized with known rate constants where available, calculated rate constants from Rissanen et al. (2014), and structure-activity relationships from Jenkin et al. (2019). For example, as a function of the initial NO mixing ratio: What fraction of O₃ reacts with C₆H₁₀ vs. NO? What fraction of the initially formed C₆H₉O₄-RO₂ undergoes unimolecular isomerization vs. reaction with NO or other RO₂? How much NO₃ is formed? What fraction of the C₆H₁₀ decay is due to reaction with NO₃ vs. O₃ vs. OH? What fraction of OH reacts with NO₂ vs. C₆H₁₀?

4. Figures 2 and 3. Molecular formulas should either be rotated 90 degrees and placed directly above the corresponding peaks or moved into the white space at the top of the figure and linked to peaks with arrows. The existing format is cluttered and difficult to interpret. Also, the y-axis should be labeled “normalized ion counts,” given the normalization to the reagent ion signals.

5. Figure 4. The discussion of Figure 4 is sparse and does not provide any information beyond that presented in relation to Figure 5. However, there are some interesting non-linear NO_x dependences displayed in Figure 4. For example, the C₁₂H₁₉NO₁₁ signal decreases between 8 and 10 ppb NO_x after increasing from 1 to 4 ppb NO_x, presum-

[Printer-friendly version](#)[Discussion paper](#)

ably due to suppression of RO₂ + RO₂ at sufficiently high NO. Additionally, the traces for C₆H₈O₇ and C₁₂H₂₀O₉ are effectively superposable, suggesting similar formation pathways. The discussion of this figure should be expanded to include interpretation of these trends.

6. Figures 5 and 6. The use of a color scale in Figure 5 that incorporates opacity makes it very difficult to determine if the degree of marker transparency for a given compound is due to the fit factor, relative signal change, or a combination of the two. Instead of transparency, perhaps different marker symbols (e.g., square, triangle, circle, diamond) could be used to denote different fit factor ranges (e.g., <0, 0.1-0.4, 0.5-0.7, 0.8-1.0). For consistency, the use of symbols to indicate fit factor ranges should also be applied to Figure 6.

7. Line 223: Dimers from O₃- and NO₃-Derived RO₂. The proposed formation of N-containing dimers via cross-reactions of O₃- and NO₃-derived RO₂ is quite interesting and possibly the first account of such chemistry. Several recent studies (Zhao et al. 2018, Kenseth et al. 2018, and Li et al. 2019) report on the analogous formation of dimers from cross-reactions of O₃- and OH-derived RO₂. A discussion of the current work as it relates to these studies and the potential importance of multi-oxidant chemistry would be useful.

8. Line 259: Model Adaptation. Rather than shifting the fit between FR and log₁₀(C*) reported for α-pinene in Peräkylä et al. (2020) by one unit to account for the difference in SOA concentration, which seems somewhat arbitrary, why not generate a new logistic fit using values for cyclohexene model compounds calculated by the ADCHAM model?

9. Line 282: Model-Measurement Agreement. Both models significantly overpredict the FR for compounds between m/z 250 and 300, including the major C₆H₈O₇ and C₆H₈O₉ HOMs. Was the FR overpredicted for all observed HOMs, suggesting an inability of the models to effectively capture HOMs volatilities? A more detailed discus-

[Printer-friendly version](#)[Discussion paper](#)

sion of the potential causes of the model-measurement disagreement (e.g., ability to accurately model contributions of key HOMs functionalities, such as hydroperoxides, to volatility) would be beneficial.

10. Line 292: Scaling Experimental Data. What assumptions were made in scaling the data from Peräkylä et al. (2020) to facilitate comparison with the results obtained in this work under different experimental conditions? Such a comparison seems somewhat contrived and the utility is not immediately clear, however, beyond stating that the scaled data are included in Figure 9 they are not discussed. The authors should either include a comparison of the elemental composition vs. volatility trends for cyclohexene- and α -pinene-derived HOMs or remove the Peräkylä et al. (2020) data from Figure 9. As an example, Figure 6 in Peräkylä et al. (2020) suggests that $m/z > 300$ is required for the FR of non-N-containing α -pinene compounds to drop below 50%, whereas Figure 8 in this work shows that for non-N-containing cyclohexene compounds $m/z > 250$ corresponds to FR values below 50%. A discussion of such differences would be informative.

Minor Comments

1. Line 29. Sentence beginning with “Through oxidation...” is awkward. Consider rephrasing.
2. Line 76. What is the detection limit of the NO_x analyzer?
3. Line 81. Sentence beginning with “Experiments with ammonium sulfate...” is unnecessary.
4. Figure 7. Why is there so much variability in the signal of the three species prior to ABS seed injection (~ 0.9 - 1.1)? What are the uncertainties in the measured signals of the detected species and their relative changes as a function of NO_x mixing ratio and ABS seed concentration?
5. Line 244. How were the modeled values “scaled to the range of observed remaining

[Printer-friendly version](#)[Discussion paper](#)

fractions”?

6. Line 250. A discussion of the suitability of using an activity coefficient of unity and the potential impacts of this assumption should be included.

7. Line 315. This sentence is contradictory. By definition, “increased termination of RO₂ radicals by NO” does not lead “to more alkoxy radicals” but rather to closed-shell alkyl nitrates. Radical propagation via reaction of RO₂ with NO will produce more RO radicals, which seems to be the intent of the sentence.

References

Jenkin, M. E., Valorso, R., Aumont, B., Rickard, A. R.: Estimation of rate coefficients and branching ratios for reactions of organic peroxy radicals for use in automated mechanism construction, *Atmos. Chem. Phys.*, 19, 7691–7717, 2019.

Kenseth, C. M., Huang, Y., Zhao, R., Dalleska, N. F., Hethcox, J. C., Stoltz, B. M., Seinfeld, J. H.: Synergistic O₃ + OH oxidation pathway to extremely low-volatility dimers revealed in b-pinene secondary organic aerosol, *Proc. Natl. Acad. Sci. U.S.A.*, 115, 8301–8306.

Li, X., Chee, S., Hao, J., Abbatt, J. P. D., Jiang, J., Smith, J. N.: Relative humidity effect on the formation of highly oxidized molecules and new particles during monoterpene oxidation, *Atmos. Chem. Phys.*, 19, 1555–1570, 2019.

Peräkylä, O., Riva, M., Heikkinen, L., Quéléver, L., Roldin, P., and Ehn, M.: Experimental investigation into the volatilities of highly oxygenated organic molecules (HOMs), *Atmos. Chem. Phys.*, 20, 649–669, 2020.

Rissanen, M. P.: NO₂ Suppression of Autoxidation – Inhibition of Gas-Phase Highly Oxidized Dimer Product Formation, *ACS Earth Space Chem.*, 2, 1211–1219, 2018.

Rissanen, M. P., Kurten, T., Sipilä, M., Thornton, J. A., Kangasluoma, J., Sarnela, N., Junninen, H., Jorgensen, S., Schallhart, S., Kajos, M. K., Taipale, R., Springer, M.,

Mentel, T. F., Ruuskanen, T., Petäjä, T., Worsnop, D. R., Kjærgaard, H. G., and Ehn, M.: The Formation of Highly Oxidized Multifunctional Products in the Ozonolysis of Cyclohexene, *J Am. Chem. Soc.*, 136, 15596–15606, 2014.

Zhao, Y., Thornton, J. A., Pye, H. O. T.: Quantitative constraints on autoxidation and dimer formation from direct probing of monoterpene-derived peroxy radical chemistry, *Proc. Natl. Acad. Sci. U.S.A.*, 115, 12142–12147.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2020-694>, 2020.

[Printer-friendly version](#)[Discussion paper](#)