



*Supplement of*

## **Influence of wind strength and direction on diffusive methane fluxes and atmospheric methane concentrations above the North Sea**

**Ingeborg Bussmann et al.**

*Correspondence to:* Ingeborg Bussmann ([ingeborg.bussmann@awi.de](mailto:ingeborg.bussmann@awi.de))

The copyright of individual parts of the supplement might differ from the article licence.

## 1 Supplement

Hydrographic parameters:

In the RV Ludwig Prandtl, the built-in FerryBox (Petersen, 2014) recorded data using a Teledyne RDI thermosalinograph (temperature and salinity), Aanderaa oxygen optode 4330 (oxygen concentration and saturation), Meinsberg pH electrode (pH values), and SCUFA submersible chlorophyll fluorometer (turbidity and fluorescence). The data were stored on board and automatically sent via a mobile phone connection to the Hereon database. The water flow in the FerryBox was approximately 12 L min<sup>-1</sup>. In the RV Mya II, the built in Ferrybox recorded data using a salinometer (FSI), and SBE38 Digital Oceanographic Thermometer (SBE38) for salinity and temperature. For pH measurements a Meinsberg pH Sensor (EGA150) was used and for oxygen an Aanderaa oxygen optode (4330). On the RV Uthörn and RV Littorina, a portable pocket FerryBox (4HJena, Germany) was used to record the hydrographic parameters using the following sensors: Seabird SBE45 thermosalinograph, Aanderaa oxygen optode 4835, Meinsberg pH electrode (EGA150), Seapoint Chlorophyll Fluorometer (SCF), and Seapoint Turbidity Meter. The water flow was between 3 and 4 L min<sup>-1</sup>. Data were saved once per minute.

Methane analysis:




Fig. S1: Regression lines with 95% confidence interval for the instruments on RV Mya and RV Littorina on Stern-5. The slope and  $r^2$  for the other calibrations were: 39.51; 20.17; 21.77 ppm to nM for Stern-3 Uthörn, Littorina and Prandtl with the respective  $r^2$  of 0.97; 0.96 and 0.97. For Stern-5 the slopes were 22.65; 26.92; 23.42 for Stern-5 Mya, Littorina and Prandtl with the respective  $r^2$  of 0.8; 0.99 and 0.98.






Figure S2: Histograms for the dissolved methane concentrations (nmol /L) for Stern-3 (left panel) and Stern-5 (right panel)



|    |                                       |           |
|----|---------------------------------------|-----------|
| #1 | coastal sea of the Elbe River         | 590000003 |
| #2 | Helgoland                             | 590000002 |
| #3 | Piep Tidal basin                      | 950000001 |
| #4 | Outer Elbe North                      | 590000006 |
| #5 | western Wadden Sea of the Elbe        | 590000005 |
| #6 | eastern Wadden Sea of the Weser River | 490000003 |
| #7 | open coastal sea of the Weser         | 490000004 |
| #8 | coastal sea of the Weser              | 490000005 |

Figure S3: Marine ecosystems of the German Bight, with the areas of eastern Wadden Sea of the Weser, open coastal sea of the Weser, coastal sea of the Weser, Helgoland, coastal sea of the Elbe, western Wadden Sea of the Elbe, Outer Elbe North and Piep Tidal basin all indicated with their respective key numbers. Modified after <https://oekosystematlas-ugr.destatis.de>






Figure S4: Concentrations of dissolved CH<sub>4</sub> (top), diffusive CH<sub>4</sub> flux (bottom) on 31.8; 1.9; 2.9 and 3.9.2020 (from left to right). Schlitzer, Reiner, Ocean Data View, [odv.awi.de](http://odv.awi.de), 2024




Fig. S5: Range of diffusive fluxes calculated with all in-situ data from 2019 (flux-1), with in-situ data and atmospheric CH<sub>4</sub> from the land station (flux-2) and within-situ data and atmospheric CH<sub>4</sub> and wind from a land stations (flux-3). The calculations were performed for the region of Bremerhaven (blue), for Cuxhaven (green) and Helgoland (red, see Fig. 2).






Figure S6: Frequency distribution (n) of the diffusive fluxes (flux-1, in  $\mu\text{mol m}^{-2} \text{d}^{-1}$ ) in September 2019




Figure S7: Frequency distribution (n) of the diffusive fluxes (flux-1, in  $\mu\text{mol m}^{-2} \text{d}^{-1}$ ) in September 2020

Table S1: Excel sheet for the calculation of the area weighted flux.

|   |   | total number of samples                      |                                   | total area (m <sup>2</sup> )                   |   |  |  |   |  |
|---|---|--|-----------------------------------|--|---|--|--|---|--|
|   |   | 3028   |                                   | 3779470000                                     |   |  |  |   |  |
|   |   |  |                                   |  |   | sum of relative flux<br>( $\mu\text{mol/d}$ )<br>(SUMME(F7:F48)) |  | sum of relative flux<br>( $\text{kmol/d}$ ) |  |
|   |   |  |                                   |  |   | 6.00217E+11  |  | 600   |  |
|   |   |  |                                   |  |   | 20747454895  |  | 21  |  |
|   |   |  |                                   |  |   |  |  | std deviation from column F                 |  |
| scores / bins range<br>( $\mu\text{mol/m}^2/\text{d}$ ) | mean diffusive flux of<br>each bin ( $\mu\text{mol/m}^2/\text{d}$ ) | Frequency (number of<br>samples in this bin) | relative frequency<br>(C7/\$C\$2) | relative area in<br>m <sup>2</sup> (D7*\$E\$2) | relative flux in<br>$\mu\text{mol/d}$ (E7*B7) |  |  |   |  |
| (0,100]   | 34  | 2026   | 0.669                             | 2528799941                                     | 86485460243                                   |  |  |   |  |
| (100,200]   | 140   | 636  | 0.210                             | 793838480.8                                    | 1.1112E+11                                    |  |  |   |  |
| (200,300]   | 233   | 156  | 0.052                             | 194715099.1                                    | 45438933283                                   |  |  |   |  |
| (300,400]   | 334   | 32   | 0.011                             | 39941558.78                                    | 13354141174                                   |  |  |   |  |
| (400,500]   | 459   | 17   | 0.006                             | 21218953.1                                     | 9730448397                                    |  |  |   |  |
| (500,600]   | 556   | 26   | 0.009                             | 32452516.51                                    | 18044021432                                   |  |  |   |  |
| (600,700]   | 662   | 21   | 0.007                             | 26211647.95                                    | 17348183034                                   |  |  |   |  |




Figure S8: Backward trajectory over 24 hours for air mass at 10 m height for 1 and 3 September 2020. The star indicates an arbitrary point in our study area.