

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Do we (need to) care about canopy radiation schemes in DGVMs? An evaluation and assessment study

A. Loew¹, P. M. van Bodegom², J.-L. Widlowski³, J. Otto⁴, T. Quaife⁵, B. Pinty³,
and T. Raddatz¹

¹Max Planck Institute for Meteorology, KlimaCampus, Hamburg, Germany

²VU University Amsterdam, Department of Ecological Science, Subdepartment of Systems Ecology, Amsterdam, the Netherlands

³European Commission, Joint Research Centre, Institute for Environment and Sustainability Joint Research Centre, Via E. Fermi 2749, 21027 Ispra (VA), Italy

⁴Laboratoire des Sciences du Climat et l'Environnement (LSCE), CEA-CNRS-UVSQ, Gif-sur-Yvette, France

⁵Department of Meteorology, University of Reading, Earley Gate, Reading, UK

Received: 28 September 2013 – Accepted: 14 October 2013 – Published: 25 October 2013

Correspondence to: A. Loew (alexander.loew@zmaw.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Dynamic Global Vegetation Models (DGVM) are an essential part of current state-of-the-art Earth System Models. In recent years, the complexity of DGVM has increased by incorporating new important processes, like e.g. nutrient cycling and land cover dynamics while biogeophysical processes, like surface radiation have been not much further developed. Canopy radiation models are however very important for the estimation of absorption and reflected fluxes and are essential for a proper estimation of surface carbon, energy and water fluxes.

The present study provides an overview about current implementations of canopy radiation schemes in a couple of state-of-the-art DGVMs and evaluates their accuracy in simulating canopy absorption and reflection for a variety of different surface conditions. Systematic deviations in surface albedo and fraction of absorbed photosynthetic active radiation (faPAR) are identified and potential impacts are assessed.

The results show clear deviations for both, absorbed and reflected, surface solar radiation fluxes. FaPAR is typically underestimated which results in an underestimation of Gross Primary Productivity (GPP) for the investigated cases. The deviation can be as large as 25 % in extreme cases. Deviations in surface albedo range between $-0.15 \leq \Delta\alpha \leq 0.36$ with slight positive bias in the order of $\Delta\alpha \approx 0.04$. Potential radiative forcing caused by albedo deviations is estimated as $-1.25 \leq RF \leq -0.8 \text{ [W m}^{-2}\text{]}$ caused by a neglecting the diurnal cycle of surface albedo.

The present study is the first one that provides an evaluation of canopy RT schemes in different currently used DGVMs together with an assessment of the potential impact of the identified deviations. The paper illustrates that there is a general need to improve the canopy radiation schemes in DGVMs and provides different perspectives for their improvement.

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Land surface models are one of the required tools for understanding land surface dynamics, land–atmosphere interactions and climate-carbon feedbacks and are an essential part of Dynamic Global Vegetation Models (DGVM). DGVMs are widely used for assessing climate change impacts on vegetation distribution and terrestrial carbon, water and energy fluxes and feedbacks of the biosphere in the Earth System as well as climatic consequences of land cover change (Friedlingstein et al., 2006; Brovkin et al., 2013a). Given the major policy implications for climate change mitigation, much attention is placed on the performance and realism of these models resulting in an increased overall complexity. At the same time, among land surface models, there is no consensus on important aspects of the carbon cycle in a future climate (Sitch et al., 2008).

DGVMs typically represent the land surface dynamics in a simplified manner. For practical reasons they represent small scale processes using large scale variables. In doing so, different models make different approximations in their representation of processes. The relevance and realism of several approximations and their potential implications for the range of projections found have been discussed elsewhere as far as it concerns the ecology (Harrison et al., 2010; Van Bodegom et al., 2012) and soil carbon dynamics (Ostle et al., 2009). However, also in their representation of exchanges of energy (radiation and heat), models differ widely and occasionally have been implemented differently for albedo and fraction of absorbed photosynthetic active radiation (faPAR) calculations. Most models are confined to one dimensional (vertical) exchange of radiation, mostly relying on solutions derived from two-stream approximations based on plane-parallel turbid media assumptions, like those of Sellers (1985).

The transfer of radiation within canopies is complicated by multiple scattering of radiation, mutual shadowing, variations in leaf orientations and crown closure, as well as variable optical properties. Sprintsin et al. (2012) have shown that neglecting e.g. the differences between sunlit and shadowed leaves in canopy radiative transfer (RT)

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



schemes results in a significant underestimation of the canopy-level Gross Primary Production (GPP). A similar study analyzing the impact of a revised canopy RT scheme in the ISBA land surface model has been documented by Carrer et al. (2013). In addition, full 3-D radiative transfer models exist for canopies as well as landscapes (Gastellu-Etchegorry et al., 2004; Kobayashi and Iwabuchi, 2008), but these cannot be used directly in DGVMs due to their high computational demand and the high number of required (vegetation structural) parameters. Numerically fast 1-D models with a limited number of required input parameters are therefore still preferably used.

These 1-D-models simplify radiative complexities caused by vegetation clumping, representing the concentration of vegetation and thus canopy scattering and absorption within a given area. This leads to so-called effective variables (Widlowski et al., 2005; Rochdi et al., 2006; Pinty et al., 2006). However, neither 1-D-approach has been incorporated consistently within the currently state-of-the-art DGVMs. Alternative 1-D-models have been developed which assume large scale canopy elements to have simple shapes (e.g. spherical crowns) and give analytical solutions for the calculation of surface fluxes using a 1-D-model (Dickinson et al., 2008; Haverd et al., 2012).

Given these complications, there is increasingly a call for systematically evaluating and benchmarking DGVMs (Abramowitz et al., 2008; Luo et al., 2012; Hagemann et al., 2013; Brovkin et al., 2013b). Benchmark analysis is essential to identify uncertainties in predictions as well as to guide priorities for further model development (Blyth et al., 2011). Despite the importance of heat and energy exchange in DGVMs as major drivers of surface temperatures and carbon productivity, current benchmark analyses do not yet account for energy budgets. Where benchmark initiatives consider albedo and faPAR, separate – and thus from an RT point of view potentially inconsistent – Earth Observation products are proposed as candidate benchmarks (Luo et al., 2012; Hagemann and Stacke, 2013).

There is thus a need to evaluate the consistency and accuracy of radiative transfer schemes in DGVMs and to assess the potential impact of uncertainties in the widely used radiative transfer models on surface temperature and carbon production esti-

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mates, which so far has not been quantified for widely used DGVMs. Such assessment requires physically consistent three-dimensional radiative transfer formulations as a reference. Such reference simulations are provided by e.g. the Radiative Transfer Model Intercomparison Initiative (RAMI) (Pinty et al., 2001; Pinty, 2004; Widlowski et al., 2007).

Here, we aim at a show case to (i) evaluate the consistency among a number of representative state-of-the-art DGVMs, employing different definitions, assumptions and temporal and spatial scales for the canopy RT formulations, using various reference RAMI4PILPS simulations, (ii) evaluate at which conditions the used canopy RT schemes (and their simplifications) lead to major errors in faPAR and/or albedo in these representative DGVMs, and (iii), importantly, assess the potential implications thereof for net irradiance and carbon productivity estimates.

Thus the overall objective of this paper is to raise awareness on the relevance of canopy radiation and surface albedo schemes in globally applied land surface schemes being used in a wide range of applications.

2 Data and methods

2.1 Representation of canopy RT in DGVMs

Global DGVM simulations are typically performed on coarse spatial resolutions with model grid cell sizes in the order of 10^2 to 10^8 km² and long time periods (decades . . . millenia). To represent surface processes – including radiative transfer – at the sub-grid scale, a tiling (mosaic) approach is widely used where surface processes are simulated on tiles of N individual plant functional types (PFT) and where results are combined to a model grid box average by area-weighted averaging. Plant functional types provide a means to use a finite set of model parameters to simulate plants with similar ecological behavior (Diaz and M., 1997; Prentice et al., 2007). The analysis of

the present study focuses on the performance of the canopy RT models and assumes that a grid cell is covered by a single land cover type only ($N = 1$).

The applications of DGVM models demand a numerically stable and fast canopy radiative transfer scheme. Simple 1-D canopy radiative transfer schemes or parametric approaches are therefore used in DGVMs instead of more computationally demanding 3-D-RT schemes. A 1-D model is understood here as a model where the surface state variables (e.g. leaf area, leaf reflectance and absorption properties) vary only along a single coordinate axis (typically z-direction). Note that this is not related to the spatial integration needed for flux calculations. When integrating over the upper and lower hemispheres, the model is called a 2-stream model. However, angular integrations can also be made only for parts of spheres, resulting then in a 4-stream model for half spheres and an N-stream solution for N numbers of subspaces. If N is large, a bidirectional reflectance model (BRF) is obtained (Gobron et al., 1997).

A turbid medium assumption is widely used for 1-D canopy simulations, where foliage elements are assumed to be point like scatterers – leaves are assumed having an infinitely small size – randomly and uniformly distributed and typically stems and branches are neglected. The gaps within canopies can be expressed by the gap probability (P_{gap}) which is defined as the probability of a beam at sun zenith angle θ to hit the ground without interacting with canopy elements. In other words, it is the ratio between the uncollided radiation flux and the incident radiation (Haverd et al., 2012) and represents the domain averaged direct transmission. In case of a horizontally homogeneous canopy with randomly distributed leaves (and no foliage), P_{gap} follows a Poisson distribution and is given by

$$P_{\text{gap}}(\theta, \Lambda) = e^{-G(\theta) \cdot \Lambda / \mu} \quad (1)$$

where $\mu = \cos(\theta)$, Λ is the leaf area index (LAI) [$\text{m}^2 \text{m}^{-2}$] and $G(\theta)$ is the mean projection of unit leaf area in the direction perpendicular to the incoming beam. Often a spherical leaf angle distribution is assumed, which results in $G(\theta) = 0.5$.

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This representation of P_{gap} is widely used in DGVMs and applies to all models investigated in this study. While Eq. (1) is a good approximation of P_{gap} for closed canopy cases, it has been recognized that it overestimates canopy absorption for open (non-uniform) canopies (Pinty et al., 2006; Chen et al., 2008; Haverd et al., 2012). To account for non-uniform distributions, a so called clumping factor $0 < \Omega < 1$ has been introduced for use in a one-dimensional radiative transfer model. The gap probability is then calculated as

$$P_{\text{gap}}(\theta, \Lambda) = e^{-\Omega(\theta)G(\theta)\cdot\Lambda/\mu} = e^{-\tilde{\Lambda}(\mu)} \quad (2)$$

where $\tilde{\Lambda}(\mu)$ is an effective leaf area index which reproduces the correct canopy RT fluxes using a 1-D formulation instead of a 3-D model (Pinty et al., 2006).

The fraction of absorbed photosynthetic radiation (faPAR) is a crucial variable in DGVM carbon flux simulations. Ignoring horizontal fluxes (Widlowski et al., 2006), faPAR is defined as the fraction of radiation absorbed by the canopy elements (foliage and woody parts) in the photosynthetic active electromagnetic spectrum (= VIS) and is defined by considering the energy balance as

$$\text{faPAR} = 1 - \frac{\text{PAR}_\alpha}{\text{PAR}_i} - (1 - \alpha_s) \frac{\text{PAR}_T}{\text{PAR}_i} \quad (3)$$

where PAR_i is the incident down-welling PAR at the top of the canopy, PAR_α is the reflected component, PAR_T is the total transmission through the canopy and α_s is the albedo of the soil. A first order approximation to faPAR is given by assuming the leaves and soil are completely black (i.e. all radiation incident on them is absorbed). In this case $\text{PAR}_\alpha = 0$ and $\text{PAR}_T/\text{PAR}_i = P_{\text{gap}}$, which yields:

$$\text{faPAR}(\theta, \Lambda) = 1 - P_{\text{gap}} = 1 - e^{-\Omega(\theta)G(\theta)\cdot\Lambda/\mu} \quad (4)$$

However, if the optical properties of the canopy elements are non-zero, then the exact solution becomes more complex as it must consider multiple scattering within

the canopy, and between the crown and the ground if the soil albedo is also non-zero. Among the various currently used faPAR approximation schemes Eq. (4) was shown to provide the least systematic bias (Widlowski, 2010).

Ideally, a 1-D-RT scheme would also consider the proportion of direct illumination and shading at different levels in the canopy as this has potentially significant impacts on photosynthesis (Mercado et al., 2009). This entails splitting the incoming solar radiation into direct and scattered components (Spitters et al., 1986) and calculating the amount of energy intercepted at each canopy layer that is from the direct component which has not previously been scattered (i.e. the sunlit fraction), and the diffuse and multiply scattered components (i.e. the shaded fraction). Multiple scattering is typically neglected in simple canopy RT schemes employed in DGVMs.

Most DGVMs use a parametric 1-D-RT scheme that is somewhere within the range of complexities described above. However, more complex 1-D-schemes are available. In general the added complexity is in the form of extra detail in the description of canopy structure. These models range from Monte-Carlo Ray Tracing which explicitly compute the path of a large number of photons through the vegetation canopy (Disney et al., 2000) to Geometric Optic techniques that treat the tree crowns as a distribution of geometric primitives and calculate the proportion of illuminated and vegetated parts of the scene (Li and Strahler, 1986). Ray tracing models are unlikely to be practical for directly embedding them in DGVMs although they are useful tools for validating simpler models. Some models based on Geometric Optics are viable however. An example is the ACTS (Analytical Clumped Two-Stream) model (Ni-Meister et al., 2010) which is derived from the GORT (Geometric Optic Radiative Transfer) model (Ni et al., 1999). The approach used is to consider the canopy as forest of randomly distributed spheroids filled with leaves. The probability of a photon entering a crown is calculated from geometric optic theory and then scattering is treated as a 1-D-RT problem similar to those described elsewhere in this section.

DGVM canopy
radiation

A. Loew et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2.2 Handling sparse vegetation (open canopies) in DGVMs

Despite the existence of concepts to account for vegetation clumping using geometric optics, more simplified approaches to handle sparse vegetated canopies have been implemented in DGVMs and are widely used. Figure 1 shows a model grid cell with unit area A which is assumed to be fully covered by a particular PFT. The area A corresponds to the reference ground area for the leaf area index of the PFT. In case of sparse vegetation types, like e.g. savannas, the vegetated area is typically covered by a dominant plant functional type (e.g. trees) and understory vegetation (e.g. shrubs, grass) or bare soil. In the process of converting land cover information, as is available from satellite products, a partitioning between these different components is made (Poulter et al., 2011). The total area A is thus defined as

$$A = f_{\text{veg}} + f_{\text{under}} \quad (5)$$

where f_{veg} is the fractional coverage of the major PFT and f_{under} can correspond to a different PFT or soil surface. It is important to recognize that f_{under} is different from the gap probability $P_{\text{gap}}(\theta = 0)$, as gaps within the dominant canopy are not considered in f_{under} , therefore $P_{\text{gap}}(\theta = 0) \geq f_{\text{under}}$. In other words, a common concept in DGVMs is to approximate P_{gap} as

$$\tilde{P}_{\text{gap}}(\Lambda) \approx f_{\text{under}} + g(\Lambda(t)) \quad (6)$$

where $g(\Lambda)$ is a function that estimates the gaps within the canopy of the dominant vegetation type and might change over time t as a function of leaf area index Λ , while f_{under} is static (examples for the investigated models are given in the following section). Note that while P_{gap} in Eq. (1) is a function of illumination conditions (θ) this is not the case for \tilde{P}_{gap} Eq. (6).

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3 Models

The present study uses three different DGVMs, which are used in renowned Earth System models which all contribute to the Coupled Model Intercomparison Projects on a regular basis (Meehl et al., 2007; Taylor et al., 2012).

2.3.1 JULES

JULES is the Joint UK Land Environment Simulator, a land surface model designed to predict the fluxes of heat, water and carbon between the land surface and the atmosphere. It originates from the Met Office Surface Exchange Scheme (MOSES) and is designed to be linked to the UK Met Office Unified Model. The fundamental equations underlying the model are in common to many land surface schemes and are described in detail elsewhere (Best et al., 2011; Clark et al., 2011). In addition, JULES allows for a coupling with a General Circulation Model (GCM) and provides optional modules that allow taking into account longer term processes such as succession of plant functional types. These allow it to be used to simulate the response of the land surface to changing climatic conditions. In particular this includes the Dynamic Vegetation Model TRIFFID (Top-down Representation of Interactive Foliage and Flora Including Dynamics) (Cox, 2001).

The canopy radiative transfer scheme in JULES is based on the two-stream approximation proposed by Sellers (1985), which translates into consistent calculations of surface albedo and canopy absorption. Sellers (1985) provides an analytical solution for the two stream model given as

$$\begin{aligned} -\bar{\mu} \frac{dI^{\uparrow}}{dL} + [1 - (1 - \beta)\omega] I^{\uparrow} - \omega\beta I^{\downarrow} &= \omega\bar{\mu}\tau\beta_0 e^{-\tau L} \\ \bar{\mu} \frac{dI^{\downarrow}}{dL} + [1 - (1 - \beta)\omega] I^{\downarrow} - \omega\beta I^{\uparrow} &= \omega\bar{\mu}\tau(1 - \beta_0) e^{-\tau L} \end{aligned} \quad (7)$$

where I^{\uparrow} (I^{\downarrow}) are the upward and downward diffuse radiation fluxes normalized by the incoming radiative flux, $\tau = G(\mu)/\mu$ is the optical depth and $\omega = r + t$ the single scattering albedo. β and β_0 are upscatter parameters for the diffuse and direct beams respectively and $\bar{\mu}$ is the average inverse optical depth per unit leaf area and L corresponds to the cumulative leaf area index. For details on the solution of Eq. (7) see Sellers (1985).

JULES is the only one of the analyzed models which does a consistent simulation of absorbed and scattered radiation. Direct and diffuse radiation fluxes are computed individually. The leaf scattering and extinction properties are prescribed by the leaf reflectance (r) and single scattering albedo (ω). JULES uses a number of layers, as defined by the user, with equally distributed LAI density to simulate the canopy radiative transfer for the direct and diffuse flux components. Here an $n = 20$ layer model has been used as default.

The scheme of (Sellers, 1985) used here is from JULES version 3.2. An earlier version (2.1) was used in the RAMI4PILPS exercise (Widlowski et al., 2011) and was found to give analogous results (not shown here).

2.3.2 JSBACH

The DGVM JSBACH (Raddatz et al., 2007; Brovkin et al., 2009; Reick et al., 2013), is implicitly coupled to ECHAM6, the atmospheric component of the Max-Planck-Institute for Meteorology Earth System model (Stevens et al., 2013). It simulates all relevant land surface water, energy and carbon fluxes. The present study uses version 2.03 of JSBACH which is comparable to the model version which was used for the Coupled Model Intercomparison Project (CMIP5) experiments (Taylor et al., 2012). A validation of global scale energy and water flux components of MPI-ESM CMIP5 simulations is given in Hagemann et al. (2013) and Brovkin et al. (2013b). The JSBACH model has two independent schemes for calculating surface albedo and canopy absorption.

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Albedo

The surface albedo of a vegetated area (α_{veg}) is calculated as a weighted sum of the leaf albedo (α_{le}) and soil background albedo (α_{s}) as

$$\alpha_{\text{veg}}(\Lambda, \alpha_{\text{s}}, \alpha_{\text{le}}) = (1 - \Omega_{\text{sky}}(\Lambda)) \alpha_{\text{le}} + \Omega_{\text{sky}}(\Lambda) \alpha_{\text{s}} \quad (8)$$

Soil albedo α_{s} depends on soil color and the soil litter content (Vamborg et al., 2011). A PFT specific leaf albedo (α_{le}) as well as the soil background albedo was derived from MODIS observations (Otto et al., 2011).

The sky-view factor (Ω_{sky}), which weights between canopy and soil albedo, is calculated assuming a random leaf-angle distribution as

$$\Omega_{\text{sky}}(\Lambda) = 1 - \left(f_{\text{vegmax}} \cdot \left(1 - e^{-0.5\Lambda} \right) \right) \quad (9)$$

where $0 \leq f_{\text{vegmax}} \leq 1$ is the fraction of a model grid cell with vegetation for $\Lambda \rightarrow \infty$ which is similar to f_{veg} in Fig. 1. Note that Ω_{sky} is independent of the sun zenith angle in this parameterization which does therefore not allow to simulate mutual shadowing in the canopy or its effect on surface albedo, neither the diurnal dependency of surface albedo.

Canopy absorption

FaPAR is calculated in JSBACH using a 1-D two-stream approximation based on Sellers (1985), similar to JULES. The faPAR is calculated for direct and diffuse radiation components separately using $n = 3$ canopy layers with equally distributed leaf densities. Further assumptions are that the leaf reflectivity and transmissivity are equal ($r = t = \text{const}$). The canopy single scattering albedo is assumed to be $\omega = \text{const} = 0.12$. Note that this is independent of the leaf albedo used for the albedo calculations. A spherical leaf angle distribution ($G(\theta) = 0.5$) is also assumed in JSBACH which is similar to JULES.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To account for sparse vegetation, faPAR is corrected for the vegetation fraction f_{veg} . This vegetated fraction is parametrized for each plant functional type as

$$f_{\text{veg}}(\Lambda_{\text{max}}) = 1 - \exp\left(-\frac{\Lambda_{\text{max}}}{\gamma}\right) \quad (10)$$

where Λ_{max} and γ are PFT specific parameters representing the maximum leaf area index for a particular PFT and an empirical clumping parameter, respectively. The calculation of $f_{\text{veg}}(\Lambda_{\text{max}})$ for faPAR calculations is therefore slightly different than for the calculation of the surface albedo but also independent of the solar zenith angle.

While the assumptions for the canopy radiative transfer simulations and technical implementations are different between JSBACH and JULES, it can be shown that both models provide nearly identical results for faPAR when parametrized in the same way.

The JSBACH RT scheme allows to explicitly account for gaps within the canopy. It is assumed that, due to canopy gaps, only a fraction $f_{\text{clump}} < 1$ of the canopy area is actually covered with leaves. The radiative transfer simulations are then calculated with an effective LAI $\Lambda_{\text{eff}} = \Lambda / f_{\text{clump}}$. Note that the leaf area index used for the canopy radiative transfer simulations is therefore larger than the original leaf area index ($\Lambda_{\text{eff}} > \Lambda$). The simulated radiation fluxes are therefore rescaled by multiplying with f_{clump} after the 1-D-RT calculations have been performed. To assess the impact of this clumping on the simulations in the present study, two different JSBACH model versions (with/without clumping) are evaluated.

2.3.3 ORCHIDEE

ORCHIDEE (ORganizing Carbon and Hydrology In Dynamic Ecosystems) is a land-surface model that simulates the energy and water cycles of soil and vegetation, the terrestrial carbon cycle, and the vegetation composition and distribution (Krinner, 2005).

ORCHIDEE is used as the land surface scheme of the IPSL Earth System model. A global model validation is described in Dufresne et al. (2013) for the CMIP5 simulation experiments. We use the version ORCHIDEE 1.9.5.2 (Revision 816). No changes

have been introduced to the calculation of albedo and faPAR since the version which was used for the CMIP5 simulations (Dufresne et al., 2013). The calculation of the light absorbed by vegetation and vegetation albedo is calculated in two independent schemes.

5 Albedo

The albedo calculation in ORCHIDEE is the same as in JSBACH (Eq. 8). Only the PFT-specific canopy albedo differs slightly compared to JSBACH. The soil background albedo has been derived from a database of soil colors in the standard setup of ORCHIDEE according to Wilson and Henderson-Sellers (1985).

10 Canopy absorption

The faPAR is not a standard output variable in ORCHIDEE. The light absorbed by the canopy is calculated by means of Beer's law assuming a constant extinction coefficient ($G(\theta) = 0.5$) for all PFTs and assuming an exponential profile of leaf area index (Λ^*) within the vertical canopy profile. FaPAR is calculated once and stored in a look-up-table. The calculation is independent of the solar zenith angle:

$$15 \text{ faPAR}(\Lambda^*) = 1 - e^{-0.5\Lambda^*} \quad (11)$$

The vertical distribution of LAI is calculated using $n = 20$ different canopy layers. The LAI for a given layer i is calculated as

$$20 \Lambda^*(i) = \Lambda_{\max} \frac{e^{0.5(i-1)} - 1}{e^{0.15n} - 1} \quad (12)$$

whereas $\Lambda_{\max} = 12 \text{ [m}^2 \text{ m}^{-2}\text{]}$, independent of the PFT used.

The actual LAI, as calculated dynamically by a phenology model, is compared to this precomputed profile to determine the number of canopy layers, which is thus not fixed

in ORCHIDEE. The LAI for this layer is then used to compute faPAR using Eq. (11). A maximum number of 20 canopy layers is reached at the theoretical LAI of $\Lambda^*(20) = 12$ [$\text{m}^2 \text{m}^{-2}$].

2.4 RAMI4PILPS experiments

The RAMI4PILPS suite of virtual experiments was designed to evaluate the accuracy and consistency of shortwave RT formulations (as used in DGVMs) under perfectly controlled experimental conditions. More specifically, RAMI4PILPS prescribed a series of virtual canopy scenarios having accurately described structural, spectral and illumination related characteristics. For these test cases, RT model simulations have been generated as a reference using a Monte Carlo approach. The Monte Carlo model in question had been extensively verified during previous RAMI phases, e.g., Widlowski et al. (2007). Models, participating in the RAMI4PILPS benchmarking exercise (Widlowski et al., 2011) had to simulate the canopy albedo, transmission and absorption in both the visible and near-infrared spectral domains. The resulting data were evaluated against the 3-D Monte Carlo reference solution.

Contrary to efforts comparing model simulations against in situ observations at specific test sites, the RAMI4PILPS approach eliminates uncertainties arising from an incomplete or erroneous knowledge of (1) the structural, spectral, and illumination related characteristics of the canopy target, and (2) the uncertainties introduced into the reference solution by calibration, sampling and upscaling errors (Fig. 2).

The complexity of the RAMI4PILPS scenes was adapted to the typical capability of available shortwave RT formulations. More specifically, RAMI4PILPS proposed two homogeneous plant canopy types, i.e., grasslands (GRA) and closed forest canopy (CFC) scenes that differ only in their predominant leaf orientations as well as in the height of the canopy. This corresponds to the idealized case of a 1-D-turbid medium, where 1-D canopy radiative transfer schemes are expected to perform best.

In addition, RAMI4PILPS proposed two heterogeneous canopy scenarios where tree crowns were approximated by woodless spheres, i.e. a shrubland (SHR) and an open

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



forest canopy (OFC) scene that differed only in their size of the spheres, the height of the canopy, and the degree of mutual shading between neighboring crowns (Fig. 2). Details of the RAMI4PILPS experiments used in the present study are summarized in Table 1 while further details of the RAMI4PILPS experiments can be found in Widowski et al. (2011). For each scenario, simulations for different leaf area and varying soil brightness are performed, assuming direct insolation for three different sun zenith angles as well as isotropic illumination conditions.

2.4.1 Interfacing with RAMI4PILPS

To enable all three models to simulate the RAMI4PILPS scenes the experiment parameters were implemented as follows:

- *Single scattering albedo*: JULES uses the vegetation single scattering albedo (ω) as an input. It was calculated from the RAMI4PILPS leaf reflectance and transmittance as $\omega = r + t = 0.1301$. JSBACH assumes ω to be constant for calculation of canopy absorption. The standard value used in JSBACH simulations ($\omega = 0.12$) was also used in the present study. The impact of this assumption was however tested by comparing JSBACH simulations with $\omega = 0.12$ and $\omega = 0.1301$. A minor impact on the results of this study was found, indicating minor importance of ω for the present study. The leaf albedo (α_{le}) and soil albedo (α_s) were both obtained from Table 1. ORCHIDEE does not use any information about the single scattering albedo.
- *Snow covered areas*: One of the RAMI4PILPS simulations assumes snow below the canopy (SNW). In these cases, the soil albedo was replaced by the snow albedo. No snow on canopy was simulated as this was not foreseen in the RAMI4PILPS experiments.
- *Open vs. closed canopies*: Open canopies consist of vegetated and non-vegetated (soil) patches as discussed before. The RAMI4PILPS experiment de-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



scription contains information on the fractional area covered by vegetation (f_{veg}) as well as the amount of uncollided radiation ($P_{\text{gap}}(\theta = 0)$).

All models rely on the assumption of a plane-parallel turbid media, which prohibits a direct application to open canopies. Open canopies are challenging to simulate for 1-D-RT models and require effective RT model parameters, as discussed in Sect. 2.1. The applied models try to mimic open canopies by correcting by the area fraction covered by vegetation as discussed before.

The vegetation fraction f_{veg} was therefore obtained from the RAMI4PILPS experiment setup (Table 1). The soil albedo α_s was used for the understory. For JSBACH we set $f_{\text{vegmax}} = f_{\text{veg}}$. The surface albedo for open canopies was calculated for JSBACH and ORCHIDEE by using Eq. (8), while for JULES the surface albedo was calculated as

$$\alpha_{\text{veg}}(\alpha_s, \alpha_{\text{le}}, f_{\text{veg}}) = f_{\text{veg}}\alpha_{\text{le}} + (1 - f_{\text{veg}})\alpha_s \quad (13)$$

where α_{le} is calculated by the JULES RT model. The total faPAR was calculated in a similar way by weighting the faPAR calculated by the RT models ($\text{faPAR}_{\text{canopy}}$) by the actual area fraction covered by vegetation as

$$\text{faPAR} = f_{\text{veg}} \cdot \text{faPAR}_{\text{canopy}} \quad (14)$$

The spectral definition in all models and in RAMI4PILPS is consistent. The solar spectrum was divided in two broad spectral bands (VIS: 400...700nm, NIR: 700...3000nm). Both bands were analyzed for albedo in this study, while only the VIS band was used for the faPAR analysis.

2.5 Impact assessment – does it matter?

This paper aims at identifying systematic deviations for albedo and faPAR for well established DGVMs and to provide guidance for estimating the impact of these deviations

on global climate simulations and carbon flux estimates. A comprehensive impact assessment would, however, require a detailed knowledge of the spatial distribution of faPAR and albedo deviations and their consideration in coupled models, with all else being the same including LAI and all other vegetation variables. This is far beyond the scope of the present paper.

It is however of major importance for further model development to raise awareness of potential impacts of the identified deviations. We will detail in the following how we performed straightforward calculations for a first assessment of potential impacts caused by deviations in RT schemes. These are likely to first affect net photosynthesis by differences in absorbed radiation through the canopy at a given LAI and second through heat exchange between the land surface and the atmosphere through albedo. Whereas DGVM behavior with respect to energy, water and carbon exchange will be affected in multiple other ways, we constrained our assessment to these two prime targets.

2.5.1 Potential impacts of RT scheme differences on the C-cycle

In order to evaluate the potential implications of differences in canopy faPAR profiles, induced by different canopy RT schemes, we determined net photosynthesis rates for each of the virtual experiments of the closed forest canopy case only (CFC). We made this constraint because the photosynthesis calculations in combination with the representation of faPAR for different layers of the canopy are likely only valid for the closed forest canopy case, given the assumptions made in the photosynthesis schemes adopted in most DGVMs. To determine net photosynthesis estimates, we chose the Farquhar scheme as described by von Caemmerer (2000), which is representative for the photosynthesis schemes in current DGVMs. According to this scheme, net photosynthesis is the minimum of CO₂-limited photosynthesis and photosynthesis as limited by the availability of ribulose bi-phosphate (RuBP), the latter being driven by absorbed radiation, corrected for dark respiration.

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We evaluated this scheme for two representative closed canopy situations at typical clear sky conditions in summer, following Widlowski et al. (2011). The two selected situations coincide with a tropical forest with a solar zenith angle at local noon (Θ_{noon}) of 15° and a boreal (presumably evergreen needleleaved) forest with $\Theta_{\text{noon}} = 50^\circ$. For these situations, and the given sun zenith angles of the virtual experiment, the total incident shortwave radiation at noon was calculated according to Wang et al. (2002). When combined with the faPAR profiles through the canopy, this provided the absorbed photosynthetic active radiation (aPAR) in each canopy layer. Coupling aPAR to the Farquhar scheme provided CO₂-limited and light-limited photosynthesis rate in each canopy layer. The minimum of these represents the net photosynthesis rate, which was summed across all layers to provide the net photosynthesis rate [$\mu\text{mol C m}^{-2} \text{s}^{-1}$] for a given virtual experiment. Absolute differences between net photosynthesis rates between the RAMI4PILPS scenes and each of the DGVMs were compared.

The Farquhar scheme was parameterized according to Sharkey et al. (2007) which currently provides the most comprehensive analysis of the variables involved. Estimates of the maximum rates of Rubisco carboxylase activity ($V_{c_{\text{max}}}$) and the maximum rates of photosynthetic electron transport (J_{max}), were however taken to represent the mean observed values at a reference temperature of 25°C for tropical broadleaved evergreen forests and boreal (needleleaved) evergreen forests, according to the TRY database (Kattge et al., 2011) and Domingues et al. (2010), as compiled by Verheijen et al. (2013). Dark respiration was scaled to $V_{c_{\text{max}}}$.

2.5.2 Surface radiation budget

Surface albedo directly affects the net surface radiation budget and thus indirectly also near surface temperature. To quantify the influence of a factor which is changing the balance of incoming and outgoing energy in an Earth-atmosphere system, the concept of radiative forcing (RF) has been developed (IPCC, 2007). Radiative forcing has been widely used to quantify global impact of regional changes in surface albedo (Hall and Qu, 2006; Pongratz et al., 2009; Lenton and Vaughan, 2009; Bright et al., 2012) and

can be calculated in various ways. Here, we applied a simple method to estimate RF (Bright et al., 2012; Cherubini et al., 2012).

Effect of albedo biases

The radiative forcing calculated in this study is understood as a disturbance of radiation fluxes caused by variations in surface albedo. The estimates calculated in this study must therefore not be compared directly with estimates of radiative forcing from different forcing agents like those used e.g. in coupled climate model assessments (IPCC, 2007). The albedo induced radiative forcing RF_{α_p} is related to a change in planetary albedo α_p as

$$RF_{\alpha_p} = -R_{\text{TOA}}^{\downarrow} \Delta\alpha_p \quad (15)$$

where $R_{\text{TOA}}^{\downarrow}$ [W m^{-2}] is the incoming solar radiation flux at the top of the atmosphere (TOA). The change in planetary albedo is linearly related to a change in surface albedo (Lenton and Vaughan, 2009). The corresponding radiative forcing is then given by

$$RF_{\alpha_s} = -R_{\text{TOA}}^{\downarrow} \Gamma_a^{\uparrow} \Delta\alpha_s \quad (16)$$

where Γ_a^{\uparrow} is the two-way atmospheric transmissivity for solar radiation which is given as the product of the downward (Γ^{\downarrow}) and upward (Γ^{\uparrow}) atmospheric transmissivities. Γ^{\uparrow} was parameterized in two ways. Similar to previous studies focusing on radiative forcing (Lenton and Vaughan, 2009; Cherubini et al., 2012), we assumed a constant value of $\Gamma^{\uparrow} = 0.854$. This global mean value is based on the assumption that the outgoing shortwave flux is leaving the atmosphere given clear-sky conditions. Alternatively, we assumed that under cloudy sky conditions, the upward and downward atmospheric transmissivities are equal ($\Gamma^{\uparrow} = \Gamma^{\downarrow}$).

The TOA solar radiation flux can be calculated for a particular point on Earth using simple formulations (Widowski et al., 2011; Bright et al., 2012). The atmospheric one-way downward transmissivity Γ_a^{\downarrow} can be derived from shortwave surface all-sky ($R_{\text{surf}}^{\downarrow}$)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and TOA flux ($R_{\text{TOA}}^{\downarrow}$) estimates as available from satellite data products (Rossow and Zhang, 1995; Posselt et al., 2012; Loeb et al., 2012) as

$$\Gamma^{\downarrow} = R_{\text{surf}}^{\downarrow} / R_{\text{TOA}}^{\downarrow} \quad (17)$$

Let us now assume that we observe a surface albedo bias ($\Delta\alpha_s$) for a particular plant functional type (PFT) and that this PFT covers a fraction f_{pft} of a model grid box, then the total radiative forcing for the Earth with surface area $A = 510 \times 10^6$ [km²] is given using Eq. (16) by integrating over the surface area as

$$\text{RF}_{\text{tot}} = -\frac{1}{A} \int_A f_{\text{pft}}(a) R_{\text{TOA}}^{\downarrow}(a) \Gamma_a^{\downarrow}(a) \Gamma_a^{\uparrow}(a) \Delta\alpha_s(a) da \quad (18)$$

where “ a ” is a spatial index. If $\Delta\alpha_s$ is assumed to be constant (time independent) for a particular PFT, then the sensitivity to surface albedo changes can be defined as

$$\frac{\partial \text{RF}_{\text{tot}}}{\partial \Delta\alpha_s} = \lambda = -\frac{1}{A} \int_A f_{\text{pft}}(a) R_{\text{TOA}}^{\downarrow}(a) \Gamma_a^{\downarrow}(a) \Gamma_a^{\uparrow}(a) da \quad (19)$$

which can be calculated for each combination of PFT fraction f_{pft} and Γ_a^{\uparrow} and Γ_a^{\downarrow} .

Table 2 provides global mean estimates of the one-way atmospheric transmissivity (Γ^{\downarrow}) as well as global means of $R_{\text{TOA}}^{\downarrow}$ and $R_{\text{surf}}^{\downarrow}$ based on various data sources over land areas. The one-way atmospheric transmissivity ranges from $0.41 \leq \Gamma^{\downarrow} \leq 0.58$, dependent on the dataset.

Figure C1 shows the mean fields of $R_{\text{surf}}^{\downarrow}$ and the estimated one-way transmissivity over land as derived from CERES EBAF (v2.6) (Loeb et al., 2012). Using Eq. (19) and values in Table 2 one can estimate the radiative forcing caused by an albedo change for a given distribution of plant functional types.

Impacts of neglecting the diurnal cycle on radiative forcing

The surface albedo can change substantially throughout the day due to its dependency on the solar zenith angle. The JSBACH and ORCHIDEE surface albedo models are representative for a wide range of albedo models applied in DGVMs. These models do not simulate the surface albedo diurnal behavior, which may affect the surface radiation budget.

The surface albedo can be expressed as a weighted average between the so called black-sky-albedo (BSA) and the albedo for complete isotropic diffuse radiation (white sky albedo, WSA) (Lewis and Barnsley, 1994; Schaepman-Strub et al., 2006) as

$$\alpha_{\text{blue}}(f_{\text{dir}}, \theta) = f_{\text{dir}}\alpha_{\text{BSA}}(\theta) + (1 - f_{\text{dir}})\alpha_{\text{WSA}} \quad (20)$$

where f_{dir} [-] corresponds to the fraction of the direct surface solar radiation flux.

The surface albedo models of JSBACH and ORCHIDEE are based on the white sky albedo (α_{WSA}). To assess the potential impacts of neglecting the diurnal cycle in surface albedo calculations, we used the difference between the albedo given by Eq. (20) and α_{WSA} which is given as

$$\Delta\alpha(f_{\text{dir}}, \theta) = \alpha_{\text{WSA}} - \alpha_{\text{blue}} = f_{\text{dir}}(\alpha_{\text{WSA}} - \alpha_{\text{BSA}}(\theta)) \quad (21)$$

Assuming clear sky-conditions, we calculated the surface solar radiation flux $R_{\text{surf}}^{\downarrow}$ for each latitude ϕ and day of the year using the MAGIC atmospheric RT code (Mueller et al., 2012; Posselt et al., 2012). Required atmospheric aerosol and water vapor content were taken from climatological mean values (Kinne et al., 2013).

The temporal average RF disturbance due to neglecting the diurnal cycle of α , averaged over a timeperiod T , is then given as

$$\text{RF}_{\text{tot}} = -\frac{1}{T} \frac{1}{A} \int_A f_{\text{pft}}(a) \int_0^T R_{\text{surf}}^{\downarrow}(t, a) \Gamma_a^{\uparrow}(t, a) \Delta\alpha(t, a) dt da \quad (22)$$

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We calculated Eq. (22) by using either $\Gamma_a^\uparrow = \text{const} = 0.854$ (Lenton and Vaughan, 2009) or by assuming that $\Gamma^\uparrow(t, a) = \Gamma^\downarrow(t, a)$, where the one-way downward transmissivity was obtained from Eq. (17) assuming clear sky conditions and as calculated by MAGIC. The fraction of direct surface radiation flux (f_{dir}) was calculated from monthly climatologies of direct and total radiation fluxes as obtained from satellite radiation products and provided by the NASA Atmospheric Data Centre (<https://eosweb.larc.nasa.gov/>). The global radiative forcing RF_{tot} was calculated for each day ($T = 24$ h) and two different PFT distributions (tree, grass) as derived from MODIS vegetation continuous fields were used (DiMiceli et al., 2011) which provide f_{pft} for each grid cell.

2.5.3 Consistency of canopy absorption and reflection calculations

The canopy radiative transfer needs to fulfill the law of conservation of energy. Following Eq. (3) this is defined as

$$1 - \alpha_{\text{veg}} - (1 - \alpha_s)\Gamma_{\text{can}} - \text{faPAR} = 0 \quad (23)$$

where $0 \leq \Gamma_{\text{can}} \leq 1$ is the one-way canopy transmission. The canopy RT schemes in the used DGVMs differ how they simulate canopy absorption and reflection as discussed before. Two models (JSBACH, ORCHIDEE) use separate approaches for the simulation of canopy absorption and reflection which might lead to a lack of energy conservation in these models.

To test the energy conservation, the canopy one-way transmissivity would be needed as model output. This is however not available from all of the investigated models. We therefore adopted a simpler approach to test in general for energy conservation of the used canopy RT schemes. From Eq. (23) it follows that

$$\Gamma_{\text{can}} = \frac{1 - \text{faPAR} - \alpha_{\text{veg}}}{1 - \alpha_s} \quad (24)$$

To test if the models are in general energy conserving, we investigated if Γ_{can} from Eq. (24) is in its physically defined limits ($0 \leq \Gamma_{\text{can}} \leq 1$), which is the most conservative approach.

3 Results

3.1 Canopy absorbed radiation

The investigated models generally underestimate the radiation absorbed in the canopy for the RAMI4PILPS experiments analyzed (Fig. 3). Especially for the open canopy cases (SHR,OFC), a strong underestimation of faPAR is observed with increasing negative bias for increasing solar zenith angle. The 1-D-canopy RT models without clumping (JULES, JSBACH-no-clumping) perform best for the closed canopy cases (CFC,GRA), as expected.

Some positive biases are observed for the isotropic (ISO) illumination conditions for JULES and JSBACH-no-clumping. In general, results of these models are almost identical due to the same basis for the canopy RT simulation code as discussed in Sect. 2.3.

3.1.1 Does it matter?

Deviations in net photosynthesis due to differences in the faPAR canopy profile for the RAMI4PILPS reference scene vs. each of the models strongly depend on the virtual experiment (Fig. 4). On average, deviations are largest for $\theta = 60^\circ$, where deviations in faPAR are considerable, while incident radiation is still high enough to support net photosynthesis (in contrast to $\theta = 83^\circ$). Moreover, the differences in net photosynthesis strongly increase with increasing LAI, when more leaf surface is available for photosynthesis. Finally, the impacts are also larger for the boreal forest ($\Theta_{\text{noon}} = 50$) slightly increase with increasing albedo, although the latter effects are minor.

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Depending on the model and even within a given model setting, differences in calculated net photosynthesis can be both negative and positive. Largest deviations occur for ORCHIDEE with both overestimations as well as underestimations of net photosynthesis rates, although overestimations by ORCHIDEE predominated. Net photosynthesis estimates according to JSBACH are on average lower than those for the RAMI4PILPS reference profiles. The deviations in net photosynthesis for JSBACH-no-clumping and JULES are minor overall, indicating a good performance of the canopy 1-D RT schemes in these cases. This is expected, because the closed canopy corresponds to the idealized case where the 1-D RT schemes are supposed to perform best. Overall, JULES performed best of all models, although even for this model substantial deviations occur.

These deviations in net photosynthesis rates can be considerable, up to $10 \mu\text{molCm}^{-2}\text{s}^{-1}$, corresponding to up to 25 % of the photosynthesis rates. These conditions (with high LAI and $\theta = 60^\circ$) commonly prevail. Therefore, the impacts for the total calculated carbon budget are likely to be similarly large, which implies that the deviations in faPAR profiles detected in the current study are of critical importance for global carbon cycle studies. At lower incident radiation conditions, e.g. for $\theta = 83^\circ$, deviations amount even up to more than 75 %. However, given the lower incident radiation and consequently lower net photosynthesis, their impacts on the global carbon balance will be less.

3.2 Surface albedo

Figure 5 shows deviations of simulated surface albedo compared to RAMI4PILPS reference solutions for the entire solar spectrum (0.3...3.0 μm). Deviations in the entire solar spectrum are of particular importance as these determine the net effect on the surface radiation budget. More details on the separate deviations in the visible and infrared spectral bands is provided in the Appendix (Figs. A1 and A2).

The closed canopy experiments (CFC, GRA) show very similar results, as expected given that they only differ in their leaf orientation (random vs. erectophile). Largest deviations to the RAMI4PILPS references solutions are found for bright surface back-

ground (SNW) for JSBACH and ORCHIDEE models. Deviations for these two models are between $0.18 \leq \Delta\alpha \leq 0.25$ for SNW experiments. In contrast, the JULES albedo simulations show much smaller deviations for all closed canopy experiments ($-0.03 \leq \Delta\alpha \leq 0.02$). Also here, largest deviations are observed for bright backgrounds.

For soils with medium albedo (MED), the deviations are smaller for JSBACH and ORCHIDEE. Positive biases are obtained for isotropic conditions and small solar zenith angles ($-0.01 \leq \Delta\alpha \leq 0.06$). For large solar zenith angles, JULES shows minor deviations, while JSBACH and ORCHIDEE show a negative bias of $\Delta\alpha = -0.06$. While the deviations for JSBACH and ORCHIDEE are becoming smaller with increasing leaf area index for bright surface, due to an increased masking of the bright background, $\Delta\alpha$ increases with increasing LAI for the medium bright soils.

Open forest canopies (OFC) and shrubland (SHR) simulations show larger deviations than the closed canopy cases. Large positive deviations ($0.07 \leq \Delta\alpha \leq 0.36$) are observed for the snow covered cases with increasing deviations with increasing solar zenith angle. The larger deviations result from a lack of the representation of canopy shadowing effects in all of the canopy RT models, resulting in an overestimation of the simulated surface albedo. For medium soils, a positive bias is observed for all models. The deviations are larger ($0.01 \leq \Delta\alpha \leq 0.06$) than for the closed canopies and an increase in $\Delta\alpha$ with increasing leaf area index is observed.

3.2.1 Does it matter?

Effect of systematic albedo biases

It was shown in the previous section that the investigated models show albedo biases of $-0.27 < \Delta\alpha < 0.36$. The extreme deviations however occur typically either at very large sun zenith angles or at open forest canopies with snow as a background. In case of snow-free and closed canopies, the typical albedo biases for isotropic illumination conditions and small solar zenith angles are in the order of $0.02 < \Delta\alpha < 0.05$ for closed canopies.

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The impact of albedo biases on the calculated radiative forcing sensitivity (λ) is summarized in Table 3. The mean radiative forcing sensitivity for trees (grassland) is -5.5 (-13.11) [W m^{-2}]. Thus a change of surface albedo of e.g. $\Delta\alpha = 0.04$ would correspond to a radiative forcing of -0.22 (-0.52) [W m^{-2}]. The use of different global radiation datasets as input has a minor impact on the radiative forcing estimates. The standard deviation of RF caused by different radiation data is $0.3 \leq \sigma \leq 0.45$ ($1.03 \leq \sigma \leq 1.41$) for trees (grassland). However, much larger differences are observed for different assumptions for Γ^\uparrow . For clear sky conditions ($\Gamma^\uparrow = \text{const}$), the radiative cooling effect is 50. . .60% larger than for cloudy skies ($\Gamma^\uparrow = \Gamma^\downarrow$). It is therefore expected that the radiative forcing for the two investigated PFT types will be somewhere in between these two extremes.

Figure 6 shows the spatial distribution of the temporal mean radiative forcing sensitivity for tree and grassland for both assumptions of Γ^\uparrow . The much larger RF sensitivity for grassland is mainly due to its larger spatial coverage globally as well as due to the higher abundance in tropical areas with large insolation. For trees, the highest radiative cooling is observed in the tropics with a secondary maximum in boreal regions which is mainly during boreal summer.

Given the similarity in albedo bias for closed forest and grassland canopies, compared to RAMI4PILPS references, the radiative forcing of the forest and grassland PFTs might be summed. The global mean RF sensitivity is therefore -9.31 [W m^{-2}], assuming equal probability for the two parameterizations of atmospheric transmissivity. For a typical albedo bias of $\Delta\alpha = 0.04$, this would result in $RF_{\text{tot}} = -0.37$ [W m^{-2}].

Diurnal effects

Figure 7 shows the seasonal cycle of the changes in radiative forcing caused by neglecting the albedo diurnal cycle for grassland and tree covered areas for different leaf areas and parameterizations of the upward atmospheric transmissivity (Γ^\uparrow).

The maximum radiative cooling effect is $-1.25 < \text{RF}_{\text{tot}} < -0.8$ [W m^{-2}]. The radiative cooling is mainly caused by large albedo biases in tropical areas and changes substantially throughout the year.

With increasing leaf-area index, the radiative forcing effect increases. In general, the radiative cooling by grassland areas is higher than for tree covered areas due to their different spatial distribution. The annual mean radiative forcing is -0.61 (-0.53) and -0.84 (-0.73) for a leaf-area index of one and four, respectively, assuming $\Gamma^{\uparrow} = \text{const}$ ($\Gamma^{\uparrow} = \Gamma^{\downarrow}$). While the LAI and seasonal cycle have the largest impacts on RF_{tot} , also a different choice in the parameterization of Γ^{\uparrow} results in deviations of $\approx 15\%$.

3.3 Consistency of surface energy fluxes

Energy conservation was evaluated for the different experiments following Appendix D. Results are illustrated in Fig. D1. The JSBACH and ORCHIDEE models are not energy conservative for high leaf area and bright background (SNW). The JSBACH model shows clear deviations for the closed canopy cases (CFC, GRA) for almost all experiments with bright background.

The JULES model is the only model which is simulating the canopy RT fluxes in a consistent manner. No obvious violation of the energy conservation is found for JULES.

It needs to be emphasized however that the energy conservation might be violated also in the cases where the calculated canopy transmission is within its physical limits ($0 \leq \Gamma_{\text{can}} \leq 1$). It can however not be quantified in this study as the canopy transmission information is not available for the different models. JSBACH and ORCHIDEE estimates for Γ_{can} often differ from those of JULES by a factor of two or more. If we assume that JULES simulations are closest to energy conservation it is rather likely these models would not be energy conserving for experiments where a large deviation from the transmission estimates of JULES is observed.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Discussion

DGVMs are widely used to simulate land surface dynamics in coupled Earth System models, to assess the role of the biosphere in the Earth System and to study global water, energy and carbon fluxes. This paper aims to raise awareness of potential problems and impacts associated with different canopy radiative transfer schemes and their impacts on carbon fluxes and global climate simulations.

Large deviations in faPAR were found. In most cases, faPAR was underestimated with up to 60 % underestimation in open canopy cases with high solar zenith angles and/or snow as a background. Only in a few instances higher faPAR than in the RAMI4PILPS reference were found (typically in closed canopy cases with low zenith angles or isotropic insolation). Moreover, deviations were strongly model-dependent. JSBACH and ORCHIDEE consistently had the highest (negative) biases. This suggests that a two-stream approximation such as proposed by Sellers (1985) or other consistent treatments of radiative transfer are essential for obtaining unbiased faPAR calculations Pinty et al. (2006).

For surface albedo, largest deviations in surface albedo were observed in case of bright background albedo (SNW), for both open and closed canopies. High deviations ($\Delta\alpha > 0.2$) were diagnosed for the simplified surface albedo schemes implemented in JSBACH and ORCHIDEE. The albedo model of JULES only shows minor deviations ($-0.03 \leq \Delta\alpha \leq -0.01$) for closed canopy cases, but also large deviations for the open canopy cases. For surfaces with a typical soil background and a closed canopy, the 1-D canopy RT model of JULES is superior compared to the simple albedo schemes of JSBACH and ORCHIDEE which have an albedo bias which is twice to five times larger than the JULES albedo bias.

The simplified surface albedo schemes of JSBACH and ORCHIDEE do not take into account the diurnal variability in surface albedo. Isotropic insolation conditions are assumed instead. The impact of this assumption on global scale was therefore analyzed. While the applied assessment of radiative forcing is simple, we use state-of-the

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



art observations to get a most realistic assessment of RF_{tot} and its uncertainties. The developed framework can be used in general to estimate radiative forcing effects of temporally invariant as well as changing surface albedo biases.

While we have identified faPAR and albedo biases by comparing to RAMI4PILPS experiments, it needs to be emphasized that the RAMI4PILPS experiments provide idealized cases of canopies and that real surfaces and vegetation patches are much more complicated. For instance, none of the analyzed canopy RT schemes is capable to take into account large scale vegetation structural effects, like e.g. mutual shadowing or the scattering between multiple canopy elements.

4.1 Do we (need to) care?

Existing representations of shortwave radiative flux estimates in DGVMs have been developed throughout the last decades. Developing a DGVM always requires prioritization of resources between different model components. Recent DGVMs become more and more complex and incorporate complex carbon fluxes and nutrient cycling. In the past, a lot of attention has been devoted towards an improvement of especially the biochemical components in DGVMs (Goll et al., 2012; Zaehle and Friend, 2010; Brovkin et al., 2012), whereas biophysical components, like the canopy RT have been neglected because no study had analyzed systematically the major caveats of existing schemes so far. While model benchmarking and evaluation studies detail particular deviations of model results compared to observations and assess a (relative) model skill (Gleckler et al., 2008; Luo et al., 2012; Brovkin et al., 2013b; Hagemann et al., 2013), they typically do not provide an assessment of potential impacts of the observed model deficits.

It was therefore one of the objectives of the present study to provide an assessment of the implications due to a choice for a particular canopy radiative transfer scheme. Answering the question “Does it matter?” is of particular importance when decisions on further DGVM development need to be made with limited resources.

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy
radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The differences in faPAR discussed in the previous section do not translate straightforwardly into differences in net photosynthesis, and depend on total radiation and the extent to which CO₂ or light is limiting photosynthesis in different canopy layers (which again depends on the maximum capacity of the light and CO₂ limited photosynthesis pathways and on faPAR profiles through the canopy). Here, we only evaluated the impacts on net photosynthesis for some of the best (i.e. least deviating) cases, i.e. for closed forest canopies. We chose this setting, because the assumptions of the implementation of the Farquhar-photosynthesis scheme in DGVMs only apply to closed (forest) canopies. In that sense, it is important to recognize that some impacts on net photosynthesis (discussed below) may be even worse for open shrub or forest canopies, let alone for situations where one PFT literally grows on top of another affecting the light regime of the one below.

Moreover, given that the differences in faPAR are situation- and model-specific, the impacts are also expected to differ for various regions and models. Thus, we selected two common closed canopy cases (boreal forests and tropical forests) which we parameterized with representative radiation at top of canopy and maximum photosynthesis capacities. In contrast to faPAR, net photosynthesis was most affected at medium zenith angles (when radiation at top of canopy was still high) and high LAI (while faPAR was almost unaffected by LAI). While in this study net photosynthesis rates and their biases were not integrated to diurnal values, conditions of medium zenith angle occur commonly and large deviations in net photosynthesis estimates are thus expected. These deviations were on average stronger for the boreal forest than for the tropical forest case.

Likely, these deviations have been partly captured by tuning maximum photosynthesis rates (which may explain their deviations from the observed means (Kattge et al., 2011; Verheijen et al., 2013). Even so, given that the magnitude and direction differ between different model experiments, tuning will not allow capturing regional biases. While other factors like biases in precipitation also likely contribute to regional differences in GPP estimates between DGVMs and flux observation-derived estimates

(Beer et al., 2010), our results suggests the importance of incorporating consistent and adequate radiative transfer schemes.

Likewise a typical surface albedo bias in the order of $\Delta\alpha = 0.04$ results in a mean radiative forcing of $RF_{\text{tot}} = -0.22(-0.52) [\text{W m}^{-2}]$ for trees (grasslands). In addition it was shown in Sect. 3.2.1 that neglecting the diurnal cycle results in a radiative cooling of between -1.25 and $-0.8 [\text{W m}^{-2}]$, dependent on the leaf area index and atmospheric opacity model chosen. While an actual assessment of the radiative forcing effect would require simulations with a coupled (land-atmosphere) climate model, the provided values are likely to provide a realistic range of RF_{tot} estimates. These radiative forcing effects correspond to a considerable cooling, which is mainly caused by tropical land areas. This emphasizes that the diurnal cycle cannot be neglected for albedo calculations. Especially in tropical areas, where the insulation conditions change strongly throughout the day and solar insulation is high, a large effect on RF_{tot} is observed.

It needs to be emphasized that these results for net photosynthesis and radiative forcing are only valid for some vegetation settings and are tailored to provide only the order of magnitude of the impact of faPAR and surface albedo biases on net photosynthesis and radiative forcing.

4.2 A way forward?!

A combination of different approaches could lead to a general improvement of the representation of canopy radiation fluxes in DGVMs.

- *Model evaluation* is essential for any model development. Only once the target is clearly defined and accurate reference solutions are available, the potential biases in canopy RT schemes can be rigorously identified. In case that tolerance criteria are specified which allow to judge the relevance of deviations between models and a defined reference, model evaluation studies can determine the urgency of new developments to improve relevant model formulations (Widlowski et al., 2013). Specific test cases, like those provided by RAMI4PILPS are required for

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that purpose as well as large scale model evaluation studies (Hagemann et al., 2013; Brovkin et al., 2013b). The current RAMI4PILPS experiments provide a very idealized setup which does not represent real canopies in a realistic manner. To approach a fully quantitative and realistic assessment of canopy RT schemes in climate models, several aspects are of particular importance: (a) snow cover strongly affects the surface albedo. Current RAMI4PILPS experiments assume snow only below the canopy, while the surface albedo of a model grid cell is largely determined by snow on top and within the canopy. It is therefore recommended to represent snow cover dynamics in a more realistic way in RAMI4PILPS. (b) To allow for an appropriate assessment of the impact of biases in surface albedo and faPAR, it is important to provide reference solutions which cover a large variety of illumination conditions (representation of diurnal cycle) and which provide information on the reflected and absorbed fluxes also within distinct canopy layers. This is required e.g. for the quantification of the effect of the faPAR bias on net photosynthesis as discussed. (c) RAMI4PILPS neglects the woody part of the vegetation (stems, branches). These have a major effect on the surface albedo in high latitudes, especially when the leaf area index is negligible and solar zenith angle is large and the ground is covered by snow. In these cases, stems contribute to a large part to shadowing of the bright background (snow). An accurate representation of this masking effect in models is important for the sensitivity of surface temperature to surface albedo. (d) Vegetation types within RAMI4PILPS that more closely resemble plant functional types as defined in DGVMs, would allow an easier translation of results.

- *Model re-parameterization* could be used to compensate (partly) for some of the biases observed between models and references. Structural deficits in canopy RT schemes are likely to be not overcome with a re-parameterization of models, but could reduce their effect on model simulations. Some of the observed faPAR and albedo biases may be reduced by simply changing model parameters like e.g. leaf albedo (α_{leaf}), canopy single scattering albedo (ω) or specific leaf area (SLA) to

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



artificially converge model simulations towards mean reference data. This might reduce considerably potential biases and thus potential impacts on e.g. global radiative forcing, although it will not allow alleviating regional biases.

- *Improved consistency of existing schemes*: the present study has clearly shown that simplified albedo schemes result in considerable biases and radiative forcing which is not negligible. As a first step, the simplified surface albedo models should be replaced by 1-D RT schemes that are consistent with those used by the same models when computing faPAR. An adaptation of the 1-D-RT model for a consistent calculation of canopy absorption and reflectance should require medium effort, would be physically consistent and could also provide more accurate calculations of surface albedo diurnal cycle.
- *Effective radiative model parameters*: a replacement of 1-D canopy RT schemes by more complex 3-D canopy RT schemes in DGVMs is not foreseeable in the future as the required canopy structural information required for 3-D models is not available as prognostic variables from DGVMs and because of the much higher computational costs of these models. Effective radiative model parameters, however, aim to obtain realistic surface flux estimates and flux partitioning by means of physics-based 1-D radiative transfer formulations (Pinty et al., 2006). Effective radiative model parameters therefore compensate for inherent differences in process descriptions between simple (1-D) and complex (3-D) radiative modelling approaches. The key challenge, however, is to develop a deterministic or empirical relationship between model state variables and radiative model effective parameters which satisfies the consistency of canopy RT fluxes in agreement with observations. Pinty et al. (2006) have shown that the 3-D canopy radiative transfer problem can be approximated by a 1-D model using effective leaf area index as well as effective parameters for the vegetation optical properties. Different approaches have been proposed to relate a leaf area index to an effective leaf area index by defining a structural parameter which accounts for vegetation clumping

DGVM canopy radiation

A. Loew et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



(Yang et al., 2001; Pinty, 2004; Ni-Meister et al., 2010; Haverd et al., 2012). New globally available dataset of effective RT model parameters as derived from satellite observations (Pinty et al., 2011) could facilitate RT model parametrizations of DGVMs for different plant functional types. Given the fact that it has been shown that 1-D models can provide appropriate RT fluxes when parametrized in the right way, these novel dataset provide a novel source for the re-parameterization of canopy RT models at the global scale.

- As an alternative, effective vegetation model parameters or structural parameters may be obtained through *re-calibration using global satellite observations*. Global satellite observations provide longterm and high resolution estimates of surface variables, like surface albedo, faPAR and canopy radiation flux partitioning (Tucker et al., 2005; Loew and Govaerts, 2010; Pinty et al., 2011). However, a direct assimilation of geophysical EO products might be complicated due to e.g. different definitions of geophysical variables among different products as well as between a DGVM and an EO based observational dataset. Unless the canopy RT schemes in DGVMs do not satisfy basic principles of energy conservation and the appropriate representation of canopy scattering processes, an assimilation of EO based surface albedo or faPAR products in DGVMs might lead to erroneous results. As an example, faPAR satellite products show a wide range of variability due to different definitions and algorithms applied for the retrieval. A comprehensive review is given by Gobron and Verstraete (2009). Dahlke et al. (2012) compared different globally available faPAR satellite products and found considerable biases between the different data products, whereas the vegetation seasonality was consistently captured by the different investigated data products. Similarly, an assimilation of satellite leaf area indices or faPAR products in DGVMs might be complicated or impossible due to different canopy radiation schemes applied. While DGVMs use 1-D-canopy radiation schemes, as discussed, some satellite based products of vegetation variables (LAI, faPAR) are based on the inversion of 3-D canopy radiation models (Myneni et al., 2002). As a consequence, it is not physically valid to

ingest the retrieved values of for instance LAI into 1-D representations such as those provided by 2-stream approaches. The usage of effective variables in data assimilation schemes is required instead.

5 Conclusions

5 Major conclusions of the present study are:

1. *Considerable biases*: currently used simplified canopy RT schemes produce considerable biases in both surface albedo and faPAR calculations. Very simple albedo schemes result in large albedo biases ($\Delta\alpha > 0.3$) in snow covered cases and also much higher albedo biases for surfaces with a typical soil albedo. Our first assessment of the impacts of these biases indicate that these deviations lead to a considerable radiative forcing when not corrected for by an appropriate model parameterization while the faPAR differences can lead to biased estimates of GPP.
- 10 2. *Albedo diurnal cycle*: considering the albedo diurnal cycle in global surface albedo schemes is of major importance. State-of-the-art DGVM and climate models should therefore implement albedo schemes that consider appropriately the surface albedo diurnal behavior.
- 15 3. *Physical inconsistency*: canopy radiation schemes that use separate approaches to simulate reflected and absorbed radiation fluxes can lead to physically meaningless results. For some of the simulated experiments, it was clearly shown that the investigated models violate the conservation law of energy for the canopy RT fluxes.
- 20 4. *RAMI4PILPS continued*: the current RAMI4PILPS experiments (Widlowski et al., 2011) should be extended to provide more realistic settings for evaluating DGVMs.

Such continued RAMI4PILPS experiments may provide the basis for a more comprehensive analysis for to assessing the impact of actual surface albedo changes on climate using simulations of a coupled Earth System model.

5 *Using two-stream based 1-D canopy radiative transfer schemes for faPAR and surface albedo calculations* is a useful approach to minimize faPAR and albedo biases and to obtain consistent estimates of absorbed and reflected canopy radiation fluxes. Such schemes for radiative transfer in canopies are quite well understood and. The implementation of such accurate canopy RT simulations in DGVMs should be comparably easy compared to much more complicated processes like e.g. carbon fluxes. The major challenge will, however, be to decide
10 how to best link model prognostic variables like e.g. LAI to radiative effective variables needed for a 1-D canopy RT model. Implementing physically consistent canopy RT schemes in DGVMs would also allow for the assimilation of satellite based observations of canopy RT fluxes at the global scale and open new perspectives for DGVM model parameterization at the regional to global scale. This
15 paper is therefore a plea for a more rigorous treatment of surface and canopy radiation fluxes in DGVMs which has been largely neglected in the past.

Appendix A

Surface albedo deviations for VIS and NIR

20 Figure 5 shows surface albedo deviations for the entire solar spectrum. The following figures show the deviations of surface albedo for the visible and near infrared shortwave bands separately.

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Appendix B

PFT distributions

See Fig. B1.

Appendix C

5 CERES radiation and one-way transmissivity

The climatological mean one-way atmospheric transmissivity was derived from CERES top-of-atmosphere and surface radiation fluxes using Eq. (17). Resulting distribution of Γ^{\downarrow} is shown in Fig. C1.

Appendix D

10 Energy conservation

The energy conservation was tested by estimating the theoretical canopy transmission (T) under the assumption of energy conservation. The following figures show where the energy conservation is clearly violated (red). Note however, that also experiments with $0 \leq T \leq 1$ can be physically inconsistent, which could be however not be tested, as the values for T are not available from the models.

15 *Acknowledgements.* Collaboration among the authors was supported by the TERRABITES COST Action ES0805 and by the International Space Science Institute (nr. 263). The main author was supported through the Cluster of Excellence “CliSAP” (EXC177), University of Hamburg, funded through the German Science Foundation (DFG) and the ESA Climate Change Initiative Climate Modelling User Group (contract: 4000100222/10/I-AM). Juliane Otto
20 was funded through ERC starting grant 242564. Atmospheric radiation data were obtained

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from the NASA Langley Research Center Atmospheric Science Data Center. Tristan Quaife was funded by the European Space Agency (contract 4000104980/11/I-LG) with additional support from the UK National Centre for Earth Observation. The authors thank Julia Pongratz for reviewing a previous version of the manuscript.

The service charges for this open access publication have been covered by the Max Planck Society.

References

- Abramowitz, G., Leuning, R., Clark, M., and Pitman, A.: Evaluating the performance of land surface models, *J. Climate*, 21, 5468–5481, doi:10.1175/2008JCLI2378.1, 2008. 16554
- Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rödenbeck, C., Arain, M. A., Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M., Luysaert, S., Margolis, H., Oleson, K. W., Rouspard, O., Veenendaal, E., Viovy, N., Williams, C., Woodward, F. I., and Papale, D.: Terrestrial gross carbon dioxide uptake: global distribution and covariation with climate, *Science*, 329, 834–838, doi:10.1126/science.1184984, 2010. 16582
- Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., and Harding, R. J.: The Joint UK Land Environment Simulator (JULES), model description – Part 1: Energy and water fluxes, *Geosci. Model Dev.*, 4, 677–699, doi:10.5194/gmd-4-677-2011, 2011. 16560
- Blyth, E., Clark, D. B., Ellis, R., Huntingford, C., Los, S., Pryor, M., Best, M., and Sitch, S.: A comprehensive set of benchmark tests for a land surface model of simultaneous fluxes of water and carbon at both the global and seasonal scale, *Geosci. Model Dev.*, 4, 255–269, doi:10.5194/gmd-4-255-2011, 2011. 16554
- Bright, R. M., Cherubini, F., and Strømman, A. H.: Climate impacts of bioenergy: inclusion of carbon cycle and albedo dynamics in life cycle impact assessment, *Environ. Impact Assess.*, 37, 2–11, doi:10.1016/j.eiar.2012.01.002, 2012. 16569, 16570, 16600, 16601

BGD

10, 16551–16613, 2013

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Brovkin, V., Raddatz, T., Reick, C. H., Claussen, M., and Gayler, V.: Global biogeophysical interactions between forest and climate, *Geophys. Res. Lett.*, 36, L07405, doi:10.1029/2009GL037543, 2009. 16561
- Brovkin, V., van Bodegom, P. M., Kleinen, T., Wirth, C., Cornwell, W. K., Cornelissen, J. H. C., and Kattge, J.: Plant-driven variation in decomposition rates improves projections of global litter stock distribution, *Biogeosciences*, 9, 565–576, doi:10.5194/bg-9-565-2012, 2012. 16580
- Brovkin, V., Boysen, L., Arora, V. K., Boisier, J. P., Cadule, P., Chini, L., Claussen, M., Friedlingstein, P., Gayler, V., van den Hurk, B. J. J. M., Hurtt, G. C., Jones, C. D., Kato, E., de Noblet-Ducoudré, N., Pacifico, F., Pongratz, J., and Weiss, M.: Effect of anthropogenic land-use and land cover changes on climate and land carbon storage in CMIP5 projections for the 21st century, *J. Climate*, 26, 6859–6881, doi:10.1175/JCLI-D-12-00623.1, 2013a. 16553
- Brovkin, V., Boysen, L., Raddatz, T., Gayler, V., Loew, A., and Claussen, M.: Evaluation of vegetation cover and land-surface albedo in MPI-ESM CMIP5 simulations, *J. Adv. Model. Earth Sys.*, 5, 48–57, doi:10.1029/2012MS000169, 2013b. 16554, 16561, 16580, 16583
- Carrer, D., Roujean, J.-L., Lafont, S., Calvet, J.-C., Boone, A., Decharme, B., Delire, C., and Gastellu-Etchegorry, J.-P.: A canopy radiative transfer scheme with explicit FAPAR for the interactive vegetation model ISBA-A-gs: impact on carbon fluxes, *J. Geophys. Res.-Biogeo.*, 118, 888–903, doi:10.1002/jgrg.20070, 2013. 16554
- Chen, Q., Baldocchi, D., Gong, P., and Dawson, T.: Modeling radiation and photosynthesis of a heterogeneous savanna woodland landscape with a hierarchy of model complexities, *Agr. Forest Meteorol.*, 148, 1005–1020, doi:10.1016/j.agrformet.2008.01.020, 2008. 16557
- Cherubini, F., Bright, R. M., and Strömman, A. H.: Site-specific global warming potentials of biogenic CO₂ for bioenergy: contributions from carbon fluxes and albedo dynamics, *Environ. Res. Lett.*, 7, 045902, doi:10.1088/1748-9326/7/4/045902, 2012. 16570
- Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C., and Cox, P. M.: The Joint UK Land Environment Simulator (JULES), model description – Part 2: Carbon fluxes and vegetation dynamics, *Geosci. Model Dev.*, 4, 701–722, doi:10.5194/gmd-4-701-2011, 2011. 16560
- Cox, P.: Description of the TRIFFID dynamic global vegetation model, Tech. rep., Hadley Centre, Met Office, London Road, Bracknell, Berks, RG122SY, UK, 2001. 16560

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dahlke, C., Loew, A., and Reick, C.: Robust identification of global greening phase patterns from remote sensing vegetation products, *J. Climate*, 25, 8289–8307, doi:10.1175/JCLI-D-11-00319.1, 2012. 16585

Diaz, S. and M., Cabido: Plant functional types and ecosystem function in relation to global change, *J. Veg. Sci.*, 8, 463–474, 1997. 16555

Dickinson, R. E., Zhou, L., Tian, Y., Liu, Q., Lavergne, T., Pinty, B., Schaaf, C. B., and Knyazikhin, Y.: A three-dimensional analytic model for the scattering of a spherical bush, *J. Geophys. Res.*, 113, D20113, doi:10.1029/2007JD009564, 2008. 16554

DiMiceli, C., Carroll, M., Sohlberg, R., Huang, C., Hansen, M., and Townshend, J.: Annual Global Automated MODIS Vegetation Continuous Fields (MOD44B) at 250 m Spatial Resolution for Data Years Beginning Day 65, 2000–2010, Collection 5 Percent Tree Cover, University of Maryland, College Park, MD, USA, 2011. 16573

Disney, M., Lewis, P., and North, P.: Monte Carlo ray tracing in optical canopy reflectance modelling, *Remote Sens. Rev.*, 18, 163–196, doi:10.1080/02757250009532389, 2000. 16558

Domingues, T. F., Meir, P., Feldpausch, T. R., Saiz, G., Veenendaal, E. M., Schrodt, F., Bird, M., Djagbletey, G., Hien, F., Compaore, H., Diallo, A., Grace, J., and Lloyd, J.: Co-limitation of photosynthetic capacity by nitrogen and phosphorus in West Africa woodlands, *Plant Cell Environ.*, 33, 959–80, doi:10.1111/j.1365-3040.2010.02119.x, 2010. 16569

Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benshila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, a., Cugnet, D., Noblet, N., Duvel, J.-P., Ethé, C., Fairhead, L., Fichet, T., Flavoni, S., Friedlingstein, P., Grandpeix, J.-Y., Guez, L., Guilyardi, E., Hauglustaine, D., Hourdin, F., Idelkadi, A., Ghattas, J., Jousaume, S., Kageyama, M., Krinner, G., Labetoulle, S., Lahellec, A., Lefebvre, M.-P., Lefevre, F., Levy, C., Li, Z. X., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N., and Vuichard, N.: Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, *Clim. Dynam.*, 40, 2123–2165, doi:10.1007/s00382-012-1636-1, 2013. 16563, 16564

Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.-G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate-carbon

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cycle feedback analysis: results from the C4MIP model intercomparison, *J. Climate*, 19, 3337–3353, doi:10.1175/JCLI3800.1, 2006. 16553

Gastellu-Etchegorry, J.-P., Martin, E., and Gascon, F.: DART: a 3-D model for simulating satellite images and studying surface radiation budget, *Int. J. Remote Sens.*, 25, 73–96, 2004. 16554

5 Gleckler, P. J., Taylor, K. E., and Doutriaux, C.: Performance metrics for climate models, *J. Geophys. Res.*, 113, D06104, doi:10.1029/2007JD008972, 2008. 16580

Gobron, N. and Verstraete, M. M.: FAPAR, available at: <http://www.fao.org/gtos/doc/ECVs/T10/T10.pdf>, 2009. 16585

10 Gobron, N., Pinty, B., Verstraete, M. M., and Govaerts, Y.: A semidiscrete model for the scattering of light by vegetation, *J. Geophys. Res.*, 102, 9431–9446, doi:10.1029/96JD04013, 1997. 16556

Goll, D. S., Brovkin, V., Parida, B. R., Reick, C. H., Kattge, J., Reich, P. B., van Bodegom, P. M., and Niinemets, Ü.: Nutrient limitation reduces land carbon uptake in simulations with a model of combined carbon, nitrogen and phosphorus cycling, *Biogeosciences*, 9, 3547–3569, doi:10.5194/bg-9-3547-2012, 2012. 16580

Hagemann, S. and Stacke, T.: Impact of the soil hydrology scheme on simulated soil moisture memory, in preparation, 2013. 16554

15 Hagemann, S., Loew, A., and Andersson, A.: Combined evaluation of MPI-ESM land surface water and energy fluxes, *J. Adv. Model. Earth Sys.*, 5, 259–286, doi:10.1029/2012MS000173, 2013. 16554, 16561, 16580, 16583

Hall, A. and Qu, X.: Using the current seasonal cycle to constrain snow albedo feedback in future climate change, *Geophys. Res. Lett.*, 33, 1–4, doi:10.1029/2005GL025127, 2006. 16569

20 Harrison, S. P., Prentice, I. C., Barboni, D., Kohfeld, K. E., Ni, J., and Sutra, J.-P.: Ecophysiological and bioclimatic foundations for a global plant functional classification, *J. Veg. Sci.*, 21, 300–317, doi:10.1111/j.1654-1103.2009.01144.x, 2010. 16553

Haverd, V., Lovell, J., Cuntz, M., Jupp, D., Newnham, G., and Sea, W.: The Canopy Semi-analytic Pgap And Radiative Transfer (CanSPART) model: formulation and application, *Agr. Forest Meteorol.*, 160, 14–35, doi:10.1016/j.agrformet.2012.01.018, 2012. 16554, 16556, 16557, 16585

25 IPCC: Climate Change 2007: synthesis report, Tech. rep., Intergovernmental Panel on Climate Change, 2007. 16569, 16570

30 Kattge, J., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Bönsch, G., Garnier, E., Westoby, M., Reich, P. B., Wright, I. J., Cornelissen, J. H. C., Violle, C., Harrison, S. P., Van bode-

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



gom, P. M., Reichstein, M., Enquist, B. J., Soudzilovskaia, N. A., Ackerly, D. D., Anand, M.,
Atkin, O., Bahn, M., Baker, T. R., Baldocchi, D., Bekker, R., Blanco, C. C., Blonder, B.,
Bond, W. J., Bradstock, R., Bunker, D. E., Casanoves, F., Cavender-Bares, J., Cham-
bers, J. Q., Chapin III, F. S., Chave, J., Coomes, D., Cornwell, W. K., Craine, J. M., Do-
brin, B. H., Duarte, L., Durka, W., Elser, J., Esser, G., Estiarte, M., Fagan, W. F., Fang, J.,
Fernández-Méndez, F., Fidelis, A., Finegan, B., Flores, O., Ford, H., Frank, D., Fres-
chet, G. T., Fyllas, N. M., Gallagher, R. V., Green, W. A., Gutierrez, A. G., Hickler, T., Hig-
gins, S. I., Hodgson, J. G., Jalili, A., Jansen, S., Joly, C. A., Kerkhoff, A. J., Kirkup, D., Ki-
tajima, K., Kleyer, M., Klotz, S., Knops, J. M. H., and Kramer, K.: TRY – a global database
of plant traits, *Glob. Change Biol.*, 17, 2905–2935, doi:10.1111/j.1365-2486.2011.02451.x,
2011. 16569, 16581

Kinne, S., O'Donnel, D., Stier, P., Kloster, S., Zhang, K., Schmidt, H., Rast, S., Giorgetta, M.,
Eck, T. F., and Stevens, B.: HAC-v1: A new global aerosol climatology for climate studies, *J.*
Adv. Model. Earth Sys., doi:10.1002/jame.20035, 2013. 16572

Kobayashi, H. and Iwabuchi, H.: A coupled 1-D atmosphere and 3-D canopy radiative transfer
model for canopy reflectance, light environment, and photosynthesis simulation in a hetero-
geneous landscape, *Remote Sens. Environ.*, 112, 173–185, 2008. 16554

Krinner, G.: A dynamic global vegetation model for studies of the coupled atmosphere-
biosphere system, *Global Biogeochem. Cy.*, 19, GB1015, doi:10.1029/2003GB002199,
2005. 16563

Lenton, T. M. and Vaughan, N. E.: The radiative forcing potential of different climate geoengi-
neering options, *Atmos. Chem. Phys.*, 9, 5539–5561, doi:10.5194/acp-9-5539-2009, 2009.
16569, 16570, 16573

Lewis, P. and Barnsley, M.: Influence of the sky radiance distribution on various formulations
of the Earth surface albedo, in: *Proceedings of the 6th ISPRS International Symposium*
on Physical Measurements and Signatures in Remote Sensing, Val d'Isère, France, 17–21
January 1994, 707–715, 1994. 16572

Li, X. and Strahler, A.: Geometric-optical bidirectional reflectance modeling of a conifer for-
est canopy, *IEEE T. Geosci. Remote*, 24, 906–919, doi:10.1109/TGRS.1986.289706, 1986.
16558

Loeb, N. G., Kato, S., Su, W., Wong, T., Rose, F. G., Doelling, D. R., Norris, J. R., and Huang, X.:
Advances in understanding top-of-atmosphere radiation variability from satellite observa-
tions, *Surv. Geophys.*, 33, 359–385, doi:10.1007/s10712-012-9175-1, 2012. 16571

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Loew, A. and Govaerts, Y.: Towards multidecadal consistent meteosat surface albedo time series, *Remote Sens.*, 2, 957–967, doi:10.3390/rs2040957, 2010. 16585
- Luo, Y. Q., Randerson, J. T., Abramowitz, G., Bacour, C., Blyth, E., Carvalhais, N., Ciais, P., Dalmonech, D., Fisher, J. B., Fisher, R., Friedlingstein, P., Hibbard, K., Hoffman, F., Huntzinger, D., Jones, C. D., Koven, C., Lawrence, D., Li, D. J., Mahecha, M., Niu, S. L., Norby, R., Piao, S. L., Qi, X., Peylin, P., Prentice, I. C., Riley, W., Reichstein, M., Schwalm, C., Wang, Y. P., Xia, J. Y., Zaehle, S., and Zhou, X. H.: A framework for benchmarking land models, *Biogeosciences*, 9, 3857–3874, doi:10.5194/bg-9-3857-2012, 2012. 16554, 16580
- Meehl, G. A., Covey, C., Taylor, K. E., Delworth, T., Stouffer, R. J., Latif, M., McAvaney, B., and Mitchell, J. F. B.: The WCRP CMIP3 multimodel dataset: a new era in climate change research, *B. Am. Meteorol. Soc.*, 88, 1383–1394, doi:10.1175/BAMS-88-9-1383, 2007. 16560
- Mercado, L. M., Bellouin, N., Sitch, S., Boucher, O., Huntingford, C., Wild, M., and Cox, P. M.: Impact of changes in diffuse radiation on the global land carbon sink, *Nature*, 458, 1014–10177, doi:10.1038/nature07949, 2009. 16558
- Mueller, R., Behrendt, T., Hammer, A., and Kemper, A.: A new algorithm for the satellite-based retrieval of solar surface irradiance in spectral bands, *Remote Sens.*, 4, 622–647, doi:10.3390/rs4030622, 2012. 16572
- Myneni, R., Hoffman, S., and Knyazikhin, Y.: Global products of vegetation leaf area and fraction absorbed PAR from year one of MODIS data, *Remote Sens.*, 83, 214–231, doi:10.1016/S0034-4257(02)00074-3, 2002. 16585
- Ni, W., Li, X., Woodcock, C., Caetano, M., and Strahler, A.: An analytical hybrid GORT model for bidirectional reflectance over discontinuous plant canopies, *IEEE T. Geosci. Remote*, 37, 987–999, doi:10.1109/36.752217, 1999. 16558
- Ni-Meister, W., Yang, W., and Kiang, N. Y.: A clumped-foliage canopy radiative transfer model for a global dynamic terrestrial ecosystem model, 1: Theory, *Agr. Forest Meteorol.*, 150, 881–894, doi:10.1016/j.agrformet.2010.02.009, 2010. 16558, 16585
- Ostle, N. J., Smith, P., Fisher, R., Ian Woodward, F., Fisher, J. B., Smith, J. U., Galbraith, D., Levy, P., Meir, P., McNamara, N. P., and Bardgett, R. D.: Integrating plant-soil interactions into global carbon cycle models, *J. Ecol.*, 97, 851–863, doi:10.1111/j.1365-2745.2009.01547.x, 2009. 16553
- Otto, J., Raddatz, T., and Claussen, M.: Strength of forest-albedo feedback in mid-Holocene climate simulations, *Clim. Past*, 7, 1027–1039, doi:10.5194/cp-7-1027-2011, 2011. 16562

DGVM canopy
radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pinty, B.: Synergy between 1-D and 3-D radiation transfer models to retrieve vegetation canopy properties from remote sensing data, *J. Geophys. Res.*, 109, D21205, doi:10.1029/2004JD005214, 2004. 16555, 16585
- Pinty, B., Gobron, N., Widlowski, J.-L., Gerstl, S. A. W., Verstraete, M. M., Antunes, M., Ba-
 5 cour, C., Gascon, F., Gastellu, J.-P., Goel, N., Jacquemoud, S., North, P., Qin, W., and Thompson, R.: Radiation transfer model intercomparison (RAMI) exercise, *J. Geophys. Res.*, 106, 11937, doi:10.1029/2000JD900493, 2001. 16555
- Pinty, B., Lavergne, T., Dickinson, R. E., Widlowski, J.-L., Gobron, N., and Verstraete, M. M.: Simplifying the interaction of land surfaces with radiation for relating remote sensing prod-
 10 ducts to climate models, *J. Geophys. Res.*, 111, D02116, doi:10.1029/2005JD005952, 2006. 16554, 16557, 16579, 16584
- Pinty, B., Andredakis, I., Clerici, M., Kaminski, T., Taberner, M., Verstraete, M. M., Gobron, N., Plummer, S., and Widlowski, J.-L.: Exploiting the MODIS albedos with the Two-stream In-
 15 version Package (JRC-TIP): 1. Effective leaf area index, vegetation, and soil properties, *J. Geophys. Res.*, 116, D09105, doi:10.1029/2010JD015372, 2011. 16585
- Pongratz, J., Raddatz, T., Reick, C. H., Esch, M., and Claussen, M.: Radiative forcing from anthropogenic land cover change since AD 800, *Geophys. Res. Lett.*, 36, L02709, doi:10.1029/2008GL036394, 2009. 16569
- Posselt, R., Mueller, R., Stöckli, R., and Trentmann, J.: Remote sensing of solar surface radia-
 20 tion for climate monitoring – the CM-SAF retrieval in international comparison, *Remote Sens. Environ.*, 118, 186–198, doi:10.1016/j.rse.2011.11.016, 2012. 16571, 16572
- Poulter, B., Ciais, P., Hodson, E., Lischke, H., Maignan, F., Plummer, S., and Zimmer-
 mann, N. E.: Plant functional type mapping for earth system models, *Geosci. Model Dev.*, 4, 993–1010, doi:10.5194/gmd-4-993-2011, 2011. 16559
- Prentice, I. C., Bondeau, A., Cramer, W., Harrison, S. P., Hickler, T., Lucht, W., Sitch, S.,
 25 Smith, B., and Sykes, M. T.: Dynamic global vegetation modeling: quantifying terrestrial ecosystem responses to large-scale environmental change, in: *Terrestrial Ecosystems in a Changing World*, edited by: Canadell, J., Pataki, D. E., and Pitelka, L. F., IGBP Book series, chap. Dynamic GI, Springer, 175–192, available at: <http://www.springerlink.com/index/r765181510882450.pdf>, 2007. 16555
- Raddatz, T. J., Reick, C. H., Knorr, W., Kattge, J., Roeckner, E., Schnur, R., Schnitzler, K.-
 30 G., Wetzell, P., and Jungclaus, J.: Will the tropical land biosphere dominate the climate-

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



carbon cycle feedback during the twenty-first century?, *Clim. Dynam.*, 29, 565–574, doi:10.1007/s00382-007-0247-8, 2007. 16561

Reick, C. H., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and anthropogenic land cover change in MPI-ESM, *J. Adv. Model. Earth Sys.*, 5, 459–482, doi:10.1002/jame.20022, 2013. 16561

Rochdi, N., Fernandes, R., and Chelle, M.: An assessment of needles clumping within shoots when modeling radiative transfer within homogeneous canopies, *Remote Sens. Environ.*, 102, 116–135, 2006. 16554

Rossow, W. B. and Zhang, Y.-C.: Calculation of surface and top of atmosphere radiative fluxes from physical quantities based on ISCCP data sets: 2. Validation and first results, *J. Geophys. Res.*, 100, 1167, doi:10.1029/94JD02746, 1995. 16571

Schaepman-Strub, G., Schaepman, M., Painter, T., Dangel, S., and Martonchik, J.: Reflectance quantities in optical remote sensing – definitions and case studies, *Remote Sens. Environ.*, 103, 27–42, doi:10.1016/j.rse.2006.03.002, 2006. 16572

Sellers, P. J.: Canopy reflectance, photosynthesis and transpiration, *Int. J. Remote Sens.*, 6, 1335–1372, doi:10.1080/01431168508948283, 1985. 16553, 16560, 16561, 16562, 16579

Sharkey, T. D., Bernacchi, C. J., Farquhar, G. D., and Singsaas, E. L.: Fitting photosynthetic carbon dioxide response curves for C(3) leaves, *Plant Cell Environ.*, 30, 1035–1040, doi:10.1111/j.1365-3040.2007.01710.x, 2007. 16569

Sitch, S., Huntingford, C., Gedney, N., Levy, P. E., Lomas, M., Piao, S. L., Betts, R., Ciais, P., Cox, P., Friedlingstein, P., Jones, C. D., Prentice, I. C., and Woodward, F. I.: Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs), *Glob. Change Biol.*, 14, 2015–2039, 2008. 16553

Spitters, C., Toussaint, H., and Goudriaan, J.: Separating the diffuse and direct component of global radiation and its implications for modeling canopy photosynthesis, Part 1. Components of incoming radiation, *Agr. Forest Meteorol.*, 38, 217–229, doi:10.1016/0168-1923(86)90060-2, 1986. 16558

Sprintsin, M., Chen, J. M., Desai, A., and Gough, C. M.: Evaluation of leaf-to-canopy upscaling methodologies against carbon flux data in North America, *J. Geophys. Res.*, 117, G01023, doi:10.1029/2010JG001407, 2012. 16553

Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., Brokopf, R., Fast, I., Kinne, S., Kornblueh, L., Lohmann, U.,

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Pincus, R., Reichler, T., and Roeckner, E.: Atmospheric component of the MPI-M Earth System Model: ECHAM6, *J. Adv. Model. Earth Sys.*, 5, 146–172, doi:10.1002/jame.20015, 2013. 16561

Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *B. Am. Meteorol. Soc.*, 93, 485–498, doi:10.1175/BAMS-D-11-00094.1, 2012. 16560, 16561

Trenberth, K. E., Fasullo, J. T., and Kiehl, J.: Earth's global energy budget, *B. Am. Meteorol. Soc.*, 90, 311–323, doi:10.1175/2008BAMS2634.1, 2009. 16600, 16601

Tucker, C. J., Pinzon, J. E., Brown, M. E., Slayback, D., Pak, E. W., Mahoney, R., V., Ermote, E., and Saleous, N.: An extended AVHRR 8-km NDVI data set compatible with MODIS and SPOT vegetation NDVI data., *Int. J. Remote Sens.*, 26, 4485–4498, 2005. 16585

Vamborg, F. S. E., Brovkin, V., and Claussen, M.: The effect of a dynamic background albedo scheme on Sahel/Sahara precipitation during the mid-Holocene, *Clim. Past*, 7, 117–131, doi:10.5194/cp-7-117-2011, 2011. 16562

Van Bodegom, P. M., Douma, J. C., Witte, J. P. M., Ordoñez, J. C., Bartholomeus, R. P., and Aerts, R.: Going beyond limitations of plant functional types when predicting global ecosystem-atmosphere fluxes: exploring the merits of traits-based approaches, *Global. Ecol. Biogeogr.*, 21, 625–636, doi:10.1111/j.1466-8238.2011.00717.x, 2012. 16553

Verheijen, L. M., Brovkin, V., Aerts, R., Bönisch, G., Cornelissen, J. H. C., Kattge, J., Reich, P. B., Wright, I. J., and van Bodegom, P. M.: Impacts of trait variation through observed trait–climate relationships on performance of an Earth system model: a conceptual analysis, *Biogeosciences*, 10, 5497–5515, doi:10.5194/bg-10-5497-2013, 2013. 16569, 16581

von Caemmerer, S.: *Biochemical Models of Leaf Photosynthesis*, CSIRO PUBLISHING, available at: <http://www.publish.csiro.au/pid/1592.htm>, 2000. 16568

Wang, S., Chen, W., and Cihlar, J.: New calculation methods of diurnal distribution of solar radiation and its interception by canopy over complex terrain, *Ecol. Model.*, 155, 191–204, doi:10.1016/S0304-3800(02)00122-9, 2002. 16569

Widowski, J.-L.: On the bias of instantaneous FAPAR estimates in open-canopy forests, *Agr. Forest Meteorol.*, 150, 1501–1522, doi:10.1016/j.agrformet.2010.07.011, 2010. 16558

Widowski, J.-L., Pinty, B., Lavergne, T., Verstraete, M., and Gobron, N.: Using 1-D models to interpret the reflectance anisotropy of 3-D canopy targets: issues and caveats, *IEEE T. Geosci. Remote*, 43, 2008–2017, doi:10.1109/TGRS.2005.853718, 2005. 16554

Widowski, J.-L., Pinty, B., Lavergne, T., Verstraete, M. M., and Gobron, N.: Horizontal radiation transport in 3-D forest canopies at multiple spatial resolutions: simulated impact on

canopy absorption, *Remote Sens. Environ.*, 103, 379–397, doi:10.1016/j.rse.2006.03.014, 2006. 16557

Widowski, J.-L., Taberner, M., Pinty, B., Bruniquel-Pinel, V., Disney, M., Fernandes, R., Gastellu-Etchegorry, J.-P., Gobron, N., Kuusk, A., Lavergne, T., Leblanc, S., Lewis, P. E., Martin, E., Mörtus, M., North, P. R. J., Qin, W., Robustelli, M., Rochdi, N., Ruiloba, R., Soler, C., Thompson, R., Verhoef, W., Verstraete, M. M., and Xie, D.: Third Radiation Transfer Model Intercomparison (RAMI) exercise: documenting progress in canopy reflectance models, *J. Geophys. Res.*, 112, D09111, doi:10.1029/2006JD007821, 2007. 16555, 16565

Widowski, J.-L., Pinty, B., Clerici, M., Dai, Y., De Kauwe, M., de Ridder, K., Kallel, A., Kobayashi, H., Lavergne, T., Ni-Meister, W., Olchev, A., Quaife, T., Wang, S., Yang, W., Yang, Y., and Yuan, H.: RAMI4PILPS: an intercomparison of formulations for the partitioning of solar radiation in land surface models, *J. Geophys. Res.*, 116, G02019, doi:10.1029/2010JG001511, 2011. 16561, 16565, 16566, 16569, 16570, 16586

Widowski, J.-L., Pinty, B., Lopatka, M., Atzberger, C., Buzica, D., Chelle, M., Disney, M., Gastellu-Etchegorry, J.-P., Gerboles, M., Gobron, N., Grau, E., Huang, H., Kallel, A., Kobayashi, H., Lewis, P. E., Qin, W., Schlerf, M., Stuckens, J., and Xie, D.: The 4th radiation transfer model intercomparison (RAMI-IV): proficiency testing of canopy reflectance models with ISO-13528, *J. Geophys. Res.-Atmos.*, 118, 6869–6890, doi:10.1002/jgrd.50497, 2013. 16582

Wilson, M. and Henderson-Sellers, A.: A global archive of land cover and soils data for use in general circulation climate models, *J. Climate*, 5, 119–143, 1985. 16564

Yang, R., Friedl, M. A., and Ni, W.: Parameterization of shortwave radiation fluxes for nonuniform vegetation canopies in land surface models, *J. Geophys. Res.*, 106, 14275–14286, doi:10.1029/2001JD900180, 2001. 16585

Zaehle, S. and Friend, A. D.: Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model description, site-scale evaluation, and sensitivity to parameter estimates, *Global Biogeochem. Cy.*, 24, GB1005, doi:10.1029/2009GB003521, 2010. 16580

DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.

Table 1. Summary of experiment parameters (see Wildowski et al., 2011, for details); different soil backgrounds are specified as BLK = black, MED = Medium, SNW = snow, difference canopy types are specified as GRA = grassland, CFC = closed forest canopy, SHR = shrubland, OFC = open forest canopies.

Experiment	GRA	CFC	SHR	OFC
Vegetation type	homog. grass	closed forest	shrubs 3-D	open forest
$LAI_{\text{canopy}} [m^3 m^{-3}]$	1/2/4	1/2/4	2.5/2.5/2.5	5.0/5.0/5.0
$LAI_{\text{scene}} [m^3 m^{-3}]$	1/2/4	1/2/4	0.25/0.50/1.00	0.5/1.5/2.5
$\alpha_{s,VIS}/\alpha_{s,NIR}$	BLK: 0., 0./MED: 0.1217, 0.2142/SNW: 0.9640, 0.5568			
$\alpha_{le,VIS}/\alpha_{le,NIR}$	0.0735/0.3912			
$t_{\text{leaf,VIS}}/t_{\text{leaf,NIR}}$	0.0566/0.4146			
sun zenith angle [deg]	27.4643/60.0/83.5289/isotropic			
f_{veg}	1/1/1	1/1/1	0.1/0.2/0.4	0.1/0.3/0.5
$1 - P_{\text{gap}}(\theta = 0)$	1/1/1	1/1/1	0.068/0.136/0.273	0.088/0.26244/0.437

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy
radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Summary of atmospheric one-way or two-way transmissivity as derived from different sources. Input fluxes used for the transmissivity estimates are provided. Values are for land areas only.

Dataset	Γ_a^{\downarrow} [–]	$R_{\text{TOA}}^{\downarrow}$ [W m ⁻²]	$R_{\text{surf}}^{\downarrow}$ [W m ⁻²]	Reference
Trenberth et al. (2009)	0.47	341	161	Trenberth et al. (2009)
ISCCP-FD	0.45	330.1	147.2	Trenberth et al. (2009)
ERA-40	0.41	330.3	134.3	Trenberth et al. (2009)
JRA	0.47	328.2	154.9	Trenberth et al. (2009)
Bright et al. (2012)	0.45	–	–	Bright et al. (2012)
CERES EBAF2.6	0.58	324.7	188.1	this study

DGVM canopy radiation

A. Loew et al.

Table 3. Radiative forcing sensitivity ($\partial\text{RF}/\partial\Delta\alpha$) for different radiation datasets as derived from Eq. (19) for different global radiation datasets used to estimate the one-way atmospheric transmissivity and different assumptions on the calculation of r^\uparrow .

Radiation data	$\lambda_{\text{tree}} [\text{W m}^{-2}]$		$\lambda_{\text{grass}} [\text{W m}^{-2}]$	
	$r^\uparrow = 0.854$	$r^\uparrow = r^\downarrow$	$r^\uparrow = 0.854$	$r^\uparrow = r^\downarrow$
(Trenberth et al., 2009)	-7.0	-4.2	-16.0	-10.3
ISCCP-FD	-6.7	-4.0	-15.4	-9.9
ERA-40	-6.1	-3.7	-14.0	-9.0
JRA	-7.0	-4.2	-16.0	-10.3
Bright et al. (2012)	-6.7	-4.0	-15.4	-9.9
CERES EBAF (v2.6)	-7.6	-4.7	-18.7	-12.4
Mean $\pm 1\sigma$	-6.86 ± 0.45	-4.13 ± 0.30	-15.91 ± 1.41	-10.30 ± 1.03
Mean per PFT	-5.50		-13.11	
Global mean (tree & grass)	-9.31			

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.




Fig. 1. Illustration of difference between vegetation fraction (f_{veg}) and gap probability ($P_{gap}(\theta = 0)$) for a model grid cell which is assumed to be totally vegetated.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy
radiation

A. Loew et al.




Fig. 2. Schematic depictions of RAMI4PILPS scenes: closed forest canopies, grasslands, 3-D shrublands and 3-D open forest canopies (left to right).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

DGVM canopy radiation

A. Loew et al.




Fig. 3. Differences of faPAR between canopy RT models and RAMI4PILPS for different experiments (model-RAMI4PILPS); columns correspond to different leaf area or vegetation fraction and soil brightness (see Table 1) while rows correspond to different illumination sun zenith angles (degree); ISO=isotropic illumination (θ); Triangles correspond to different models: left = ORCHIDEE, right = JULES, top = JSBACH, bottom = JSBACH without clumping.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.




Fig. 4. Deviations in GPP for closed forest canopies as derived from differences in canopy radiation profiles between RAMI4PILPS and canopy RT schemes. Columns correspond to different leaf area and soil brightness, while rows correspond to different sun zenith angles for tropical ($\Theta_{\text{noon}} = 15$) and boreal ($\Theta_{\text{noon}} = 50$) conditions. Triangles correspond to different models.

DGVM canopy radiation

A. Loew et al.




Fig. 5. Deviations of simulated surface albedo from RAMI4PILPS reference solutions ($\alpha_{\text{model}} - \alpha_{\text{ref}}$) for the solar spectrum (0.3...4.0 μm). Details of figure are the same as for Fig. 3.

DGVM canopy radiation

A. Loew et al.




Fig. 6. Impact of a systematic albedo difference of $\Delta \alpha = 0.04$ on radiative forcing for tree cover (top) and grassland (bottom), using CERES solar radiation flux ($R_{\text{surf}}^\downarrow$) and different upward atmospheric one-way atmospheric transmissivity: $\Gamma^\uparrow = \text{const}$ (left), $\Gamma^\uparrow = \Gamma^\downarrow$ (right).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 7. Difference in radiative forcing (cooling) due to neglecting the diurnal cycle for surface albedo schemes for global tree and grass cover and different leaf area (columns). Different rows correspond to different parameterizations of the shortwave upward one-way atmospheric transmissivity (Γ^1). Values in parentheses correspond to global mean values.




Fig. A1. Deviations of simulated surface albedo from RAMI4PILPS reference solutions ($\alpha_{\text{model}} - \alpha_{\text{ref}}$) for the VIS spectrum (0.3...0.7 μm). Details of figure structure are the same as Fig. 3.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy radiation

A. Loew et al.




Fig. A2. Deviations of simulated surface albedo from RAMI4PILPS reference solutions ($\alpha_{\text{model}} - \alpha_{\text{ref}}$) for the NIR spectrum (0.7..3.0 μm). Details of figure structure are the same as Fig. 3.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I ◀

▶ I

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DGVM canopy
radiation

A. Loew et al.




Fig. B1. PFT distribution for grassland and forests.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

DGVM canopy radiation

A. Loew et al.




Fig. D1. Canopy transmission as calculated using Eq. (24) by assuming energy conservation. Red labels indicate a clear inconsistency in the canopy flux estimates. Details of figure structure are the same as Fig. 3.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

