

Response to comments

Paper #: *bg-2017-291*

Title: *Gross changes in forest area shape the future carbon balance of tropical forests*

Journal: *Biogeosciences*

Reviewer #1:

General Comments:

Comment #1

This is an interesting study, pointing out the importance of using gross instead of net land use transitions, distinguish between clearing of primary vs. secondary forest and to define a specific and reasonable time horizon when making land-based mitigation policies. Three main steps were taken: 1) the comparison between different response curves, 2) calculating different theoretical scenarios with a bookkeeping model to show the importance of considering gross forest area change and finding critical and 3) applying the ratio to real net land cover transitions from satellite data. Thereby, step 2 clearly takes the highest priority and consideration.

Response #1

We thank the reviewer for the comments and suggestions. Please see the detailed point-by-point responses below.

Comment #2

Still, some revision is needed: the abstract is very long which makes it partly difficult to get the main message of the study. Also, the gap in current research is not carved out very well (page 2, ln 16 ff, says that other models have already implemented gross transitions) and the objectives should become clearer. In the introduction a two-fold purpose of the study is mentioned, what about the 3rd step? What was its objective? The 3rd point cannot be found in the method section, it is just roughly described in the results. Thereby some steps remain unclear: e.g. the model considers LC transition to take place at time = t_0 , but the satellite covers a time series of 12 years. Are all the transitions during these years threatened as if they took place at one time $t=0$ and then the results for the different time horizons of 20, 50 and 100 years are calculated based on that? Or is the exact time of each transition considered and the time horizons starts to be calculated after the last transition took place? Or do the gross transitions in this case refer not to time (i.e. shift of one LC to another LC and back) but instead refer to transitions within the calculated gridcells of 0.5° resolution, as the satellite data was mentioned to have a 30 m resolution?

Response #2

We will shorten the abstract in the revised manuscript (reproduced in **Response #4**).

We will add sentences on **P2L21** to make the research gap and objectives more clear: “[However, uncertainties in the simulated \$E_{LULCC}\$ by grid-based DGVMs arise from the translation of the original LULCC datasets into plant functional type \(PFT\) maps and different processes comprised in different models \(Arneth et al., 2017; Li et al., 2017\). Although DGVMs are spatially and temporally explicit and include detailed physiological processes, the simulations using these models are time consuming and require long spin-up simulations, small time step calculations of biophysical effects and carbon fluxes, including processes less relevant to \$E_{LULCC}\$. Thus, DGVMs are not appropriate to perform, for instance, sensitivity tests for the assessment of LULCC carbon fluxes.](#)”

As suggested in **Comment #3**, we will include the 3rd step in the objectives in Introduction and also mention it in the Methods.

We will add sentences in the Methods to explain how we processed the 30 m forest cover data from Hansen et al. (2013): “[Forest cover data from Hansen et al. \(2013\) comprise three layers at 30 m](#)

resolution: tree cover fraction (0-100% in each pixel) in year 2000, forest area loss (each pixel labeled with a loss year) during 2000-2012, and forest gain during 2000-2012 (not specifying the gain year). Attributing the forest gain to a specific year is challenging because of the difficulty in detecting young forests from satellite reflectance measurements (Hansen et al., 2013). In this study, we used the forest loss and forest gain layers to calculate the ratios of gross-to-net area changes (γ_{Anet}^{Agross}) at a $0.5^\circ \times 0.5^\circ$ resolution, and γ_{Anet}^{Agross} represents the average values during 2000-2012 rather than for a single year since the year of forest gain is not reported. The gross changes at the 0.5° level were calculated by summing the absolute areas of forest loss and gain at the 30 m level during 2000-2012 in each 0.5° grid cell, while the net changes were the sum of gross forest loss (negative) and gross forest gain (positive).”

Comment #3

One possibility to handle the dominance of point 2 would be to make it to the only objective, and shift point 1 to the method section – the comparison seems to be anyhow just a plot of the different curves that justifies the usage of response curves based on Poorter instead of those from Houghton and Hansis. Another possibility would be to include 3 in the method section and give point 1 and 3 more weight - e.g. by calculating the critical gross to net ratios based on the Houghton and Hansis functions, applying it to the same grid cells and comparing it then with the results based on the Poorter function. This would be interesting outcome and extend the first objective of comparing the different response curves to more than just a simple plot of the different curves in the same graph. Further, it would be very interesting to not only know whether each gridcell was a sink or source but also to quantify the ELUC and sum it to total number – one if everything was primary forest at the first transition, and one as if all was secondary, and the same if the other response curves were used.

Response #3

We will include the 3rd step in the method section as suggested.

We calculated the critical ratios based on the exponential response curves from Hansis et al. (2015) (see **Figure R1** below) and compared the number of grid cells above the critical ratios with that using curves from Poorter et al. (2016) (see **Table R1** below). We didn't show the results from Houghton (1999) because the parameters of the exponential functions from Hansis et al. (2015) were already calibrated from the linear function of Houghton (1999).

As show in **Figure 1**, the equilibrium of secondary forest vegetation density with the recovery curve of Hansis et al. (2015) is higher than with Poorter et al. (2016) and we assumed that the same density of primary forest for both, and thus $L_{Hansis,primary}(t) = L_{Poorter,primary}(t)$, $L_{Hansis,secondary}(t) > L_{Poorter,secondary}(t)$ and $G_{Hansis}(\infty) > G_{Poorter}(\infty)$. Note that a positive value of carbon flux indicates carbon emission to the atmosphere. Combined with **Eq (7)** in the manuscript, $\gamma_{Anet}^{Agross} = \frac{L(t)-G(t)}{L(t)+G(t)}$, the different equilibrium states of secondary forest vegetation can explain the differences of critical ratios over time between Hansis et al. (2015) and Poorter et al. (2016) in **Figure R1**. Accordingly, a higher critical ratio leads to smaller number of $0.5^\circ \times 0.5^\circ$ grid cells with γ_{Anet}^{Agross} beyond the critical ratio (**Table R1**).

Figure R1 The critical value of γ_{Anet}^{Agross} at which $\Sigma E_{LULCC, gross}$ is zero, going from a net source to a net sink with different time horizons, using the biomass recovery response curves from Poorter et al. (2016) (solid, same as **Figure 4c**) and from Hansis et al. (2015) (dashed). Values larger than this critical value indicate that the initial forest area change has the net cumulative effect to emit CO₂ at a given time-horizon on the x-axis. Note the different y-axis scale. The lower critical ratio values in the case of primary forest initial loss is because primary forests have a larger biomass, so that a small gross-to-net initial loss in forest area will legate a source at a given horizon than if secondary forests are initially lost.




Table R1 Number of $0.5^\circ \times 0.5^\circ$ grid cells with γ_{Anet}^{Agross} above the critical ratio for which the system is a net cumulative source of CO₂ to the atmosphere, for different time horizons. The calculation was done using the biomass recovery response curves from Hansis et al. (2015) and Poorter et al. (2016) in Latin America. The values of γ_{Anet}^{Agross} were calculated based on high-resolution net and gross forest area change data from Hansen et al. (2013) during 2000-2012. Secondary-to-secondary represents a net forest gain with gross secondary forest loss by assuming that all lost forests were secondary, and primary-to-secondary represents a net forest gain with gross primary forest loss by assuming that all lost forests were primary.

		20 yr		50 yr		100 yr	
		Critical ratio	Grid cell number	Critical ratio	Grid cell number	Critical ratio	Grid cell number
secondary-to-secondary	Poorter et al.	7.2	102	22.5	42	-	-
	Hansis et al.	4.2	143	15.3	57	97.4	9
primary-to-secondary	Poorter et al.	2.4	199	3.1	175	3.7	155
	Hansis et al.	2.5	198	4.1	147	5.0	126

As suggested, we also calculated the total amount of ΣE_{LULCC} of the grid cells beyond the critical ratios with different time horizons in **Figure 5**. The numbers are given in the **Table R2** below and we will incorporate it in the revised manuscript.

Taking C1 secondary-to-secondary at 20 yr horizon for example, using net transitions results in a carbon sink of 12 Tg C but using gross transitions results in a carbon emission of 21 Tg C in the grid cells with $\gamma_{Anet}^{Agross} > 7.2$ (**Figure 5**).

Table R2 Cumulative carbon flux (Tg C) using gross transitions ($\Sigma E_{LULCC, gross}$) and net transitions ($\Sigma E_{LULCC, net}$) in the grid cells with γ_{Anet}^{Agross} beyond the critical ratios at different time horizons. The gross

and net forest area changes are based on the data from Hansen et al. (2013). Positive value of carbon flux indicates carbon emission to the atmosphere. Secondary-to-secondary represents a net forest gain with gross secondary forest loss (C1) by assuming that all lost forests were secondary, and primary-to-secondary represents a net forest gain with gross primary forest loss (C2) by assuming that all lost forests were primary.

Time horizon	C1: secondary-to-secondary			C2: primary-to-secondary		
	Critical ratio	$\Sigma E_{LULCC, gross}$	$\Sigma E_{LULCC, net}$	Critical ratio	$\Sigma E_{LULCC, gross}$	$\Sigma E_{LULCC, net}$
20 yr	7.2	21	-12	2.4	162	-38
50 yr	22.5	3	-2	3.1	125	-39
100 yr	-	-	-	3.7	99	-36

We will add these new analyses suggested by the reviewer in the revised manuscript.

Specific Comments:

Comment #4

Abstract: The abstract should be shortened to better focus on the findings, which would make it easier to read and understand. E.g. is the 3rd sentence really relevant for the findings of this study? Especially also from line 19 to 27 there might be possibilities to shorten, summarize and simplify. Where shapes of the three different curves relevant for the finding? The finding here is difficult to understand, the sentences a bit complicated and several sentences basically say the same: You found and show critical values of gross to net forest area change above which ELUC of a net a net forest area gain switches from CO₂ sink to source.

Response #4

As suggested, we will shorten the Abstract (160 words less) as follows: “

Bookkeeping models are used to estimate land-use and land-cover change (LULCC) carbon fluxes (E_{LULCC}). The uncertainty of bookkeeping models partly arises from data used to define response curves (usually from local data) and their representativeness for application to large regions. Here, we compare biomass recovery curves derived from a recent synthesis of secondary forest plots in Latin America by Poorter et al. (2016) with the curves used previously in bookkeeping models from Houghton (1999) and Hansis et al. (2015). We find that the two latter models overestimate the long-term (100 years) biomass carbon density of secondary forest by about 25%. We also use idealized LULCC scenarios combined with these three different response curves to demonstrate the importance of considering gross forest area changes instead of net forest area changes for estimating regional E_{LULCC} . In the illustrative case of a net gain in forest area composed of a large gross loss and a large gross gain occurring during a single year, the initial gross loss has an important legacy effect on E_{LULCC} so that the system can be a net source of CO₂ to the atmosphere long after the initial forest area change. We show the existence of critical values of the ratio of gross area change over net area change ($\gamma_{Anet}^{A_{gross}}$) above which cumulative E_{LULCC} is a net CO₂ source rather than a sink for a given time horizon after the initial perturbation. These theoretical critical ratio values derived from simulations of a bookkeeping model are compared with real-world observations from the 30 m resolution Landsat TM data of gross and net forest area change in the Amazon. This allows us to diagnose areas where current forest gains with a large land turnover will still legate LULCC carbon emissions in 20, 50 and 100 years.

”

Comment #5

Introduction: Page 2, ln 8: that is for DGVMs the sub-grid transitions that sum up to net changes, here a reference to e.g. Bayer et al. 2017: doi:10.5194/esd-8-91-2017 could be nice, who focused on the problematic of sub-grid transitions.

Response #5

We will add this reference in the revised manuscript.

Comment #6

3.4 Page 7, ln 14: “we pose the question whether such ratios can be observed in the real world” – but this is not what you are answering with your approach. As far as I understood you just calculate using your rates, whether the regions are a sink or a source.

Response #6

This sentence on **P7L14** will be revised as: “...we further combined such ratios with the land use and land cover change datasets to determine whether a region is a carbon sink or source at a given time horizon.”

Comment #7

Page 9, ln 22: “With a too high rotation rate of forests, i.e. a large gross to net area change ratio, a net forest gain could still legate a net carbon source over a long period in the future.” I don’t agree, as I think long rotation secondary forests should have other response curves than short rotation forest, as short rotation forest don’t store as much carbon that can be lost afterwards.

Response #7

This sentence on **P9L22** will be revised as: “With a large gross to net area change ratio, a net forest gain could still legate a net carbon source over a long period in the future.”

Comment #8

4. Discussion: You state that the response curves used in bookkeeping models from Houghton (1999) and Hansis et al. (2015) overestimate carbon density – that implies that Porters values are true, while Houghton and Hansis are wrong. But also Houghton and Hansis are based on measurements, right? Maybe just not in the right region? It would be helpful to mention in the discussion where the measurements for Houghton and Hansis models were located.

Response #8

We showed the differences in biomass recovery curves between Poorter et al. (2016) and Houghton (1999) and Hansis et al. (2015), but we didn’t say that it implies “Porters values are true, while Houghton and Hansis are wrong.”. We only argued that it may bias in this particular region where Poorter et al.’s field survey covers. The reasons for these differences could be the assumptions made for secondary forest by Houghton et al. (1983), the number of field sites and the different locations where field measurements were conducted, as the reviewer said.

We will add some sentences on **P8L9** to clarify it: “The biomass recovery curves of Neotropical secondary forests from Poorter et al. (2016) are lower 20 years after the initial perturbation than those used in the bookkeeping models of Houghton (1999) and Hansis et al. (2015), implying that these models simulate different LULCC carbon fluxes in Latin America from those using the recovery curves of Poorter et al. (2016). The carbon density in undisturbed forests in the bookkeeping models of Houghton (1999) and Hansis et al. (2015) were essentially based on Whittaker and Likens (1973), multiplied by a factor of 0.75 to approximate the lower carbon density of secondary forests (Houghton et al., 1983). The carbon density data from Whittaker and Likens (1973) are subject to two sources of uncertainty. First, these values represent biomass in the 1950s (Woodwell et al., 1978) rather than present days, and second, they were compiled from very limited field measurements for tropical forests. In fact, Whittaker and Likens (1973) claimed in their study that data “for tropical communities are very meager” and the mean biomass density is “a subjectively chosen intermediate value based on very few measurements” to avoid extreme values.”

Reference

- Houghton, R. A., Hobbie, J. E., Melillo, J. M., Moore, B., Peterson, B. J., Shaver, G. R. and Woodwell, G. M.: Changes in the Carbon Content of Terrestrial Biota and Soils between 1860 and 1980: A Net Release of CO₂ to the Atmosphere, *Ecol. Monogr.*, 53(3), 235–262, doi:10.2307/1942531, 1983.
- Whittaker, R. H. and Likens, G. E.: Carbon in the biota., in *Brookhaven symposia in biology*, pp. 281–302., 1973.
- Woodwell, G. M., Whittaker, R. H., Reiners, W. a, Likens, G. E., Delwiche, C. C. and Botkin, D. B.: The biota and the world carbon budget, *Science*, 199(4325), 141–146, doi:10.1126/science.199.4325.141, 1978.
-

Comment #9

Is the extent to which gross versus net transitions affect ELUC comparable what other studies investigating gross versus net transitions studied? You mention several studies about this issue which were performed with carbon models – they should also appear in the discussion, showing how your results compare with what they found.

Response #9

We will add some discussion about the impacts of gross and net transitions on E_{LULCC} on **P8L13**: “Some DGVMs (Shevliakova et al., 2009; Stocker et al., 2014; Wilkenskield et al., 2014; Yue et al., 2017; Bayer et al., 2017) as well as a bookkeeping model (Hansis et al., 2015) have implemented gross land use and land cover transitions, and thus simulated a higher E_{LULCC} than using net transitions. Arneth et al. (2017) reviewed the “missing processes” in LULCC modeling by DGVMs and found that ignoring gross LULCC could underestimate the global cumulative E_{LULCC} by 36 Pg C on average over the historical period (1901-2014).”. However, the E_{LULCC} from land carbon models cannot be directly compared with E_{LULCC} from bookkeeping models, because of the different processes in models and the different definitions of E_{LULCC} . In addition, our study focused on the ratios of gross-to-net changes rather than the estimates of E_{LULCC} , and thus it is difficult and not necessary to compare with E_{LULCC} from land carbon models.

Comment #10

Technical corrections

General: In all figures and the text, it might be useful to replace the term “biomass carbon density” by “vegetation carbon density”, as biomass is less well defined and includes in some disciplines also dead biomass and soil microbial biomass which would count here to the soil pool.

Response #10

We will revise it accordingly.

Comment #11

Page 1, ln 26: “critical value” should be plural

Response #11

We will revise it accordingly.

Comment #12

Page 2, ln 10: “Gross LUC occurs in tropical regions with shifting cultivation (Hurtt et al., 2011) but also everywhere forests are cut and new plantations created at the same time” Is here a “where” after “everywhere” missing?

Response #12

This sentence on **P2L10** will be revised as: “Gross LULCC occurs in tropical regions with shifting cultivation (Hurtt et al., 2011) and also in other regions where forests are cut and new plantations created at the same time.”

Comment #13

Table 1: The table caption uses gamma gross to net but in the table heading and the text gamma Agross to Anet is used.

Response #13

We will revise it accordingly.

Comment #14

Fig 1: biomass from primary forest: reference missing; Legend for a) and b) shows biomass, which can only be found in plot c) that has an own legend.

Response #14

We will add the reference in the legend and remove biomass from the legend in (a) and (b).

Fig 3: include dashed and solid line in legend. In figure description logarithmic asymptotic should be removed or referred to both – solid and dashed, so it becomes more clear that there is no difference in the response curve between solid and dashed, but just which systems are transformed.

Response #15

We will add it in the legend and revise the caption.

Comment #15

Fig 4 is a bit difficult to understand. The difference between plot a) and b) is hidden in the middle of the figure description in the end of a sentence. Would be better to have it in the description directly following a) resp. b), whereas “net forest gain at $t=0$ ” which is true for both plots should either be in the end or before the separation in a) and b). Or/ additionally it could be mentioned as title in each plot whether it is primary to secondary or secondary to secondary. The axis title is only in plot a) but not in b) whereas the legend can be found in both plots. Please add the axis title to b) or remove the legend from a) or do both and set the legend a bit aside, which would also help the reader to not confuse it with a second y-axis title at a first glance.


Please extend “Exponential carbon loss curve from (Hansis et al., 2015) and logarithmic gain curve from (Poorter et al., 2016) are used in this example” to something like “Exponential curve from Hansis et al., (2015) for carbon loss in all pools and gain in soil pool and logarithmic curve from Poorter et al., (2016) for gain in biomass pool are used in this example, which corresponds to the combinations C1 and C2 from Table 2 for a) and b) respectively.”

Response #16


As suggested, we will revise the caption and re-plot the figure (reproduced below).

Figure 4 Time evolution of cumulative carbon flux ($\Sigma E_{LULCC, gross}$) after an initial forest area change involving gross forest area changes followed by no forest area change. The three panels show results of our bookkeeping model for three case studies (a) a net forest gain at $t = 0$ with initial secondary forest loss followed by secondary forest regrowth (secondary-to-secondary, C1 in Table 2), (b) the same net area gain at $t = 0$ with initial primary forest loss followed by secondary forest regrowth (primary-to-secondary C2 in Table 2), and (c) the critical value of γ_{Anet}^{Agross} at which $\Sigma E_{LULCC, gross}$ is zero, going from a net source to a net sink for different time horizon in the x-axis. The colored curves in (a) and (b) have the same net area change ($A_{net} = +1$ ha) at $t = 0$ but variable values of the initial gross-to-net area change ratios (γ_{Anet}^{Agross}). The red line in (a) and (b) is the zero line, defining the time after initial forest area change at which the system reaches a neutral carbon balance. The light and dark green lines in (c) represent the critical ratios for a net initial forest gain scenario with secondary-to-secondary (a) and primary-to-secondary (b) gross forest area change, respectively. Values larger than this critical value indicate that the initial forest area change has the net effect to emit CO_2 for a given time horizon in the x-axis. Exponential curve from Hansis et al. (2015) for carbon loss in all pools and gain in soil pool and logarithmic curve from Poorter et al. (2016) for gain in biomass pool are used in this example.


(a) secondary-to-secondary



(b) primary-to-secondary



(c)



Reviewer #2:

General Comments:

Comment #1

I read this manuscript with much interest, and found it to have novel elements which provide new and useful information. However, it could benefit from some revisions.

Response #1

We thank the reviewer for the comments and suggestions. Please see the detailed point-by-point responses below.

Comment #2

The first part of the paper is about land use changes, from forest to agriculture. However changes from primary and secondary forest to plantations are also discussed (and the abbreviation LUC is used). Harvesting in rotation is not generally considered land use change (but a land cover change), so this abbreviation might be better changed to Land cover change (LCC) which would encompass both the forest gain scenario and land use change (loss of forest to agriculture). The paper uses land cover data (Hansen), which further confuses the reader, when land use is mainly used. The authors could check the consistency of these terms (land use and land cover) in the paper.

Response #2

We agree that the satellite data from Hansen et al. (2013) we used in the case study is land cover change rather than land use change, and the idealized scenarios are more land use change although also a land cover change, as described on **P3L12** “The land-use changes considered in this study are forest loss (tropical moist forest transformed to cropland) and forest gain (cropland abandonment to secondary tropical moist forest) in Latin America.”. We will change the term into “land-use and land-cover change (LULCC)” throughout in the text to be consistent.

Comment #3

One of the concerns in the paper is the methods, which could be expanded to clarify some points. The analysis of the Hansen data, for example is not included. For example, what forest cover threshold did you use in the analysis? Did you for example mask out those pixels with loss or gain but with <10%, or another appropriate canopy cover threshold for the region? Or is it exactly following the Poorters map? How was the change of grid cell to 0.50 done? For example, pixels only partially within the area of interest are included or not? I wonder if the choice of grid cell size would impact the results? Was 0.50 chosen for a specific reason?

Response #3

We will add some sentences to clarify the forest cover change data from Hansen et al. (2013) in the revised manuscript: “Forest cover data from Hansen et al. (2013) comprise three layers at 30 m resolution: tree cover fraction (0-100% in each pixel) in year 2000, forest area loss (each pixel labeled with a loss year) during 2000-2012, and forest gain during 2000-2012 (not specifying the gain year). Attributing the forest gain to a specific year is challenging because of the difficulty in detecting young forests from satellite reflectance measurements (Hansen et al., 2013). In this study, we used the forest loss and forest gain layers to calculate the ratios of gross-to-net area changes (γ_{Anet}^{Agross}) at a $0.5^\circ \times 0.5^\circ$ resolution, and γ_{Anet}^{Agross} represents the average values during 2000-2012 rather than for a single year since the year of forest gain is not reported. The gross changes at the 0.5° level were calculated by summing the absolute areas of forest loss and gain at the 30 m level during 2000-2012 in each 0.5° grid cell, while the net changes were the sum of gross forest loss (negative) and gross forest gain (positive).” Thus, we didn’t use the tree cover fraction threshold because we didn’t use the tree fraction data.

It is not necessary to be exactly the same region of the Poorter et al.’s map because the biomass recovery estimates from Poorter et al. (2016) are based on forest sites and forest plots and thus represent a rough (not precise) Latin America region. Thus there is no such issue of partially

overlapped pixels. We gave the latitudes and longitudes of the region we used from the map of Hansen et al. (2013) in **Figure 5**.

The gross changes compared to net changes essentially is a matter of resolution. For example, if the source data is at 30 m spatial resolution and all the models are run at 30 m resolution, there would be no difference between gross and net changes. The differences between gross and net changes only emerge when aggregating high-resolution data into a coarser resolution. The reason for choosing the 0.5° resolution was described on **P7L18**: “[The spatial resolution of 0.5° is a typical resolution of DGVMs when they simulate global E_{LULCC}](#).” Because the 30 m spatial resolution from Hansen et al.’s data is relatively high, using other grid cell size like 0.1° or 1° would be expected to give similar patterns as using 0.5° in **Figure 5**.

Comment #4

Figure 5 is also not clear to me, for example (if I understand correctly), those pixels in blue reached the threshold for the secondary forest clearing (and also the primary forest clearing) and those in green reached the threshold for the primary forest clearing only? This would be useful information to include in the caption.

Response #4

Yes, that is correct. We will add it in the caption as suggested: “[The blue grid cells represent a cumulative carbon emission in 20 years no matter whether the lost forest is primary or secondary. The green ones represent a cumulative carbon emission only if the cleared forests are primary forests.](#)”

Comment #5

The results for the soil carbon change are also interesting and useful to include, but I find the discussion about this lacking. Indeed, there is a huge amount of uncertainty related to changes in soil carbon (see for example Don et al. 2011 Impact of tropical land-use change on soil organic carbon stocks – a meta-analysis). Incorporating some aspect of uncertainties related to this could have been helpful, and indeed, uncertainties are missing in all findings of the paper.

Response #5

We will revise the sentences on **P8L9**: “[Differences may also exist for soil carbon dynamics after LULCC. There are a great number of meta-analyses or reviews \(e.g. Davidson & Ackerman, 1993; Post & Kwon, 2000; Conant et al., 2001; Paul et al., 2002; Davis & Condon, 2002; Guo & Gifford, 2002; Murty et al., 2002; West et al., 2004; Laganière et al., 2010; Poeplau et al., 2011; Powers et al., 2011; Don et al., 2011; Li et al., 2012; Marín-Spiotta & Sharma, 2013; Wei et al., 2014; Kurganova et al., 2014\) on the soil carbon change after LULCC based on field measurement data \(mostly paired sites and chronosequences\). These studies may generally agree with the directions of soil carbon change after LULCC \(e.g. soil carbon loss after forest clearing for cropland\), but the magnitudes and temporal dynamics of soil carbon changes remain highly uncertain because, among other things, of the limited site number and the diversity of soil properties. Field measurements at site level may be unrepresentative of the whole region because the distribution of biophysical conditions like soil texture, precipitation and temperature may not match the distribution of the whole set of such factors in the LULCC areas in a given region \(Powers et al., 2011\).](#)”

Specific Comments:

Comment #6

Page 7, line 14-16. There are a number of datasets which you could use, and the data also do not limit the work to small scale analysis, so this sentence seems not to be useful.

Response #6

This sentence will be deleted.

Comment #7

Page 3 line 29/30. I would include here or somewhere appropriate, some numbers related to the total biomass used in the paper from Poorter.

Response #7

The number related to the ratio of aboveground to total biomass is only mentioned in the supporting information in Poorter et al. (2016). The ratio Poorter et al. (2016) used is from FAO FRA, which is 0.82, basically the same as we used (0.81). We will revise the sentence about on **P3L30**: “For both response curves, a ratio of 0.81 (Liu et al., 2015; Peacock et al., 2007; Saatchi et al., 2011) was used to convert aboveground biomass reported by Poorter et al. (2016) to total biomass, and this ratio is consistent with the one (0.82) that Poorter et al. (2016) used based on FAO FRA (2010).”

Comment #8

Page 9 line 9 – the “new planted forest in rotation practice”- it is not clear what you mean, and do you have a reference for this?

Response #8

We will revise this sentence as: “Forest management practices like wood harvest and thinning extract carbon from the ecosystem and release it to the atmosphere (Houghton et al., 2012), while recovering secondary forest from past deforestation and logging (Pan et al., 2011) and even old-growth forests (Luyssaert et al., 2008) can act as carbon sinks.”

Reference

Luyssaert, S., Schulze, E.-D., Börner, A., Knohl, A., Hessenmöller, D., Law, B. E., Ciais, P. and Grace, J.: Old-growth forests as global carbon sinks, *Nature*, 455(7210), 213–215, doi:10.1038/nature07276, 2008.
Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L., Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W., McGuire, A. D., Piao, S., Rautiainen, A., Sitch, S. and Hayes, D.: A large and persistent carbon sink in the world’s forests., *Science*, 333(6045), 988–993, doi:10.1126/science.1201609, 2011.

Comment #9

You refer to “idealized scenarios”. I am not sure about the choice of term here. Ideal for what?

Response #9

We think this is a matter of English here. “idealized” refers to “conceptual” while “ideal” is more like “optimal”. Because we want to demonstrate the difference between gross and net changes on E_{LULCC} and determine the critical gross-to-net change ratio, we used these idealized scenarios that are simple and representative, and may not the case in reality.

Page 1 line 28. “compared against” could be changed to “compared to”. Landsat is more commonly referred to as medium resolution (rather than high resolution), although the global maps are termed high resolution global maps. I would remove the term or would specify the resolution in m.

Response #10

We will revise it accordingly.

Comment #10

Page 2 line 7. Why “so-called”?

Response #11

We will delete it accordingly.

Comment #11

Page 8 line 5. could be rephrased: ”are lower 20 years after the initial LULCC” or in another way.

Response #12

We will revise it accordingly.

Comment #12

Page 7 Line 24. Is it necessary to describe a map as ‘spatial’?

Response #13

We will delete it accordingly.

Comment #13

Page 7 line 14. Instead of “real world”, “in a case study” or similar?

Response #14

This sentence on **P7L14** will be revised as: “...we further combined such ratios with the land use and land cover change datasets to determine whether a region is a carbon sink or source at a given time horizon.”

Gross changes in forest area shape the future carbon balance of tropical forests

Wei Li¹, Philippe Ciais¹, Chao Yue¹, Thomas Gasser², Shushi Peng³, Ana Bastos¹

¹Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

²International Institute for Applied Systems Analysis (IIASA), A-2361 Laxenburg, Austria

³Sino-French Institute for Earth System Science, College of Urban and Environmental Sciences, Peking University, Beijing 100871, China

Correspondence to: Wei Li (wei.li@lsce.ipsl.fr)

10 **Abstract.** Bookkeeping models are used to estimate land-use and land-cover change (LULCC) carbon fluxes (E_{LULCC}). The
uncertainty of bookkeeping models partly arises from data used to define response curves (usually from local data) and their
representativeness for application to large regions. Here, we compare biomass recovery curves derived from a recent synthesis
of secondary forest plots in Latin America by Poorter et al. (2016) with the curves used previously in bookkeeping models
from Houghton (1999) and Hansis et al. (2015). We find that the two latter models overestimate the long-term (100 years)
15 vegetation carbon density of secondary forest by about 25%. We also use idealized LULCC scenarios combined with these
three different response curves to demonstrate the importance of considering gross forest area changes instead of net forest
area changes for estimating regional E_{LULCC} . In the illustrative case of a net gain in forest area composed of a large gross loss
and a large gross gain occurring during a single year, the initial gross loss has an important legacy effect on E_{LULCC} so that the
system can be a net source of CO_2 to the atmosphere long after the initial forest area change. We show the existence of critical
20 values of the ratio of gross area change over net area change ($\gamma_{\text{Gross Anet}}^{\text{Amazon}}$), above which cumulative E_{LULCC} is a net CO_2 source rather
than a sink for a given time horizon after the initial perturbation. These theoretical critical ratio values derived from simulations
of a bookkeeping model are compared with real-world observations from the 30 m resolution Landsat TM data of gross and
net forest area change in the Amazon. This allows us to diagnose areas where current forest gains with a large land turnover
will still legate LULCC carbon emissions in 20, 50 and 100 years.

25 ▼

Formatted: Subscript

Deleted: Bookkeeping models are used to estimate land-use change (LUC) carbon fluxes (E_{LUC}). These models combine time series of areas subject to different LUC types with response curves of carbon pools in ecosystems and harvested products after a unit change of land use. The level of detail of bookkeeping models depends on the number of response curves used for different regions, the carbon pools they represent, and the diversity of LUC types considered. The uncertainty of bookkeeping models arises from data used to define response curves (usually local data) and their representativeness of large regions. Here, we compare biomass recovery curves derived from a recent synthesis of secondary forest plots data by Poorter et al. (2016) with the curves used in bookkeeping models from Houghton (1999) and Hansis et al. (2015) in Latin America. We find that both Houghton (1999) and Hansis et al. (2015) overestimate the long-term (100 years) biomass carbon density of secondary forest, by about 25%. We also show the importance of considering gross forest area change in addition to the net forest area change for estimating regional E_{LUC} . To do so, simulations are constructed with a bookkeeping model calibrated with three different sets of response curves (linear, exponential and logarithmic) to study E_{LUC} created by a pulse of net forest area change, with different gross-to-net forest area change ratios ($\gamma_{\text{Gross Anet}}$). Following the initial pulse of forest area change, E_{LUC} is subsequently calculated over 100 years. Considering a region subject to a net gain in forest area during one year, different values of gross forest area changes that sum up to this initial net gain can change the magnitude and even the sign of E_{LUC} with a given time horizon after the initial forest area change. In other words, in the case of a net gain in forest area composed of a large gross loss and a large gross gain, the initial gross loss has an important legacy effect that the system can be a net source of CO_2 to the atmosphere. We show the existence of a critical value of $\gamma_{\text{Gross Anet}}$ above which E_{LUC} switches from CO_2 sink to source with a given time horizon after the initial forest area change. This critical ratio derived from the structure of the bookkeeping model is compared against real-world high resolution Landsat TM observations of gross forest area change in the Amazon to distinguish areas where current forest land turnover will legate LUC carbon emissions or sinks in 20 years, 50 years and 100 years in the future. -

1 Introduction

The global carbon flux from land-use and land-cover change (E_{LULCC}) represents a net source of carbon to the atmosphere of $0.9 \pm 0.5 \text{ Gt C yr}^{-1}$ during the last decade (Ciais et al., 2013; Le Quéré et al., 2015). E_{LULCC} is usually estimated by bookkeeping models (Hansis et al., 2015; Houghton, 2003), dynamic global vegetation models (DGVMs) (Le Quéré et al., 2015; Sitch et al., 2015) or compact earth system models (Gasser et al., 2017). Most DGVMs (e.g. in the TRENDY project, Sitch et al., 2015) estimate emissions due only to net area changes between different land-use / land-cover types in a grid cell. At the moment, efforts are being made to incorporate gross land-use and land-cover change (LULCC) in these models, that is for DGVMs the sub-grid transitions that sum up to net changes (Bayer et al., 2017). The bookkeeping model of Houghton (1999) includes emissions from both net area changes and gross LULCC from shifting cultivation, previously at the scale of large regions (Houghton, 2003), and more recently for each country (Houghton and Nassikas, 2017). Gross LULCC occurs in tropical regions with shifting cultivation (Hurtt et al., 2011) and also in other regions where forests are cut and new plantations created at the same time. For example, consider a region with co-existing forest and crops where 20% of the land is converted from primary forest to crops while 20% sees crop abandonment to forest in the same period. The net change corresponds to a stable forest area, but the large carbon loss from primary forest is not compensated by the small carbon gain of the new plantations. In this example, the region will be a net source of CO_2 during several years. Because of the non-symmetrical dynamics of CO_2 fluxes between forest loss and gain, E_{LULCC} differs between net and gross area changes. Arneeth et al. (2017) recently reviewed this issue using DGVMs and concluded that considering gross LULCC significantly increased the simulated E_{LULCC} at global scale. Gross land-use area transition datasets including e.g. shifting cultivation practice (Hurtt et al., 2011) and reconstructions using empirical ratios between gross and net transitions (Fuchs et al., 2015) are now available and have been implemented in a bookkeeping model (Hansis et al., 2015) as well as in some DGVMs to improve the estimate of E_{LULCC} (Fuchs et al., 2016; Shevliakova et al., 2009; Stocker et al., 2014; Wilkenskeld et al., 2014; Yue et al., 2017). However, uncertainties in the simulated E_{LULCC} by grid-based DGVMs arise from the translation of the original LULCC datasets into plant functional type (PFT) maps and different processes comprised in different models (Arneeth et al., 2017; Li et al., 2017). Although DGVMs are spatially and temporally explicit and include detailed physiological processes, the simulations using these models are time consuming and require long spin-up simulations, small time step calculations of biophysical effects and carbon fluxes, including processes less relevant to E_{LULCC} . Thus, DGVMs are not appropriate to perform, for instance, sensitivity tests for the assessment of LULCC carbon fluxes. Bookkeeping models use response curves for biomass and soil carbon stocks consecutive to LULCC disturbance and time-series of LULCC areas to estimate E_{LULCC} (Hansis et al., 2015; Houghton, 1999). Response curves can be linear (Houghton, 1999, 2003), exponential (Hansis et al., 2015) or of other types. The carbon densities of different land-use types are derived from field measurements (Houghton et al., 1983). Even though carbon densities have a high spatial variability in the real world, the same response curve measured at one location is often applied in bookkeeping models over large regions. A recent study of the biomass resilience of secondary forests in the Neotropics provides new biomass recovery curves from 45 secondary

Deleted: LUC

Deleted: so-called

Deleted: LUC

Deleted: Gross LUC occurs in tropical regions with shifting cultivation (Hurtt et al., 2011) but also everywhere forests are cut and new plantations created at the same time.

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: land carbon models

Deleted: LUC

Formatted: Subscript

Formatted: Subscript

Deleted: LUC

Deleted: LUC

Deleted: LUC

forest sites (Poorter et al., 2016). These new data are valuable to revisit the response curves for the regrowth of secondary forest in the Amazon area, an important region with a large E_{vULCC} .

In this study, we first aim to compare the recent biomass regrowth curves from Poorter et al. (2016) with the ones used in two bookkeeping models (Hansis et al., 2015; Houghton, 1999) for their implications in E_{vULCC} . Second, we demonstrate that because of the asymmetry between carbon loss from deforestation and carbon gains from regrowth, even when the net forest area change is positive, a large initial gross forest area change can still cause E_{vULCC} to be a source of CO_2 to the atmosphere on multi-decadal horizons. Last, we apply our conceptual calculation to the satellite forest data to diagnose areas with net forest gains but cumulative LULCC carbon emissions.

Based on E_{vULCC} calculated using a bookkeeping approach and several idealized scenarios constructed to have different gross forest area changes but with the same net area change (Section 3.2), we show the existence of a critical ratio of gross-to-net forest area change above which cumulative E_{vULCC} remains a net source after initial JULCC, because carbon losses from deforestation are not compensated by carbon gains from secondary forest growth (Section 3.3). The theoretical value of this ratio derived from the idealized scenarios is then compared with actual estimates of gross-to-net forest area change over the Amazon derived from high-resolution (30 m) Landsat satellite imagery over the period of 2000-2012 (Hansen et al., 2013). This allows us to identify sensitive regions where the current turnover of forest is too large, and may result in an emission source of CO_2 to the atmosphere over different time horizons in the future.

2 Methods

The land-use and land-cover changes considered in this study are forest loss (tropical moist forest transformed to cropland) and forest gain (cropland abandonment to secondary tropical moist forest) in Latin America. We construct a bookkeeping model to simulate the carbon balance of simultaneous forest loss and gain in the same region. This model is similar to those developed by Houghton (1999) and Hansis et al. (2015) for global applications. After forest area loss, carbon density changes are calculated for biomass, two soil organic carbon pools (rapid and slow) and two products pools with turnover times of 1 and 10 years respectively. After the establishment of a secondary forest, carbon density changes in biomass and soil pools are considered. Only one slow soil pool is used in the regrowth of secondary forest, similar to Houghton (1999) and Hansis et al. (2015).

Both the linear response curves from Houghton (1999) and the exponential ones from Hansis et al. (2015) are used to simulate the dynamics of each carbon pool consecutive to initial JULCC (Figure 1). For re-growing secondary forest, we also used two curves for biomass recovery based on a collection of field measurements by Poorter et al. (2016). The first one is a logarithmic equation describing aboveground biomass carbon as a function of stand age from Poorter et al. (2016), the parameters of which are derived using the average aboveground biomass recovery from multiple stands after 20 years. It should be noted that with a logarithmic curve, no asymptotic value is reached even after an infinite time, which is not realistic for estimating long-term budgets, as it would mean permanent carbon gains. To overcome this problem of the logarithmic curve, we define a fixed time

Deleted: LUC

Deleted: The purpose of this study is thus two-fold.

Deleted: First,

Deleted: LUC

Deleted: will

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: ,

Deleted: ed

Deleted: LUC

horizon of 100 years after ~~JULCC~~ at which biomass becomes constant. The second biomass carbon gain curve is an exponential curve obtained by fitting the data from Poorter et al. (2016) with a saturating exponential function like in Hansis et al. (2015). This equation avoids the infinite increase of biomass after ~~JULCC~~ in the logarithmic curve. For both response curves, a ratio of 0.81 (Liu et al., 2015; Peacock et al., 2007; Saatchi et al., 2011) is used to convert aboveground biomass reported by Poorter et al. (2016) to total biomass, and this ratio is consistent with the one (0.82) that Poorter et al. (2016) used based on FAO FRA report (FAO, 2010).

Deleted: LUC

Deleted: LUC

Deleted: was

To model the sensitivity of the carbon balance of a typical region in Latin America to different ratios of gross-to-net forest area change during initial pulse of forest area change followed by no-change in forest area, we construct five idealized scenarios (Table 1). These scenarios are: S0 with no net but gross area changes; S1 with a net forest area loss being the sum of small gross area changes; S2 with the same net forest area loss as S1, but being a sum of large gross area changes; and S3 and S4, similar to S1 and S2 but with a net forest area gain, instead of a net loss. An example of small versus large gross area changes with the same net area change is illustrated in Figure 2.

Deleted: s

Deleted: .

In each scenario, ~~JULCC~~ is applied as a pulse of forest area change at time $t = 0$, and we evaluate carbon changes over the following 100 years. The parameter $\gamma_{A_{net}}^{A_{gross}}$ is the ratio of gross forest change area (A_{gross}) to net forest change area (A_{net}) applied at $t = 0$.

Deleted: LUC

$$\gamma_{A_{net}}^{A_{gross}} = \frac{A_{gross}}{A_{net}} \quad (1)$$

where:

$$A_{gross} = |A_{loss}| + A_{gain} \quad (2)$$

$$A_{net} = A_{loss} + A_{gain} \quad (3)$$

By convention, $A_{loss} (<0)$ and $A_{gain} (>0)$ are the gross forest loss and gain areas applied at $t = 0$. A positive value of A_{net} is an increase in forest area. For instance, the illustrative scenario S3 described in Table 1 explores the effects of a large positive value of $\gamma_{A_{net}}^{A_{gross}}$ on ~~F_{JULCC}~~. ~~F_{JULCC}~~ is then simulated for contrasting A_{gross} and A_{net} transitions with the bookkeeping model as the sum of changes in all carbon pools over the area that was disturbed at $t = 0$. ~~$\Sigma F_{JULCC,net}$~~ is the cumulative ~~JULCC~~ carbon flux up to a time horizon t , calculated using only net area changes (A_{net}) and ignoring gross area changes. ~~$\Sigma F_{JULCC,gross}$~~ is the cumulative carbon flux using gross forest area change, which has two component fluxes: the cumulative emissions (~~$\Sigma F_{JULCC,loss}$~~) from gross forest loss and the carbon sink (~~$\Sigma F_{JULCC,gain}$~~) from secondary forest regrowth. This is given by:

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

$$\Sigma E_{LUC,gross} = \Sigma E_{LUC,loss} + \Sigma E_{LUC,gain} \quad (4)$$

$$\Sigma E_{LUC,loss} = -A_{loss} \times L(t) \quad (5)$$

$$\Sigma E_{LUC,gain} = A_{gain} \times G(t) \quad (6)$$

where $L(t)$ and $G(t)$ stand for the cumulative carbon density change in all carbon pools up to time t . Positive values of carbon fluxes indicate a loss of land carbon to the atmosphere.

For each scenario in Table 1, we test different loss and gain response curves in our bookkeeping model, namely, linear or exponential carbon loss and linear, logarithmic or exponential increase for forest gain. In the case of gross forest area loss, we

considered two options, either a primary forest (primary-to-secondary) or a secondary forest (secondary-to-secondary) being cleared (Table 2, also see an illustration in Figure 2). This gives a total of eight combinations (C1 to C8 in Table 2) to calculate legacy F_{LULCC} after a forest area disturbance. Note that one basic principle of bookkeeping models is that the same equilibrium vegetation carbon density is assumed between a secondary forest being lost and a secondary forest having fully recovered.

Deleted: LUC

Deleted: biomass

5 Therefore, the equilibrium biomass density of secondary forest being lost at $t=0$ in C1, C3 and C5 is set to be the same as that of the fully recovered (100 years) secondary forest in Poorter et al. (2016).

We use Global Forest Change data from Hansen et al. (2013) to apply our conceptual calculation to the real-world gross and net forest changes. Forest cover data from Hansen et al. (2013) comprise three layers at 30 m resolution: tree cover fraction (0-100% in each pixel) in year 2000, forest area loss (each pixel labeled with a loss year) during 2000-2012, and forest gain during 2000-2012 (not specifying the gain year). As noted in Hansen et al. (2013), attributing the forest gain to a specific year is challenging because of the difficulty in detecting young forests from satellite reflectance measurements. In this study, we use the forest loss and forest gain layers to calculate the ratios of gross-to-net area changes ($\gamma_{\text{Area}}^{\text{Gross}}$) at a $0.5^\circ \times 0.5^\circ$ resolution, and thus $\gamma_{\text{Area}}^{\text{Gross}}$ represents the average values during 2000-2012 rather than for a single year since the year of forest gain is not reported. The gross changes at the 0.5° level are calculated by summing the absolute areas of forest loss and gain at the 30 m level during 2000-2012 in each $0.5^\circ \times 0.5^\circ$ grid cell, while the net changes are the sum of gross forest loss (negative) and gross forest gain (positive).

Formatted: Font color: Auto

Formatted: Not Highlight

Formatted: Not Highlight

3 Results

3.1 Response curves and comparison with field measurements

The response curves of tropical moist forest from bookkeeping models of Houghton (1999) and Hansis et al. (2015) and from Poorter et al. (2016) for Latin America used in this study (Section 2) are displayed in Figure 1. The curves of Houghton (1999) (linear) and Hansis et al. (2015) (exponential) are similar (Figure 1) because the parameters of the exponential function were calibrated from the linear one (Hansis et al., 2015). Due to the higher carbon density of primary compared to secondary forest and the identical time at which both loss curves reach zero in Houghton (1999) and Hansis et al. (2015), the loss curves for a cleared primary forest are steeper than those for a cleared secondary forest (Figure 1a, b). This implies that clearing a primary forest instead of a secondary one leads to larger legacy emissions. The fast decay of the rapid soil carbon pool in Figure 1a and 1b is due to the fact that a fraction of the initial biomass is assigned to this pool after forest clearing (Hansis et al., 2015; Houghton, 1999).

Deleted: gain

The logarithmic recovery curve (lime dashed lines in Figure 1c) from Poorter et al. (2016) has an initial faster biomass growth rate up to 20 years than in the curves used in previous bookkeeping models. After 20 or 30 years, however, the recovery curves of Houghton (1999) and Hansis et al. (2015) surpass the one of Poorter et al. (2016), leading to a higher equilibrium biomass of mature secondary forests (Figure 1c). More precisely, the 100-year biomass of a secondary forest in Houghton (1999) and Hansis et al. (2015) is $\approx 25\%$ higher than in Poorter et al. (2016). The median time to recover 90% of the maximum biomass

is 66 years in Poorter et al. (2016), compared to only 44 years in Houghton (1999) and 55 years in Hansis et al. (2015) (Figure 1c). The exponential recovery curve fit to the data from Poorter et al. (2016) (lime dash-dotted line in Figure 1c) has lower biomass than the logarithmic curve in the first 40 years but reaches a similar density after 100 years (by construction). The exponential curve from Poorter et al. (2016) agrees well with the linear curve of Houghton (1999) during the first 20 years (Figure 1c).

3.2 Temporal change of cumulative carbon fluxes in different LULCC scenarios

We calculated cumulative carbon fluxes for the five idealized forest area change scenarios (Table 1) with the eight combinations of response curves (Table 2), giving an ensemble of 40 simulations. Results for each simulation are shown in Figure S1. We compare here the response curve combination C1 (exponential secondary forest loss and logarithmic biomass recovery) and C2 (exponential primary forest loss and logarithmic biomass recovery) as examples in Figure 3 (see annual fluxes in Figure S2) to illustrate the effect of different gross forest area change with the same net area change on cumulative carbon flux, i.e., the impact of $\gamma_{\text{Anet}}^{\text{Agross}}$ on the F_{LULCC} . Other combinations provide similar conclusions as C1 and C2. For example, F_{LULCC} for C5 and C6 using linear curves for forest loss are very similar to C1 and C2 in Figure S1.

In the scenario S0 with initial secondary forests and no net forest area change, $\Sigma F_{\text{LULCC,net}}$ is zero when calculated based on net area change (Figure 3a) but the gross carbon flux ($\Sigma F_{\text{LULCC,gross}}$) is distinct from zero. In the variant of the S0 scenario with initial primary forest (C2), due to the lower equilibrium carbon density of the secondary forest, $\Sigma F_{\text{LULCC,gross}}$ is a large source after 100 years (red dashed lines in Figure 3a). In the secondary forest loss and gain case (C1), $\Sigma F_{\text{LULCC,gross}}$ is a carbon source in the initial period and gradually becomes carbon neutral with the compensation effects of secondary forest regrowth (red solid lines in Figure 3a).

Both S1 and S2 scenarios have the same net forest area loss ($A_{\text{net}} = -1$ ha) but different gross forest area changes ($\gamma_i^{1,2} = -1.2$ and $\gamma_i^{201} = -201$ for S1 and S2 respectively, Table 1). In S1 with a small gross area change ($A_{\text{gross}} = 1.2$ ha), $\Sigma F_{\text{LULCC,gross}}$ is close to $\Sigma F_{\text{LULCC,net}}$ (Figure 3b), starting with either primary and secondary initial forests. By contrast, the difference between $\Sigma F_{\text{LULCC,gross}}$ and $\Sigma F_{\text{LULCC,net}}$ in S2 is large and positive, indicating a cumulative carbon loss much higher than S1 due to its large gross area change (Figure 3c).

The scenarios S3 versus S4 with a net forest gain ($A_{\text{net}} = +1$ ha) but different ratios of gross-to-net area changes ($\gamma_{\text{Anet}}^{\text{Agross}}$) present a similar behavior as S1 versus S2. However, the sign of $\Sigma F_{\text{LULCC,gross}}$ is reversed, from a sink in S3 (red lines in Figure 3d) to a source in S4 (red lines in Figure 3e). Especially for the gross primary forest loss, $\Sigma F_{\text{LULCC,gross}}$ exhibits a large source even after 100 years (red dashed lines in Figure 3d,e). This implies that despite the net initial forest gain, the rate of gross area change determines the sign of ΣF_{LULCC} over a certain time horizon after the pulse of forest area change. More generally, this shows that, while long term cumulative land use change emissions are determined only by the net land use area change (e.g. Gasser and Ciais, 2013), short term cumulative emissions are determined by the gross area change.

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: s.

Deleted: s.

Deleted: LUC

Deleted: LUC

Deleted: LUC

3.3 Change of $\Sigma E_{\text{LULCC, gross}}$ with the same net forest gain but different gross area changes

The comparison of $\Sigma E_{\text{LULCC, gross}}$ (Figure 3) for the idealized scenarios (Table 1) illustrates the fact that different values of $\gamma_{\text{Anet}}^{\text{Agross}}$ have a large impact on the magnitude and the sign of cumulative JULCC emissions depending on the time elapsed after the initial pulse of forest area change. We thus calculated the difference between $\Sigma E_{\text{LULCC, gross}}$ and $\Sigma E_{\text{LULCC, net}}$ by varying $\gamma_{\text{Anet}}^{\text{Agross}}$ in a systematic manner in a net forest gain scenario (Figure 4).

When $\gamma_{\text{Anet}}^{\text{Agross}}$ is increased, i.e., with more forest land turnover at $t = 0$ for the same initial net forest area gain ($A_{\text{net}} = +1$ ha), the time for $\Sigma E_{\text{LULCC, gross}}$ to become a net carbon sink becomes longer (Figure 4a). With initial primary forest being cut at $t = 0$, the cumulative JULCC carbon flux is still a source of CO_2 to the atmosphere after 100 years, even in simulations where the net forest area was increased at $t = 0$ (Figure 4b). This highlights that the different initial carbon density of primary forest from secondary forest can lead to very long-term legacy emissions.

The critical value of $\gamma_{\text{Anet}}^{\text{Agross}}$ that reverses the sign of $\Sigma E_{\text{LULCC, gross}}$ from carbon source to sink increases as a function of the time-horizon considered after the initial forest area change (Figure 4c). The two cases with initial secondary and primary forest loss show a different trajectory of this ratio along time. In the former, $\gamma_{\text{Anet}}^{\text{Agross}}$ increases slowly in the beginning and then sharply, while in the latter $\gamma_{\text{Anet}}^{\text{Agross}}$ increases quickly at the initial stage and then at a smaller rate. In fact, if $\Sigma E_{\text{LULCC, gross}}$ can reach zero (the point of sign changed, let $\Sigma E_{\text{LULCC, gross}} = 0$), combining with equations (1) to (6), the critical value of $\gamma_{\text{Anet}}^{\text{Agross}}$ can be expressed as:

$$\gamma_{\text{Anet}}^{\text{Agross}} = \frac{L(t) - G(t)}{L(t) + G(t)} \quad (7)$$

This critical value of $\gamma_{\text{Anet}}^{\text{Agross}}$ is independent of the initial forest area but determined by the carbon density changes at a given time consecutive to a change of forest area. Thus, for a secondary forest loss and gain at $t = 0$, the long-term $L(t) + G(t)$ tends to zero and $\gamma_{\text{Anet}}^{\text{Agross}}$ goes to infinite. For a primary forest loss and secondary forest gain at $t = 0$, the long-term $L(t) + G(t)$ is the difference in the equilibrium carbon densities between primary and secondary forest, and therefore $\gamma_{\text{Anet}}^{\text{Agross}}$ approaches a constant value at $t = \text{infinite}$. Furthermore, it should be noted that our approach of analyzing the critical value of $\gamma_{\text{Anet}}^{\text{Agross}}$ is not limited to net forest gain scenarios or to JULCC transitions between forest and cropland. The framework of $\gamma_{\text{Anet}}^{\text{Agross}}$ can also be extended to other JULCC scenarios, including lower, higher, and equal equilibrium biomass density between two land-use types. For example, if a re-growing forest can achieve a higher equilibrium carbon density than the initial one, there is also a critical $\gamma_{\text{Anet}}^{\text{Agross}}$ for the net forest loss scenario, for which the gross carbon emission becomes a sink at a certain time after initial forest area change. This situation may happen in reality, if the deforested forests are replaced by more productive species or under active management like fertilization and irrigation. Even in the field measurements by Poorter et al. (2016), some Neotropical secondary forests show very high biomass resilience, i.e., reaching to a higher biomass than pre-deforestation.

We also calculated the critical ratios over time based on the exponential biomass response curves from Hansis et al. (2015) in comparison with the response curves from Poorter et al. (2016) (Figure S3). As show in Figure 1, the equilibrium of secondary forest vegetation density with the recovery curve of Hansis et al. (2015) is higher than with Poorter et al. (2016) and we assumed that the same density of primary forest for both, and thus $L_{\text{Hansis, primary}}(t) = L_{\text{Poorter, primary}}(t)$, $L_{\text{Hansis, secondary}}(t) >$

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: densities

Deleted: between

Deleted: and

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

$L_{\text{Poorter,secondary}}(t)$ and $G_{\text{Hansis}}(\infty) > G_{\text{Poorter}}(\infty)$. Note that a positive value of carbon flux indicates carbon emission to the atmosphere. Combined with Eq (7), the different equilibrium states of secondary forest vegetation can explain the differences of critical ratios over time between Hansis et al. (2015) and Poorter et al. (2016) in Figure S3.

3.4 Ratios in Latin America from satellite imagery

- 5 Based on the theoretical evidence for the existence of a critical value of the gross-to-net forest area change ratio ($\gamma_{\text{Anet}}^{\text{Agross}}$), which determines the sign and magnitude of $\Sigma E_{\text{LULCC, gross}}$ at a given time after an initial net forest area change, we further combined such ratios with the land-cover change dataset to determine whether a region is a carbon sink or source at a given time horizon. Using the 30 m resolution forest area change data of Hansen et al. (2013) between 2000 and 2012, we calculated the ratios ($\gamma_{\text{Anet}}^{\text{Agross}}$) at a spatial resolution of $0.5^\circ \times 0.5^\circ$ in the same region of Latin America as Poorter et al. (2016). The spatial resolution
- 10 of $0.5^\circ \times 0.5^\circ$ is a typical resolution of DGVMs when they simulate global E_{LULCC} . We set a future time horizon of 20 years as that is close to the targeted year in the Nationally Determined Contributions (NDCs) (Grassi et al., 2017). From Figure 4c, the critical values of $\gamma_{\text{Anet}}^{\text{Agross}}$ at 20 years after an initial change in forest area are 7.2 and 2.4 respectively for secondary-to-secondary and primary-to-secondary initial transitions. For a longer time horizon of 50 years, the critical values are 22.5 and 3.1, respectively. After 100 years of the initial forest area change, while the critical value of $\gamma_{\text{Anet}}^{\text{Agross}}$ for secondary-to-secondary
- 15 transition goes to infinite, it approaches a constant value of 3.7 for primary-to-secondary forest change (Figure 4c). The map of $\gamma_{\text{Anet}}^{\text{Agross}}$ diagnosed from the 30 m Landsat forest cover data in grid cells of $0.5^\circ \times 0.5^\circ$ is shown in Figure 5. Note that here we focus only on the grid cells with a net forest gain. The number of $0.5^\circ \times 0.5^\circ$ grid cells where $\gamma_{\text{Anet}}^{\text{Agross}} > 7.2$, that is grid cells where current forest area change will lead to a source of CO₂ over a 20-year horizon, is 102 in our domain (Figure 5a), which accounts for 35% of the total number of grid cells where a net forest gain is observed between 2000 and 2012. In these
- 20 102 grid cells, the $\Sigma E_{\text{LULCC, gross}}$ is simulated to be a cumulative carbon emission in 20 years, no matter whether the lost forest is primary or secondary. If primary forests are cleared in grid cells with $2.4 < \gamma_{\text{Anet}}^{\text{Agross}} < 7.2$ (33% of the total forest gain grid cells, Figure 5a), the 20-year $\Sigma E_{\text{LULCC, gross}}$ is also a carbon source rather than a sink. We note that it is not possible to separate the primary and secondary forest in the forest cover data of Hansen et al. (2013), so we cannot say whether these grid cells with $2.4 < \gamma_{\text{Anet}}^{\text{Agross}} < 7.2$ are carbon source or sink in the real world. For a time horizon of 50 years, the fractions of grid cells
- 25 with $\gamma_{\text{Anet}}^{\text{Agross}} > 22.5$ and with $3.1 < \gamma_{\text{Anet}}^{\text{Agross}} < 22.5$ in total net forest gain grid cells are 14% and 46% respectively (Figure 5c). The 100-year $\Sigma E_{\text{LULCC, gross}}$ in grid cells with $\gamma_{\text{Anet}}^{\text{Agross}} > 3.7$ (53% of total) is also possible to be a carbon source if lost forest is primary in these grid cells (Figure 5d). The grid cells with $\gamma_{\text{Anet}}^{\text{Agross}}$ greater than the critical values are mainly distributed in Southeast Brazil (Figure 5b,c,d).
- By comparison, we also calculated the number of grid cells with $\gamma_{\text{Anet}}^{\text{Agross}}$ above the critical ratio for the biomass response curves
- 30 from Hansis et al. (2015) (Table S1). Because of the differences in the critical values of $\gamma_{\text{Anet}}^{\text{Agross}}$ over time (Figure S3) between curves from Poorter et al. (2016) and Hansis et al. (2015), a higher critical ratio leads to smaller number of $0.5^\circ \times 0.5^\circ$ grid cells with $\gamma_{\text{Anet}}^{\text{Agross}}$ beyond the critical ratio (Table S1).

Deleted: LUC

Deleted: we pose the question whether such ratios can be observed in the real world.

Deleted: The availability of 30 m resolution forest cover change data from satellites (Hansen et al., 2013) makes it possible to calculate these ratios for a small region of the neo-tropics, where both gross forest loss and gain are going on.

Deleted: high

Deleted: LUC

Deleted: spatial

Deleted: LUC

Deleted: LUC

Deleted: (Hansen et al., 2013)

Deleted: cumulative

Deleted: LUC

In addition to the number of grid cells with $\gamma_{\text{Anet}}^{\text{Anet}}$ above the critical ratio, we further showed the differences between the cumulative carbon flux using gross transitions ($\Sigma E_{\text{LULCC,gross}}$) and net transitions ($\Sigma E_{\text{LULCC,net}}$) in these grid cells (Table S2). Taking C1 (secondary-to-secondary) at 20 yr horizon for example, using net transitions results in a carbon sink of 12 Tg C but using gross transitions results in a carbon emission of 21 Tg C (Table S2) in the grid cells with $\gamma_{\text{Anet}}^{\text{Anet}} > 7.2$ (Figure 5b).

5 4 Discussion

The biomass recovery curves of Neotropical secondary forests from Poorter et al. (2016) are lower 20 years after the initial perturbation than those used in the bookkeeping models of Houghton (1999) and Hansis et al. (2015), implying that these models simulate different LULCC carbon fluxes in Latin America from those using the recovery curves of Poorter et al. (2016). The carbon density in undisturbed forests in the bookkeeping models of Houghton (1999) and Hansis et al. (2015) were essentially based on Whittaker and Likens (1973), multiplied by a factor of 0.75 to approximate the lower carbon density of secondary forests (Houghton et al., 1983). The carbon density data from Whittaker and Likens (1973) are subject to two sources of uncertainty. First, these values represent biomass in the 1950s (Woodwell et al., 1978) rather than present days, and second, they were compiled from very limited field measurements for tropical forests. In fact, Whittaker and Likens (1973) claimed in their study that data “for tropical communities are very meager” and the mean biomass density is “a subjectively chosen intermediate value based on very few measurements” to avoid extreme values.

Differences may also exist for soil carbon dynamics after LULCC. There are a great number of meta-analyses or reviews (Conant et al., 2001; Davidson and Ackerman, 1993; Davis and Condrón, 2002; Don et al., 2011; Guo and Gifford, 2002; Kurganova et al., 2014; Laganière et al., 2010; Li et al., 2012; Marín-Spiotta and Sharma, 2013; Murty et al., 2002; Paul et al., 2002; Poeplau et al., 2011; Post and Kwon, 2000; Powers et al., 2011; Wei et al., 2014; West et al., 2004) on the soil carbon change after LULCC based on field measurement data (mostly paired sites and chronosequences). These studies may generally agree with the directions of soil carbon change after LULCC (e.g. soil carbon loss after forest clearing for cropland), but the magnitudes and temporal dynamics of soil carbon changes remain highly uncertain because, among other things, of the limited site number and the diversity of soil properties. Field measurements at site level may be unrepresentative of the whole region because the distribution of biophysical conditions like soil texture, precipitation and temperature may not match the distribution of the whole set of such factors in the LULCC areas in a given region (Powers et al., 2011).

Some DGVMs (Bayer et al., 2017; Shevliakova et al., 2009; Stocker et al., 2014; Wilkenskjeld et al., 2014; Yue et al., 2017) as well as a bookkeeping model Hansis et al. (2015) have implemented gross land use and land cover transitions, and thus simulated a higher E_{LULCC} than using net transitions. Armeth et al. (2017) reviewed the “missing processes” in LULCC modeling by DGVMs and found that ignoring gross LULCC could underestimate the global ΣE_{LULCC} by 36 Pg C on average over the historical period (1901-2014). In this study, we used a bookkeeping method to quantify the difference in LULCC emissions calculated using net versus gross forest area transitions, and to show the existence of critical ratios of gross-to-net forest area changes above which land use action will cause a reversed sign of cumulative carbon flux. Evidently, the choice of

Formatted: Not Highlight
Deleted: (
Deleted: ,
Formatted: Not Highlight
Formatted: Not Highlight

Deleted: The biomass recovery curves of Neotropical secondary forests from Poorter et al. (2016) are lower after 20 years since the initial LUC than those used in the bookkeeping models of Houghton (1999) and Hansis et al. (2015), implying that these models may bias the temporal changes of LUC carbon fluxes in Latin America. The biomass recovery curves of the bookkeeping models were based on estimates of global or regional vegetation biomass from a synthesis of field measurements and based on prescribed recovery time depending on vegetation types and regions (Houghton, 1999).

Deleted: Differences may also exist for soil carbon dynamics after LUC, because the distribution of biophysical conditions like soil texture, precipitation and temperature from the sites that are measured may not match the distribution of the whole set of such factors in the LUC areas in a given region (Powers et al., 2011).
Formatted: Not Highlight
Formatted: Not Highlight
Formatted: Subscript
Deleted: LUC

a time horizon to assess the carbon balance of a system after an initial pulse of forest area change influences the value of the critical ratio $\gamma_{\text{Anet}}^{\text{gross}}$. The desirable target time lengths could be different depending on specific mitigation projects or land-use reduction policies, and thus critical values of the gross-to-net forest area change ratio are different (Figure 4c). On the other hand, because of the temporal evolution of legacy carbon fluxes after initial land disturbance, it is important to define a specific and reasonable time horizon when making land-based mitigation policies.

As a conceptual analysis, the assumptions we made raise uncertainties. First, the logarithmic biomass recovery curve adopted in Poorter et al. (2016) does not seem to be appropriate for J_{LULCC} emission modelling because it does not reach an equilibrium state. We thus fitted the data from Poorter et al. (2016) with an exponential saturating curve to avoid this issue. Second, we used a median biomass recovery rate for the whole tropical moist forest region in Latin America. In reality, however, due to the different climate, soils and other ecosystem conditions, recovery rates vary, and thus spatially explicit recovery rates should better depict regional patterns of secondary forest regrowth and net J_{LULCC} emissions. In the dry tropics, the critical ratio values may be smaller because of the slower biomass recovery rates. Third, the biomass and soil carbon densities in initial vegetation and the equilibrium vegetation after J_{LULCC} are also spatially different in the real world. The distinction between primary and secondary forest being lost at t = 0 is a typical example of how different initial carbon density impacts the legacy J_{LULCC} carbon flux and thus the determined critical gross-to-net ratio values. In fact, a large spatial gradient of biomass exists from Northeast to Southwest Amazon region (Saatchi et al., 2007, 2011). One possible approach to account for the spatial variations of both biomass recovery rate and biomass density would be to reconstruct spatially explicit biomass–age curves using relationship between regrowth rates and climate (Poorter et al., 2016) and to combine with observation-based biomass densities (Baccini et al., 2012; Saatchi et al., 2011) and satellite-based forest cover change (Hansen et al., 2013). However, uncertainties arise in the up-scaling of biomass recovery rates and lack of information on annually resolved forest gain from Hansen et al. (2013). In addition, spatially explicit soil carbon density maps are also uncertain.

The effect of gross-versus-net forest area change on legacy J_{LULCC} emissions certainly differs across forest ecosystems and other J_{LULCC} transition types (e.g. transitions between grassland and cropland). The concept of critical ratios of gross-to-net J_{LULCC} affecting legacy carbon balance can be extended in other regions where forest management practice is critical (e.g. North America and Europe). Forest management practices like wood harvest and thinning extract carbon from the ecosystem and release it to the atmosphere (Houghton et al., 2012), while recovering secondary forest from past deforestation and logging (Pan et al., 2011) and even old-growth forests (Luyssaert et al., 2012) can act as carbon sinks. In theory, likewise, a critical ratio value should exist to balance the bi-directional carbon fluxes in forest management practices. An advantage of this concept of critical ratio is that it can be directly measured with satellite observations, which provides a quick guide for local land use management practice through near-real-time forest cover change data (e.g. Global Forest Watch <http://www.globalforestwatch.org/>).

Accurate estimates of J_{LULCC} carbon fluxes in the Neotropical forests are increasingly important for climate mitigation policy with the progressive implementation of Reducing Emissions from Deforestation and forest Degradation (REDD+) programs under the United Nations Framework Convention on Climate Change (UNCCC) (Angelsen et al., 2009; Magnago et al., 2015).

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Deleted: Forest management practices like wood harvest and thinning extract carbon from the ecosystem and release it to the atmosphere (Houghton et al., 2012), while the new planted forest in rotation practice and even old-growth forests can act as carbon sinks (Luyssaert et al., 2008)

Deleted: LUC

Furthermore, forest-based climate mitigation has been taken as a key option in the Nationally Determined Contributions (NDCs) proposed by some countries to the Paris Climate Agreement, accounting for about one-fourth of total intended emission reductions from a pre-defined baseline (Grassi et al., 2017). Brazil contributes about one-third of the global forest-based emission reduction in the NDCs (Grassi et al., 2017). Based on the results of this study, we argue that it will be important to carefully distinguish the amount of gross vs. net forest changes and clearing of primary vs. secondary forest when assessing national forest-based mitigation pledges. **With a large gross to net area change ratio, a net forest gain could still legate a net carbon source over a long period in the future.** Our work has the potential to be extended to country-level and other J_{LULCC} types as long as information on vegetation and soil carbon densities changes after J_{LULCC} is available, and a critical value of $\gamma_{\text{Anet}}^{\text{Agross}}$ can be estimated as a guideline to evaluate land-based mitigation policies for each region. More observation-based data on land-use area change and carbon loss and gain curves will definitely help to extend the range of applications of the critical gross-to-net area ratio concept.

Deleted: With a too high rotation rate of forests, i.e. a large gross to net area change ratio, a net forest gain could still legate a net carbon source over a long period in the future.

Deleted: LUC

Deleted: LUC

5 Conclusions

Using only net J_{LULCC} transitions instead of gross values can bias the magnitude of estimated J_{LULCC} carbon fluxes, to the point of estimating a sink instead of a source in reality if high gross forest area change occurs. We used idealized scenarios to demonstrate different aspects of the discrepancy between net and gross forest changes, defining the $\gamma_{\text{Anet}}^{\text{Agross}}$ metric as the ratio of gross area change to net area change. Our S0 experiment shows even that there is no net forest change, J_{LULCC} may actually lead to a carbon source, depending on the gross forest change area. S1 and S2 show that with the same net forest loss, different ratios of gross-to-net forest change ($\gamma_{\text{Anet}}^{\text{Agross}}$) alter the magnitude of differences between net and gross cumulative carbon fluxes. Similarly, S3 and S4 show that with the same amount of net forest gain area, different $\gamma_{\text{Anet}}^{\text{Agross}}$ can even change the directions of carbon fluxes, i.e. from a gross carbon sink to source even that net forest area increases. We further determined the critical ratios in net forest gain grid cells ($\gamma_{\text{Anet}}^{\text{Agross}} = 7.2$ and 2.4 respectively for secondary and primary forest clearing), above which the gross cumulative carbon fluxes show a reversed sign than the net ones at 20 years after J_{LULCC} occurred. These analyses reveal the importance of using gross J_{LULCC} transitions rather than net J_{LULCC} transitions in both bookkeeping models and DGVMs. The concept of critical ratio can be also implemented in other J_{LULCC} transitions in other regions and used as a guide for carbon balance estimation in forest management.

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Deleted: LUC

Acknowledgements

W.L., P.C., T.G. and S.P. acknowledge support from the European Research Council through Synergy grant ERC-2013-SyG-610028 “IMBALANCE-P”. W.L. and C.Y. are supported by the European Commission-funded project LUC4C (No. 603542).

References

- Angelsen, A., Brown, S., Loisel, C., Peskett, L., Streck, C. and Zarin, D.: Reducing Emissions from Deforestation and Forest Degradation (REDD): An Options Assessment Report., 2009.
- Armeth, A., Sitch, S., Pongratz, J., Stocker, B. D., Ciais, P., Poulter, B., Bayer, A. D., Bondeau, A., Calle, L., Chini, L. P., Gasser, T., Fader, M., Friedlingstein, P., Kato, E., Li, W., Lindeskog, M., Nabel, J. E. M. S., Pugh, T. A. M., Robertson, E., Viovy, N., Yue, C. and Zaehle, S.: Historical carbon dioxide emissions caused by land-use changes are possibly larger than assumed, *Nat. Geosci.*, 10(2), 79–84, doi:10.1038/ngeo2882, 2017.
- Baccini, A., Goetz, S. J., Walker, W. S., Laporte, N. T., Sun, M., Sulla-Menashe, D., Hackler, J., Beck, P. S. A., Dubayah, R., Friedl, M. A., Samanta, S. and Houghton, R. A.: Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps, *Nat. Clim. Chang.*, 2(3), 182–185, doi:10.1038/nclimate1354, 2012.
- Bayer, A. D., Lindeskog, M., Pugh, T. A. M., Anthoni, P. M., Fuchs, R. and Armeth, A.: Uncertainties in the land-use flux resulting from land-use change reconstructions and gross land transitions, *Earth Syst. Dyn.*, 8(1), 91–111, doi:10.5194/esd-8-91-2017, 2017.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Quéré, C. Le, Myeni, R. B., Piao, S., Thornton, P., France, P. C., Willem, J., Friedlingstein, P. and Munhoven, G.: 2013: Carbon and Other Biogeochemical Cycles, *Clim. Chang. 2013 Phys. Sci. Basis. Contrib. Work. Gr. I to Fifth Assess. Rep. Intergov. Panel Clim. Chang.*, 465–570, doi:10.1017/CBO9781107415324.015, 2013.
- Conant, R. T., Paustian, K. and Elliott, E. T.: Grassland management and conversion into grassland: Effects on soil carbon, *Ecol. Appl.*, 11(2), 343–355, doi:10.2307/3060893, 2001.
- Davidson, E. A. and Ackerman, I. L.: Changes in soil carbon inventories following cultivation of previously untilled soils, *Biogeochemistry*, 20(3), 161–193, doi:10.1007/BF00000786, 1993.
- Davis, M. R. and Condon, L. M.: Impact of grassland afforestation on soil carbon in New Zealand: A review of paired-site studies, *Aust. J. Soil Res.*, 40(4), 675–690, doi:10.1071/SR01074, 2002.
- Don, A., Schumacher, J. and Freibauer, A.: Impact of tropical land-use change on soil organic carbon stocks - a meta-analysis, *Global Chang. Biol.*, 17(4), 1658–1670, doi:10.1111/j.1365-2486.2010.02336.x, 2011.
- FAO: Global Forest Resources Assessment 2010, 2010.
- Fuchs, R., Herold, M., Verburg, P. H., Clevers, J. G. P. W. and Eberle, J.: Gross changes in reconstructions of historic land cover/use for Europe between 1900 and 2010, *Global Chang. Biol.*, 21(1), 299–313, doi:10.1111/gcb.12714, 2015.
- Fuchs, R., Schulp, C. J. E., Hengeveld, G. M., Verburg, P. H., Clevers, J. G. P. W., Schelhaas, M.-J. and Herold, M.: Assessing the influence of historic net and gross land changes on the carbon fluxes of Europe, *Global Chang. Biol.*, 22(7), 2526–2539, doi:10.1111/gcb.13191, 2016.
- Gasser, T. and Ciais, P.: A theoretical framework for the net land-to-atmosphere CO₂ flux and its implications in the definition of “emissions from land-use change,” *Earth Syst. Dyn.*, 4(1), 171–186, doi:10.5194/esd-4-171-2013, 2013.

- Gasser, T., Ciais, P., Boucher, O., Quilcaille, Y., Tortora, M., Bopp, L. and Hauglustaine, D.: The compact Earth system model OSCAR v2.2: description and first results, *Geosci. Model Dev.*, 10(1), 271–319, doi:10.5194/gmd-10-271-2017, 2017.
- Grassi, G., House, J., Dentener, F., Federici, S., den Elzen, M. and Penman, J.: The key role of forests in meeting climate targets requires science for credible mitigation, *Nat. Clim. Chang.*, 7(3), 220–226, 2017.
- 5 Guo, L. B. and Gifford, R. M.: Soil carbon stocks and land use change: A meta analysis, *Global Chang. Biol.*, 8(4), 345–360, doi:10.1046/j.1354-1013.2002.00486.x, 2002.
- Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. a, Tyukavina, a, Thau, D., Stehman, S. V., Goetz, S. J., Loveland, T. R., Kommareddy, a, Egorov, a, Chini, L., Justice, C. O. and Townshend, J. R. G.: High-resolution global maps of 21st-century forest cover change., *Science*, 342(6160), 850–3, doi:10.1126/science.1244693, 2013.
- 10 Hansis, E., Davis, S. J. and Pongratz, J.: Relevance of methodological choices for accounting of land use change carbon fluxes, *Global Biogeochem. Cycles*, 29(8), 1230–1246, doi:10.1002/2014GB004997, 2015.
- Houghton, R. A.: The annual net flux of carbon to the atmosphere from changes in land use 1850-1990, *Tellus B*, 51(2), 298–313, doi:10.1034/j.1600-0889.1999.00013.x, 1999.
- Houghton, R. A.: Revised estimates of the annual net flux of carbon to the atmosphere from changes in land use and land management 1850-2000, *Tellus Ser. B-Chemical Phys. Meteorol.*, 55(2), 378–390, doi:10.1034/j.1600-0889.2003.01450.x, 2003.
- 15 Houghton, R. A., Hobbie, J. E., Melillo, J. M., Moore, B., Peterson, B. J., Shaver, G. R. and Woodwell, G. M.: Changes in the Carbon Content of Terrestrial Biota and Soils between 1860 and 1980: A Net Release of CO² to the Atmosphere, *Ecol. Monogr.*, 53(3), 235–262, doi:10.2307/1942531, 1983.
- 20 Houghton, R. a., House, J. I., Pongratz, J., Van Der Werf, G. R., Defries, R. S., Hansen, M. C., Le Quéré, C. and Ramankutty, N.: Carbon emissions from land use and land-cover change, *Biogeosciences*, 9(12), 5125–5142, doi:10.5194/bg-9-5125-2012, 2012.
- Houghton, R. A. and Nassikas, A. A.: Global and regional fluxes of carbon from land use and land cover change 1850-2015, *Global Biogeochem. Cycles*, 31(3), 456–472, doi:10.1002/2016GB005546, 2017.
- 25 Hurtt, G. C., Chini, L. P., Frolking, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P., Hibbard, K., Houghton, R. A., Janetos, A., Jones, C. D., Kindermann, G., Kinoshita, T., Klein Goldewijk, K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D. P. and Wang, Y. P.: Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands, *Clim. Change*, 109(1–2), 117–161, doi:10.1007/s10584-011-0153-2, 2011.
- 30 Kurganova, I., Lopes de Gerenyu, V., Six, J. and Kuzyakov, Y.: Carbon cost of collective farming collapse in Russia, *Global Chang. Biol.*, 20(3), 938–947, doi:10.1111/gcb.12379, 2014.
- Laganière, J., Angers, D. A. and Paré, D.: Carbon accumulation in agricultural soils after afforestation: A meta-analysis, *Global Chang. Biol.*, 16(1), 439–453, doi:10.1111/j.1365-2486.2009.01930.x, 2010.
- Li, D., Niu, S. and Luo, Y.: Global patterns of the dynamics of soil carbon and nitrogen stocks following afforestation: A meta-

- analysis, *New Phytol.*, 195, 172–181, doi:10.1111/j.1469-8137.2012.04150.x, 2012.
- Li, W., Ciais, P., Peng, S., Yue, C., Wang, Y., Thurner, M., Saatchi, S. S., Arno, A., Avitabile, V., Carvalhais, N., Harper, A. B., Kato, E., Koven, C., Liu, Y. Y., Nabel, J. E. M. S., Pan, Y., Pongratz, J., Poulter, B., Pugh, T. A. M., Santoro, M., Sitch, S., Stocker, B. D., Viovy, N., Wiltshire, A., Yousefpour, R. and Zaehle, S.: Land-use and land-cover change carbon emissions between 1901 and 2012 constrained by biomass observations, *Biogeosciences Discuss.*, 1–25, doi:10.5194/bg-2017-186, 2017.
- 5 Liu, Y. Y., van Dijk, A. I. J. M., de Jeu, R. A. M., Canadell, J. G., McCabe, M. F., Evans, J. P. and Wang, G.: Recent reversal in loss of global terrestrial biomass, *Nat. Clim. Chang.*, 5(5), 470–474, doi:10.1038/nclimate2581, 2015.
- Luysaert, S., Abril, G., Andres, R., Bastviken, D., Bellassen, V., Bergamaschi, P., Bousquet, P., Chevallier, F., Ciais, P., Corazza, M., Dechow, R., Erb, K.-H., Etiope, G., Fortems-Cheiney, A., Grassi, G., Hartmann, J., Jung, M., Lathière, J., Lohila, A., Mayorga, E., Moosdorf, N., Njakou, D. S., Otto, J., Papale, D., Peters, W., Peylin, P., Raymond, P., Rödenbeck, C., Saarnio, S., Schulze, E.-D., Szopa, S., Thompson, R., Verkerk, P. J., Vuichard, N., Wang, R., Wattenbach, M. and Zaehle, S.: The European land and inland water CO₂, CO, CH₄ and N₂O balance between 2001 and 2005, *Biogeosciences*, 9(8), 3357–3380, doi:10.5194/bg-9-3357-2012, 2012.
- 10 Magnago, L. F. S., Magrach, A., Laurance, W. F., Martins, S. V., Meira-Neto, J. A. A., Simonelli, M. and Edwards, D. P.: Would protecting tropical forest fragments provide carbon and biodiversity cobenefits under REDD+?, *Global Chang. Biol.*, 21(9), 3455–3468, doi:10.1111/gcb.12937, 2015.
- Marin-Spiotta, E. and Sharma, S.: Carbon storage in successional and plantation forest soils: A tropical analysis, *Global Ecol. Biogeogr.*, 22(1), 105–117, doi:10.1111/j.1466-8238.2012.00788.x, 2013.
- Murty, D., Kirschbaum, M. U. F., McMurtrie, R. E. and McGilvray, H.: Does conversion of forest to agricultural land change soil carbon and nitrogen? a review of the literature, *Global Chang. Biol.*, 8(2), 105–123, doi:10.1046/j.1354-1013.2001.00459.x, 2002.
- 20 Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L., Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W., McGuire, A. D., Piao, S., Rautiainen, A., Sitch, S. and Hayes, D.: A large and persistent carbon sink in the world's forests., *Science*, 333(6045), 988–993, doi:10.1126/science.1201609, 2011.
- 25 Paul, K. I., Polglase, P. J., Nyakuengama, J. G. and Khanna, P. K.: Change in soil carbon following afforestation, *Forest Ecol. Manag.*, 168(1–3), 241–257, doi:10.1016/S0378-1127(01)00740-X, 2002.
- Peacock, J., Baker, T. R., Lewis, S. L., Lopez-Gonzalez, G. and Phillips, O. L.: The RAINFOR database: monitoring forest biomass and dynamics, *J. Veg. Sci.*, 18(4), 535–542, doi:10.1111/j.1654-1103.2007.tb02568.x, 2007.
- Poeplau, C., Don, A., Vesterdal, L., Leifeld, J., Van Wesemael, B., Schumacher, J. and Gensior, A.: Temporal dynamics of soil organic carbon after land-use change in the temperate zone - carbon response functions as a model approach, *Global Chang. Biol.*, 17(7), 2415–2427, doi:10.1111/j.1365-2486.2011.02408.x, 2011.
- 30 Poorter, L., Bongers, F., Aide, T. M., Almeyda Zambrano, A. M., Balvanera, P., Becknell, J. M., Boukili, V., Brancalion, P. H. S., Broadbent, E. N., Chazdon, R. L., Craven, D., de Almeida-Cortez, J. S., Cabral, G. A. L., de Jong, B. H. J., Denslow, J. S., Dent, D. H., DeWalt, S. J., Dupuy, J. M., Durán, S. M., Espirito-Santo, M. M., Fandino, M. C., César, R. G., Hall, J. S.,

- Hernandez-Stefanoni, J. L., Jakovac, C. C., Junqueira, A. B., Kennard, D., Letcher, S. G., Licona, J.-C., Lohbeck, M., Marin-Spiotta, E., Martínez-Ramos, M., Massoca, P., Meave, J. A., Mesquita, R., Mora, F., Muñoz, R., Muscarella, R., Nunes, Y. R. F., Ochoa-Gaona, S., de Oliveira, A. A., Orihuela-Belmonte, E., Peña-Claros, M., Pérez-García, E. A., Piotta, D., Powers, J. S., Rodríguez-Velázquez, J., Romero-Pérez, I. E., Ruiz, J., Saldarriaga, J. G., Sanchez-Azofeifa, A., Schwartz, N. B.,
5 Steininger, M. K., Swenson, N. G., Toledo, M., Uriarte, M., van Breugel, M., van der Wal, H., Veloso, M. D. M., Vester, H. F. M., Vicentini, A., Vieira, I. C. G., Bentos, T. V., Williamson, G. B. and Rozendaal, D. M. A.: Biomass resilience of Neotropical secondary forests., *Nature*, 530(7589), 211–214, doi:10.1038/nature16512, 2016.
- Post, W. M. and Kwon, K. C.: Soil carbon sequestration and land-use change: Processes and potential, *Global Chang. Biol.*, 6(3), 317–327, doi:10.1046/j.1365-2486.2000.00308.x, 2000.
- 10 Powers, J. S., Corre, M. D., Twine, T. E. and Veldkamp, E.: Geographic bias of field observations of soil carbon stocks with tropical land-use changes precludes spatial extrapolation., *Proc. Natl. Acad. Sci. USA*, 108(15), 6318–6322, doi:10.1073/pnas.1016774108, 2011.
- Le Quéré, C., Moriarty, R., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Friedlingstein, P., Peters, G. P., Andres, R. J., Boden, T. A., Houghton, R. A., House, J. I., Keeling, R. F., Tans, P., Arneeth, A., Bakker, D. C. E., Barbero, L., Bopp,
15 L., Chang, J., Chevallier, F., Chini, L. P., Ciais, P., Fader, M., Feely, R. A., Gkritzalis, T., Harris, I., Hauck, J., Ilyina, T., Jain, A. K., Kato, E., Kitidis, V., Klein Goldewijk, K., Koven, C., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lima, I. D., Metzl, N., Millero, F., Munro, D. R., Murata, A., Nabel, J. E. M. S., Nakaoka, S., Nojiri, Y., O'Brien, K., Olsen, A., Ono, T., Pérez, F. F., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Rödenbeck, C., Saito, S., Schuster, U., Schwinger, J., Séférian, R., Steinhoff, T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van der Laan-Luijckx, I. T., van der Werf,
20 G. R., van Heuven, S., Vandemark, D., Viovy, N., Wiltshire, A., Zaehle, S. and Zeng, N.: Global Carbon Budget 2015, *Earth Syst. Sci. Data*, 7(2), 349–396, doi:10.5194/essd-7-349-2015, 2015.
- Saatchi, S., Houghton, R. A., Dos Santos Alvalá, R. C., Soares, J. V. and Yu, Y.: Distribution of aboveground live biomass in the Amazon basin, *Global Chang. Biol.*, 13(4), 816–837, doi:10.1111/j.1365-2486.2007.01323.x, 2007.
- Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T. A., Salas, W., Zutta, B. R., Buermann, W., Lewis, S. L.,
25 Hagen, S., Petrova, S., White, L., Silman, M. and Morel, A.: Benchmark map of forest carbon stocks in tropical regions across three continents., *Proc. Natl. Acad. Sci. USA*, 108(24), 9899–9904, doi:10.1073/pnas.1019576108, 2011.
- Shevliakova, E., Pacala, S. W., Malyshev, S., Hurtt, G. C., Milly, P. C. D., Caspersen, J. P., Sentman, L. T., Fisk, J. P., Wirth, C. and Crevoisier, C.: Carbon cycling under 300 years of land use change: Importance of the secondary vegetation sink, *Global Biogeochem. Cycles*, 23(2), n/a-n/a, doi:10.1029/2007GB003176, 2009.
- 30 Sitch, S., Friedlingstein, P., Gruber, N., Jones, S. D., Murray-Tortarolo, G., Ahlström, A., Doney, S. C., Graven, H., Heinze, C., Huntingford, C., Levis, S., Levy, P. E., Lomas, M., Poulter, B., Viovy, N., Zaehle, S., Zeng, N., Arneeth, A., Bonan, G., Bopp, L., Canadell, J. G., Chevallier, F., Ciais, P., Ellis, R., Gloor, M., Peylin, P., Piao, S. L., Le Quéré, C., Smith, B., Zhu, Z. and Myneni, R.: Recent trends and drivers of regional sources and sinks of carbon dioxide, *Biogeosciences*, 12(3), 653–679, doi:10.5194/bg-12-653-2015, 2015.

- Stocker, B., Feissli, F., Strassmann, K. and Physics, E.: Past and future carbon fluxes from land use change, shifting cultivation and wood harvest, *Tellus B*, 1, 1–15, doi:10.3402/tellusb.v66.23188, 2014.
- Wei, X., Shao, M., Gale, W. and Li, L.: Global pattern of soil carbon losses due to the conversion of forests to agricultural land., *Sci. Rep.*, 4, 4062, doi:10.1038/srep04062, 2014.
- 5 West, T. O., Marland, G., King, A. W., Post, W. M., Jain, A. K. and Andrasko, K.: Carbon Management Response curves: estimates of temporal soil carbon dynamics., *Environ. Manage.*, 33(4), 507–18, doi:10.1007/s00267-003-9108-3, 2004.
- Whittaker, R. H. and Likens, G. E.: Carbon in the biota., in *Brookhaven symposia in biology*, pp. 281–302., 1973.
- Wilkenskjeld, S., Kloster, S., Pongratz, J., Raddatz, T. and Reick, C. H.: Comparing the influence of net and gross anthropogenic land-use and land-cover changes on the carbon cycle in the MPI-ESM, *Biogeosciences*, 11(17), 4817–4828, doi:10.5194/bg-11-4817-2014, 2014.
- 10 Woodwell, G. M., Whittaker, R. H., Reiners, W. a, Likens, G. E., Delwiche, C. C. and Botkin, D. B.: The biota and the world carbon budget, *Science*, 199(4325), 141–146, doi:10.1126/science.199.4325.141, 1978.
- Yue, C., Ciais, P., Luysaert, S., Li, W., McGrath, M., Chang, J. and Peng, S.: Representing anthropogenic gross land use change, wood harvest and forest age dynamics in a global vegetation model ORCHIDEE-MICT (r4259), *Geosci. Model Dev. Discuss.*, 2017.
- 15

Table 1 Illustrative scenarios with different ratios of gross-to-net forest area changes impacting legacy LULCC emissions after a pulse disturbance of forest area at $t = 0$. A_{net} , A_{gross} , A_{loss} , A_{gain} and $\gamma_{net}^{A_{gross}}$ are the applied net forest area change, gross forest area change, gross forest loss area, gross forest gain area and the ratio of A_{gross} to A_{net} at $t = 0$. Positive value of an area change is an increase of forest area.

Scenario	$\gamma_{A_{net}}^{A_{gross}}$	A_{net} (ha)	A_{gross} (ha)	A_{loss} (ha)	A_{gain} (ha)
S0	$\gamma_0^2 = \infty$	0	2	-1	1
S1	$\gamma_{i,1}^{1,2} = -1.2$	-1	1.2	-1.1	0.1
S2	$\gamma_{i,1}^{201} = -201$	-1	201	-101	100
S3	$\gamma_{i,1}^{1,2} = 1.2$	1	1.2	-0.1	1.1
S4	$\gamma_{i,1}^{201} = 201$	1	201	-100	101

5

Table 2 Different combinations of response curves to calculate E_{LULCC} .

Combination	Forest loss		Forest gain		
	forest type	response curve in all carbon pools	forest type	response curve for biomass	response curve for soil
C1	secondary	exponential, Hansis	secondary	logarithmic, Poorter	exponential, Hansis
C2	primary	exponential, Hansis	secondary	logarithmic, Poorter	exponential, Hansis
C3	secondary	exponential, Hansis	secondary	exponential, Poorter	exponential, Hansis
C4	primary	exponential, Hansis	secondary	exponential, Poorter	exponential, Hansis
C5	secondary	linear, Houghton	secondary	logarithmic, Poorter	exponential, Hansis
C6	primary	linear, Houghton	secondary	logarithmic, Poorter	exponential, Hansis
C7	secondary	exponential, Hansis	secondary	exponential, Hansis	exponential, Hansis
C8	primary	exponential, Hansis	secondary	exponential, Hansis	exponential, Hansis

Deleted: LUC

Deleted: g

Deleted: LUC

Figure 1 Response curves for tropical moist forest in bookkeeping models and from a recent field study. Solid and dotted lines indicate the linear (Houghton, 1999) and exponential (Hansis et al., 2015) curves, respectively. Lime dashed and dash-dotted lines are the logarithmic and exponential curves from forest plots (Poorter et al., 2016). **Vegetation carbon density in primary forest (Houghton, 1999)** is also shown as a star in (c) for comparison.

Deleted: Biomass




Figure 2 An illustration of different gross forest area changes with the same net area change. (a) Net forest gain with small gross secondary forest area changes (secondary-to-secondary), thus low $\gamma_{A_{net}}^{A_{gross}}$. (b) Same net forest gain as (a) but with large gross secondary forest area changes (secondary-to-secondary), thus high $\gamma_{A_{net}}^{A_{gross}}$. (c) Same as (a) but with gross primary forest loss (primary-to-secondary) instead of gross secondary loss.

5




Figure 3 Cumulative carbon flux ($\sum E_{LULCC}$) after an initial forest area change at $t = 0$ followed by no change in forest area, for the different scenarios S0 to S4 in Table 1 with different net and gross initial forest area changes. The response curves used in those bookkeeping model simulations are C1 in solid lines (Table 2) with a secondary-to-secondary forest change at $t = 0$, and a logarithmic biomass recovery curve with an asymptote, and C2 in the dashed lines (primary-to-secondary forest change at $t = 0$ and a logarithmic biomass recovery curve with an asymptote). The dotted line is the zero line. Positive value of carbon flux indicates carbon emission to the atmosphere.

Deleted: LUC

Deleted: ,





Figure 4 Time evolution of cumulative carbon flux ($\Sigma E_{LU,CC, gross}$) after an initial forest area change involving gross forest area changes followed by no forest area change. The three panels show results of our bookkeeping model for three case studies (a) a net forest gain at $t = 0$ with initial secondary forest loss followed by secondary forest regrowth (secondary-to-secondary, C1 in Table 2), (b) the same net area gain at $t = 0$ with initial primary forest loss followed by secondary forest regrowth (primary-to-secondary C2 in Table 2), and (c) the critical value of γ_{Anet}^{Agross} at which $\Sigma E_{LU,CC, gross}$ is zero, going from a net source to a net sink for different time horizon in the x-axis. The colored curves in (a) and (b) have the same net area change ($A_{net} = +1$ ha) at $t = 0$ but variable values of the initial gross-to-net area change ratios (γ_{Anet}^{Agross}). The red line in (a) and (b) is the zero line, defining the time after initial forest area change at which the system reaches a neutral carbon balance. The light and dark green lines in (c) represent the critical ratios for a net initial forest gain scenario with secondary-to-secondary (a) and primary-to-secondary (b) gross forest area change, respectively. Values larger than this critical value indicate that the initial forest area change has the net effect to emit CO_2 for a given time horizon in the x-axis. Exponential curve from Hansis et al. (2015) for carbon loss in all pools and gain in soil pool and logarithmic curve from Poorter et al. (2016) for gain in biomass pool are used in this example (Table 2).



Formatted: Font color: Auto

Field Code Changed

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Deleted: Time evolution of cumulative carbon flux ($\Sigma E_{LU,CC, gross}$) after an initial forest area change involving gross forest area changes followed by no forest area change. The colored curves have the same net area change ($A_{net} = +1$ ha) at $t = 0$ but variable values of the initial gross-to-net area change ratios (γ_{Anet}^{Agross}). The three panels show results of our bookkeeping model for (a) a net forest gain at $t = 0$ scenario with gross secondary forests (secondary-to-secondary), (b) the same net area gain at $t = 0$ with gross primary forest loss (primary-to-secondary), and (c) the critical value of γ_{Anet}^{Agross} at which $\Sigma E_{LU,CC, gross}$ is zero, going from a net source to a net sink with different time horizon. Values larger than this critical value indicate that the initial forest area change has the net effect to emit CO_2 . Exponential carbon loss curve from (Hansis et al., 2015) and logarithmic gain curve from (Poorter et al., 2016) are used in this example. The red line in (a) and (b) is the zero line, defining the time after initial disturbance at which the system reaches a neutral carbon balance. The light and dark green lines in (c) represent the critical ratios for a net initial forest gain scenario with secondary-to-secondary (a) and primary-to-secondary (b) gross forest area change, respectively.

Deleted: .

Formatted: Justified, Line spacing: 1.5 lines

Figure 5 Ratios of gross-to-net forest area change (γ_{Anet}^{Agross}) in $0.5^\circ \times 0.5^\circ$ grid cells in Latin America (same region as Poorter et al. (2016)) calculated from the high-resolution forest cover change map (Hansen et al., 2013). Grid cells with $\gamma_{Anet}^{Agross} < 2.4$ are masked. (b) is the zoom-in area of 20-30° S and 40-60° W in (a) (red rectangle) and grid cells with $\gamma_{Anet}^{Agross} > 7.2$ and with $2.4 < \gamma_{Anet}^{Agross} < 7.2$ are shown as blue and green respectively to indicate those beyond the critical ratios with a time horizon of 20 years. (c) and (d) are similar to (b) but indicate a time horizon of 50 and 100 years respectively. The blue grid cells in (b) and (c) represent a cumulative carbon emission in 20 years no matter whether the lost forest is primary or secondary. The green ones in (b), (c) and (d) represent a cumulative carbon emission only if the cleared forests are primary forests.

Deleted: (

Deleted: ,

