



Evidence of a prolonged drought ca. 4200 yr BP correlated with prehistoric settlement abandonment from the Gueldaman GLD1 Cave, Northern Algeria

J. Ruan^{1,5}, F. Kherbouche², D. Genty¹, D. Blamart¹, H. Cheng^{3,4}, F. Dewilde¹, S. Hachi², R. L. Edwards⁴, E. Régnier¹, and J.-L. Michelot⁵

¹Laboratoire des Sciences du Climat et de l'Environnement, Gif-sur-Yvette, France

²Centre National de Recherches Préhistoriques, Anthropologiques et Historiques, Algiers, Algeria

³Institute of Global Environmental Change, Xi'an Jiaotong University, Xi'an, China

⁴Department of Geological Sciences, University of Minnesota, Minnesota, USA

⁵Laboratoire Géosciences Paris Sud, UMR 8148, Université Paris-Sud, Orsay, France

Correspondence to: J. Ruan (jiaoyangruan@gmail.com)

Received: 12 June 2015 – Published in Clim. Past Discuss.: 3 July 2015

Revised: 3 December 2015 – Accepted: 14 December 2015 – Published: 15 January 2016

Abstract. Middle Holocene cultures have been widely studied around the Eastern-Mediterranean basin in the last 30 years and past cultural activities have been commonly linked with regional climate changes. However, in many cases such linkage is equivocal, in part due to existing climatic evidence that has been derived from areas outside the distribution of ancient settlements, leading to uncertainty from complex spatial heterogeneity in both climate and demography. A few high-resolution well-dated paleoclimate records were recently established using speleothems in the Central and Eastern-Mediterranean basin, however, the scarcity of such records in the western part of the Mediterranean prevents us from correlating past climate evolutions across the basin and deciphering climate–culture relation at fine timescales.

Here we report the first decadal-resolved Mid-Holocene climate proxy records from the Western-Mediterranean basin based on the stable carbon and oxygen isotopes analyses of two U/Th dated stalagmites from the Gueldaman GLD1 Cave in Northern Algeria. Comparison of our records with those from Italy and Israel reveals synchronous (multi) centennial dry phases centered at ca. 5600, ca. 5200 and ca. 4200 yr BP across the Mediterranean basin. New calibrated radiocarbon dating constrains reasonably well the age of rich anthropogenic deposits (e.g., faunal remains, pottery, charcoal) excavated inside the cave, which allows the comparison

between in situ evidence of human occupation and of climate change. This approach shows that the timing of a prolonged drought at ca. 4400–3800 yr BP blankets the onset of cave abandonment shortly after ca. 4403 cal yr BP, supporting the hypothesis that a climate anomaly may have played a role in this cultural disruption.

1 Introduction

As drought in Northwestern Africa is a recurring phenomenon and prolonged dry conditions exert a significant impact on local social systems, it becomes important to accurately document the role of drought conditions on the area. For instance, the most recent drought in Algeria began in 1998, as part of a widespread pattern of drying in the Northern Hemisphere, and brought considerable loss in regards to water resource and agricultural yields (Hoerling and Kumar, 2003). Increasingly dry sub-tropical conditions are predicted as one potential consequence of anthropogenic climate change, but current general circulation models do not completely capture the magnitude and spatial extent of observed drought conditions (Seager et al., 2007). To help understand recent climate anomalies, paleoclimate studies are crucial to characterize the range of potential natural variability in the past and to improve our understanding of the links between regional drought and large-scale forcing. Instrumental data




Figure 1. The Gueldaman GLD1 Cave ($36^{\circ}26' \text{ N}$, $4^{\circ}34' \text{ E}$, 507 m a.s.l.). The top left shows the location of the Gueldaman GLD1 Cave, in the Northern-Algerian part of the Western-Mediterranean basin; the bottom left shows a photo of cave entrance and local vegetation cover; the right panel shows maps of inner cave where stalagmites and archaeological deposits are collected.

from weather stations in Northwestern Africa report less than 100 years. Tree ring based drought reconstructions in Algeria and Tunisia have been extended back to the last 9 centuries, which reveals large spatial heterogeneity of past climate evolutions in Northwestern Africa and concludes that the climate anomaly 1998–2002 appears to be the most severe in the last millennium (Touchan et al., 2008, 2011). Holocene paleoclimate studies in other regions, however, have suggested larger oscillations at centennial to millennial timescales highlighting the need for new records from this area (Mayewski et al., 2004; Wanner et al., 2008).

A significant climate excursion ca. 4200 yr BP has been widely reported and is considered as an ideal case to study the causes and effects of a large-scale climate anomaly that occurred against background conditions similar to those of today (Berkelhammer et al., 2013; Booth et al., 2005; Dixit et al., 2014; Roland, 2012). The climatic expression of the 4200 yr BP event differs around the world. For example, it has been documented as droughts in much of mid-to-low latitudes, across Africa, Asia and North America, wet and stormy in Northern Europe and cooler in North Atlantic (Booth et al., 2005; Roland, 2012). More recently, this climatic anomaly was characterized by extreme dry conditions on high-resolved speleothem isotope records from the Central (Drysdale et al., 2006; Zanchetta et al., 2014) and Eastern-Mediterranean basin (Bar-Matthews and Ayalon, 2011), but, until now, such records have not been available in the Western Mediterranean which prevents us from

having access to the correlation of past climate anomalies across the basin.

Aside from its climatic interest, such an episode likely influenced numerous human cultures. Major societal changes have been observed across the Mediterranean basin during the Mid-Holocene, and in particular, a catastrophic desiccation ca. 4200 yr BP has been suggested to be the trigger of the collapse of the Akkadian Empire in Mesopotamia, the Old Kingdom in Egypt and the Early Bronze Age civilizations of Greece and Crete (Weiss and Bradley, 2000; Weiss et al., 1993; Wiener, 2014). These studies have been stimulating an increasing number of debates on climate–culture relationship (e.g., Coombes and Barber, 2005). Uncertainty regarding the societal impact of such an event is still large, due in part that climatic evidence, in many cases, has been derived from regions far from the distribution of ancient settlements (e.g., Cullen et al., 2000). Although the 4200 yr BP dry event has been observed in several mid-latitude sites, the database remains incomplete and conflicting observations of climatic conditions between seemingly adjacent regions exist (Magny et al., 2013; Staubwasser and Weiss, 2006). Additionally, a recent study demonstrated that the climatic impact on many agricultural settlements in ancient Near East was diverse even within spatially limited cultural units (Riehla et al., 2014).

In Northern Algeria, the extinction of large mammal species (e.g., *Syncerus antiquus*) during the Mid-Holocene was correlated with regional climate aridity, likely due to

the competition with pastoralists and livestock for increasingly scarce water (Faith, 2014). Similarly, the evidence of the aridity (i.e., the termination of the African Humid Period) that provoked this extinction has been derived from the Sahara and its surroundings (deMenocal et al., 2000), which is several hundred kilometers away, leaving this assertion ambiguous and stimulating the search for new high-resolution paleoclimate records in the area.

In this study, we document the Mid-Holocene climate history in the Western Mediterranean by decadal-resolved stable carbon and oxygen isotopes analyses of two U/Th dated stalagmites from the Gueldaman GLD1 Cave of Northern Algeria. We compare the records with those established earlier in the Central and Eastern-Mediterranean basin. In addition, we describe archaeological deposits layers inside the cave whose ages have been reasonably well constrained due to new radiocarbon dating. Finally we test the links between cultural changes and climate anomalies with a particular emphasis on the 4200 yr BP event.

2 Samples and methods

2.1 Study site

Gueldaman GLD1 Cave is one of a series of karstic caves formed within the southeastward slope of the Adrar Gueldaman ridge, western part of Babor mountains in Northern Algeria (Kherbouche et al., 2014). It is located close to the large Soummam River, 5–6 km from the Akbou town, and approximately 65 km southern inland from the Western-Mediterranean Sea (36°26' N, 4°34' E, 507 m a.s.l.; Fig. 1). Gueldaman GLD1 is a relatively short cave (total extension of ~80 m) that developed in Jurassic limestone. The entrance, facing to the SE, is a semi-circular ~6 m large arch, leading to a dome-shaped ~10 m high and 6 m wide corridor which ends with the main chamber “Grande Salle” at a depth of 30–40 m. Previous archaeological excavation and field investigation suggest that Gueldaman GLD1 Cave has probably been open since at least ca. 7 ka ago (Kherbouche et al., 2014). The area is covered by a thin layer (<10 cm) of soil derived from the limestone bedrock, wind-blown silicate dust, and organic matter from local vegetation such as *Pistacia lentiscus*, *Quercus ilex*, *Buxus sempervirens*, typical Mediterranean *Garrigue* type plant assemblage (C3 dominated).

Local climate is Mediterranean semi-arid type, characterized by hot-dry summers and mild-wetter winters. From the ERA-interim reanalysis data between 1979 and 2013 (<http://apps.ecmwf.int/datasets/>) the annual total rainfall is 516 mm, and the annual mean temperature is 17.2 °C. Rainfall occurs rarely in the summer (37 mm) but relatively evenly through the autumn (155 mm), winter (178 mm) and spring (147 mm). Gueldaman GLD1 Cave is well ventilated with the outside atmosphere due to its larger opening and shorter extension. Hobo logger data at 10 min resolution from Novem-




Figure 2. U/Th dating of stalagmites GLD1-stm2 and GLD1-stm4 from the Gueldaman GLD1 Cave. U/Th dates and 2σ errors are shown next to sampling positions.

ber 2013 to April 2015 shows significant variations in cave temperature ranging from 13.7 to 19.5 °C. The relative humidity varies from 56 to 94 %. Carbon dioxide has not been measured, but it is likely to be close to the atmospheric value.

2.2 Stalagmites Analyses

Two stalagmites and three modern calcites samples were collected in 2012 and 2013 from the main chamber of Gueldaman GLD1 Cave. Both stalagmites were laid on the cave floor during collection. Stalagmite GLD1-stm2 is 350 mm long and 100–200 mm wide; stalagmite GLD1-stm4 is 203 mm long with a diameter of 50–120 mm (Fig. 2). They were halved and polished along the longitudinal axis. It was noticed from the sectioned sample that stalagmite GLD1-stm2 was broken at the depth of 5 mm from the top and covered by the calcite deposited at a certain time after that point. The top 5 mm was not analyzed in this study. Both stalagmites show well-marked laminae with several shifts in the drip apex of the lower parts (Fig. 2). Black bandings, with visible incorporations of charcoal particles, are found throughout both stalagmite profiles.

U/Th dating – Seventeen powder samples were drilled from the two stalagmites and dated by a multi-collector inductively coupled plasma mass spectrometer. The procedure to separate uranium and thorium was referred to in Edwards et al. (1987) and Cheng et al. (2013). The dating work was carried out at the University of Minnesota (USA) and the Xi’an Jiaotong University (China). One dating from the base part of stalagmite GLD1-stm2, for the exploration of preliminary age frame, was done at the Laboratoire des Sciences du Climat et de l’Environnement (LSCE, France). The U/Th dates were reported in years before 2000 AD. (Fig. 2, Table 1). The age model for both stalagmites was devel-

Table 1. U/Th dates from MC-ICP-MS analyses of stalagmites GLD1-stm2 and GLD1-stm4 from the Gueldaman GLD1 Cave. Analytical errors are 2σ of the mean. U decay constants: $\lambda_{238} = 1.55125 \times 10^{-10}$ (Jaffey et al., 1971) and $\lambda_{234} = 2.82206 \times 10^{-6}$ (Cheng et al., 2013). Th decay constant: $\lambda_{230} = 9.1705 \times 10^{-6}$ (Cheng et al., 2013). * $\delta^{234}\text{U} = ([^{234}\text{U}/^{238}\text{U}]_{\text{activity}} - 1) \times 1000$. ** $\delta^{234}\text{U}_{\text{initial}}$ was calculated based on ^{230}Th age (T), i.e., $\delta^{234}\text{U}_{\text{initial}} = \delta^{234}\text{U}_{\text{measured}} \times e^{\lambda_{234} \times T}$. Corrected ^{230}Th ages assume the initial $^{230}\text{Th}/^{232}\text{Th}$ atomic ratio of $4.4 \pm 2.2 \times 10^{-6}$. Those are the values for a material at secular equilibrium, with the bulk earth $^{232}\text{Th}/^{238}\text{U}$ value of 3.8. The errors are arbitrarily assumed to be 50 %. *** BP stands for “Before Present” where the “Present” is defined as the year 2000 AD.

Sample Number	^{238}U (ppb)	^{232}Th (ppt)	$^{230}\text{Th}/^{232}\text{Th}$ (atomic $\times 10^{-6}$)	$d^{234}\text{U}$ * (measured)	$^{230}\text{Th}/^{238}\text{U}$ (activity)	^{230}Th Age (yr) (uncorrected)	^{230}Th Age (yr) (corrected)	$d^{234}\text{U}_{\text{initial}}$ ** (corrected)	^{230}Th Age (yr BP)*** (corrected)	Laboratory
GLD1-stm2-7	169 ± 0.1	396 ± 0	136 ± 1	863 ± 2.3	0.105 ± 0.001	6297 ± 40	6228 ± 72	863 ± 2.3	6228 ± 72	LSCE
GLD1-stm2-36	154 ± 0.2	1922 ± 39	137 ± 3	910 ± 2.2	0.103 ± 0.000	6036 ± 29	5848 ± 136	925 ± 2.3	5835 ± 136	UM
GLD1-stm2-98	150 ± 0.2	69 ± 2	3077 ± 73	776 ± 2.4	0.086 ± 0.000	5374 ± 29	5366 ± 29	788 ± 2.5	5353 ± 29	UM
GLD1-stm2-180	152 ± 0.2	161 ± 3	1157 ± 25	644 ± 2.0	0.074 ± 0.000	5036 ± 30	5018 ± 33	653 ± 2.1	5005 ± 33	UM
GLD1-stm2-192	162 ± 0.1	182 ± 4	1158 ± 24	807 ± 1.7	0.079 ± 0.000	4858 ± 16	4840 ± 20	818 ± 1.7	4827 ± 20	UM
GLD1-stm2-213	175 ± 0.2	547 ± 11	390 ± 8	697 ± 2.0	0.074 ± 0.000	4841 ± 33	4788 ± 50	707 ± 2.1	4775 ± 50	UM
GLD1-stm2-286	169 ± 0.2	207 ± 4	980 ± 21	756 ± 1.8	0.073 ± 0.000	4601 ± 25	4581 ± 28	765 ± 1.9	4568 ± 28	UM
GLD1-stm2-320	195 ± 0.3	384 ± 8	575 ± 12	759 ± 2.2	0.069 ± 0.000	4321 ± 26	4288 ± 34	769 ± 2.2	4276 ± 34	Xi'an U
GLD1-stm2-340	194 ± 0.3	354 ± 7	612 ± 13	800 ± 2.9	0.068 ± 0.000	4172 ± 20	4142 ± 29	809 ± 2.9	4129 ± 29	UM
GLD1-stm4-10	105 ± 0.1	283 ± 6	415 ± 11	322 ± 1.5	0.068 ± 0.001	5734 ± 90	5675 ± 99	327 ± 1.6	5662 ± 99	UM
GLD1-stm4-24	126 ± 0.1	983 ± 20	135 ± 3	347 ± 1.9	0.064 ± 0.001	5311 ± 49	5143 ± 128	352 ± 1.9	5131 ± 128	Xi'an U
GLD1-stm4-30	106 ± 0.1	1362 ± 27	80 ± 2	298 ± 1.9	0.062 ± 0.001	5321 ± 57	5035 ± 210	302 ± 1.9	5023 ± 210	Xi'an U
GLD1-stm4-47	96 ± 0.1	1104 ± 22	88 ± 2	290 ± 1.9	0.062 ± 0.001	5341 ± 63	5082 ± 194	294 ± 1.9	5070 ± 194	Xi'an U
GLD1-stm4-70	224 ± 0.4	749 ± 15	254 ± 5	334 ± 2.5	0.051 ± 0.000	4282 ± 27	4210 ± 58	338 ± 2.6	4197 ± 58	UM
GLD1-stm4-113	155 ± 0.2	130 ± 3	910 ± 20	363 ± 2.1	0.046 ± 0.000	3761 ± 29	3743 ± 32	367 ± 2.1	3730 ± 32	UM
GLD1-stm4-152	180 ± 0.3	582 ± 12	226 ± 5	373 ± 2.0	0.045 ± 0.000	3590 ± 25	3521 ± 54	376 ± 2.0	3508 ± 54	UM
GLD1-stm4-195	175 ± 0.1	929 ± 19	131 ± 4	369 ± 1.7	0.042 ± 0.001	3403 ± 74	3290 ± 108	372 ± 1.8	3277 ± 108	UM




Figure 3. Age models of stalagmites GLD1-stm2 and GLD1-stm4 from the Gueldaman GLD1 Cave. The age models were calculated using the StalAge program (Scholz and Hoffmann, 2011). Note that the U/Th date of sample GLD1-stm4-47 was detected as a major outlier and not used in the final age model of stalagmite GLD1-stm4. The 2σ analytical uncertainty of each U/Th date (dot) is represented by the error bars, whereas the 95 % uncertainty assessed from the model simulation is represented by thin curves.

oped using the StalAge program (Scholz and Hoffmann, 2011) where a linear interpolation between depth and age is made through each progressive triplet of adjacent U/Th dates (Fig. 3). This procedure provides a quantitative estimate of age uncertainty continuously along the record despite having analytical constraints only at locations where the U/Th dates exist. Stalagmite growth rates were calculated based on the StalAge age model (Fig. 4).

Stable isotopes – Four hundred and thirty samples were drilled every 1 to 2 mm along the stalagmite central growth axis (Fig. 2). Stable carbon and oxygen isotopes compositions of both stalagmites and modern calcites were measured using a VG-OPTIMA mass spectrometer at the LSCE. For each analysis, 60 to 80 μg calcite powder is reacted with phosphoric acid at 90 °C, and the resultant CO_2 is measured relative to a reference gas that has been calibrated against a series of isotopic standards. Duplicates were run every 10 to 20 samples to check replicability. All values are reported in ‰ relative to the V-PDB (Fig. 4). The error is 0.08 ‰ for $\delta^{18}\text{O}$ and 0.05 ‰ for $\delta^{13}\text{C}$.

2.3 Archaeological analyses

Archaeological excavations were carried out at two sectors S2 and S3 inside the Gueldaman GLD1 Cave during the 2010–2012 campaign (Fig. 1). This work consisted mainly in collecting, identifying, and referencing the archaeological materials found in stratigraphic layers (refer to Kherbouche et al., 2014 for details). More than 7000 anthropogenic remains were collected, consisting mainly of faunal remains, ceramic, and lithic and bone tools. Besides, all sediments were water screened through 1.5 and 4 mm mesh and subjected systematically to flotation with collection in a 250 μm mesh yielding a huge amount of charcoals. Initial radiocarbon dating of upper stratigraphic sequences from S2 and S3 gave the median ages ranging from ca. 6800 to 1500 cal yr BP (Kherbouche et al., 2014). In order to refine the chronology of these deposits, in this study, six new charcoal samples were collected from the key archaeological layers in excavation area MN 47/48 of S2. These samples were dated using the AMS radiocarbon method at the CEA Saclay (France).

Table 2. Radiocarbon dates from AMS analyses of charcoals from excavation sector S2 inside the Gueldaman GLD1 Cave. * Kherbouche et al. (2014). Ages are reported in years before 2000 AD.

Depth Z (cm)	Square	Lab no. (SacA#)	Material	^{14}C Age ($\pm 2\sigma$; yr)	Median age (yr)	Cal. interval (2σ ; yr)	Note
60	M48	39408	Charcoal	1600 ± 30	1482	1385–1604	This study
65	N48	29731	Charcoal	1610 ± 25	1484	1415–1547	*
84	N48	39410	Charcoal	4020 ± 30	4495	4411–4785	This study
86	N48	39411	Charcoal	3975 ± 30	4416	4290–4569	This study
91	M48	39409	Charcoal	3945 ± 30	4403	4244–4522	This study
108	N47	36982	Charcoal	4355 ± 30	4918	4851–5032	This study
124	L48	23883	Charcoal	5250 ± 35	6003	5924–6178	*
132	L48	23884	Charcoal	4260 ± 30	6025	5933–6178	*
147	M47	36981	Charcoal	6120 ± 35	7002	6907–7157	This study

Detailed procedures of the chemical preparation and the dating in the lab were referred to Cottereau et al. (2007). The dates were calibrated using the IntCal13 data set (Reimer et al., 2013) and reported in years before 2000 AD (Table 2).

3 Results

3.1 Stalagmites U/Th dates and growth rates

The uranium contents of measured stalagmites samples are relatively high ranging from 95 to 225 ppb (Table 1). The 2 sigma U/Th errors vary from 20 to 210 years with an average of 77 years (1.6%). The U/Th date (5070 ± 194 yr BP) of sample GLD1-stm4-47 was detected as a major outlier by the StalAge program, thus, it was not used to calculate the final age model. Calculated StalAge age model for stalagmite GLD1-stm4 shows large errors up to 500 years during ca. 4900–4200 yr BP (Fig. 3). This may partly be due to relatively large errors of adjacent U-series dates and/or inappropriate hypotheses applied in the algorithm. Based on individual StalAge age model, stalagmite GLD1-stm2 grew from ca. 6200 to 4000 yr BP (4100 yr BP if excluding the top 5 mm), whereas stalagmite GLD1-stm4 grew from ca. 5800 to 3200 yr BP (Fig. 3).

Stalagmite GLD1-stm2 shows high and variable growth rates (mean = $180 \mu\text{m yr}^{-1}$) with higher values $\sim 400 \mu\text{m yr}^{-1}$ at ca. 4800–4600 yr BP; whereas stalagmite GLD1-stm4 shows relatively lower and less variable growth rates (mean = $120 \mu\text{m yr}^{-1}$) with higher values $\sim 200 \mu\text{m yr}^{-1}$ at ca. 3800–3200 yr BP (Fig. 4).

3.2 Stable carbon and oxygen isotopes

The isotopic compositions of modern calcite vary from -5.40 to -5.56 ‰ for the $\delta^{18}\text{O}$ and from -8.43 to -10.34 ‰ for the $\delta^{13}\text{C}$. The $\delta^{18}\text{O}$ values from stalagmites GLD1-stm2 and GLD1-stm4 range from -7.8 to -2.8 ‰ and from -7.3 to -0.6 ‰, respectively; the $\delta^{13}\text{C}$ values range from -10.6 to -3.3 ‰ and from -11.9 to -0.6 ‰,




Figure 4. The $\delta^{18}\text{O}$, $\delta^{13}\text{C}$ and growth rate of stalagmites GLD1-stm2 and GLD1-stm4 from the Gueldaman GLD1 Cave. U/Th dates with 2σ errors are presented at the top. The isotopic ranges of modern calcites are also shown on the left (rectangles). Growth rates are calculated from the StalAge age model. Note that the extraordinarily high growth at ca. 4600 yr BP in stalagmite GLD1-stm2 and ca. 3800 yr BP in stalagmite GLD1-stm4 are likely attributed to artificial simulations by the StalAge program and thus are not fully discussed in terms of climate in the text.

respectively. The $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ significantly correlate in both stalagmites: $R = 0.87$, $P < 0.01$ for GLD1-stm2 and $R = 0.92$, $P < 0.01$ for GLD1-stm4. Albeit the different amplitudes, the isotopic profiles of the two stalagmites show similarities during their common development of ca. 5800–

4000 yr BP: relatively elevated isotope values are found at ca. 5700–5400, ca. 5200, and ca. 4500 yr BP (Fig. 4). Two other isotopically enriched periods in stalagmite GLD1-stm2 are found at ca. 6200 and ca. 4900 yr BP (Fig. 4). There is a common isotopic enrichment trend since ca. 4800–4600 yr BP (depending on individual age model; abrupt in stalagmite GLD1-stm4 whereas more gradual in GLD1-stm2). Toward the end of this trend, the most prominent anomaly occurs in stalagmite GLD1-stm4 at ca. 4400–3800 yr BP during which the $\delta^{18}\text{O}$ values are enriched by approximately 3.5‰ relative to the background values of that time as well as the modern calcite values for a period of ~ 500 years (Fig. 4). Specifically within this anomalous period, there is a mild isotopic depletion at ca. 4200–4000 yr BP, blanketed by two significant enrichments at ca. 4400–4200 yr BP and ca. 4000–3800 yr BP (Fig. 4). The last part, ca. 3800–3200 yr BP, of GLD1-stm4, is characterized by a $\delta^{18}\text{O}$ recovery of about -3 ‰, synchronous with increased growth rates (Fig. 4).

3.3 Anthropogenic deposits and ^{14}C dates

Excavations inside Gueldaman GLD1 Cave revealed a large variety of archeological remains and, among them, are numerous precious macro charcoals that have been used for establishing the chronology of the deposits. In the $\sim 7\text{ m}^2$ total excavated area of S2, more than 7000 archaeological objects were identified and consisted of faunal remains, lithic artifacts and grinding equipment, potteries, bone tools, ornaments, and ochre. In addition, a fragment of a human mandible and two isolated teeth were found during the excavation 2010–2012 in Gueldaman GLD1 Cave (Kherbouche et al., 2014). These deposits belong mainly to the Neolithic; only the top level of the sequence contains potsherds of the historic period. In the lower Neolithic levels, identified domestic species (i.e. sheep and goats) represented $\sim 25\%$ of total faunal assemblages ($N = 2378$) suggesting a partly pastoral based economy. The potteries ($N = 825$) are mostly related to cooking vessels of 25–40 cm rim diameter. Hundreds of black charcoals ($> 1\text{ cm}$) were found and always associated with ceramic concentrations suggesting evidence of cooking activities.

Determined radiocarbon dates give the median ages of the sequence between 7002 and 1482 cal yr BP, with their 2σ -error intervals varying from 132 to 374 years (Table 2). These dates provide a first chronology for the archaeological deposits excavated from sector S2 (Fig. 6; Kherbouche et al., 2014): anthropogenic remains (i.e. charcoals, bones, teeth and potteries) are numerous during ca. 7002–6003 cal yr BP (depths of ~ 150 –120 cm), decreased at ca. 6003–4918 cal yr BP (depths of ~ 120 –105 cm), most abundant in the period of ca. 4918–4403 cal yr BP (depths of 105–75 cm), significantly diminished during the long interval of ca. 4403–1484 cal yr BP (depths of 75–60 cm), and finally, numerous again from ca. 1484 cal yr BP (depths of ~ 60 –50 cm). With an overall decrease in archeological materials,




Figure 5. Comparison of high-resolution Mid-Holocene stalagmite $\delta^{18}\text{O}$ records across the Mediterranean basin. From the top to bottom are stalagmite records from the Gueldaman GLD1 Cave in Northern Algeria of the Western-Mediterranean basin (this study), the Corchia Cave (Zanchetta et al., 2014) and the Renella Cave (Drysedale et al., 2006) in Central Italy of the Central Mediterranean basin, and the Soreq Cave (Bar-Matthews and Ayalon, 2011; Kaufman et al., 1998) in Israel of Eastern-Mediterranean basin. Different stalagmites from each area are represented in distinct colors. U/Th dates with 2σ errors are shown at the top of each curve. Ages are reported in years before 2000 AD.

there are two levels clearly marked by their poverty in charcoal and pottery during the periods of ca. 6003–4918 and ca. 4403–1484 cal yr BP (Fig. 6).

4 Discussions

4.1 Climatic significance of stalagmites proxies

Under isotopic equilibrium precipitation, stalagmite calcite $\delta^{18}\text{O}$ depends mainly on the temperature of calcite-water fractionation and on the $\delta^{18}\text{O}$ of drip water that is controlled by local rainfall $\delta^{18}\text{O}$ (Genty et al., 2014; Lachniet, 2009). Observations from the IAEA network show that the rainfall $\delta^{18}\text{O}$ at many Mediterranean stations (including one in Algiers, Algeria) are partly controlled by the amount of rainfall (IAEA, 2005), which is coherent with previous studies that stalagmite $\delta^{18}\text{O}$ records from the Mediterranean regions were interpreted to primarily reflect changes in rainfall amount (inversely correlated; e.g., Bar-Matthews and Ay-




Figure 6. Radiocarbon dating of anthropogenic deposits layers in excavation sector S2 inside the Gueldaman GLD1 Cave. From the left to right are ^{14}C dates of charcoal samples, anthropogenic deposit distribution, and a photo at depth across $\sim 75\text{--}88$ cm showing a transition of layer from rich to rare anthropogenic deposits. The gray color highlights two phases with diminished anthropogenic remains (especially pottery and charcoal) at ca. 4403–1484 cal yr BP (depths of $\sim 75\text{--}60$ cm) and ca. 6003–4918 cal yr BP (depths of $\sim 120\text{--}105$ cm).

alon, 2011; Bar-Matthews et al., 2003, 1997; Drysdale et al., 2004, 2006; Zanchetta et al., 2014). Therefore, higher stalagmite $\delta^{18}\text{O}$ values are expected during periods of decrease in rainfall (drier). The temperature effect on calcite-water fractionation, on the other hand, is partly counteracted by the condensation temperature effect on rainfall $\delta^{18}\text{O}$ (Drysdale et al., 2006). We notice that this interpretation may particularly hold true for the present study because the regional temperature seems to have been relatively constant since the Mid-Holocene (Martrat et al., 2004).

Stalagmite $\delta^{13}\text{C}$ variations have several potential causes, the most likely, considering the studied location and time interval, being variations in soil CO_2 input and water flow rate (Genty et al., 2001a; McDermott, 2004). Despite the fact that soil biogenic CO_2 production varies according to both temperature and moisture level, moisture is likely to be a major controlling factor due to low temperature variability of the considered time interval and limited water availability under semi-arid climates. Moisture also influences water flow rate and thus the CO_2 loss during the prior calcite precipitation (Fairchild et al., 2000). Longer residence time due to lower flow rate, under diminished moisture condition, enhances CO_2 loss and preferential ^{12}C removal from solution causing enrichments in stalagmite $\delta^{13}\text{C}$ (Johnson et al., 2006; Mickler et al., 2004). Eventually, atmospheric rainfall largely determines the moisture level and controls the $\delta^{13}\text{C}$ variations. Therefore, the significant correlation of stalagmite $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ may suggest a common control of rainfall.

The rainfall signal imprinted in the Gueldaman stalagmite isotopes (inversely correlated) is probably amplified by

two other processes – evaporation and disequilibrium isotopic fractionation (Mickler et al., 2006) that are very likely to have occurred at the Gueldaman GLD1 Cave due to its large entrance. Longer residence time during drier (lower rainfall) periods would allow extended evaporative and non-equilibrium fractionations, which drives stalagmite isotopes further higher (Mickler et al., 2004). It has recently been observed that evaporation in semi-arid caves could cause 4–5 ‰ $\delta^{18}\text{O}$ enrichments of a wide range of drip waters (Cuthbert et al., 2014). The Hendy test (i.e. studying the isotopic variation in contemporaneous laminae; Hendy, 1971) carried out at three different depths in stalagmite GLD1-stm2 show that the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ significantly correlate and simultaneously increase by up to ~ 1 ‰ from the apex to the edge probably due to the presence of progressive kinetic fractionations when feeding waters flow outwards from the apex, suggesting that the stalagmite formed out of isotopic equilibrium (see the Supplement). These two processes may partly account for the significant correlation of the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ profiles in both stalagmites ($R = 0.87$ for GLD1-stm2; $R = 0.92$ for GLD1-stm4). Comparisons of two stalagmites show different amplitudes in their isotopic profiles. This might be due to the following: (1) having been fed by reservoirs that smooth rainfall signal differently, and (2) having been suffered from variable evaporative and kinetic enrichments associated with different recharge features. Despite this discrepancy the isotopic profiles of two stalagmites broadly show similar patterns. Consequently, synchronous $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ variations in the two stalagmites can be interpreted in terms of humidity change. A prolonged severe drought is inferred from the most ele-

vated isotopes values during ca. 4400–3800 yr BP together with drier periods from higher values at ca. 6200, ca. 5700–5400, ca. 5200, and ca. 4500 yr BP (Fig. 4), while wetter periods are suggested from depleted isotopes at ca. 4600–5100 and ca. 5300 yr BP.

Moreover, stalagmite growth phase has long been regarded as an environmental indicator (e.g., Genty et al., 2001b; Stoll et al., 2013) and could be used to complement and/or test the climatic interpretation from isotope records. Water availability is an essential factor for stalagmite growth in arid-to-semi-arid areas, as shown by Vaks et al. (2013) who correlated growth periods well with periods of effective rainfall regimes. The growth cessation of stalagmite GLD1-stm2 at ca. 4000 yr BP may suggest a phase of increased aridity, which is consistent with the dryness inferred from elevated isotopes (Fig. 4). However, the continuous growth of stalagmite GLD1-stm4 at that time suggests that the growth cessation might also have been caused by shifts in dripping position under deteriorating climates (Fairchild and Baker, 2012). Moreover, fast stalagmite growths together with wide diameters usually are associated with high drip rates suggesting humid conditions. A wet period ca. 4800–4600 yr BP is indicated by the high growth rates of stalagmite GLD1-stm2, which is broadly in line with the humid period inferred from depleted isotopes (Fig. 4). Another wetter period ca. 3800–3200 yr BP is suggested by faster growths and relatively depleted isotopes of stalagmite GLD1-stm4 (Fig. 4). The discrepancy in the growth rate profiles of the two stalagmites is possibly due in part to (1) the lack of substantial U/Th dating especially for stalagmite GLD1-stm4 between 5023 and 4197 yr BP, and/or (2) site-specific processes due to different reservoirs play a key role in controlling stalagmite growth.

Finally, the modern calcite isotopic values fall in the range of two stalagmite records (Fig. 4), which suggests that the current humidity condition in Northern Algeria seems to be within the range of its Mid-Holocene variability. The average $\delta^{18}\text{O}$ value during ca. 4400–3800 yr BP in stalagmite GLD1-stm4 is more enriched by about 3.5‰ relative to the modern values (Fig. 4), therefore, the 4400–3800 yr BP climate anomaly may be considered analogous to end numbers of the most recent and ongoing drying.

4.2 Mid-Holocene climate anomalies across the Mediterranean basin and their dynamic implications

High-resolution absolute-dated Mid-Holocene climate records are rare in the Western-Mediterranean basin, however, there are number of paleoenvironment studies using sediment cores that document large oscillations in vegetation ecology and provide clues of climate change during the Mid-Holocene. A drying trend from ca. 4600 cal yr BP onwards was inferred, based on the decreasing pollen ratio of deciduous broad-leaf vs. evergreen sclerophyllous taxa at Capestang in the Mediterranean, Southern France (Jalut et al., 2000). At a nearby site in Northeastern Spain, the most

arid Mid-Holocene condition at ca. 4800–4000 cal yr BP was interpreted using maximum salinity values, more positive organic carbon isotope values, and decreased algal productivity in Estanya Lake (Morrellon et al., 2009). In the Mediterranean, Southern Spain, desertification phases at ca. 5200 and ca. 4100 cal yr BP were inferred using multiple paleoecological indicators including pollen, microcharcoal, spores of terrestrial plants, fungi, non-siliceous algae, and other microfossils in Siles Lake (Carrión, 2002). Similar environment changes have also been observed in the Central Mediterranean basin, as shown by a synthesis study of lacustrine palynological data which suggested a dryness peaking at ca. 4000 cal yr BP (Sadori et al., 2011). Evaluated carbonate oxygen isotopes from Lake Shkodra were argued to be an indicator of dryness at ca. 4100–4000 cal yr BP (Zanchetta et al., 2012b). Increasing aridity at ca. 5000–4000 cal yr BP was suggested to explain the increases in non-tree pollen percentage and micro charcoal content in the Lago di Pergusa Lake, Sicily (Roberts et al., 2011; Sadori and Giardini, 2007; Sadori and Narcisi, 2001). Closer to our site, a study at Preola Lake, Eastern Sicily, documented a significant low stand lake level at ca. 4500–4000 cal yr BP suggesting extreme aridity (Magny et al., 2011). Moreover, six tephra layers were carefully studied to correlate climate anomalies at ca. 4500–3800 cal yr BP from different archives in the central Mediterranean (Zanchetta et al., 2012a). In the east, Finné et al. (2011) reviewed the climate history of the Eastern Mediterranean over the last 6000 years and concluded with much evidence of drying conditions at ca. 4600–3800 yr BP. Although the sampling and dating resolutions in most of the above studies are low, they are in good agreement with the present study regarding the 5200 yr BP dry event, the drying trend from 4800–4600 yr BP onward, and the 4400–3800 yr BP drought.

Recently, high frequency Mid-Holocene climate change has been well documented by multiple detailed speleothem studies from the Central and Eastern-Mediterranean basin (Fig. 5). Drysdale et al. (2006) demonstrated a severe drought peaked at 4400–3800 yr BP through multiproxy analyses on a U/Th-dated flowstone from Renella Cave, Central Italy. This finding is supported by following work at nearby Corchia Cave. Zanchetta et al. (2007) established a high-resolution speleothem isotope record for the entire Holocene and interpreted it as an indicator of rainfall change, in which a drier period persisting from ca. 4800 to 3800 yr BP could be inferred from relatively elevated $\delta^{18}\text{O}$ values. Elemental analysis on the same stalagmite by Regattieri et al. (2014) presented more clear signals of dryness peaked at ca. 4700–4200 yr BP. In the Eastern-Mediterranean basin, Bar-Matthews and Ayalon (2011) explicitly discussed the Mid-Holocene climate change by high-resolution dating and isotopic analyses on speleothems from Soreq Cave in Israel; they identified multiple dry events at 6250–6180, 5700–5600 and 5250–5170 yr BP as well as a long drying trend since ca. 4700 yr BP peaked at 4200–4050 yr BP. Zanchetta

Table 3. Summary of the features of climate and human activity in different climate periods and occupation phases.

Climate period	Age (^{230}Th yr)	Climate condition	Occupation phase	Age (^{14}C cal yr)	Human activity
–	–	–	0	~ 7002–6003	Permanent and intensive occupation
1	~ 6200–5100	Wet & oscillatory	1	~ 6003–4918	Permanent but less intensive occupation
2	~ 5100–4400	Wettest, ending with a dramatic shift in the last ~ 200 years	2	~ 4918–4403	Permanent and most intensive occupation
3	~ 4400–3800	Extremely dry	3	~ 4403–1484	Abandonment of the cave/occasional visit
4	~ 3800–3200	Relatively wetter	4	~ 1484–	Re-occupation of the cave

et al. (2014) carefully compared the speleothem isotope records from Corchia Cave and Soreq Cave in the Central and Eastern-Mediterranean basin, and found two coeval dry events at ca. 5600 and ca. 5200 yr BP from the comparable $\delta^{18}\text{O}$ enrichments in two speleothems. Detailed comparisons of these speleothem records with our new records from Guedaman GLD1 Cave reveal many consistencies. In particular, periods with elevated $\delta^{18}\text{O}$ values at ca. 5700–5400, 5200 and 4400–3800 yr BP observed in Guedaman stalagmites are all identifiable in the speleothems from Corchia, Renella and Soreq Cave (Fig. 5), suggesting that anomalous dryness at these periods synchronously developed across the Mediterranean basin. These observations indicate that climates in the Mediterranean basin might have been under an identical regional scale atmospheric regime during the Mid-Holocene.

It has been suggested that climate change in mid-latitude Europe and the Mediterranean might arise from a perturbation of the Westerlies from a high-latitude trigger (i.e. the North Atlantic; Bond et al., 2001; Drysdale et al., 2006; Zanchetta et al., 2014) or from dynamics within the tropics (Booth et al., 2005; Hoerling and Kumar, 2003). The three dry periods in the Mediterranean are broadly in phase with the ice rafting events in the subpolar North Atlantic (Bond et al., 2001), which suggests some links with the North Atlantic circulation. Based on the coincidence with the elevated wind strength in Iceland (Jackson et al., 2005), Zanchetta et al. (2014) argued that the dry events at ca. 5700–5400 and ca. 5200 yr BP might be caused by reduction of vapor advection into the Mediterranean, due to the intensification and northward displacement of the North Atlantic Westerlies. However, lacking evidence of strengthened wind in the fourth millennium BP argues for a different forcing of the 4400–3800 yr BP drought. The considerably lower amplitude of the Bond ice rafting event at ca. 4200 yr BP than at the fifth millennium BP also indicates a varied ocean-atmosphere circulation state. The modern mid-latitude droughts (1998–2002) have been linked to the increased warmth in equatorial oceans (Booth et al., 2005). During this event, SST changes lead to persistent high pressure over the Northern Hemisphere's mid-latitudes, caus-

ing widespread synchronous drought (Hoerling and Kumar, 2003). However, the challenge in applying the dynamics under the 1988–2002 drought towards an understanding of the 4400–3800 yr BP climate anomaly is that while the mechanism operates effectively on short timescales, it has never been tested as to whether they could produce an anomalous climate mode for several centuries (Berkelhammer et al., 2013). General circulation model simulations that begin with realistic boundary conditions and are perturbed with a variety of forcings have been successfully undertaken to understand potential mechanisms that lead to the 8200 yr BP event (Tindall and Valdes, 2011). Similar efforts would be a useful starting point to produce hypotheses for the dynamical underpinnings of the 4200 yr BP event.

4.3 Possible relations between climate anomaly and cultural change

A regional drought ca. 4200 yr BP has been widely linked to ancient cultural changes in the Eastern Mediterranean and Asia (Staubwasser and Weiss, 2006), though, in many cases, climatic inferences have been derived from sites that are distant to these human settlements. For instance, evidence of reduced precipitation from elevated $\delta^{18}\text{O}$ of Soreq stalagmites (Israel) and increased dust input into the Gulf of Oman sediment core has been suggested to contribute to the collapse of the Akkadian empire in Mesopotamia (Bar-Matthews and Ayalon, 2011; Cullen et al., 2000; Weiss et al., 1993). Similarly, a dry period inferred from reduced discharge of the Indus river and elevated $\delta^{18}\text{O}$ of a Northeast-Indian stalagmite has been linked with the Indus Valley deurbanization (Berkelhammer et al., 2013; Staubwasser et al., 2003; Staubwasser and Weiss, 2006). A recent study in the ancient Near East, however, revealed that the regional impact of the drought on ancient civilizations, being influenced by geographic factors and human technology, were highly diverse even within spatially limited cultural units (Riehla et al., 2014). This highlights the need for caution when linking evidence of human activities from a site to evidence of climate oscillations from another one.




Figure 7. Comparison between evidence of ancient human occupation and of past climate change from the Gueldaman GLD1 Cave. The definition of occupation phases 0–4 and climate periods 1–4 are referred to the text.

The present study in the Gueldaman GLD1 Cave provides an opportunity to test climate–culture relations by comparing in situ archaeological sequences and high-resolution paleoclimate records, thereby avoiding the uncertainty of inter-site correlation arising from complex spatial heterogeneity in climate and demography.

To facilitate the comparison, stalagmite-inferred climate changes at the cave site during ca. 6200–3200 yr BP are separated into four periods 1–4 (Table 3):

- Period 1 (~ 6200–5100 yr BP): wet, superimposed by several centennial-scale drier events;
- Period 2 (~ 5100–4400 yr BP): wettest, ending with a ~ 200-year long shift from the wettest to extreme dry conditions;
- Period 3 (~ 4400–3800 yr BP): a drought-like climatic anomaly;
- Period 4 (~ 3800–3200 yr BP): relatively wetter.

In parallel, from the abundance of archaeological remains (especially bone, charcoal and pottery; Fig. 6), the temporal evolution of past cave occupations can be separated into five phases 0–4 (Table 3):

- Phase 0 (~ 7002–6003 cal yr BP): permanent and intensive occupation;
- Phase 1 (~ 6003–4918 cal yr BP): permanent but less intensive occupation;

- Phase 2 (~ 4918–4403 cal yr BP): permanent and most intensive occupation;
- Phase 3 (~ 4403–1484 cal yr BP): abandonment of the cave/occasional visits;
- Phase 4 (~ 1484 cal yr BP): re-occupation of the cave.

Correlations can be identified when comparing the climatic and archaeological records, though this does not necessarily mean that occupation of the cave depends merely on climate (Table 3; Fig. 7). When the climate was wet and variable ca. 6200–5100 yr BP (Period 1), the Gueldaman GLD1 Cave preserved a few bones and rare charcoals and potteries (Phase 1; Fig. 7). When the climate was wettest ca. 5100–4400 yr BP (Period 2), the most abundance of bones, charcoals and potteries suggests a permanent and more intensive occupation of the cave (Phase 2; Fig. 7). More striking is the drought-like climatic anomaly that has been establishing ca. 4400–3800 yr BP (Period 3), from which the cave was abandoned for ca. 3000 years (indicated by a dramatic decrease in anthropogenic remains, especially charcoal and pottery; Phase 3; Fig. 7). The rarer bones seen in this period imply that the cave might have been occasionally visited until its re-occupation at ca. 1484 cal yr BP (Fig. 7). These observations argue for links between climate and settlement activity especially during the 4200 yr BP climate anomaly. Water availability was likely crucial to maintain the Neolithic community at Gueldaman, Northern Algeria and the prolonged severe drought ca. 4400–3800 yr BP might have played a role in triggering the settlement abandonment, indicating that the

pastoral economy may not be as resistant, as commonly assumed, to climate anomaly in semi-arid area.

Moreover, the sole piece of the bone from large ungulate (supposed to be elephant or rhinoceros) found at the depth of ~ 110 cm (Kherbouche et al., 2014) was anchored by two calibrated ^{14}C dates from the present study between 6003 and 4913 cal yr BP, which is in line with the latest survival of large mammal species (e.g., *S. antiquus*) at the proximate sites of Northern Algeria during the Mid-Holocene (Faith, 2014 and references therein). The extinction of the Mid-Holocene large mammal in Northern Algeria was attributed to the competition with pastoralists and livestock for increasingly scarce water, corresponding with an abrupt climatic shift toward extreme aridity in the Sahara region ca. 5500 cal yr BP (i.e. the end of the Humid Africa Period (deMenocal et al., 2000; Faith, 2014). Recently, the timing of this climatic transition was refined to ca. 4900 yr BP \pm 200 yrs (McGee et al., 2013). In addition, a paleoenvironmental study in the Sahara revealed that the Mid-Holocene deteriorations of terrestrial ecosystem and climate culminated at ca. 4200–3900 cal yr BP (e.g., Kröpelin et al., 2006). Therefore, it is more likely based on evidence from the Gueldaman GLD1 Cave and the proximate sites (Faith, 2014) that extinction of large mammal/ungulate in Northern Algeria occurred during the prolonged drought ca. 4400–3800 yr BP.

5 Conclusions

It is increasingly clear based on a growing number of records spanning across much of mid-to-low latitudes, Northern Europe, and the Atlantic ocean that there was a significant large-scale climate anomaly at around 4200 yr BP (Booth et al., 2005). The 4200 yr BP aridity that had been suggested to affect the Early Bronze Age populations from the Aegean to ancient Near East was recently characterized by high-resolution speleothem records from the Central and Eastern-Mediterranean basin (Bar-Matthews and Ayalon, 2011; Drysdale et al., 2006; Zanchetta et al., 2014). The new record presented here from the Gueldaman GLD1 Cave in Northern Algeria provides increased evidence of a prolonged severe drought ca. 4400–3800 yr BP, which suggests that the Mid-Holocene dryness spread to the Western Mediterranean of Northern Africa.

Radiocarbon dating made on charcoals constrains reasonably well the age of archaeological deposits excavated inside the cave (Kherbouche et al., 2014) and reveals significant changes in human occupation during the last ca. 7000 years. Comparison of the stalagmite record with in situ archaeological sequence suggests synchronicity between climate and settlement activity. Relatively wet periods coincide with the periods of intensive human occupation. Particularly, the timing of the prolonged drought at ca. 4400–3800 yr BP blanket the onset of the cave abandonment event shortly after ca. 4403 cal yr BP, which argues a possible role of climate

anomaly in this societal disruption. Further work on pollen-based reconstruction of environment change from the excavation sequence and on refinement of the chronology of transitions between different occupation phases would potentially uncover the intrinsic relations among climate, environment and settlement. It is suggested that the methodology and the findings from the present study at the Gueldaman GLD1 Cave be applied and tested at other sites.

The Supplement related to this article is available online at doi:10.5194/cp-12-1-2016-supplement.

Acknowledgements. Radiocarbon dating was analyzed by Jean-Pierre Dumoulin at ARTEMIS (LMC14, Saclay). We thank Edwige Pons-Branchu, Monique Pierre for the U/Th dating of the base part of stalagmite GLD1-stm2 during the earlier stage of this study. We also thank Lijuan Sha at the Xi'an Jiaotong University for assistance with the U/Th dating. We appreciate Russell N. Drysdale, Giuliano Zanchetta and Miryam Bar-Matthews for sharing the isotopic data shown in Fig. 5. Thanks to Cecilia Garrec for editing assistance. We are thankful to Giuliano Zanchetta and an anonymous referee for their constructive comments which improved this paper. This work was funded by the CNRS INSU program PALEOMEX-ISOMEX, the NSFC grant 41230524 and the CSC scholarship.

Edited by: N. C. Nebout

References

- Bar-Matthews, M. and Ayalon, A.: Mid-Holocene climate variations revealed by high-resolution speleothem records from Soreq Cave, Israel and their correlation with cultural changes, *Holocene*, 21, 163–171, 2011.
- Bar-Matthews, M., Ayalon, A., and Kaufman, A.: Late Quaternary Paleoclimate in the Eastern Mediterranean Region from Stable Isotope Analysis of Speleothems at Soreq Cave, Israel, *Quat. Res.*, 47, 155–168, 1997.
- Bar-Matthews, M., Ayalon, A., Gilmour, M., Matthews, A., and Hawkesworth, C. J.: Sea–land oxygen isotopic relationships from planktonic foraminifera and speleothems in the Eastern Mediterranean region and their implication for paleorainfall during interglacial intervals, *Geochim. Cosmochim. Ac.*, 67, 3181–3199, 2003.
- Berkehamer, M., Sinha, A., Stott, L., Cheng, H., Pausata, F. S. R., and Yoshimura, K.: An abrupt shift in the Indian Monsoon 4000 years ago, in: *Climates, Landscapes, and Civilizations*, edited by: Giosan, L., Fuller, D. Q., Nicoll, K., Flad, R. K., and Clift, P. D., American Geophysical Union, Washington, DC, 75–87, 2013.
- Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M. N., Showers, W., Hoffmann, S., Lotti-Bond, R., Hajdas, I., and Bonani, G.: Persistent Solar Influence on North Atlantic Climate During the Holocene, *Science*, 294, 2130–2136, 2001.

- Booth, R. K., Jackson, S. T., Forman, S. L., Kutzbach, J. E., E.A. Bettis, I., Kreig, J., and Wright, D. K.: A severe centennial-scale drought in midcontinental North America 4200 years ago and apparent global linkages, *Holocene*, 15, 321–328, 2005.
- Carrión, J. S.: Patterns and processes of Late Quaternary environmental change in a montane region of southwestern Europe, *Quat. Sci. Rev.*, 21, 2047–2066, 2002.
- Cheng, H., Edwards, L. R., Shen, C. C., Polyak, V. J., Asmerom, Y., Woodhead, J., Hellstrom, J., Wang, Y., Kong, X., Spötl, C., Wang, X., and Alexander Jr, E. C.: Improvements in ^{230}Th dating, ^{230}Th and ^{234}U half-life values, and U-Th isotopic measurements by multi-collector inductively coupled plasma mass spectrometry, *Earth Planet. Sci. Lett.*, 371–372, 82–91, 2013.
- Coombes, P. and Barber, K.: Environmental determinism in Holocene research: causality or coincidence?, *Area*, 37, 303–311, 2005.
- Cottreau, E., Arnold, M., Moreau, C., Baque, D., Bavay, D., Caffy, I., Comby, C., Dumoulin, J.-P., Hain, S., Perron, M., Salomon, J., and Setti, V.: Artemis, the New 14C AMS14 in Saclay, France, *Radiocarbon*, 49, 291–299, 2007.
- Cullen, H. M., deMenocal, P. B., Hemming, S., Hemming, G., Brown, F. H., Guilderson, T., and Sirocko, F.: Climate change and the collapse of the Akkadian empire: Evidence from the deep sea, *Geology*, 28, 379–382, 2000.
- Cuthbert, M. O., Baker, A., Jex, C. N., Graham, P. W., Treble, P. C., Andersen, M. S., and Acworth, R. I.: Drip water isotopes in semi-arid karst: Implications for speleothem paleoclimatology, *Earth Planet. Sci. Lett.*, 395, 194–204, 2014.
- deMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L., and Yarusinsky, M.: Abrupt onset and termination of the African Humid Period: rapid climate responses to gradual insolation forcing, *Quat. Sci. Rev.*, 19, 347–361, 2000.
- Dixit, Y., Hodell, D. A., and Petrie, C. A.: Abrupt weakening of the summer monsoon in northwest India ~4100 yr ago, *Geology*, 42, 339–342, 2014.
- Drysdale, R., Zanchetta, G., Hellstrom, J., Maas, R., Fallick, A., Pickett, M., Cartwright, I., and Piccini, L.: Late Holocene drought responsible for the collapse of Old World civilizations is recorded in an Italian cave flowstone, *Geology*, 34, 101–104, 2006.
- Drysdale, R. N., Zanchetta, G., Hellstrom, J. C., Fallick, A. E., Zhao, J.-x., Isola, I., and Bruschi, G.: Palaeoclimatic implications of the growth history and stable isotope ($\delta^{18}\text{O}$ and $\delta^{13}\text{C}$) geochemistry of a Middle to Late Pleistocene stalagmite from central-western Italy, *Earth Planet. Sci. Lett.*, 227, 215–229, 2004.
- Edwards, R. L., Chen, J. H., and Wasserburg, G. J.: ^{238}U - ^{234}U - ^{230}Th - ^{232}Th systematics and the precise measurements of time over the past 500 000 years., *Earth Planet. Sci. Lett.*, 81, 175–192, 1987.
- Fairchild, I. J. and Baker, A.: *Speleothem science – from process to past environments*, John Wiley & Sons, Ltd, Chichester, UK, 2012.
- Fairchild, I. J., Borsato, A., Tooth, A. F., Frisia, S., Hawkesworth, C. J., Huang, Y. M., McDermott, F., and Spiro, B.: Controls on trace element (Sr–Mg) compositions of carbonate cave waters: implications for speleothem climatic records, *Chem. Geol.*, 166, 255–269, 2000.
- Faith, J. T.: Late Pleistocene and Holocene mammal extinctions on continental Africa, *Earth-Sci. Rev.*, 128, 105–121, 2014.
- Finné, M., Holmgren, K., Sundqvist, H. S., Weiberg, E., and Lindblom, M.: Climate in the eastern Mediterranean, and adjacent regions, during the past 6000 years – A review, *J. Archaeol. Sci.*, 38, 3153–3173, 2011.
- Genty, D., Baker, A., Massault, M., Proctor, C., Gilmour, M., and Pons-Branchu, E.: Dead carbon in stalagmites: Carbonate bedrock paleodissolution vs. ageing of soil organic matter. Implications for ^{13}C variations in speleothems, *Geochim. Cosmochim. Ac.*, 65, 3443–3457, 2001a.
- Genty, D., Baker, A., and Vokal, B.: Intra- and inter-annual growth rate of modern stalagmites, *Chem. Geol.*, 176, 191–212, 2001b.
- Genty, D., Labuhn, I., Hoffmann, G., Danis, P. A., Mestre, O., Bourges, F., Wainer, K., Massault, M., Régnier, E., Orengo, P., Falourd, S., and Minster, B.: Rainfall and cave water isotopic relationships in two South-France sites, *Geochim. Cosmochim. Ac.*, 131, 323–343, 2014.
- Hendy, C. H.: The isotopic geochemistry of speleothems-I, The calculation of the effects of different modes of formation on the isotopic composition of speleothems and their applicability as palaeoclimatic indicators, *Geochim. Cosmochim. Ac.*, 35, 801–824, 1971.
- Hoerling, M. and Kumar, A.: The perfect ocean for drought, *Science*, 299, 691–694, 2003.
- IAEA: Isotopic composition of precipitation in the Mediterranean Basin in relation to air circulation patterns and climate: final report of a coordinated research project, 2000–2004, International Atomic Energy Agency, Vienna, Austria, 2005.
- Jackson, M. G., Oskarsson, N., Tronnes, R. G., McManus, J. F., Oppo, D. W., Gronvold, K., Hart, S. R., and Sachs, J. P.: Holocene loess deposition in Iceland: evidence for millennial-scale atmosphere-ocean coupling in the North Atlantic, *Geology*, 33, 509–512, 2005.
- Jaffey, A. H., Flynn, K. F., Glendenin, L. E., Bentley, W. C., and Essling, A. M.: Precision measurement of half-lives and specific activities of ^{235}U and ^{238}U , *Phys. Rev. C*, 4, 1889–1906, 1971.
- Jalut, G., Esteban Amat, A., Bonnet, L., Gauquelin, T., and Fontugne, M.: Holocene climatic changes in the Western Mediterranean, from south-east France to south-east Spain, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 160, 255–290, 2000.
- Johnson, K. R., Hu, C. Y., Belshaw, N. S., and Henderson, G. M.: Seasonal trace-element and stable-isotope variations in a Chinese speleothem: The potential for high-resolution paleomonsoon reconstruction, *Earth Planet. Sci. Lett.*, 244, 394–407, 2006.
- Kaufman, A., Wasserburg, G. J., Porcelli, D., Bar-Matthews, M., Ayalon, A., and Halicz, L.: U-Th isotope systematics from the Soreq cave, Israel and climatic correlations, *Earth Planet. Sci. Lett.*, 156, 141–155, 1998.
- Kherbouche, F., Hachi, S., Abdessadok, S., Sehil, N., Merzoug, S., Sari, L., Benchernine, R., Chelli, R., Fontugne, M., Barbaza, M., and Roubet, C.: Preliminary results from excavations at Guel-daman Cave GLD1 (Akbou, Algeria), *Quat. Int.*, 320, 109–124, 2014.
- Kröpelin, S., Verschuren, D., Lézine, A.-M., Eggermont, H., Cocquyt, C., Francus, P., Cazet, J.-P., Fagot, M., Rumes, B., Russell, J. M., Darius, F., Conley, D. J., Schuster, M., Suchodoletz, H. v., and Engstrom, D. R.: Climate-Driven Ecosystem Succession in the Sahara: The Past 6000 Years, *Science*, 320, 765–768, 2006.

- Lachniet, M. S.: Climatic and environmental controls on speleothem oxygen-isotope values, *Quat. Sci. Rev.*, 28, 412–432, 2009.
- Magny, M., Vanni re, B., Calo, C., Millet, L., Leroux, A., Peyron, O., Zanchetta, G., La Mantia, T., and Tinner, W.: Holocene hydrological changes in south-western Mediterranean as recorded by lake-level fluctuations at Lago Preola, a coastal lake in southern Sicily, Italy, *Quat. Sci. Rev.*, 30, 2459–2475, 2011.
- Magny, M., Combourieu-Nebout, N., de Beaulieu, J. L., Bout-Roumazeilles, V., Colombaroli, D., Desprat, S., Francke, A., Joannin, S., Ortu, E., Peyron, O., Revel, M., Sadori, L., Siani, G., Sicre, M. A., Samartin, S., Simonneau, A., Tinner, W., Vanni re, B., Wagner, B., Zanchetta, G., Anselmetti, F., Brugiapaglia, E., Chapron, E., Debret, M., Desmet, M., Didier, J., Essallami, L., Galop, D., Gilli, A., Haas, J. N., Kallel, N., Millet, L., Stock, A., Turon, J. L., and Wirth, S.: North-south palaeohydrological contrasts in the central Mediterranean during the Holocene: tentative synthesis and working hypotheses, *Clim. Past*, 9, 2043–2071, doi:10.5194/cp-9-2043-2013, 2013.
- Martrat, B., Grimalt, J. O., Lopez-Martinez, C., Cacho, I., Sierro, F. J., Flores, J. A., Zahn, R., Canals, M., Curtis, J. H., and Hodell, D. A.: Abrupt temperature changes in the Western Mediterranean over the past 250 000 years, *Science*, 306, 1762–1765, 2004.
- Mayewski, P. A., Rohling, E. E., Stager, J. C., Karl n, W., Maasch, K. A., Meeker, L. D., Meyerson, E. A., Gasse, F., Kreveld, S. V., Holmgren, K., Lee-Thorp, J., Rosqvist, G., Rack, F., Staubwasser, M., Schneider, R. R., and Steig, E. J.: Holocene climate variability, *Quat. Res.*, 62, 243–255, 2004.
- McDermott, F.: Palaeo-climate reconstruction from stable isotope variations in speleothems: a review, *Quat. Sci. Rev.*, 23, 901–918, 2004.
- McGee, D., deMenocal, P. B., Winckler, G., W.Stuut, J. B., and Bradtmiller, L. I.: The magnitude, timing and abruptness of changes in North African dust deposition over the last 20 000 yr, *Earth Planet. Sci. Lett.*, 371–372, 163–176, 2013.
- Mickler, P. J., Banner, J. L., Stern, L., Asmeron, Y., Edwards, R. L., and Ito, E.: Stable isotope variations in modern tropical speleothems: Evaluating equilibrium vs. kinetic isotope effects, *Geochim. Cosmochim. Ac.*, 68, 4381–4393, 2004.
- Mickler, P. J., Stern, L. A., and Banner, J. L.: Large kinetic isotope effects in modern speleothems, *Geol. Soc. Am. Bull.*, 118, 65–81, 2006.
- Morrellon, M., zvalero-Garces, B., Vegas-Vilarrubia, T., Gonzalez-Samperiz, P., Romero, O., Delgado-Huertas, A., Mata, P., Moreno, A., Rico, M., and Corella, J. P.: Lateglacial and Holocene paleohydrology in the western Mediterranean region: the Lake Estanya record (NE Spain), *Quat. Sci. Rev.*, 28, 2582–2599, 2009.
- Regattieri, E., Zanchetta, G., Drysdale, R. N., Isola, I., Hellstrom, J. C., and Dallai, L.: Lateglacial to Holocene trace element record (Ba, Mg, Sr) from Corchia Cave (Apuan Alps, central Italy): paleoenvironmental implications, *J. Quat. Sci.*, 29, 381–392, 2014.
- Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. J., Bronk Ramsey, C., Buck, C. E., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hafflidason, H., Hajdas, I., Hatt , C., Heaton, T. J., Hoffmann, D. L., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B., Manning, S. W., Niu, M., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Staff, R. A., Turney, C. S. M., and van der Plicht, J.: IntCal13 and Marine13 radiocarbon age calibration curves 0–52 000 years cal BP, *Radiocarbon*, 55, 1869–1887, 2013.
- Riehla, S., Pustovoytov, K. E., Weippert, H., Klett, S., and Hole, F.: Drought stress variability in ancient Near Eastern agricultural systems evidenced by $\delta^{13}\text{C}$ in barley grain, *P. Natl. Acad. Sci. USA*, 111, 12348–12353, 2014.
- Roberts, N., Brayshaw, D., Kuzucuoglu, C., Perez, R., and Sadori, L.: The mid-Holocene climatic transition in the Mediterranean: Causes and consequences, *Holocene*, 21, 3–13, 2011.
- Roland, T. P.: Was there a “4.2 kyr event” in Great Britain and Ireland? Evidence from the peatland record, PhD, University of Exeter, Exeter, UK, 2012.
- Sadori, L. and Narcisi, B.: The postglacial record of environmental history from Lago di Pergusa, Sicily, *Holocene*, 11, 655–672, 2001.
- Sadori, L. and Giardini, M.: Charcoal analysis, a method to study vegetation and climate of the Holocene: The case of Lago di Pergusa (Sicily, Italy), *Geobios*, 40, 173–180, 2007.
- Sadori, L., Jahns, S., and Peyron, O.: Mid-Holocene vegetation history of the central Mediterranean, *Holocene*, 21, 117–129, 2011.
- Scholz, D. and Hoffmann, D. L.: StalAge – An algorithm designed for construction of speleothem age models, *Quat. Geochronol.*, 6, 369–382, 2011.
- Seager, R., Ting, M., Held, I., Kushnir, Y., Lu, J., Vecchi, G., Huang, H. P., Harnik, N., Leetmaa, A., Lau, N. C., Li, C., Velez, J., and Naik, N.: Model projections of an imminent transition to a more arid climate in southwestern North America, *Science*, 316, 1181–1184, 2007.
- Staubwasser, M. and Weiss, H.: Holocene climate and cultural evolution in late prehistoric–early historic West Asia, *Quat. Res.*, 66, 372–378, 2006.
- Staubwasser, M., Sirocko, F., Grootes, P., and Segl, M.: Climate change at the 4.2 ka BP termination of the Indus valley civilization and Holocene south Asian monsoon variability, *Geophys. Res. Lett.*, 30, 1–4, 2003.
- Stoll, H. M., Moreno, A., Mendez-Vicente, A., Gonzalez-Lemos, S., Jimenez-Sanchez, M., Dominguez-Guesta, M. J., Edwards, R. L., Cheng, H., and Wang, X.: Paleoclimate and growth rates of speleothems in the northwestern Iberian peninsula over the last two glacial cycles, *Quat. Res.*, 80, 284–290, 2013.
- Tindall, J. C. and Valdes, P. J.: Modeling the 8.2 ka event using a coupled atmosphere-ocean GCM, *Global Planet. Change*, 79, 312–321, 2011.
- Touchan, R., Anchukaitis, K. J., Meko, D. M., Attalah, S., Baisan, C., and Aloui, A.: Long term context for recent drought in northwestern Africa, *Geophys. Res. Lett.*, 35, L13705, doi:10.1029/2008GL034264, 2008.
- Touchan, R., Anchukaitis, K. J., Meko, D. M., Sabir, M., Attalah, S., and Aloui, A.: Spatiotemporal drought variability in northwestern Africa over the last nine centuries, *Clim. Dyn.*, 37, 237–252, 2011.
- Vaks, A., Woodhead, J., Bar-Matthews, M., Ayalon, A., Cliff, R. A., Zilberman, T., Matthews, A., and Frumkin, A.: Pliocene–Pleistocene climate of the northern margin of Saharan–Arabian Desert recorded in speleothems from the Negev Desert, Israel, *Earth Planet. Sci. Lett.*, 368, 88–100, 2013.
- Wanner, H., Beer, J., B tikofer, J., Crowley, T. J., Cubasch, U., Fl ckiger, J., Goosse, H., Grosjean, M., Joos, F., Kaplan, J. O., K ttel, M., M ller, S. A., Prentice, C. I., Solomina, O., Stocker,

- T. F., Tarasov, P., Wagner, M., and Widmann, M.: Mid- to Late Holocene climate change: an overview, *Quat. Sci. Rev.*, 27, 1791–1828, 2008.
- Weiss, H. and Bradley, R. S.: What Drives Societal Collapse?, *Science*, 291, 609–610, 2000.
- Weiss, H., Courty, M.-A., Wetterstrom, W., Guichard, F., Senior, L., Meadow, R., and Curnow, A.: The genesis and collapse of third millennium North Mesopotamian civilization, *Science*, 261, 995–1004, 1993.
- Wiener, M. H.: The interaction of climate change and agency in the collapse of civilizations ca. 2300–2000 BC, *Radiocarbon*, 56, S1–S16, doi:10.2458/azu_rc.56.18325, 2014.
- Zanchetta, G., Drysdale, R. N., Hellstrom, J. C., Fallick, A. E., Isola, I., Gagan, M. K., and Paresch, M. T.: Enhanced rainfall in the Western Mediterranean during deposition of sapropel S1: stalagmite evidence from Corchia cave (Central Italy), *Quat. Sci. Rev.*, 26, 279–286, 2007.
- Zanchetta, G., Giraudi, C., Sulpizio, R., Magny, M., Drysdale, R. N., and Sadori, L.: Constraining the onset of the Holocene “Neoglacial” over the central Italy using tephra layers, *Quat. Res.*, 78, 236–247, 2012a.
- Zanchetta, G., van Welden, A., Baneschi, I., Drysdale, R. N., Sadori, L., Roberts, N., Giardini, M., Beck, C., and Pascucci, V.: Multi-proxy record for the last 4500 years from Lake Shkodra (Albania/Montenegro), *J. Quat. Sci.*, 27, 780–789, 2012b.
- Zanchetta, G., Bar-Matthews, M., Drysdale, R. N., Lionello, P., Ayalon, A., Hellstrom, J. C., Isola, I., and Regattieri, E.: Coeval dry events in the central and eastern Mediterranean basin at 5.2 and 5.6 ka recorded in Corchia (Italy) and Soreq caves (Israel) speleothems, *Global Planet. Change*, 122, 130–139, 2014.