

Optimizing the terrestrial ecosystem gross primary productivity using carbonyl sulfide (COS) within a “two-leaf” modeling framework

Huajie Zhu^{1†}, Xiuli ~~Xing~~^{1,2}~~Xing~~^{2†}, Mousong Wu^{1*}, Weimin Ju¹, Fei Jiang^{1,3}

1 International Institute for Earth System Science, Nanjing University, Nanjing, China

5 ~~2-currently at:2~~ Department of Environmental Science and Engineering, Fudan University, Shanghai, China

3 Frontiers Science Center for Critical Earth Material Cycling, Nanjing University, Nanjing, China

Correspondence to: Mousong Wu (mousongwu@nju.edu.cn)

†they contribute equally to this work

10 **Abstract.** Accurately modeling gross primary productivity (GPP) is of great importance in diagnosing terrestrial carbon-climate feedbacks. Process-based terrestrial ecosystem models are often subject to substantial uncertainties, primarily attributed to inadequately calibrated parameters. Recent attention has identified carbonyl sulfide (COS) as a promising proxy of GPP, due to the close linkage between leaf exchange of COS and carbon dioxide (CO₂) through their shared pathway of stomatal diffusion. However, most of the current modeling approaches for COS and CO₂ ~~did not~~ not explicitly consider the

15 vegetation structural impacts, i.e., the differences between the ~~sun-shade and~~ sunlit and shaded leaves in COS uptake. This study used ecosystem COS fluxes ~~data~~ from 7 sites to optimize GPP estimation across various ecosystems with the ~~Boreal Ecosystem Productivity~~ Biosphere-atmosphere Exchange Process Simulator (BEPS), which was further developed for simulating the leaf COS uptake under its state-of-the-art “two-leaf” framework. Our results demonstrated the substantial improvement in GPP simulation across various ecosystems through the fusion data assimilation of COS ~~data flux~~ into the “two-

20 leaf” model, with the ensemble mean of root mean square error (RMSE) for simulated GPP reduced by ~~18.99~~ 20.16 % to ~~66.64~~ 12 %. Notably, we also shed light on the remarkable identifiability of key parameters within the BEPS model, including the maximum carboxylation rate of Rubisco at 25 °C (V_{cmax25}), minimum stomatal conductance (b_{H_2O}), and leaf nitrogen content (N_{leaf}), despite intricate interactions among COS-related parameters. Furthermore, our global sensitivity analysis delineated both shared and disparate sensitivities of COS and GPP to model parameters and suggested the unique treatment of

25 parameters for each site in COS and GPP modeling. In summary, our study deepened insights into the sensitivity, identifiability, and interactions of parameters related to COS, and showcased the efficacy of COS in reducing uncertainty in GPP simulations.

Keywords: carbonyl sulfide, ~~photosynthesis, model gross primary productivity (GPP),~~ data fusion assimilation, ecosystem modeling, parameter optimization

1 Introduction

30 Over the past five decades, terrestrial ecosystems have been absorbing about 30 % of anthropogenic carbon dioxide (CO₂) emissions, playing a crucial role in mitigating climate change (Friedlingstein et al., 2022). Driven by the photosynthesis of terrestrial vegetation, ~~the~~ gross primary productivity (GPP) is the largest terrestrial carbon flux and plays an important role in understanding terrestrial carbon-climate feedbacks (Luo, 2007; Wang et al., 2021). However, as the direct observations of GPP using atmospheric CO₂ observations are confounded by respiration (Hilton et al., 2017), and the modeling of GPP ~~are~~ is affected by a range of uncertainties, such as the poorly calibrated parameters (Macbean et al., 2022), the precise quantification of GPP in terrestrial ecosystems has been a major challenge (Canadell et al., 2000; Yuan et al., 2007).

~~In the past decade, carbonyl sulfide (COS)~~ Carbonyl sulfide (COS) is the most abundant sulfur-containing trace gas in the atmosphere with a lifetime of about 2 years (Montzka et al., 2007; Karu et al., 2023). The tropospheric atmospheric mole fraction of COS is approximately 500 parts per trillion (ppt), exhibiting a typical seasonal amplitude of ~ 100–200 ppt (Montzka et al., 2007; Ma et al., 2021; Hu et al., 2021; Remaud et al., 2022; Remaud et al., 2023; Ma et al., 2023). In the past decade, COS has emerged as a promising tracer for terrestrial photosynthesis (Stimler et al., 2010; Asaf et al., 2013; ~~Billesbach et al., 2014; Maseyk et al., 2014;~~ Launois et al., 2015; ~~Hilton et al., 2017; Yang et al., 2018;~~ Kooijmans et al., 2019; ~~Spielmann et al., 2019; Hu et al., 2021; Kohonen et al., 2022~~) and stomatal conductance (Commane et al., 2015; Wehr et al., 2017; Sun et al., 2022; ~~Cho et al., 2023~~) as the leaf exchange of COS and carbon dioxide (CO₂) are tightly coupled through stomata (45 ~~Goldan et al., 1988;~~ Sandoval-Soto et al., 2005; Seibt et al., 2010; Wohlfahrt et al., 2012; Whelan et al., 2018). Unlike CO₂, which is emitted back to the atmosphere via leaf respiration (Sun et al., 2022), COS is completely destroyed by a hydrolysis reaction catalyzed by carbonic anhydrase (Protoschill-Krebs et al., 1996) without back-flux in leaves under normal conditions (Stimler et al., 2010). Consequently, the measurement of COS flux is able to provide a direct and independent way to estimate GPP (Sandoval-Soto et al., 2005; Abadie et al., 2023).

50 In most of the early studies, GPP was directly estimated by scaling measurement of plant COS uptake with the empirically derived leaf relative uptake (LRU) approach or its extensions that ~~take~~ incorporate the effects of temperature, humidity, light and CO₂ concentration on stomatal conductance ~~into account~~ (Kohonen et al., 2022; Sun et al., 2022; Abadie et al., 2023) because of the simplicity of this approach and the sufficiency of it in many cases (Sandoval-Soto et al., 2005; Whelan et al., 2018). In contrast, the process-based model that mechanistically simulates COS plant uptake by incorporating stomatal transport processes were also developed and widely evaluated (Maignan et al., 2021; Kooijmans et al., 2021). Concurrently, the significance of soil COS exchange has also been recognized, leading to the development of a suite of empirical or mechanistic COS soil exchange models (Kesselmeier et al., 1999; Berry et al., 2013; Launois et al., 2015; Sun et al., 2015; Ogée et al., 2016; Whelan et al., 2022). Process-based COS plant uptake model and soil exchange ~~model~~ models have been integrated into land surface models (LSMs) (Berry et al., 2013; Maignan et al., 2021; Kooijmans et al., 2021; ~~Chen et al.,~~ 60 ~~2023~~). Consequently, by constraining the model parameters of LSMs with COS through data assimilation, not only the model

variables like GPP are expected to be ~~optimized~~improved, but also our understanding of ecosystem processes is expected to be significantly ~~deepened~~enhanced.

Currently, several studies have been ~~endeavored~~conducted to refine the model parameters of LSMs through assimilating the COS data, and thereby ~~optimized~~optimizing the modeling of water-carbon fluxes (Chen et al., 2023; Abadie et al., 2023; Zhu et al., 2023). Within a big-leaf framework, Abadie et al. (2023) demonstrated COS could provide mechanistic constraint on stomatal diffusion, and the joint assimilation of COS and GPP ~~can also~~is able to improve the model performance of GPP and latent heat. Ecosystem carbon, water and energy processes are ~~interacted~~interacting and nonlinear, the changes in one process could induce variations in the other processes. While COS assimilation has proven effective in constraining COS-related model parameters and optimizing GPP, there remains a gap of systematic understanding of the ability of COS to optimize model parameters from different processes, ~~e.g.~~ For example, how effective is the assimilation of COS in reducing model prediction uncertainty of GPP as well as the relevant ecosystem processes in different ecosystems?

~~In this context, we have further explored the capacity of COS to constrain the model parameters of LSM and to optimize GPP, with the aims to answering the following questions:~~

~~Which~~Due to the dissimilar illumination conditions, there is the significant variability of leaf photosynthesis between sunlit and shaded leaves (Chen et al., 1999; Pignon et al., 2017; Wang et al., 2018; Bao et al., 2022). It is now clearly recognized that big-leaf models are conceptually flawed and practically inaccurate and sunlit-shaded leaf stratification is necessary to make accurate canopy-level photosynthesis estimation (Chen et al., 1999; Chen et al., 2012; Luo et al., 2018). Consequently, in the process-based LSM that simulates COS plant uptake and photosynthesis in a coupled manner (Ball et al., 1987; Berry et al., 2013), the application of the two-leaf model shows promise for providing accurate simulation of plant COS uptake. In this context, we have further explored the capacity of COS to constrain the model parameters of a LSM and to optimize GPP within the two-leaf modeling framework.

~~Our goal is to address the following questions:~~

~~To which~~ parameters is the COS simulation ~~is~~ sensitive ~~to~~, and what are the differences in parameter sensitivities between COS and GPP?

85 How effective is COS assimilation in ~~enhancing~~improving model prediction and reducing prediction uncertainty of GPP?

Which process parameters can be well identified by the assimilation of COS?

How do process parameters interact in COS modeling across diverse ecosystems?

To address these questions, we utilized ~~the~~ ecosystem COS flux data to optimize GPP across various ecosystems based on the ~~combination~~coupling of COS modeling with the ~~Boreal Ecosystem Productivity~~two-leaf based Biosphere-atmosphere Exchange Process Simulator (BEPS). ~~Specifically~~Through Monte Carlo simulations, we investigated~~conducted a global parameter sensitivity analysis to explore the usefulness~~sensitivity of COS ~~in constraining the ecosystem processes~~and GPP simulations to model parameters related ~~to~~ not only to photosynthesis, but also ~~the~~to water and energy. ~~Furthermore, parameter uncertainties as well as the uncertainties in the optimization results~~The interaction and identifiability of these parameters were

95 quantified using Monte Carlo optimized parameter sets. Additionally, the effectiveness of COS in constraining model uncertainty in simulated COS and GPP is evaluated.

2 Materials and methods

2.1 Model description

2.1.1 BEPS basic model

The BEPS model (Liu et al., 1997; Chen et al., 1999; Chen et al., 2012) used in this study is a process-based diagnostic model driven by remotely sensed vegetation parameters, including leaf area index (LAI), clumping index, and land cover type, as well as meteorological and soil data (Chen et al., 2019). With the ~~consideration of~~ coupling among terrestrial carbon, water, and nitrogen cycles (He et al., 2021), it simulates photosynthesis, energy balance, and hydrological and soil biogeochemical processes at hourly time steps (Ju et al., 2006; Liu et al., 2015). For photosynthesis, it stratifies whole canopies into sunlit and shaded leaves and ~~calculated~~calculates GPP for each group of leaves by scaling Farquhar's leaf biochemical model (Farquhar et al., 1980) up to canopy-level with a temporal and spatial scaling scheme (Chen et al., 1999). In this study, the BEPS model stratifies the soil profile into five layers, and the model implicitly solves the soil water content values for these layers (Ju et al., 2010). Over the last few decades, the BEPS model has been continuously improved and has been used in a wide variety of terrestrial ecosystems (Schwalm et al., 2010; Liu et al., 2015). This study ~~used~~uses the BEPS model that simulates water, carbon and energy processes at hourly interval which enables the detection of diel variations of model variables (Xing et al., 2023).

2.1.2 The ~~two-leaf~~² scheme for GPP and COS modeling

The BEPS model simulates the canopy photosynthesis for the sunlit and ~~sun-shaded~~shaded leaves separately,

$$GPP = GPP_{sunlit}LAI_{sunlit} + GPP_{shaded}LAI_{shaded} \quad (1)$$

where- GPP_{sunlit} and GPP_{shaded} denote the GPP per unit area for sunlit and shaded leaves, LAI_{sunlit} and LAI_{shaded} represent the LAI values of sunlit and shaded leaves, respectively. ~~The detailed descriptions about the photosynthesis~~ LAI_{sunlit} and LAI_{shaded} are calculated as (Chen et al., 1999):

$$LAI_{sunlit} = 2 \cos \theta \left(1 - e^{\left(1 - \frac{0.5\Omega LAI_{total}}{\cos \theta}\right)} \right) \quad (2)$$

$$LAI_{shaded} = LAI_{total} - LAI_{sunlit} \quad (3)$$

Where θ is the solar zenith angle, LAI_{total} is the total leaf area index of the canopy, and Ω is the clumping index.
120 GPP values of sunlit and shaded leaves are calculated using the Farquhar's model (Farquhar et al., 1980) with consideration of the large difference in incident solar irradiance between these two-leaf groups (Chen et al., 2012; Chen et al., 2019). Stomatal conductances of sunlit and shaded leaves are determined separately according to photosynthesis rates of these leaves.

atmospheric CO₂ concentration, relative humidity and soil moisture (Ball et al., 1987; Ju et al., 2010). The detailed descriptions about the photosynthesis and stomatal conductance modeling approach of BEPS are illustrated in Appendix A1.

125 The ecosystem COS flux includes both plant COS uptake $F_{\text{ecos,plant}} F_{\text{COS,plant}}$ and soil COS flux exchange $F_{\text{ecos,soil}} F_{\text{COS,soil}}$ (Whelan et al., 2016). In this work, the canopy-level COS plant uptake $F_{\text{ecos,plant}} F_{\text{COS,plant}}$ (pmol m⁻² s⁻¹) was calculated by upscaling the resistance analog model of COS uptake (Berry et al., 2013) with the ‘two-leaf’ upscaling scheme (Chen et al., 1999). Considering the different responses of foliage to diffuse and direct solar radiation (Gu et al., 2002), $F_{\text{COS,plant}}$ is calculated as:

$$130 \quad F_{\text{COS,plant}} = F_{\text{COS,sunlit}} LAI_{\text{sunlit}} + F_{\text{COS,shaded}} LAI_{\text{shaded}} \quad (4)$$

where $F_{\text{COS,sunlit}}$ and $F_{\text{COS,shaded}}$ denote the leaf-level COS uptake rate (pmol m⁻² s⁻¹) for sunlit and shaded leaves. The leaf-level COS uptake rate $F_{\text{COS,leaf}}$ is determined by the formula (Berry et al., 2013):

$$F_{\text{COS,leaf}} = COS_a \left(\frac{1.94}{g_{sw}} + \frac{1.56}{g_{bw}} + \frac{1}{g_{\text{COS}}} \right)^{-1} \quad (5)$$

135 ~~$F_{\text{COS,plant}} = F_{\text{COS,sunlit}} LAI_{\text{sunlit}} + F_{\text{COS,shaded}} LAI_{\text{shaded}} \quad (2)$~~
 ~~g_{sw} and g_{bw}~~
~~are the stomatal conductance and leaf laminar boundary layer conductance to water vapor (H₂O). The factors 1.94 and 1.56 account for the smaller diffusivity of COS with respect to H₂O. g_{COS} indicates the apparent conductance for COS uptake from the intercellular airspaces, which combined the mesophyll conductance (Evans et al., 1994) and the biochemical reaction rate of COS and carbonic anhydrase (Badger and Price, 1994). It can be calculated as :~~

$$140 \quad g_{\text{COS}} = \alpha V_{\text{cmax}} \quad (6)$$

Where α is a parameter that is calibrated to observations of simultaneous measurements of COS and CO₂ uptake (Stimler et al., 2012). V_{cmax} is the maximum carboxylation rate of Rubisco. Analysis of these measurements yield estimates of α of ~1400 for C3 and ~7500 for C4 species. With reference the COS modelling scheme of the Simple biosphere model (version 4.2) (Haynes et al., 2020), g_{COS} can be calculated as

$$145 \quad g_{\text{COS}} = 1.4 * 10^3 * (1.0 + 5.33 * F_{\text{C4}}) * 10^{-6} F_{\text{APAR}} f_w V_{\text{cmax}} \quad (7)$$

where F_{C4} denotes the C4 plant flag, taking the value of 1 for C4 plants and 0 otherwise. f_w is a soil moisture stress factor describing the sensitivity of g_{sw} to soil water availability (Ju et al., 2006). F_{APAR} is the scaling factor for leaf radiation (Smith et al., 2008), calculated as:

$$150 \quad F_{\text{APAR}} = 1 - e^{(-0.45 LAI)} \quad (8)$$

The soil COS fluxes are simulated by considering the abiotic and biotic components separately, as by Whelan et al. (2016). We took the soil COS modeling scheme including the parameterizations from ~~Abadie et al. (2022)~~ Whelan et al. (2016) and Whelan et al. (2022) in this study (see Appendix A2) given that our focus is the COS and GPP relationships and the previous studies have verified this approach over multiple sites with measurements.

155 2.2 Site description

The model was evaluated on seven sites distributed on the Eurasian and American continents in boreal, temperate and subtropical regions based on field observations collected from several studies. Those sites were representative of different climate regions and land cover types (in the model represented by plant function types, and soil textures, as depicted in Table 1).

160 **Table 1.** Site characteristics. Site identification includes the country initials and a three-letter name for each site. ~~locations~~Locations of the sites are provided by the latitude (Lat) and longitude (Lon). PFT stands for plant functional type. ENF and DBF denote evergreen needleleaf forest and deciduous broadleaf forest respectively.

Site name	Lat (°N)	Lon (°E)	PFT	Soil texture	Year	References
AT-Neu	47.12	11.32	C3 grass	Sandy loam	2015	Spielmann et al. (2019)
DK-Sor	55.49	11.64	DBF	Sandy loam	2016	Spielmann et al. (2019)
ES-Lma	39.94	-5.77	C3 grass	Sandy loam	2016	Spielmann et al. (2019)
FI-Hyy	61.85	24.29	ENF	Sandy loam	2013-2017	Vesala et al. (2022), Sun et al. (2018)
IT-Soy	45.87	13.08	C3 crop	Silt clay	2017	Spielmann et al. (2019), Abadie et al. (2022)
US-Ha1	42.54	-72.17	DBF	Sandy loam	2012-2013	Wehr et al. (2017), Commans et al. (2015)
US-Wrc	45.82	-121.95	ENF	Loam	2014	Rastogi et al. (2018), Shaw et al. (2004)

2.3 Data

Data used in this study include LAI, land cover type, meteorological and soil data, as well as CO₂ and COS mole fraction data.

165 The CO₂ and COS mole fractions in the bulk air were assumed to be spatially invariant over the globe but to vary annually. The CO₂ mole fraction data in this study are taken from the Global Monitoring Laboratory (<https://gml.noaa.gov/ccgg/trends/global.html>). For the COS mole fraction, we utilized the average of observations from sites SPO (South Pole) and MLO (Mauna Loa, United States) to drive the model. These data are publicly available online at: <https://gml.noaa.gov/hats/gases/OCS.html>.

170 2.3.1 LAI dataset

~~The satellite LAI data that best matched the field measurements was selected from three candidate LAI products to drive the model.~~The LAI dataset used here are the GLOBMAP global leaf area index product (Version 3) (see [GLOBMAP global Leaf Area Index since 1981 | Zenodo](#)) and the Global Land Surface Satellite (GLASS) LAI product (Version 3) (acquired from <ftp://ftp.glcf.umd.edu/>). They represent ~~Leaf area index~~LAI at a spatial resolution of 8 km (Liu et al., 2012) and 1 km (Xiao et al., 2016) respectively, and a temporal resolution of 8-day. With reference to the observed LAI at these sites (Wehr et al., 2017; Rastogi et al., 2018; Spielmann et al., 2019; Kohonen et al., 2022), ~~the GLASS LAI was adopted to DK-Sor and IT-Soy, and the GLOBMAP LAI was utilized for the remaining five sites.~~we used GLOBMAP products to drive the BEPS model at most sites (5/7) due to its good agreement with the observed LAI. Specifically, as the GLOBMAP product had considerably

underestimated LAI at DK-Sor and was not consistent with the vegetation phenology at ES-Lma during the measurement
180 campaign (Spielmann et al., 2019). GLASS LAI was used at these two sites. In addition, these LAI products were interpolated
into daily values by the nearest neighbor method for the simulation.

2.3.2 Meteorological dataset

Meteorological data required to force the BEPS model include air temperature, shortwave radiation, precipitation, relative
humidity and wind speed. As the simulations were conducted at the site scale, we utilized the FLUXNET2015 data (see
185 <https://fluxnet.org> for AT-Neu, DK-Sor and ES-Lma ~~and~~, FI-Hyy and US-Ha1) and the AmeriFlux data (see
<https://ameriflux.lbl.gov> for US-Ha1 and US-Wrc). ~~Specifically, as~~ As FLUXNET2015 meteorological data for AT-Neu were
only accessible for the period 2002-2012, we conducted a linear fit between its ERA5 ~~data~~ ~~(European Centre for Medium-~~
~~Range Weather Forecasts (ECMWF) Reanalysis v5) data~~ ([https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-
single-levels?tab=overview](https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-
single-levels?tab=overview)) and FLUXNET2015 meteorological data for the corresponding period. Then, we used the fitted
190 parameters to adjust the ERA5 data ~~for~~ for 2015, thereby obtaining downscaling information for the meteorological data. In
addition, we utilized the FLUXNET data in 2012, and Ameriflux data and ERA5 shortwave radiation data ~~of~~ in 2013 to drive
the BEPS model at US-Ha1 ~~were derived from ERA5,~~ due to the absence of FLUXNET data in 2013 and the lack of
available shortwave radiation data in AmeriFlux of Ameriflux.


2.3.3 COS and GPP datasets

195 The hourly ecosystem COS flux observations were utilized to perform optimization and to evaluate the optimization results.
They were derived from existing studies with pre-processing with regard to the data quality check, as listed in ~~Table 1~~ Table 1.
Specifically, following the recommendations regarding the standardized processing of eddy covariance flux measurements of
COS by Kohonen et al. (2020), both the measured and gap-filled COS flux observations are provided in Vesala et al. (2022),
and the latter were utilized in this study. To assess the model performance of GPP, the GPP observations were also collected
200 from FLUXNET (DK-Sor, ES-Lma ~~and~~, FI-Hyy and US-Ha1 in 2012), AmeriFlux (US-Ha1 in 2013), ~~and US-Wrc~~, ~~and~~
existing studies (Spielmann et al. (2019) ~~(AT-Neu and IT-Soy)~~ for AT-Neu and IT-Soy and Rastogi et al. (2018) for US-Wrc).
Given that only CO₂ turbulent flux (FC) or net ecosystem exchange (NEE) data were available at AT-Neu, IT-Soy, ~~US-Ha1~~
and US-~~Wrc~~ Ha1 in 2013, a night flux partitioning model (Reichstein et al., 2005) was employed to derive GPP. This model
assumes that nighttime NEE represents ecosystem respiration R_{eco} , and thus partitions FC or NEE into GPP and R_{eco} based
205 on the semi-empirical models of respiration, which use air temperature as a driver (Lloyd and Taylor, 1994; Lasslop et al.,
2012).

2.4 The Monte Carlo-based parameter calibration-optimization approach

To evaluate the sensitivity, equifinality and interaction of model parameters, ~~and the uncertainty of model outputs~~, the Monte
Carlo-based parameter calibration methodology optimization approach was employed here. ~~(Figure 1)~~. The methodology calls

210 for rejecting the concept of a unique global optimum parameter set within some particular model structure, instead recognizing the “equifinality” of parameter sets that exhibit similarly good performance in producing the observed data (Beven and Freer, 2001). ~~Thus, the result is depicted with an ensemble of acceptable or behavioral parameter sets~~In a Monte Carlo simulation framework, a large number of random sets of parameters are derived across specified parameter ranges (Staudt et al., 2010) and employed to drive the model. Subsequently, model realizations are grouped into behavioral and non-behavioral model runs and associated parameter sets based on the values of the single or multiple performance measures and the predefined threshold value (Houska et al., 2014). The former describes acceptable model realizations conditioned on the available observational data (Blasone et al., 2008; Beven and Binley, 2014). ~~This method~~The latter describes parameter sets that produce behavior inconsistent with observed behavior. Given the gradual transition of performance measures between behavioral and non-behavioral model runs within the Monte Carlo framework, the threshold value used to distinguish between behavioral and non-behavioral parameter sets was often determined by an acceptable samples rate, i.e., ranking model runs and taking the top X% as behavioral (Beven and Binley, 2014). In the past few decades, this approach has been extensively used in ecosystem modeling with multiple parameters to be calibrated and shown high ability in constraining multiple ecosystem processes (Tonkin and Doherty, 2009; Houska et al., 2014; He et al., 2016; Wu et al., 2019; Wu et al., 2020; Xing et al., 2023).



225 **Figure 1.** Flowchart of the Monte Carlo-based parameter optimization approach used in this study.

2.4.1 Parameter selection and sampling strategy

Based on current understanding of COS exchange (Wohlfahrt et al., 2012; Berry et al., 2013; Whelan et al., 2016; Whelan et al., 2018; Cho et al., 2023) and photosynthesis processes (Ball et al., 1987; Raines, 2003; Blankenship, 2021) and related parameter sensitivity studies (Liu et al., 2011; Chen et al., 2012; Chen et al., 2023; Xing et al., 2023; [Abadie et al., 2023](#); [Zhu et al., 2023](#)), 9 parameters were selected to be calibrated in this study (for details see Table [B42](#)). These parameters are related to formulas describing four processes: 1) photosynthesis (V_{cmax25} , VJ_{slope} , N_{leaf}), 2) soil hydrology ($Ksat_{scalar}$, b_{scalar} , r_{decay}), 3) stomatal gas exchange (b_{H_2O} , m_{H_2O}), and 4) energy balance (f_{leaf}). Specifically, $Ksat_{scalar}$ and b_{scalar} are scaling factors designed to optimize the saturated hydraulic conductivity (Ksat) and the Campbell parameter (b) for each soil layer in the BEPS model. The [default prior](#) values and prior ranges for these parameters (Table [B42](#)) were chosen based on literature ([Jackson et al., 1996](#); [Medlyn et al., 1999](#); [Kattge et al., 2009](#); [Miner et al., 2017](#); [Ryu et al., 2018](#)) and default model settings. Uniform distributions were assigned to all parameters, and 20,000 sets of parameters were generated through random sampling.

Table 2. Descriptions of the 9 parameters that were selected to be calibrated. The prior values and prior ranges (in parentheses) of these parameters are given for each plant function type (PFT) or for each soil texture or globally according to the parameter dependency. ENF and DBF denote evergreen needleleaf forest and deciduous broadleaf forest respectively.

Parameter	Description	Dependent	Prior value and prior range			
			ENF/ Sand loam	DBF/Slit loam	C3 grass/ Loam	C3 crop
b_{H_2O}	The intercept of the Ball-Berry model ($\text{mol m}^{-2}\text{s}^{-1}$)	PFT	0.0175 (0-1)	0.0175 (0-1)	0.0175 (0-1)	0.0175 (0-1)
b_{scalar}	The scaling factor of Campbell parameter b (unitless)	Texture	1 (0.25-1.75)	1 (0.25-1.75)	1 (0.25-1.75)	1 (0.25-1.75)
f_{leaf}	Ratio of photosynthetically active radiation to shortwave radiation (unitless)	Global	0.466 (0.42-0.51)			
$Ksat_{scalar}$	The scaling factor of saturated hydraulic conductivity Ksat (unitless)	Texture	1 (0.25-1.75)	1 (0.25-1.75)	1 (0.25-1.75)	1 (0.25-1.75)
m_{H_2O}	The slope of the Ball-Berry model (unitless)	PFT	8 (2-14)	8 (2-14)	8 (2-14)	8 (2-14)
N_{leaf}	Leaf nitrogen content ($\text{m}^2 \text{g}^{-1}$)	PFT	3.10 (0.40-5.80)	1.74 (0.32-3.16)	1.75 (0.23-3.27)	1.62 (0.40-2.84)
r_{decay}	Decay rate of root distribution (unitless)	PFT	0.95 (0.80-0.99)	0.97 (0.80-0.99)	0.96 (0.80-0.99)	0.95 (0.80-0.99)
V_{cmax25}	Maximum carboxylation rate of Rubisco at 25 °C ($\mu\text{mol m}^{-2}\text{s}^{-1}$)	PFT	62.5 (13.1-111.9)	57.7 (15.3-100.1)	78.2 (16-140.4)	100.7 (27.5-173.9)
VJ_{slope}	Slope of the V_{cmax} and J_{max} (maximum electron transport rate) relationship (unitless)	PFT	2.39 (1-4)	2.39 (1-4)	2.39 (1-4)	2.39 (1-4)

240 2.4.2 Selection of behavioral simulations

To measure the agreement between model simulations and observations, a variety of performance metrics have been proposed and utilized in ~~in~~ previous studies (Beven and Binley, 1992; Moradkhani et al., 2005; Staudt et al., 2010). In this study, we employed the root mean square error (RMSE) to distinguish between behavioral and non-behavioral simulations.

$$RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (obs_i - sim_i)^2} \quad (3)$$

245 where N is the total number of observations. “obs” and “sim” denote the observations and simulations, respectively. sim_i denotes the simulation corresponding to the i th observation obs_i .

~~In specific,~~ Specifically, here we chose an acceptable samples rate of 0.5%, i.e., the top 0.5% 100 model runs with the lowest RMSE values for COS as behavioral simulations. Thus, the deterministic model prediction is given by the ensemble mean of the 100 behavioral simulations.

250 2.4.35 Uncertainty quantification

The model prediction limits or uncertainty bounds can be determined by forming the cumulative density function (CDF) of the ensemble of simulations (Beven and Binley, 2014), normally chosen at the 5 % and 95 % confidence level in most of the previous studies (Blasone et al., 2008). Similarly, we chose the 5 % and 95 % quantiles of the 20,000 simulations and the 100 behavioral simulations to quantify the model output uncertainty in this study.

255 2.56 Parameter sensitivity

In order to take full advantage of the Monte Carlo ~~sampled parameter set~~ simulations, a density-based global sensitivity analysis approach (Plischke et al., 2013) was used to investigate the sensitivity of COS and GPP simulations to the selected model parameters via the Sensitivity Analysis Library (SALib) (Iwanaga et al., 2022). This approach aims at assessing the influence of the entire input distribution on the entire output distribution without reference to a particular moment of the output
260 (Borgonovo, 2007). According to Borgonovo (2007), the sensitivity index (δ) is always between 0 and 1, ~~-it~~ equals 0 if the output is not dependent upon the model parameter, and it equals unity if all model parameters ~~is~~ are considered.

2.67 Parameter uncertainty

Due to the functional and structural complexity of ~~ecosystem~~ ecosystems, ecosystem models often require a substantial number of parameters to realize the modeling of various ecosystem processes, and some parameters are compensating ~~with~~ each other
265 (Mo et al., 2008). While the parameter interactions related to photosynthesis have been systematically studied (Tang and Zhuang, 2009; Lu et al., 2013; Wu et al., 2019; Xing et al., 2023), the parameter interactions related to COS flux simulation have not been reported. Based on the Monte Carlo-based methodology, the numerous behavioral parameter sets around the

“optimum” (Beven and Freer, 2001) provide us with the opportunity to analyze the interactions between the selected parameters. In this study, the Pearson correlation coefficient and the confidence level were employed to identify the parameter interactions.

Parameter identifiability (PI) is the concept of whether uncertain parameters can be correctly estimated from the observed data (Yi et al., 2019). The failure in PI is supposed to be caused by ‘over-parameterization’ and parameter interactions (due to high nonlinearity of model equations) (Gan et al., 2014). Inspired by Yi et al. (2019) who used likelihood confidence interval as a measure of PI, here we used parameter distribution range for the same purpose. Taking into account the influence of the prior distribution of the behavioral parameter sets, the PI is defined as the reduction of the parameter range width. Hence, a large value of PI indicates that the parameter is well identified in the calibration optimization process.


3 Results

3.1 Parameter sensitivity

The sensitivity indexes of COS and GPP simulations to the model parameters for the seven sites are illustrated in Fig. 1. It can be seen that both COS and GPP simulations exhibit high sensitivity to leaf nitrogen content (N_{leaf}) and the maximum carboxylation rate of Rubisco at 25 °C (V_{cmax25}), while showing low sensitivity to energy balance related parameter f_{leaf} as well as soil hydrology related parameters (including b_{scalar} , $Ksat_{scalar}$ and r_{decay}). With the average values of sensitivity index of 0.14, 0.1211 and 0.0810, the photosynthesis related parameters f_{leaf} and parameter V_{Jslope} as well as stomata conductance related parameter m_{H_2O} can significantly impact the simulation of GPP. However, those parameters do not exhibit high sensitivity in the modeling of COS. Furthermore, our results also highlight the crucial role of the intercept of the Ball-Berry model (b_{H_2O}) in the modeling of COS, yet its impact on the simulation of GPP is limited. In summary, our results suggest that the simulated COS and GPP share some similarities in their sensitivities to parameters, but there are also notable differences. Specifically, the parameters including f_{leaf} , m_{H_2O} and V_{Jslope} strongly influence GPP modeling simulations but have minimal impacts on COS modeling simulations. Conversely, the parameter b_{H_2O} plays a more crucial role in COS simulation.

With the mean values of 0.3033, 0.29 and 0.09 respectively, the sensitivity indexes of COS simulations to N_{leaf} , V_{cmax25} , N_{leaf} and b_{H_2O} are much larger than those of GPP simulations. However, the patterns of the sensitivity of these parameters for COS and GPP simulations are very similar across these sites. Specifically, our results reveal that the simulated COS and GPP are more sensitive to N_{leaf} , while less sensitive to influenced by V_{cmax25} at evergreen needleleaf forest sites (FI-Hyy and US-Wrc). In contrast, at comparison to other sites, the crop site (IT-Soy), role of V_{cmax25} plays a more crucial role in the simulation of COS and GPP, whereas N_{leaf} does the opposite. We also found at IT-Soy is less significant. Additionally, we observed that m_{H_2O} is much more important holds greater importance in the modeling of GPP at US-Wrc than at other sites

and f_{leaf} has a smaller effect on GPP simulations in the savanna site (ES-Lma). Furthermore, Moreover, our results suggest that the modeling of GPP at deciduous broadleaf forest sites (DK-Sor and US-Ha1) are more sensitive to VJ_{slope} while less sensitive to m_{H_2O} than at other sites.






Figure 1.- Sensitivity indexes of the modelled ecosystem COS fluxes (a) and GPP (b) to model parameters.

3.2 Posterior parameter distributions

305 The cumulative frequency distribution as well as the boxplots of each of the parameters for the 0.5 % best runs were plotted in Fig. 42, with a comparison to the uniform parameter distributions and the ~~default parameters prior parameter values~~. As shown in Fig. 42, the posterior distributions of these parameters differ significantly, indicating that the response of these parameters to the assimilation of COS is quite different. Our results demonstrated that COS fluxes have similar constraining effects to the same parameters in different ecosystems although the posterior distributions of the same parameter at different sites depicted variations. In general, parameters related to plant growth and stomatal conductance were strongly constrained by the assimilation of COS, while the parameters related to energy balance as well as soil hydrology were inadequately constrained.

With distinct shape and remarkably narrow range of the cumulative frequency curves, b_{H_2O} (the intercept of the Ball-Berry model, representing minimum stomatal conductance) was strongly ~~constraint~~ constrained by the assimilation of COS in this study. For most sites (AT-Neu, DK-Sor, FI-Hyy, IT-Soy and US-Ha1), the values of b_{H_2O} were confined to a very limited range of 0 to 0.0908 mol m⁻² s⁻¹, ~~with~~. At these five sites, the average value values of the posterior b_{H_2O} ~~less than 0.04~~ all located from 0.01 to 0.03 mol m⁻² s⁻¹. ~~For DK Sor and ES Lma, aligning well with the posterior range width default value~~

of b_{H_2O} are much larger ($0.0175 \text{ mol m}^{-2} \text{ s}^{-1}$) for the BEPS model. By contrast, with values of $0.39 \text{ mol m}^{-2} \text{ s}^{-1}$ and $0.26 \text{ mol m}^{-2} \text{ s}^{-1}$ respectively. With posterior b_{H_2O} values ranging from 0.2403 to 0.9918 and 0.03 to $0.91 \text{ mol m}^{-2} \text{ s}^{-1}$, US-Wrc is the only site where the default value of b_{H_2O} ($0.0175 \text{ mol m}^{-2} \text{ s}^{-1}$) for the BEPS model was rejected by the assimilation of COS at ES-Lma and US-Wrc. Despite the broad distribution of posterior b_{H_2O} at US-Wrc, the cumulative frequency curve still indicates that b_{H_2O} is well-constrained at this site, with 8580 % of the posterior b_{H_2O} located in a narrow range of 0.2415 to $0.50 \text{ mol m}^{-2} \text{ s}^{-1}$. Overall, our results are reasonable as literature-documented values of b_{H_2O} are highly variable and they align well with the compilation provided by Miner et al. (2017), in which more than $83\%5/6$ of the b_{H_2O} values are located between 0 and $0.4518 \text{ mol m}^{-2} \text{ s}^{-1}$, and about half are located between 0 and $0.04 \text{ mol m}^{-2} \text{ s}^{-1}$. Moreover, the mean values of posterior b_{H_2O} for almost ($5/7$) sites are larger than the default b_{H_2O} value of the BEPS model, suggesting that the current b_{H_2O} value utilized in BEPS may be underestimated.


Identified as the most sensitive parameters in COS modeling, the plant growth related parameter V_{cmax25} and N_{leaf} were generally well constrained in this study. However, unlike b_{H_2O} , which is strongly constrained at all sites, the posterior cumulative frequency curves of V_{cmax25} and N_{leaf} exhibit considerable variation across sites. Except for the evergreen needleleaf forest sites US-Ha1 and US-Wrc, the posterior V_{cmax25} and N_{leaf} were mostly distributed in the upper half of the parameter range. Particularly, all of the lower half values of V_{cmax25} and N_{leaf} were excluded by the behavioral parameter sets at DK-Sor and ES-Lma. In contrast, at FI-Hyy, both the posterior cumulative frequency curves of V_{cmax25} and N_{leaf} deviated slightly from the original uniform distribution at US-Ha1, indicating that they are not well-constrained by the assimilation of COS. Notably, the assimilation of COS effectively excluded the upper half As for US-Wrc, both the largest 7% and smallest 4% values of N_{leaf} in US-Wrc, while not for V_{cmax25} , are effectively excluded by the assimilation of COS.

Another stomatal conductance-related parameter, m_{H_2O} , demonstrated effective constraint through COS assimilation at specific sites (AT-Neu, DK-Sor, ES-Lma, FI-Hyy, and IT-Soy), with parameter range width reductions comparable to V_{cmax25} and N_{leaf} . However, at the remaining sites US-Ha1 and US-Wrc, the posterior cumulative frequency curves of m_{H_2O} show minimal deviation from the original uniform distribution. Nevertheless, the optimization of m_{H_2O} is generally achievable through COS assimilation, as supported by our results in good agreement with the compilation of Miner et al. (2017), in which the average historical values of m_{H_2O} grouped by PFT (referred to as the PFT-grouping values below) are provided. As indicated in Table 23, the average absolute bias between the default m_{H_2O} and the PFT-grouping value reached as high as 2.87 for these sites. Through COS assimilation, the mean absolute bias was reduced to 2.43 . In particular, the disparities between the mean posterior m_{H_2O} and the grouping value are consistently smaller than those between the default value and the grouping value, with the exception of DK-Sor and US-Wrc. However, it is noteworthy that our optimization results for DK-Sor and US-Wrc remain plausible, as they fall within one standard deviation (SD) of the grouping values 59.

Table 23. Mean posterior m_{H_2O} values for seven study sites in comparison with the default values and the PFT ~~grouping values (mean \pm standard deviation) (Miner et al., 2017) grouping values (mean \pm standard deviation) in Miner et al. (2017).~~ Within the compilation of Miner et al. (2017), FI-Hyy and US-Wrc are classified under the PFT of evergreen gymnosperm tree, while DK-Sor and US-Ha1 fall under the PFT of deciduous angiosperm tree.

Site name	AT-Neu	DK-Sor	ES-Lma	FI-Hyy	IT-Soy	US-Ha1	US-Wrc
default Default	8	8	8	8	8	8	8
This study	8.266.41	11.469.53	10.6937	5.8913	9.6333	8.6500	8.557.76
Miner 2017	13.3 \pm 3.1	8.7 \pm 5.1	13.3 \pm 3.1	6.7 \pm 2.5	13.5 \pm 3.1	8.7 \pm 5.1	6.7 \pm 2.5

The photosynthesis-related parameters VJ_{slope} and f_{leaf} also influence COS simulation. ~~In behavioral parameter sets for DK-Sor and ES-Lma, excessively small values (less than 0.2) for f_{leaf} were effectively rejected.~~ However, the posterior distributions of f_{leaf} resemble the original uniform distribution, suggesting that it is not a crucial parameter for COS simulations. The posterior cumulative frequency curve of VJ_{slope} also generally deviates slightly from the uniform distribution. Yet, at DK-Sor and US-Ha1, more than ~~three-quarters~~two-thirds of the posterior VJ_{slope} values are situated in the upper half of the parameter range, indicating that VJ_{slope} can also be well-constrained by the assimilation of COS in specific cases. Among these seven sites, the soil hydrology-related parameters $Ksat_{scalar}$ ~~and b_{scalar} and r_{decay}~~ did not exhibit a strong response during the assimilation of COS. However, the posterior cumulative frequency curves of ~~these parameters~~ r_{decay} show notable deviations from the uniform distribution in certain cases. ~~Specifically, at FI-Hyy~~At US-Wrc, higher values of ~~b_{scalar} , $Ksat_{scalar}$, and r_{decay}~~ are more prevalent within the behavioral parameter sets, leading to the posterior ~~means for these parameters that are all~~mean of r_{decay} much greater than the prior ~~means~~mean. Moreover, the largest 14 % values of r_{decay} are effectively excluded by the assimilation of COS at IT-Soy.






Figure 2-3. Cumulative frequency distributions and boxplots for the posterior model parameters- obtained by COS assimilation. The grey area represents a uniform parameter distribution and the black square represents distributions, while the colored areas denote posterior CDF

distributions, with parameters for different sites represented using different colors. The box extends from the default first quartile to the third quartile of the parameter values, with a line at the median. "x" markers denote outliers, and the whiskers represent the lowest or highest parameter values excluding any outliers. The black squares represent the prior parameter values, and the axis ranges denote the prior ranges of the parameters. PI denotes parameter identifiability, defined as the reduction of the parameter range width.

3.3 The optimization performance in COS fluxes

The posterior simulated COS fluxes were evaluated against the prior simulations and observations. Table 34 lists the mean RMSEs and range widths of the prior and posterior simulated COS fluxes for all the sites. The $RMSE_{mean}$ of the posterior COS simulations are smaller than the prior, and the mean RMSE reduction for all sites is $32.09\% \pm 13.49\%$ (mean \pm SD). At the same time, the simulation range widths of COS fluxes are also well constrained, with a mean reduction of $76.42\% \pm 13.31\% \pm 7.32\%$ from the prior. The reduction of $RMSE_{mean}$ and range width is particularly significant in US-Wrc, with a value of 61.39% and 96.77% , respectively.

In Fig. 24, the daily and monthly variations of COS during the observation period at each site are shown. It can be observed that both the prior and posterior simulations are able to accurately capture the daily variation or the seasonal cycle of COS across these sites, with the exception of IT-Soy. As IT-Soy is a temporary observatory with no continuous in-situ meteorological observations available, the ERA-5 meteorological data were used to drive the model for this site, resulting in the simulation not being able to characterize the COS changes very well. Although the simulations perform well in modeling the variations of COS for other sites, our results also suggest that they tend to underestimate the magnitude of COS fluxes at both ends of the growing season (e.g. Fig. 24d). Furthermore, the model markedly underestimates the magnitude of COS during rainy days (DOY 126-134) at ES-Lma (Fig. 3e4c). These findings suggest substantial deficiencies in modeling the mechanistic process of COS exchange. Nevertheless, it can be stated that the fusion of COS observations with the BEPS model has the capacity in constraining the predictive uncertainty of COS, as evidenced by significantly reduced uncertainty bounds that largely encapsulate observations.

The prior simulations ~~tend to~~ significantly underestimate the COS fluxes at ~~DK-Sor and~~ ES-Lma, with the ensemble mean of prior simulations being only about ~~half one-third of~~ that of the observations. After ~~calibration optimization~~, the simulated COS fluxes show a substantial increase and generally align with the observations. However, some observed peaks are still not included in the posterior simulation uncertainty bounds. In contrast, the prior simulations tend to overestimate COS fluxes at ~~the evergreen needleleaf forest sites, especially site FI-Hyy, US-Wrc, where Ha1 and US-Wrc. At US-Wrc and US-Ha1,~~ the ensemble ~~mean means~~ of prior simulations are 78.43% and 64.81% larger than the observations. The assimilation of COS ~~also~~ effectively corrected the overestimation but, at the same time, led to a ~~further underestimates~~ light underestimation of the ~~magnitude of variation in the~~ simulated COS for US-Wrc. ~~In addition, with~~ With the down-regulation of COS simulations, the model-observation difference at both ends of the growing season for FI-Hyy further increased. Particularly, significant underestimation is found in the posterior simulations in 2017 for FI-Hyy, despite ~~remarkable improvement is attached by~~ the posterior simulations showing a remarkable improvement in reproducing COS fluxes over the entire period (2013-2017). As the prior simulations neither noticeably overestimate nor underestimate, there is little difference between the ensemble mean

of the prior and posterior simulations at the remaining three sites (AT-Neu, ~~DK-Sor~~ and IT-Soy ~~and US-Ha1~~). Nevertheless, the assimilation of COS resulted in a remarkable reduction in both $RMSE_{mean}$ and uncertainty bounds for COS simulations at these sites, with mean reductions of ~~24.25~~23.93 % and ~~79.43~~75.11 %, respectively.


405 Overall, there are considerable uncertainties in the prior simulations, with the uncertainty bounds comparable to or much larger than the uncertainties of observations, and the ensemble mean ~~deviate remarkable~~strongly deviates from observations in some sites, i.e., ~~DK-Sor~~ES-Lma. Our results suggest that significant improvement in both the ensemble mean and predictive uncertainty of COS simulations can be achieved through the addition of the information provided by the COS observations with the Monte Carlo-based parameter ~~calibration~~optimization approach, especially for evergreen needleleaf forest sites.

410 However, limited by various factors, such as uncertainty in model-driven data and model structure (Cho et al., 2023), currently the model often underestimates the simulation at both ends of the growing season, and lacks proficiency in modeling the magnitude of COS during rainy days.

Table 34. Comparison of model performance indices for the prior and posterior COS simulations. The ~~$RMSE_{mean}$~~ $RMSE_{mean}$ of the prior and posterior simulations are the mean values of the RMSEs of 20,000 prior COS simulations and 100 behavioral COS simulations with

415 COS observations, respectively. The range widths of the prior and posterior COS simulations are defined as the mean values of the difference between the 95th and 5th percentile of the prior and posterior simulations, respectively. The reduction (%) of $RMSE_{mean}$ and range width is calculated as $(1 - \text{posterior}/\text{prior}) \times 100$.

Site name	$RMSE_{mean}$ (pmol m ⁻² s ⁻¹)			Range width (pmol m ⁻² s ⁻¹)		
	Prior	Posterior	Reduction (%)	Prior	Posterior	Reduction (%)
AT-Neu	16.01 <u>24.10</u>	11.69 <u>14.30</u>	27.01 <u>40.67</u>	22.79 <u>46.40</u>	5.59 <u>6.79</u>	75.45 <u>85.36</u>
DK-Sor	32.03 <u>369</u>	22.09 <u>24.07</u>	31.04 <u>26.36</u>	23.85 <u>45.41</u>	10.54 <u>13.33</u>	55.81 <u>70.64</u>
ES-Lma	12.96 <u>17.10</u>	10.42 <u>14.66</u>	19.62 <u>14.26</u>	6.85 <u>10.35</u>	2.76 <u>75</u>	59.68 <u>73.47</u>
FI-Hyy	17.35 <u>15.87</u>	10.4387	39.84 <u>31.52</u>	35.85 <u>20.96</u>	5.59 <u>3.88</u>	84.40 <u>81.50</u>
IT-Soy	16.8449	13.22 <u>11.35</u>	21.53 <u>31.16</u>	22.58 <u>27.26</u>	4.60 <u>5.12</u>	79.61 <u>81.21</u>
US-Ha1	20.24 <u>30.08</u>	17.02 <u>15.34</u>	24.22 <u>13.44</u>	43.88 <u>50.47</u>	7.35 <u>8.42</u>	83.2431
US-Wrc	36.1276	13.95 <u>14.28</u>	61.3914	70.17 <u>78.04</u>	2.27 <u>4.13</u>	96.77 <u>94.70</u>



420




Figure 34. Comparison of prior and posterior simulated ecosystem COS fluxes. The ensemble means of the prior (red) and posterior (blue) simulations are plotted around the uncertainty bounds (5th and 95th quantile). The mean observed COS and its uncertainty (± 1 estimated by the standard deviation) are represented by black dots with error bars. The means and uncertainties of these hourly observations and simulations are calculated and plotted on a daily or monthly scale. Particularly, error bars are not plotted when more than 3/4 of the observations are missing. The subplot numbers are assigned based on the alphabetical order of the site names.

425

3.4 The performance of simulated GPP

The mean RMSEs and range widths of both prior and posterior simulated GPP for all sites are presented in Table 45. With reduction ratios of $RMSE_{mean}$ ranging from ~~18.99~~20.16 % to ~~66.64~~12 %, the assimilation of COS effectively enhanced the model performance of GPP to varying degrees. Concurrently, the range widths of GPP simulations were well confined, exhibiting a mean reduction ratio of ~~51.83 % ± 65.81 % ± 6.77 %~~17.02 %. ~~Notably, the~~ maximum reduction in ~~both~~ $RMSE_{mean}$ ~~and range width~~ for GPP occurred at US-Wrc, aligning with the substantial improvement observed in the posterior simulated COS at this site. In contrast, a relatively limited impact on improving the prediction of GPP was observed at ~~AT-~~
~~Neu~~FI-Hyy, as evidenced by both the smaller reduction in $RMSE_{mean}$ and range width of GPP simulations.


The BEPS model demonstrated excellent performance in capturing the daily variation and seasonal cycle of GPP, as illustrated in Fig. 35. However, similar to the COS simulations, the ensemble averages of the prior simulated GPP notably deviated from observations at several sites. ~~Specifically~~For example, at DK-Sor, and ES-Lma, the ensemble averages of the prior simulated GPP were only approximately half of the observations. After the assimilation of COS, GPP simulations exhibited a significant increase, aligning well with observations at DK-Sor and ES-Lma. Conversely, substantial overestimation in prior GPP simulations was effectively corrected through the assimilation of COS at US-Wrc, resulting in a remarkable enhanced modeling performance in both RMSE and range width. For FI-Hyy and US-Ha1, minimal differences were observed between the ensemble mean of prior and posterior simulations, as the ensemble mean of prior simulated GPP had already consistently fit the observations. Nevertheless, our results highlight notable enhancements in the predictive uncertainty of GPP through COS assimilation: at these two sites. In Fig. ~~34~~5d, it is evident that, likely due to the absence of in situ meteorological data at IT-Soy, GPP trends are not well represented, although the ensemble averages of the GPP simulations are very close to the observations in magnitude. However, with a reduction of range width as high as ~~60.39~~74.72 %, our finding ~~suggests~~suggests that the assimilation of COS can significantly reduce the predictive uncertainty of GPP, despite the presence of substantial meteorological data uncertainty.

Table 45. Comparison of model performance indices for the prior and posterior GPP simulations. The $RMSE_{mean}$ of the prior and posterior simulations are the mean values of the RMSEs of 20,000 prior GPP simulations and 100 behavioral GPP simulations with GPP observations, respectively. The range widths of the prior and posterior GPP simulations are defined as the mean values of the difference between the 95th and 5th percentiles of the prior and posterior simulations, respectively. The reduction (%) of $RMSE_{mean}$ and range width is calculated as $(1 - \text{posterior}/\text{prior}) * 100$.

Site name	$RMSE_{mean}$ ($\mu\text{mol}\mu\text{mol m}^{-2} \text{ s}^{-1}$)			Range width ($\mu\text{mol}\mu\text{mol m}^{-2} \text{ s}^{-1}$)		
	Prior	Posterior	Reduction (%)	Prior	Posterior	Reduction (%)
AT-Neu	11.48 <u>13.52</u>	9.30 <u>48</u>	18.99 <u>29.90</u>	17.52 <u>27.14</u>	11.84 <u>8.41</u>	32.40 <u>69.01</u>
DK-Sor	16.51 <u>15.39</u>	7.51 <u>08</u>	54.52 <u>00</u>	18.04 <u>19.65</u>	12.14 <u>7.19</u>	32.71 <u>63.39</u>
ES-Lma	6.70 <u>7.35</u>	5.27 <u>4.63</u>	21.43 <u>37.06</u>	10.55 <u>11.99</u>	7.09 <u>4.75</u>	32.75 <u>60.39</u>
FI-Hyy	5.49 <u>14</u>	3.60 <u>4.10</u>	34.34 <u>20.16</u>	27.50 <u>8.51</u>	9.65 <u>3.98</u>	64.92 <u>53.20</u>
IT-Soy	10.93 <u>98</u>	7.59 <u>19</u>	30.60 <u>34.57</u>	20.54 <u>92</u>	8.14 <u>5.29</u>	60.39 <u>74.72</u>

US-Ha1	<u>6.748.05</u>	<u>5.304.50</u>	<u>21.4344.14</u>	<u>31.6412.09</u>	<u>10.713.51</u>	<u>66.1470.97</u>
US-Wrc	<u>19.0817.76</u>	6.37	<u>66.64.12</u>	<u>41.8940.98</u>	<u>11.0912.70</u>	<u>73.5369.00</u>

455






Figure 45. Comparison of prior and posterior simulated GPP. The ensemble means of the prior (red) and posterior (greenblue) simulations are plotted around the uncertainty bounds (5th and 95th quantile). The mean observed GPP and its uncertainty (± 1 -estimated by the standard deviation) are represented by black dots with error bars. The means and uncertainties of these hourly observations and simulations are calculated and plotted on a daily or monthly scale. Particularly, error bars are not plotted when more than 3/4 of the observations are missing. The subplot numbers are assigned based on the alphabetical order of the site names.

460

4 Discussion

4.1 Parameter sensitivity

465 As mentioned before, here we utilize the conductance analog model proposed by Berry et al. (2013) to simulate COS plant uptake. Thus, it is not ~~surprised~~surprising that both the stomatal conductance related parameter b_{H_2O} and m_{H_2O} would impact the modeling of COS flux. ~~Specifically, considering~~Considering the stress of soil moisture on stomatal conductance, the stomatal conductance was calculated by a modified version (Woodward et al., 1995; Ju et al., 2010) of the Ball-Berry model (Ball et al., 1987) based on the close relationship of stomatal conductance and photosynthesis rate. Consequently, both the soil
470 hydrology related parameters and the photosynthesis related parameters can ultimately play roles in the simulation of COS plant uptake by ~~influence~~influencing the modeling of the stomatal conductance.

It has been recognized that the photosynthetic capacity simulated by terrestrial ecosystem models is highly sensitive to V_{cmax} , J_{max} , and light conditions (Zaehle et al., 2005; Bonan et al., 2011; Rogers, 2014; Sargsyan et al., 2014; Koffi et al., 2015; Rogers et al., 2017; Xing et al., 2023). Our study corroborates these findings, highlighting the pronounced sensitivity of
475 simulated GPP to V_{cmax25} , followed by VJ_{slope} and f_{leaf} . ~~Specifically, our results reveal that GPP simulations exhibit lower sensitivity to f_{leaf} at ES Lma, where ample light is available but vegetation is sparse (El Madany et al., 2018; Spielmann et al., 2019), whereas at DK Sor and US Ha1, GPP simulations are more sensitive to VJ_{slope} , attributed to the dense vegetation cover (Wehr et al., 2017; Braendholt et al., 2018; Spielmann et al., 2019). Furthermore~~Moreover, our results reveal that the COS simulations are not notably sensitive to f_{leaf} and VJ_{slope} while V_{cmax25} plays a crucial role in the modeling of COS. It is
480 because V_{cmax25} not only affects the estimation of stomatal conductance through photosynthesis, but also is used to characterize the apparent conductance for COS uptake from the intercellular airspaces, as both mesophyll conductance and carbonic anhydrase activity tend to scale with V_{cmax} (Badger and Price, 1994; Evans et al., 1994; Berry et al., 2013). Yet, as the hydrolysis reaction of COS by carbonic anhydrase is not dependent on light, VJ_{slope} and f_{leaf} do not play any roles in the modeling of apparent conductance and thus have little effect on the simulation of COS.


485 As the COS plant uptake and photosynthesis are tightly coupled through stomata, one would naturally expect similar sensitivity in simulated COS and GPP to stomatal conductance related parameters m_{H_2O} and b_{H_2O} . However, the relationship between COS and stomatal conductance significantly differs from that ~~of~~between GPP and stomatal conductance within the model (e.g., Eq. ~~A185~~ and the Ball-Berry model). Consequently, a notable difference in sensitivity between simulated GPP and COS to m_{H_2O} and b_{H_2O} was identified in this study. Specifically, m_{H_2O} exhibited more pronounced effects on photosynthesis, while
490 b_{H_2O} played a crucial role in the simulation of COS.

Given that a significant portion of nitrogen is invested in the photosynthetic machinery (Mu and Chen, 2021), there exists a close association between leaf nitrogen content and leaf photosynthetic capacity (Sage and Percy, 1987). Additionally, the well-established relationship between leaf nitrogen content and carboxylation capacity (Kattge et al., 2009; Lu et al., 2022) further emphasizes this connection. ~~In specific~~Specifically, carboxylation capacity in leaf scale is assumed to be linearly related

495 to leaf nitrogen content in the BEPS model (Medlyn et al., 1999; Chen et al., 2012). Consequently, both V_{cmax25} and N_{leaf} play crucial roles in influencing carboxylation capacity, thus having a substantial impact on the simulation of COS. The soil hydrology related parameters can also affect the simulation of COS plant flux as we take the stress effect of soil moisture on both stomatal conductance and mesophyll conductance into account (Ju et al., 2010; Knauer et al., 2020). These parameters also affect the modeling of COS soil exchange since soil moisture is a significant factor in COS soil biotic flux (Whelan et al., 2016). However, given the comparatively smaller magnitude of soil COS exchange compared to plant uptake (Whelan et al., 2018) and the minimal impact of soil moisture stress on photosynthetic capacity (Ma et al., 2022), (Ju et al., 2010; Knauer et al., 2020). These parameters also affect the modeling of COS soil exchange since soil moisture is a significant factor in COS soil biotic flux (Whelan et al., 2016). However, given the smaller magnitude of soil COS exchange compared to plant uptake (Whelan et al., 2018) and the minimal impact of soil moisture stress on photosynthetic capacity (Ma et al., 2022), these soil hydrology relevant parameters do not significantly influence the modeling of COS.

4.2 Parameter interactions

For all seven sites, Pearson correlation coefficients and ~~confidence~~ confidence levels between the selected parameters were calculated, as depicted in Fig. 5 and Fig. C1.6. Generally, each site exhibits approximately 43 to 108 parameter combinations with significant correlations ($p < 0.05$). A total of 148 parameter combinations demonstrate significantly correlated significant correlations at more than one site, while 1011 parameter combinations exhibit significant correlations at only one site. Specifically, with a mean correlation coefficient of -0.4855 ± 0.2114 (negative value representing a negative correlation), the correlations between V_{cmax25} and N_{leaf} are very significant ($p < 0.01$) at almost all sites (6/7), indicating a robust interaction between them. In addition to V_{cmax25} and N_{leaf} , four parameter combinations show highly significant correlations ($p < 0.01$) at a minimum of two sites, they are b_{H_2O} and f_{leaf} , b_{H_2O} and m_{H_2O} , m_{H_2O} and V_{cmax25} , m_{H_2O} and VJ_{slope} , N_{leaf} and VJ_{slope} , respectively. Such results indicate the strong interactions among plant growth related parameters, even if some of them do not significantly impact the modeling of COS. In contrast, with a maximum absolute value of 0.26 for the correlation coefficient, soil hydrology related parameters do not exhibit highly significantly correlations with any parameters (except at DK Sor), indicating their weak equivalence (Wu et al., 2019). m_{H_2O} , m_{H_2O} and V_{cmax25} , m_{H_2O} and N_{leaf} , b_{scalar} and VJ_{slope} respectively. Such results indicate the strong interactions among parameters related to stomatal conductance, photosynthesis as well as soil hydrology, even if some of them do not significantly impact the modeling of COS.






Figure 56. Parameter correlation matrix plots with significance levels between the parameters of the behavioral parameter sets **at AT-Neu (a) and DK-Sor (b)**. Correlation coefficients are shown in red font when the confidence level is greater than 95 % ($p < 0.05$), with a superscript "*" indicating the confidence level greater than 99 % ($p < 0.01$).

We ~~also~~ observed substantial variations in parameter interactions across different sites. For instance, at ~~IT-Soy, b_{H_2O}~~ AT-Neu, ~~m_{H_2O} and $f_{leaf}N_{leaf}$~~ exhibited a highly significant negative correlation with a correlation coefficient as high as ~~-0.4945~~. However, these two parameters seemed irrelevant at US-WreDK-Sor with a correlation coefficient of only ~~-0.0203~~. As for soil hydrology related parameters, none of them showed significant correlations with any parameter at IT-Soy, US-Ha+FI-Hyy and ~~US-WreHa1~~, yet there were four parameter combinations related to them significantly correlated at DK-Sor (Fig. ~~5b~~ 6b). Furthermore, while V_{cmax25} is highly correlated with N_{leaf} at all sites, the correlation coefficients varied considerably, ranging from ~~-0.2441~~ to ~~-0.9387~~.

We also observed interactions not only between two parameters but also among several parameters (e.g.: ~~b_{H_2O} - f_{leaf} - m_{H_2O} - b_{scalar} - V_{cmax25} - V_{Jslope}~~ in Fig. ~~5a~~). ~~To provide deeper insights into these interactions and highlight significantly correlated parameter combinations, we generated~~ 6a). Fig. ~~6~~. ~~This figure not only elucidates~~ B1 showcases the intricate interactions among multiple parameters relevant to COS simulations and illustrates the distribution of behavioral parameter sets ~~but also captures the intricate interactions among corresponding parameters. Specifically, the~~ The parameter combinations depicted in Fig. ~~6B1~~ are particularly representative as they originate from diverse sites and include nearly all highly significant correlated combinations. Overall, since ~~at~~ the six plant growth related parameters used in this study are positively correlated with the simulation of COS, they consistently constrain each other, demonstrating a negative correlation, as shown in Fig. 6a, Fig. 6c, and Fig. 6d. However, due to the nonlinearity of the model, there is not a simple linear relationship between these parameters. For example, at ES-LmaAT-Neu, where the COS observations ~~significantly surpass~~ notably exceed the ensemble mean of prior simulations, V_{cmax25} and V_{Jslope} ~~are not linearly correlated with each other~~ exhibit a non-linear correlation, but ~~are~~ both well confined tend to the region around be distributed near their upper limits (Fig. ~~6b~~ B1).




Figure 6. Scatter plots showing the behavioral parameter sets in 3D parameter space at AT-Neu (a), ES-Lma (b) and US-Ha1 (c) and US-Wre.

4.3 Parameter identifiability

As the parameter identifiability is quantified based on the range of the behavioral parameter, its results were presented in Fig. 23 along with the plots of the cumulative likelihood distributions of the behavioral parameters. These results underscore the remarkable ability of COS assimilation to identify b_{H_2O} , with a mean PI of b_{H_2O} as high as 0.7681 ± 0.2528 . Identified as the most sensitive parameters for COS modeling, V_{cmax25} and N_{leaf} also exhibit remarkable identifiability, with mean PIs of 0.39 ± 0.25 and 0.4229 ± 0.19 and 0.26 ± 0.22 respectively. The other three plant growth related parameters (m_{H_2O} , VJ_{slope} and f_{leaf}) m_{H_2O} demonstrate varying levels of identifiability, with PIs ranging from 0.0601 to 0.42 , 0.03 to 0.30 , and 0 to 0.09 , respectively. While these 28. In contrast, the light reaction related parameters VJ_{slope} and f_{leaf} are not well identified, with the

maximum value of PI of only 0.04. The soil hydrology related parameters are well identified at some sites, they prove almost b_{scalar} , $Ksat_{scalar}$ and r_{decay} are also generally unidentifiable at others. Notably, VJ_{slope} achieves optimal identification at DK-Sor, where r_{decay} is well identified at IT-Soy, in which its PI value (0.14) is approximately eightseven times that of the other sites.


560 The In this study, the identifiability of a parameter with input data closely ~~correlates with~~ related to the sensitivity of the input data COS simulations to the parameter, although it ~~can be known to~~ is influenced by model over-parameterization and parameter interactions (Gan et al., 2014). For instance, our results demonstrate at ES-Lma, where the orders of COS simulations exhibited the greatest sensitivity to N_{leaf} and V_{cmax25} , these parameters were also found to have the highest identifiability for V_{cmax25} and N_{leaf} are almost consistent at the same site (Fig. 22a and Fig. 3). Given the high sensitivity of COS modeling to V_{cmax25} ,
565 N_{leaf} and b_{H_2O} , it is unsurprising that these parameters can be effectively identified by the assimilation of COS. However, our findings indicate that the sensitivity of V_{cmax25} , N_{leaf} is much greater than that of b_{H_2O} , yet the latter is much more identifiable. This outcome can be attributed to the highly significant correlation between V_{cmax25} and N_{leaf} , as parameter interaction is a primary contributor to parameter unidentifiability (Gan et al., 2014).

In Sect. 3.1, it was demonstrated that the modeling of COS exhibits a low sensitivity to f_{leaf} , m_{H_2O} and VJ_{slope} . Consequently,
570 it is reasonable that the assimilation of COS may did not effectively identify f_{leaf} , m_{H_2O} and VJ_{slope} at certain sites (Fig. 23). However, due to their significant correlations with other plant growth-related parameters, effective identification is possible in specific cases. Notably, combinations such as f_{leaf} - N_{leaf} , m_{H_2O} - V_{cmax25} - b_{H_2O} and m_{H_2O} - V_{cmax25} - VJ_{slope} are very significantly correlated (Fig. 5b6b), and both N_{leaf} - b_{H_2O} and V_{cmax25} were effectively identified at DK-Sor FI-Hyy. As a result, f_{leaf} , m_{H_2O} and VJ_{slope} also attain highly identifiability at this site.

575 It has been previously demonstrated that soil hydrology-related parameters exert a minimal impact on COS simulations (Fig. 2) and cannot be effectively constrained through COS assimilation in general (Fig. 3). Consequently, these parameters exhibit low identifiability, although significant combinations of correlations associated with soil hydrology-related parameters were observed at certain sites (e.g., DK-Sor).

4.4 Relationship between COS and GPP simulation ~~performance~~ performances

580 In this study, we identified the top 100 parameter sets, whose corresponding simulations displayed the smallest RMSE concerning COS observations, as the behavioral parameter sets. Subsequently, these behavioral parameter sets were employed to derive the posterior simulated COS and GPP, and to estimate prediction uncertainty. Therefore, it is necessary to investigate how the distribution of RMSEs for COS simulations and GPP simulations respond to RMSE, and to understand the relationship between the model performance of COS and that of GPP.



585

Figure 7. Comparison of RMSE for COS ($RMSE_{COS}$) and GPP ($RMSE_{GPP}$) in the Monte Carlo simulations. Each data point represents a parameter set, with color indicating data density. The gray dashed line represents the RMSE threshold for COS ~~simulations~~, calculated as the mean of the 100th and 101st smallest values of the RMSE.

590

In Fig.7, scatter plots of RMSEs for COS and GPP are presented. It can be observed that at most sites, where the scatters are most densely distributed, there tend to be relatively small RMSEs for both COS and GPP. These results indicate that the current model is generally capable to simulate COS and GPP well at the same time. ~~However, given the uncertainties of model parameters, structure and driving data etc., instances like at ES Lma arise where numerous parameter combinations~~

simultaneously exhibit a small RMSE for COS, but their RMSE values for GPP show a significant range of variation. Moreover, due to the noticeable differences in the mechanisms of COS and GPP, such as that the photosynthetic rate in the Farquhar photosynthesis model (Farquhar et al., 1980) is limited either by the RuBP (ribulose 1,5 bisphosphate) carboxylation or by the RuBP regeneration (Hikosaka et al., 2006), while the current modeling of COS is insensitive to the parameters related to RuBP regeneration (e.g. $V_{j\text{stompe}}$ and f_{leaf}), numerous parameter sets of these sites. However, given the distinct mechanisms of COS and GPP as well as the uncertainties of model structure and driving data etc., there are also numerous parameter sets that perform well for either COS or GPP but exhibit significant discrepancies with the observations of the other. More specifically, for example, the model runs with the 3 % highest RMSE for GPP instead exhibit good performance in terms of COS at ES-Lma, with their mean RMSE values ($15.42 \text{ pmol m}^{-2} \text{ s}^{-1}$) less than that of the prior ($17.10 \text{ pmol m}^{-2} \text{ s}^{-1}$). Overall, our results suggest that these behavioral parameter sets, which demonstrate good performance in COS simulation, also generally perform well in modeling GPP. However, the parameter sets with relatively good GPP simulation results exhibit significant variability in the performance of COS modeling.

4.5 Caveats and implication

Compared to the big leaf model, the two-leaf model has been demonstrated to better describe the canopy radiation distribution, GPP, and stomatal conductance (Luo et al., 2018). In this study, we take the advantage of two-leaf model to simulate COS fluxes from plant and soil based on the BEPS model within the two-leaf framework. Ecosystem COS flux data were used to calibrate the model parameters belong to BEPS and to optimize GPP simulations among diverse ecosystems within the Monte Carlo-based methodology. Our results demonstrate that COS not only improves the accuracy of GPP simulations but also reduces GPP simulation uncertainty. However, due to the lack of in-situ COS concentration and flux data, as well as BEPS model driving data (e.g., meteorological data, LAI data and clumping index data), we are still facing challenges in evaluating the performance of the two-leaf model compared to other models in COS simulation. issue. Therefore, there is an urgent need for more in situ meteorological data, vegetation canopy structural parameters, as well as COS observations.

The spatial and temporal variation in atmospheric COS concentrations has a considerable influence on the COS plant uptake (Ma et al., 2021; Kooijmans et al., 2021) due to the linear relationship between the two (Stimler et al., 2010). With the lack of continuous ground-based COS concentration observations, COS concentrations in the bulk air are assumed to be spatially invariant over the globe but to vary annually in this study, which may lead to significant biases in COS simulations. Currently, several recent studies have simulated COS vegetation fluxes based on atmospheric transport model-derived COS concentration data within the big-leaf framework (Kooijmans et al., 2021; Maignan et al., 2021; Abadie et al., 2023). These COS fluxes simulated based on big-leaf models were in turn used to drive atmospheric transport models (Remaud et al., 2023; Ma et al., 2023). Within an atmosphere inversion framework, recent studies indicate an underestimation of the biosphere COS sink in high-latitude regions of the Northern Hemisphere (NH) (Ma et al., 2021; Remaud et al., 2023). Larger underestimations of ecosystem COS exchange based on big-leaf model at high latitudes have also been confirmed at the site scale, and the underestimations of COS are consistent with biases in GPP for some sites (Kooijmans et al., 2021). Interestingly, Luo et al.

(2018) demonstrated that the reason for the underestimation of GPP by the big-leaf model is that it fails to accurately describe the instantaneous radiation distribution in the canopy, and the underestimation increases with the increase of LAI. The NH high-latitude regions have relatively high LAI (Fang et al., 2019), therefore the deficiency of the big leaf model in simulating radiation distribution may contribute to the existence of the missing COS sink in the NH high latitude in summer, and this deficiency is amplified by the larger LAI. In fact, the spatial distribution of LAI (i.e., GLOBMAP LAI) retrieved through remote sensing not only in NH high-latitude regions but also in central Africa aligns with the spatial distribution of the missing sink revealed by the “objective” inversion conducted by Ma et al. (2021) (as illustrated in Figure 7 in Ma et al. (2021)), which further validates the reasonableness of this hypothesis. Therefore, conducting COS simulations under the two-leaf framework at a global scale holds the promise of providing insights into the global COS vegetation sink and benefiting the simulation of the spatial and temporal distribution of COS concentrations. Thus, it is necessary to conduct regional and global COS simulations within the two-leaf model framework in the future.

Taking advantage of the Monte Carlo-based parameter optimization approach, we analyzed the global sensitivity, identifiability as well as interactions of COS-related parameters in this study. Furthermore, we quantified the uncertainty in simulated COS and GPP, thereby revealing the capacity of COS to constrain the uncertainty in GPP simulations. However, the Monte Carlo-based parameter optimization approach is subject to controversy (Sambridge and Mosegaard, 2002) due to the numerous subjective decisions involved in its implementation, such as the selection of parameter range, sample size and performance metric, etc. Further research is needed to investigate the impact of these factors on the parameter optimization results related to COS and the assessment of model prediction uncertainty.

645 5 Conclusions

In this study, carbonyl sulfide flux data were utilized to calibrate the ecosystem model parameters and to optimize GPP simulations among various ecosystems within the Monte Carlo-based methodology base on the coupling of approach using COS modeling and the within BEPS model. Global. A global parameter sensitivity analysis was conducted to identify the most sensitive ones among a set of 9 pre-selected parameters for COS and GPP modeling, the. The identifiability and interaction of model parameters were investigated by the behavioral parameter sets. The effectiveness of COS in improving the model performance of GPP was evaluated. The major findings are as follows:

- (1) Similar to GPP, we found the modeling of COS is sensitive to parameters V_{cmax25} and N_{leaf} and b_{H_2O} , while insensitive to soil hydrology related parameters, as well as the energy related parameter f_{leaf} . Unlike GPP, COS is also sensitive to b_{H_2O} while being insensitive to m_{H_2O} and the light reaction related parameters VJ_{slope} and $f_{leaf}VJ_{slope}$.
- (2) The assimilation of COS within the Monte Carlo-based approach effectively improved model performance of GPP and significantly reduced the model predictive uncertainty, with a mean RMSE reduction of $35.42\% \pm 17.01$ $40.56\% \pm 13.77\%$ and a mean range width reduction as high as $51.83\% \pm 17.02$ $65.81\% \pm 6.77\%$.

(3) Complex and significant two-parameter or multi-parameter interactions ~~exists~~ exist between the model parameters. ~~In particularly~~ Particularly, V_{cmax25} and N_{leaf} ~~are significantly correlated~~ show highly significant correlations ($p < 0.0501$) at all
660 sites.

(4) Generally, b_{H_2O} , V_{cmax25} and N_{leaf} can be well identified through the assimilation of COS, especially b_{H_2O} , whereas the soil hydrology related parameters ~~and the light-reaction related parameters~~ cannot be identified effectively. ~~m_{H_2O} , $V_{J_{stop}}$ and f_{leaf} can also be effectively identified due to strong parameter interactions.~~

665 Appendix A:

A1- BEPS photosynthesis and stomatal conductance modeling approach

In the BEPS model, the net photosynthesis rate (A) is calculated using the Farquhar model (Farquhar et al., 1980; Chen et al., 1999):

$$A = \min(A_i, A_j) - R_d \quad (A1)$$

$$A_c = V_{cmax} \frac{C_i - \Gamma_i^*}{C_i + K_c \left(1 + \frac{O_i}{K_o}\right)} \quad (A2)$$

$$A_j = J \frac{C_i - \Gamma_i^*}{4(C_i - 2\Gamma_i^*)} \quad (A3)$$

where A_i and A_j are Rubisco-limited and RuBP-limited gross photosynthetic rates ($\mu\text{mol m}^{-2}\text{s}^{-1}$), respectively. R_d is leaf dark respiration ($\mu\text{mol m}^{-2}\text{s}^{-1}$). V_{cmax} is the maximum carboxylation rate of Rubisco ($\mu\text{mol m}^{-2}\text{s}^{-1}$); J is the electron transport rate ($\mu\text{mol m}^{-2}\text{s}^{-1}$); C_i and O_i are the intercellular carbon dioxide (CO_2) and oxygen (O_2) concentrations
675 (mol mol^{-1}), respectively; K_c and K_o are Michaelis–Menten constants for CO_2 and O_2 (mol mol^{-1}), respectively.

The electron transport rate, J, is dependent on incident photosynthetic photon flux density (PPFD, $\mu\text{mol m}^{-2}\text{s}^{-1}$) as:

$$J = \frac{J_{max} I}{I + 2.1J_{max}} \quad (A4)$$

where J_{max} is the maximum electron transport rate ($\mu\text{mol m}^{-2}\text{s}^{-1}$), I is the incident PPFD calculated from the incident shortwave radiation R_{SW} (W m^{-2}):

$$I = \beta R_{SW} f_{leaf} \quad (A5)$$

where $\beta = 4.55$ is the energy – quanta conversion factor ($\mu\text{mol J}^{-1}$), f_{leaf} is the ratio of photosynthesis active radiation to the shortwave radiation (unitless).

~~The maximum carboxylation rate of Rubisco V_{cmax} was calculated according the Arrhenius temperature function and the maximum carboxylation rate of Rubisco at 25 °C (V_{cmax25}). V_{cmax} is generally proportional to leaf nitrogen content. Considering both the fractions of sunlit and shaded leaf areas to the total leaf area and the leaf nitrogen content vary with the depth into the canopy, the V_{cmax} values of sunlit ($V_{cmax,sun}$) and shaded ($V_{cmax,sh}$) leaves can be obtained through vertical integrations with respect to leaf area index (L) (Chen et al., 2012):~~

$$V_{cmax,sun} = V_{cmax} \chi_n N_{leaf} \frac{k[1 - e^{-(k_n+k)L}]}{(k_n + k)(1 - e^{-kL})} \quad (A6)$$

$$V_{cmax,sh} = V_{cmax} \chi_n N_{leaf} \frac{\frac{1}{k_n}[1 - e^{-k_n L}] - \frac{1}{k_n+k}[1 - e^{-(k_n+k)L}]}{L - \frac{1}{k}(1 - e^{-kL})} \quad (A7)$$

690 The maximum carboxylation rate of Rubisco, V_{cmax} , was calculated according to the modified Arrhenius temperature function (Medlyn et al., 2002) and the maximum carboxylation rate of Rubisco at 25 °C (V_{cmax25}). V_{cmax} is generally proportional to leaf nitrogen content. Considering both the fractions of sunlit and shaded leaf areas to the total leaf area and the leaf nitrogen content vary with the depth into the canopy, the V_{cmax} values of sunlit ($V_{cmax,sunlit}$) and shaded ($V_{cmax,shaded}$) leaves can be obtained through vertical integrations with respect to canopy depth (Chen et al., 2012; De Pury and Farquhar, 1997):

$$695 \quad V_{cmax,sunlit} = V_{cmax} \chi_n N_{leaf} \frac{k[1 - e^{-(k_n+k)L}]}{(k_n + k)(1 - e^{-kL})} \quad (A6)$$

$$V_{cmax,shaded} = V_{cmax} \chi_n N_{leaf} \frac{\frac{1}{k_n}[1 - e^{-k_n L}] - \frac{1}{k_n+k}[1 - e^{-(k_n+k)L}]}{L - \frac{1}{k}(1 - e^{-kL})} \quad (A7)$$

where χ_n ($m^2 g^{-1}$) is the relative change of V_{cmax} to leaf nitrogen content; N_{leaf} ($g m^{-2}$) is the leaf nitrogen content at the top of the canopy; k_n (unitless) is the leaf nitrogen content decay rate with increasing depth into the canopy, taken as 0.3; L is the canopy depth described in total LAI. k is calculated as:

$$700 \quad k = G(\theta)\Omega \cos(\theta) \quad (A8)$$

where $G(\theta)$ is the projection coefficient, taken as 0.5; Ω is the clumping index, and θ is the solar zenith angle.

After V_{cmax} values for the representative sunlit and shaded leaves are obtained, the maximum electronic transport rate for the sunlit and shaded leaves are obtained from Medlyn et al. (1999):

$$J_{max} = VJ_{slope} V_{cmax} - 14.2 \quad (A9)$$

705 where VJ_{slope} (unitless) is the slope of the relationship of V_{cmax} and J_{max} .

The leaf stomatal conductance to water vapor (g_{sw} in $\mu mol m^{-2} s^{-1}$) is estimated using a modified version of Ball-Berry (BB) empirical model (Ball et al., 1987) following Woodward et al. (1995):

$$g_{sw} = b_{H_2O} + \frac{m_{H_2O} A R_h f_w}{C_a} \quad (A10)$$

710 where b_{H_2O} is the intercept of the Ball-BerryBB model, representing the minimum g_{sw} ($mol m^{-2} s^{-1}$), m_{H_2O} is the empirical slope parameter in the BB model (unitless), R_h is the relative humidity at the leaf surface (unitless), f_w is a soil moisture stress factor describing the sensitivity of g_{sw} to soil water availability (Ju et al., 2006), C_a is the atmospheric CO_2 concentration ($\mu mol mol^{-1}$).

Soil water availability factor $f_{w,i}$ in each layer i is calculated as:

$$f_{w,i} = \frac{1.0}{f_i(\psi_i) f_i(T_{s,i})} \quad (A11)$$

715 where $f_i(\psi_i)$ is a function of matrix suction ψ_i (m) (Zierl, 2001), $f_i(T_{s,i})$ is a function describing the effect of soil temperature ($T_{s,i}$ in °C) on soil water uptake (Bonan, 1991).

To consider the variable soil water potential at different depths, the scheme of Ju et al. (2006) was employed to calculate the weight of each layer (w_i) to f_w :

$$w_i = \frac{R_i f_{w,i}}{\sum_{i=1}^n R_i f_{w,i}} \quad (\text{A12})$$

720 where n is the number of soil ~~layer~~layers (five were used in this study) of the BEPS model, R_i is the root fraction in layer i , calculated as:

$$R_i = \begin{cases} 1 - r_{decay}^{100cd_i} & i = 1 \\ r_{decay}^{100cd_{i-1}} - r_{decay}^{100cd_i} & 1 < i < n \\ r_{decay}^{100cd_{i-1}} & i = n \end{cases} \quad (\text{A13})$$

where cd_i is the cumulative ~~thickness~~depth (m) of layer i . In this study, each soil layer depth (from top to bottom) of the BEPS model is 0.05 m, 0.10 m, 0.20 m, 0.40 m and 1.25 m, respectively.

725 The overall soil water availability f_w is then calculated as:

$$f_w = \sum_{i=1}^n f_{w,i} w_i \quad (\text{A14})$$

The hydraulic conductivity of each soil layer K_i (m s^{-1}) is expressed as:

$$K_i = Ksat_i \left(\frac{SWC_i}{\theta_{s,i}} \right)^{2b_i+3} \quad (\text{A15})$$

730 where $Ksat_i$ is the saturated hydrological conductivity of soil layer i (m s^{-1}); SWC_i is the volumetric liquid soil water content of soil layer i ($\text{m}^3 \text{m}^{-3}$); $\theta_{s,i}$ is the porosity of soil layer i (unitless); b_i is the Campbell parameter for soil layer i , determining the change rate of hydraulic conductivity with SWC (unitless). In this study, $Ksat_i$ and b_i are expressed as:

~~$$Ksat_i = Ksat_{scalar} Ksat_{df,i} \quad (\text{A16})$$~~

~~$$b_i = b_{scalar} b_{df,i} \quad (\text{A17})$$~~

$$Ksat_i = Ksat_{scalar} Ksat_{df,i} \quad (\text{A16})$$

$$735 \quad b_i = b_{scalar} b_{df,i} \quad (\text{A17})$$

where $Ksat_{df,i}$ and $b_{df,i}$ are the default values of $Ksat_i$ and b_i respectively.

A2 -BEPS leafsoil COS modeling approach

The leaf level COS uptake rate $F_{COS,leaf}$ is determined by the formula:

~~$$F_{COS,leaf} = COS_a \left(\frac{1.94}{g_{sw}} + \frac{1.56}{g_{bw}} + \frac{1}{g_{COS}} \right)^{-1} \quad (\text{A18})$$~~

740 ~~g_{sw} are the stomatal conductance and leaf laminar boundary layer conductance to H_2O vapor. g_{COS} indicates the apparent~~

conductance for COS uptake from the intercellular airspaces, which combined the mesophyll conductance and the biochemical reaction rate of COS and carbonic anhydrase. It can be calculated as:

$$g_{COS} = 1.4 * 10^3 * (1.0 + 5.33 * F_{LAI}) * 10^{-6} * (1 - e^{(-0.45 * LAI)}) f_w V_{emax} \quad (A19)$$

f_w is a parameter describing soil water stress.

745 The total soil COS flux $F_{COS,soil}$ is the sum of abiotic COS flux $F_{COS,abiotic}$ and biotic COS flux $F_{COS,biotic}$.

$$F_{COS,soil} = F_{COS,abiotic} + F_{COS,biotic} \quad (A20)$$

$$F_{COS,soil} = F_{COS,abiotic} + F_{COS,biotic} \quad (A18)$$

Here, we take the approach developed in Whelan et al. (2016) for the modeling of $F_{COS,soil}$. In this approach, $F_{COS,abiotic}$ is described as an exponential function of the temperature of soil T_{soil} ($^{\circ}C$).

750
$$F_{COS,abiotic} = \alpha e^{\beta T_{soil}} \quad (A21)$$

where α and β were parameters determined using the least squares fitting approach. We assigned the values of α and β to BEPS according to the parameterizations scheme of Abadie et al. (2022).

$$F_{COS,biotic} = \alpha e^{\beta T_{soil}} \quad (A19)$$

755 where α and β were parameters determined using the least-squares fitting approach. We assigned the values of α and β to BEPS according to the parameterizations scheme of Whelan et al. (2016).

$F_{COS,biotic}$ is described as the product of a power function and an exponential function.

$$F_{COS,biotic} = F_{opt} \left(\frac{SWC}{SWC_{opt}} \right)^{-a} e^{-a \left(\frac{SWC}{SWC_{opt}} - 1 \right)} \quad (A22)$$

$$a = \ln \left(\frac{F_{opt}}{F_{SWC_g}} \right) \left(\ln \left(\frac{SWC_{opt}}{SWC_g} \right) + \left(\frac{SWC_g}{SWC_{opt}} - 1 \right) \right)^{-1} \quad (A23)$$

760
$$F_{COS,biotic} = F_{opt} \left(\frac{SWC}{SWC_{opt}} \right)^{-a} e^{-a \left(\frac{SWC}{SWC_{opt}} - 1 \right)} \quad (A20)$$

$$a = \ln \left(\frac{F_{opt}}{F_{SWC_g}} \right) \left(\ln \left(\frac{SWC_{opt}}{SWC_g} \right) + \left(\frac{SWC_g}{SWC_{opt}} - 1 \right) \right)^{-1} \quad (A21)$$

Here a is the curve shape constant. The maximum biotic COS uptake F_{opt} and the biotic COS uptake F_{SWC_g} are the COS fluxes ($pmol\ m^{-2}\ s^{-1}$) at optimum soil water content SWC_{opt} and a secondary soil water content SWC_g , and $SWC_g > SWC_{opt}$.


765 A more detailed description of the modeling of $F_{COS,biotic}$ and the parameterization scheme adopted in this study can be found in Whelan et al. (2022).

Appendix B: Additional [tablefigure](#)

Table B1. Descriptions of the 9 parameters were selected to be calibrated. The default values and prior ranges (in parentheses) of these parameters are given for each plant function type (PFT) or for each soil texture or globally according to the parameter dependent.

Parameter	Description	Dependent	Default value and prior range				
			ENE/Loamy sand	DBF/Loam	Grass/Silty loam	Crop/Sandy clay-loam	Silty clay
b_{H_2O}	The intercept of the Ball-Berry model ($\text{mol m}^{-2}\text{s}^{-1}$)	PFT	0.0175 (0.004375-1)	0.0175 (0.004375-1)	0.0175 (0.004375-1)	0.0175 (0.004375-1)	
$b_{scat_{aF}}$	The scaling factor of Campbell parameter b	Texture	1 (0.25-0.75)	1 (0.25-0.75)	1 (0.25-0.75)	1 (0.25-0.75)	1 (0.25-0.75)
f_{leaf}	Ratio of photosynthetically active radiation to shortwave radiation	Global			0.5 (0.125-1)		
$K_{sat_{scat_{aF}}}$	The scaling factor of saturated hydraulic conductivity K_{sat}	Texture	1 (0.25-0.75)	1 (0.25-0.75)	1 (0.25-0.75)	1 (0.25-0.75)	1 (0.25-0.75)
m_{H_2O}	The slope of the Ball-Berry model	PFT	8 (2-14)	8 (2-14)	8 (2-14)	8 (2-14)	
N_{leaf}	Leaf nitrogen content ($\text{m}^2 \text{g}^{-1}$)	PFT	4.45 (1-7.7875)	2.45 (1-4.2875)	2.695 (1-4.71625)	2.375 (1-4.15625)	
r_{decay}	Decay rate of root distribution	PFT	0.95 (0.2375-0.99)	0.97 (0.2425-0.99)	0.96 (0.24-0.99)	0.95 (0.2375-0.99)	
$V_{em_{ax25}}$	Maximum carboxylation rate of Rubisco at 25 °C ($\mu\text{mol m}^{-2}\text{s}^{-1}$)	PFT	62.5 (15-75)	57.7 (15-75)	48 (15-75)	84.5 (15-120)	
VJ_{slope}	Slope of the $V_{em_{ax}}$ and J_{max} (maximum electron transport rate) relationship	PFT	2.39 (1-5)	2.39 (1-5)	2.39 (1-5)	2.39 (1-5)	

770 **Appendix C: Additional figure**






Figure C1. Parameter correlation matrix B1. Scatter plots with significance levels between the parameters of showing the behavioral parameter sets at ES-Lma in 3D parameter space at AT-Neu (a), FI-Hyy (b), IT-Soy (c), and US-Ha1 (d) and US-Wrc. The scatter colors represent the magnitude of the corresponding parameters using RGB values. The projection of the scatter is illustrated with smaller markers. (e). Correlation coefficients are shown in red font when the correlation confidence level is greater than 95 % ($p < 0.05$), with a superscript "*" indicating a confidence level greater than 99 % ($p < 0.01$).

Data availability. Measured eddy covariance carbonyl sulfide fluxes data can be found at <https://zenodo.org/record/3406990> for AT-Neu, DK-Sor, ES-Lma and IT-Soy, <https://zenodo.org/record/6940750> for FI-Hyy, and from the Harvard Forest Data Archive under record HF214 (<https://portal.edirepository.org/nis/mapbrowse?packageid=knb-lter-hfr.214.4>) for US-Ha1. The raw COS concentration data of US-Wrc can be obtained at <https://zenodo.org/record/1422820>. The meteorological data can be obtained from the FLUXNET database (<https://fluxnet.org/>) for AT-Neu, DK-Sor, ES-LMa, FI-Hyy and US-Ha1; from the

AmeriFlux database (<https://ameriflux.lbl.gov/>) for US-Ha1 (except shortwave radiation data) and US-Wrc; from the ERA5 dataset (<https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels?tab=overview>) for AT-Neu, IT-Soy and US-Ha1. The GPP data can be obtained from the FLUXNET database for DK-Sor, ES-LMa, FI-Hyy and US-Ha1; from the AmeriFlux database for US-Ha1 ~~and US-Wrc; and;~~ from <https://zenodo.org/record/6940750> for AT-Neu and IT-Soy. ~~The;~~ and from <https://zenodo.org/record/1422820> for US-Wrc. The GLASS LAI is available at <ftp://ftp.glcfc.umd.edu/> and the GLOBMAP LAI is available at <https://zenodo.org/record/4700264#.YzvSYnZBxD8%2F>. ~~the GLASS LAI is available at~~ <ftp://ftp.glcfc.umd.edu/>. ~~the MODIS LAI product is available at <https://lpdaac.usgs.gov/products/mod15a2hv006/>.~~

Author contributions.

MW designed the experiments and developed the model. XX improved the model and performed the Monte Carlo simulations. HZ made the analysis and wrote the original manuscript ~~and made the analysis~~. All the authors contributed to the writing of the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This study was supported by the National Natural Science Foundation of China (42371486, 42111530184), the Research Funds for the Frontiers Science Center for Critical Earth Material Cycling, Nanjing University (Grant No: 090414380031, 020914380115). We acknowledge Prof. Tim Moore from McGill University to help us in improving the language.

Reference

References

- 805 Abadie, C., Maignan, F., Remaud, M., Kohonen, K. M., Sun, W., Kooijmans, L., Vesala, T., Seibt, U., Raoult, N., and Bastrikov, V.: Carbon and water fluxes of the boreal evergreen needleleaf forest biome constrained by assimilating ecosystem carbonyl sulfide flux observations, *Journal of Geophysical Research: Biogeosciences*, e2023JG007407, 2023.
- Abadie, C., Maignan, F., Remaud, M., Ogée, J., Campbell, J. E., Whelan, M. E., Kitz, F., Spielmann, F. M., Wohlfahrt, G., and Wehr, R.: Global modelling of soil carbonyl sulfide exchanges, *Biogeosciences*, 19, 2427-2463, 2022.
- 810 Asaf, D., Rotenberg, E., Tatarinov, F., Dicken, U., Montzka, S. A., and Yakir, D.: Ecosystem photosynthesis inferred from measurements of carbonyl sulphide flux, *Nature Geoscience*, 6, 186-190, 2013.
- Badger, M. R. and Price, G. D.: The role of carbonic anhydrase in photosynthesis, *Annual review of plant biology*, 45, 369-392, 1994.
- Ball, J. T., Woodrow, I. E., and Berry, J. A.: A model predicting stomatal conductance and its contribution to the control of photosynthesis under different environmental conditions, *Progress in photosynthesis research: volume 4 proceedings of the VIIth international congress on photosynthesis providence, Rhode Island, USA, august 10–15, 1986*, 221-224,
- 815 [Bao, S., Ibrom, A., Wohlfahrt, G., Koirala, S., Migliavacca, M., Zhang, Q., and Carvalhais, N.: Narrow but robust advantages in two-big-leaf light use efficiency models over big-leaf light use efficiency models at ecosystem level, *Agricultural and Forest Meteorology*, 326, 109185, 2022.](#)

- 820 Berry, J., Wolf, A., Campbell, J. E., Baker, I., Blake, N., Blake, D., Denning, A. S., Kawa, S. R., Montzka, S. A., and Seibt, U.: A coupled model of the global cycles of carbonyl sulfide and CO₂: A possible new window on the carbon cycle, *Journal of Geophysical Research: Biogeosciences*, 118, 842-852, 2013.
- Beven, K. and Binley, A.: The future of distributed models: model calibration and uncertainty prediction, *Hydrological processes*, 6, 279-298, 1992.
- 825 Beven, K. and Binley, A.: GLUE: 20 years on, *Hydrological processes*, 28, 5897-5918, 2014.
- Beven, K. and Freer, J.: Equifinality, data assimilation, and uncertainty estimation in mechanistic modelling of complex environmental systems using the GLUE methodology, *Journal of hydrology*, 249, 11-29, 2001.
- ~~Billesbach, D., Berry, J., Seibt, U., Maseyk, K., Torn, M., Fischer, M., Abu Naser, M., and Campbell, J.: Growing season eddy covariance measurements of carbonyl sulfide and CO₂ fluxes: COS and CO₂ relationships in Southern Great Plains winter wheat, *Agricultural and Forest Meteorology*, 184, 48-55, 2014.~~
- 830 Blankenship, R. E.: *Molecular mechanisms of photosynthesis*, John Wiley & Sons 2021.
- Blasone, R.-S., Vrugt, J. A., Madsen, H., Rosbjerg, D., Robinson, B. A., and Zyvoloski, G. A.: Generalized likelihood uncertainty estimation (GLUE) using adaptive Markov Chain Monte Carlo sampling, *Advances in Water Resources*, 31, 630-648, 2008.
- 835 Bonan, G. B.: A biophysical surface energy budget analysis of soil temperature in the boreal forests of interior Alaska, *Water Resources Research*, 27, 767-781, 1991.
- Bonan, G. B., Lawrence, P. J., Oleson, K. W., Levis, S., Jung, M., Reichstein, M., Lawrence, D. M., and Swenson, S. C.: Improving canopy processes in the Community Land Model version 4 (CLM4) using global flux fields empirically inferred from FLUXNET data, *Journal of Geophysical Research: Biogeosciences*, 116, 2011.
- 840 Borgonovo, E.: A new uncertainty importance measure, *Reliability Engineering & System Safety*, 92, 771-784, 2007.
- ~~Braendholt, A., Ibrom, A., Larsen, K. S., and Pilegaard, K.: Partitioning of ecosystem respiration in a beech forest, *Agricultural and Forest Meteorology*, 252, 88-98, 2018.~~
- Canadell, J., Mooney, H., Baldocchi, D., Berry, J., Ehleringer, J., Field, C., Gower, S. T., Hollinger, D., Hunt, J., and Jackson, R. B.: Commentary: carbon metabolism of the terrestrial biosphere: a multitechnique approach for improved understanding, *Ecosystems*, 3, 115-130, 2000.
- 845 Chen, B., Wang, P., Wang, S., Ju, W., Liu, Z., and Zhang, Y.: Simulating canopy carbonyl sulfide uptake of two forest stands through an improved ecosystem model and parameter optimization using an ensemble Kalman filter, *Ecological Modelling*, 475, 110212, 2023.
- Chen, J., Liu, J., Cihlar, J., and Goulden, M.: Daily canopy photosynthesis model through temporal and spatial scaling for remote sensing applications, *Ecological modelling*, 124, 99-119, 1999.
- 850 Chen, J. M., Ju, W., Ciais, P., Viovy, N., Liu, R., Liu, Y., and Lu, X.: Vegetation structural change since 1981 significantly enhanced the terrestrial carbon sink, *Nature communications*, 10, 4259, 2019.
- Chen, J. M., Mo, G., Pisek, J., Liu, J., Deng, F., Ishizawa, M., and Chan, D.: Effects of foliage clumping on the estimation of global terrestrial gross primary productivity, *Global Biogeochemical Cycles*, 26, 2012.
- 855 Cho, A., Kooijmans, L. M., Kohonen, K.-M., Wehr, R., and Krol, M. C.: Optimizing the carbonic anhydrase temperature response and stomatal conductance of carbonyl sulfide leaf uptake in the Simple Biosphere model (SiB4), *Biogeosciences*, 20, 2573-2594, 2023.
- Commane, R., Meredith, L. K., Baker, I. T., Berry, J. A., Munger, J. W., Montzka, S. A., Templer, P. H., Juice, S. M., Zahniser, M. S., and Wofsy, S. C.: Seasonal fluxes of carbonyl sulfide in a midlatitude forest, *Proceedings of the National Academy of Sciences*, 112, 14162-14167, 2015.
- 860 ~~El Madany, T. S., Reichstein, M., Perez Priego, O., Carrara, A., Moreno, G., Martín, M. P., Pacheco Labrador, J., Wohlfahrt, G., Nieto, H., and Weber, U.: Drivers of spatio-temporal variability of carbon dioxide and energy fluxes in a Mediterranean savanna ecosystem, *Agricultural and Forest Meteorology*, 262, 258-278, 2018.~~
- ~~De Pury, D. and Farquhar, G.: Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf models, *Plant, Cell & Environment*, 20, 537-557, 1997.~~
- 865 Evans, J. R., Caemmerer, S., Setchell, B. A., and Hudson, G. S.: The relationship between CO₂ transfer conductance and leaf anatomy in transgenic tobacco with a reduced content of Rubisco, *Functional Plant Biology*, 21, 475-495, 1994.
- ~~Fang, H., Baret, F., Plummer, S., and Schaepman-Strub, G.: An overview of global leaf area index (LAI): Methods, products, validation, and applications, *Reviews of Geophysics*, 57, 739-799, 2019.~~

- 870 Farquhar, G. D., von Caemmerer, S. v., and Berry, J. A.: A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species, *planta*, 149, 78-90, 1980.
- Friedlingstein, P., O'sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Le Quéré, C., Luijkx, I. T., Olsen, A., and Peters, G. P.: Global carbon budget 2022, *Earth System Science Data Discussions*, 2022, 1-159, 2022.
- 875 Gan, Y., Duan, Q., Gong, W., Tong, C., Sun, Y., Chu, W., Ye, A., Miao, C., and Di, Z.: A comprehensive evaluation of various sensitivity analysis methods: A case study with a hydrological model, *Environmental modelling & software*, 51, 269-285, 2014.
- ~~Goldan, P. D., Fall, R., Kuster, W. C., and Fehsenfeld, F. C.: Uptake of COS by growing vegetation: A major tropospheric sink, *Journal of Geophysical Research: Atmospheres*, 93, 14186-14192, 1988.~~
- 880 Gu, L., Baldocchi, D., Verma, S. B., Black, T., Vesala, T., Falge, E. M., and Dowty, P. R.: Advantages of diffuse radiation for terrestrial ecosystem productivity, *Journal of Geophysical Research: Atmospheres*, 107, ACL 2-1-ACL 2-23, 2002.
- ~~Haynes, K., Baker, I., and Denning, S.: Simple biosphere model version 4.2 (SiB4) technical description, Colorado State University: Fort Collins, CO, USA, 2020.~~
- 885 He, H., Jansson, P.-E., Svensson, M., Meyer, A., Klemedtsson, L., and Kasimir, Å.: Factors controlling Nitrous Oxide emission from a spruce forest ecosystem on drained organic soil, derived using the CoupModel, *Ecological Modelling*, 321, 46-63, 2016.
- He, Q., Ju, W., Dai, S., He, W., Song, L., Wang, S., Li, X., and Mao, G.: Drought risk of global terrestrial gross primary productivity over the last 40 years detected by a remote sensing-driven process model, *Journal of Geophysical Research: Biogeosciences*, 126, e2020JG005944, 2021.
- 890 ~~Hikosaka, K., Ishikawa, K., Borjigidai, A., Muller, O., and Onoda, Y.: Temperature acclimation of photosynthesis: mechanisms involved in the changes in temperature dependence of photosynthetic rate, *Journal of experimental botany*, 57, 291-302, 2006.~~
- Hilton, T. W., Whelan, M. E., Zumkehr, A., Kulkarni, S., Berry, J. A., Baker, I. T., Montzka, S. A., Sweeney, C., Miller, B. R., and Elliott Campbell, J.: Peak growing season gross uptake of carbon in North America is largest in the Midwest USA, *Nature Climate Change*, 7, 450-454, 2017.
- 895 Houska, T., Multsch, S., Kraft, P., Frede, H.-G., and Breuer, L.: Monte Carlo-based calibration and uncertainty analysis of a coupled plant growth and hydrological model, *Biogeosciences*, 11, 2069-2082, 2014.
- Hu, L., Montzka, S. A., Kaushik, A., Andrews, A. E., Sweeney, C., Miller, J., Baker, I. T., Denning, S., Campbell, E., and Shiga, Y. P.: COS-derived GPP relationships with temperature and light help explain high-latitude atmospheric CO₂ seasonal cycle amplification, *Proceedings of the National Academy of Sciences*, 118, e2103423118, 2021.
- 900 Iwanaga, T., Usher, W., and Herman, J.: Toward SALib 2.0: Advancing the accessibility and interpretability of global sensitivity analyses, *Socio-Environmental Systems Modelling*, 4, 18155-18155, 2022.
- ~~Jackson, R. B., Canadell, J., Ehleringer, J. R., Mooney, H. A., Sala, O. E., and Schulze, E.-D.: A global analysis of root distributions for terrestrial biomes, *Oecologia*, 108, 389-411, 1996.~~
- 905 Ju, W., Gao, P., Wang, J., Zhou, Y., and Zhang, X.: Combining an ecological model with remote sensing and GIS techniques to monitor soil water content of croplands with a monsoon climate, *Agricultural Water Management*, 97, 1221-1231, 2010.
- Ju, W., Chen, J. M., Black, T. A., Barr, A. G., Liu, J., and Chen, B.: Modelling multi-year coupled carbon and water fluxes in a boreal aspen forest, *Agricultural and Forest Meteorology*, 140, 136-151, 2006.
- 910 ~~Karu, E., Li, M., Ernle, L., Brenninkmeijer, C. A., Lelieveld, J., and Williams, J.: Carbonyl Sulfide (OCS) in the upper troposphere/Lowermost stratosphere (UT/LMS) region: Estimates of lifetimes and fluxes, *Geophysical Research Letters*, 50, e2023GL105826, 2023.~~
- Kattge, J., Knorr, W., Raddatz, T., and Wirth, C.: Quantifying photosynthetic capacity and its relationship to leaf nitrogen content for global-scale terrestrial biosphere models, *Global Change Biology*, 15, 976-991, 2009.
- 915 Kesselmeier, J., Teusch, N., and Kuhn, U.: Controlling variables for the uptake of atmospheric carbonyl sulfide by soil, *Journal of Geophysical Research: Atmospheres*, 104, 11577-11584, 1999.
- ~~Knauer, J., Zaehle, S., De Kauwe, M. G., Haverd, V., Reichstein, M., and Sun, Y.: Mesophyll conductance in land surface models: effects on photosynthesis and transpiration, *The Plant Journal*, 101, 858-873, 2020.~~

- 920 Koffi, E., Rayner, P., Norton, A., Frankenberg, C., and Scholze, M.: Investigating the usefulness of satellite-derived fluorescence data in inferring gross primary productivity within the carbon cycle data assimilation system, *Biogeosciences*, 12, 4067-4084, 2015.
- Kohonen, K.-M., [Kolari, P., Kooijmans, L. M., Chen, H., Seibt, U., Sun, W., and Mammarella, I.: Towards standardized processing of eddy covariance flux measurements of carbonyl sulfide, *Atmospheric Measurement Techniques*, 13, 3957-3975, 2020.](#)
- 925 [Kohonen, K.-M., Dewar, R., Tramontana, G., Mauranen, A., Kolari, P., Kooijmans, L. M., Papale, D., Vesala, T., and Mammarella, I.: Intercomparison of methods to estimate gross primary production based on CO₂ and COS flux measurements, *Biogeosciences*, 19, 4067-4088, 2022.](#)
- Kooijmans, L. M., Sun, W., Aalto, J., Erkkilä, K.-M., Maseyk, K., Seibt, U., Vesala, T., Mammarella, I., and Chen, H.: Influences of light and humidity on carbonyl sulfide-based estimates of photosynthesis, *Proceedings of the National Academy of Sciences*, 116, 2470-2475, 2019.
- 930 Kooijmans, L. M. J., Cho, A., Ma, J., Kaushik, A., Haynes, K. D., Baker, I., Luijckx, I. T., Groenink, M., Peters, W., Miller, J. B., Berry, J. A., Ogee, J., Meredith, L. K., Sun, W., Kohonen, K. M., Vesala, T., Mammarella, I., Chen, H., Spielmann, F. M., Wohlfahrt, G., Berkelhammer, M., Whelan, M. E., Maseyk, K., Seibt, U., Commane, R., Wehr, R., and Krol, M.: Evaluation of carbonyl sulfide biosphere exchange in the Simple Biosphere Model (SiB4), *Biogeosciences*, 18, 6547-6565, 10.5194/bg-18-6547-2021, 2021.
- 935 Launois, T., Peylin, P., Belviso, S., and Poulter, B.: A new model of the global biogeochemical cycle of carbonyl sulfide–Part 2: Use of carbonyl sulfide to constrain gross primary productivity in current vegetation models, *Atmospheric Chemistry and Physics*, 15, 9285-9312, 2015.
- Liu, J., Chen, J., Cihlar, J., and Park, W.: A process-based boreal ecosystem productivity simulator using remote sensing inputs, *Remote sensing of environment*, 62, 158-175, 1997.
- 940 Liu, Y., Liu, R., and Chen, J. M.: Retrospective retrieval of long-term consistent global leaf area index (1981–2011) from combined AVHRR and MODIS data, *Journal of Geophysical Research: Biogeosciences*, 117, 2012.
- Liu, Y., Xiao, J., Ju, W., Zhou, Y., Wang, S., and Wu, X.: Water use efficiency of China's terrestrial ecosystems and responses to drought, *Scientific reports*, 5, 13799, 2015.
- Liu, Z., Zhou, Y., Ju, W., and Gao, P.: Simulation of soil water content in farm lands with the BEPS ecological model, *Transactions of the Chinese Society of Agricultural Engineering*, 27, 67-72, 2011.
- 945 Lu, X., Wang, Y.-P., Ziehn, T., and Dai, Y.: An efficient method for global parameter sensitivity analysis and its applications to the Australian community land surface model (CABLE), *Agricultural and forest meteorology*, 182, 292-303, 2013.
- Lu, X., Croft, H., Chen, J. M., Luo, Y., and Ju, W.: Estimating photosynthetic capacity from optimized Rubisco–chlorophyll relationships among vegetation types and under global change, *Environmental Research Letters*, 17, 014028, 2022.
- 950 Luo, ~~Y.: Terrestrial carbon cycle feedback to climate warming, *Annu. Rev. Ecol. Evol. Syst.*, 38, 683-712, 2007.~~ [X., Chen, J. M., Liu, J., Black, T. A., Croft, H., Staebler, R., He, L., Arain, M. A., Chen, B., and Mo, G.: Comparison of big-leaf, two-big-leaf, and two-leaf upscaling schemes for evapotranspiration estimation using coupled carbon-water modeling, *Journal of Geophysical Research: Biogeosciences*, 123, 207-225, 2018.](#)
- [Luo, Y.: Terrestrial carbon cycle feedback to climate warming, *Annu. Rev. Ecol. Evol. Syst.*, 38, 683-712, 2007.](#)
- 955 [Ma, J., Kooijmans, L. M., Cho, A., Montzka, S. A., Glatthor, N., Worden, J. R., Kuai, L., Atlas, E. L., and Krol, M. C.: Inverse modelling of carbonyl sulfide: implementation, evaluation and implications for the global budget, *Atmospheric Chemistry and Physics*, 21, 3507-3529, 2021.](#)
- [Ma, J., Remaud, M., Peylin, P., Patra, P., Niwa, Y., Rodenbeck, C., Cartwright, M., Harrison, J. J., Chipperfield, M. P., and Pope, R. J.: Intercomparison of Atmospheric Carbonyl Sulfide \(TransCom-COS\): 2. Evaluation of Optimized Fluxes Using Ground-Based and Aircraft Observations, *Journal of Geophysical Research: Atmospheres*, 128, e2023JD039198, 2023.](#)
- 960 [Ma, R., Xiao, J., Liang, S., Ma, H., He, T., Guo, D., Liu, X., and Lu, H.: Pixel-level parameter optimization of a terrestrial biosphere model for improving estimation of carbon fluxes with an efficient model–data fusion method and satellite-derived LAI and GPP data, *Geoscientific Model Development*, 15, 6637-6657, 2022.](#)
- 965 MacBean, N., Bacour, C., Raoult, N., Bastrikov, V., Koffi, E., Kuppel, S., Maignan, F., Ottlé, C., Peaucelle, M., and Santaren, D.: Quantifying and reducing uncertainty in global carbon cycle predictions: lessons and perspectives from 15 years of

- data assimilation studies with the ORCHIDEE Terrestrial Biosphere Model, *Global Biogeochemical Cycles*, 36, e2021GB007177, 2022.
- 970 Maignan, F., Abadie, C., Remaud, M., Kooijmans, L. M., Kohonen, K.-M., Commane, R., Wehr, R., Campbell, J. E., Belviso, S., and Montzka, S. A.: Carbonyl sulfide: comparing a mechanistic representation of the vegetation uptake in a land surface model and the leaf relative uptake approach, *Biogeosciences*, 18, 2917-2955, 2021.
- ~~Maseyk, K., Berry, J. A., Billesbach, D., Campbell, J. E., Torn, M. S., Zahniser, M., and Seibt, U.: Sources and sinks of carbonyl sulfide in an agricultural field in the Southern Great Plains, *Proceedings of the National Academy of Sciences*, 111, 9064-9069, 2014.~~
- 975 Medlyn, B. E., Badeck, F. W., De Pury, D., Barton, C., Broadmeadow, M., Ceulemans, R., De Angelis, P., Forstreuter, M., Jach, M., and Kellomäki, S.: Effects of elevated [CO₂] on photosynthesis in European forest species: a meta-analysis of model parameters, *Plant, Cell & Environment*, 22, 1475-1495, 1999.
- ~~Medlyn, B. E., Dreyer, E., Ellsworth, D., Forstreuter, M., Harley, P., Kirschbaum, M., Le Roux, X., Montpied, P., Strassmeyer, J., and Walcroft, A.: Temperature response of parameters of a biochemically based model of photosynthesis. II. A review of experimental data, *Plant, Cell & Environment*, 25, 1167-1179, 2002.~~
- 980 Miner, G. L., Bauerle, W. L., and Baldocchi, D. D.: Estimating the sensitivity of stomatal conductance to photosynthesis: a review, *Plant, Cell & Environment*, 40, 1214-1238, 2017.
- Mo, X., Chen, J. M., Ju, W., and Black, T. A.: Optimization of ecosystem model parameters through assimilating eddy covariance flux data with an ensemble Kalman filter, *Ecological Modelling*, 217, 157-173, 2008.
- 985 ~~Montzka, S., Calvert, P., Hall, B., Elkins, J., Conway, T., Tans, P., and Sweeney, C.: On the global distribution, seasonality, and budget of atmospheric carbonyl sulfide (COS) and some similarities to CO₂, *Journal of Geophysical Research: Atmospheres*, 112, 2007.~~
- Moradkhani, H., Hsu, K. L., Gupta, H., and Sorooshian, S.: Uncertainty assessment of hydrologic model states and parameters: Sequential data assimilation using the particle filter, *Water resources research*, 41, 2005.
- 990 Mu, X. and Chen, Y.: The physiological response of photosynthesis to nitrogen deficiency, *Plant Physiology and Biochemistry*, 158, 76-82, 2021.
- Ogée, J., Sauze, J., Kesselmeier, J., Genty, B., Van Diest, H., Launois, T., and Wingate, L.: A new mechanistic framework to predict OCS fluxes from soils, *Biogeosciences*, 13, 2221-2240, 2016.
- ~~Pignou, C. P., Jaiswal, D., McGrath, J. M., and Long, S. P.: Loss of photosynthetic efficiency in the shade. An Achilles heel for the dense modern stands of our most productive C4 crops?, *Journal of Experimental Botany*, 68, 335-345, 2017.~~
- 995 Plischke, E., Borgonovo, E., and Smith, C. L.: Global sensitivity measures from given data, *European Journal of Operational Research*, 226, 536-550, 2013.
- Protoschill-Krebs, G., Wilhelm, C., and Kesselmeier, J.: Consumption of carbonyl sulphide (COS) by higher plant carbonic anhydrase (CA), *Atmospheric Environment*, 30, 3151-3156, 1996.
- 1000 Raines, C. A.: The Calvin cycle revisited, *Photosynthesis research*, 75, 1-10, 2003.
- Rastogi, B., Berkelhammer, M., Wharton, S., Whelan, M. E., Itter, M. S., Leen, J. B., Gupta, M. X., Noone, D., and Still, C. J.: Large uptake of atmospheric OCS observed at a moist old growth forest: Controls and implications for carbon cycle applications, *Journal of Geophysical Research: Biogeosciences*, 123, 3424-3438, 2018.
- 1005 Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C., Buchmann, N., Gilmanov, T., and Granier, A.: On the separation of net ecosystem exchange into assimilation and ecosystem respiration: review and improved algorithm, *Global change biology*, 11, 1424-1439, 2005.
- ~~Remaud, M., Chevallier, F., Maignan, F., Belviso, S., Berchet, A., Parouffe, A., Abadie, C., Bacour, C., Lennartz, S., and Peylin, P.: Plant gross primary production, plant respiration and carbonyl sulfide emissions over the globe inferred by atmospheric inverse modelling, *Atmospheric Chemistry and Physics*, 22, 2525-2552, 2022.~~
- 1010 ~~Remaud, M., Ma, J., Krol, M., Abadie, C., Cartwright, M. P., Patra, P., Niwa, Y., Rodenbeck, C., Belviso, S., and Kooijmans, L.: Intercomparison of atmospheric carbonyl sulfide (TransCom-COS; part one): Evaluating the impact of transport and emissions on tropospheric variability using ground-based and aircraft data, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037817, 2023.~~
- Rogers, A.: The use and misuse of V_c, max in Earth System Models, *Photosynthesis research*, 119, 15-29, 2014.

- 1015 Rogers, A., Medlyn, B. E., Dukes, J. S., Bonan, G., Von Caemmerer, S., Dietze, M. C., Kattge, J., Leakey, A. D., Mercado, L. M., and Niinemets, Ü.: A roadmap for improving the representation of photosynthesis in Earth system models, *New Phytologist*, 213, 22-42, 2017.
- 1020 [Ryu, Y., Jiang, C., Kobayashi, H., and Detto, M.: MODIS-derived global land products of shortwave radiation and diffuse and total photosynthetically active radiation at 5 km resolution from 2000, *Remote Sensing of Environment*, 204, 812-825, 2018.](#)
- Sage, R. F. and Percy, R. W.: The nitrogen use efficiency of C3 and C4 plants: II. Leaf nitrogen effects on the gas exchange characteristics of *Chenopodium album* (L.) and *Amaranthus retroflexus* (L.), *Plant physiology*, 84, 959-963, 1987.
- [Sambridge, M. and Mosegaard, K.: Monte Carlo methods in geophysical inverse problems, *Reviews of Geophysics*, 40, 3-13-29, 2002.](#)
- 1025 Sandoval-Soto, L., Stanimirov, M., Von Hobe, M., Schmitt, V., Valdes, J., Wild, A., and Kesselmeier, J.: Global uptake of carbonyl sulfide (COS) by terrestrial vegetation: Estimates corrected by deposition velocities normalized to the uptake of carbon dioxide (CO₂), *Biogeosciences*, 2, 125-132, 2005.
- Sargsyan, K., Safta, C., Najm, H. N., Debusschere, B. J., Ricciuto, D., and Thornton, P.: Dimensionality reduction for complex models via Bayesian compressive sensing, *International Journal for Uncertainty Quantification*, 4, 2014.
- 1030 Schwalm, C. R., Williams, C. A., Schaefer, K., Anderson, R., Arain, M. A., Baker, I., Barr, A., Black, T. A., Chen, G., and Chen, J. M.: A model-data intercomparison of CO₂ exchange across North America: Results from the North American Carbon Program site synthesis, *Journal of Geophysical Research: Biogeosciences*, 115, 2010.
- Seibt, U., Kesselmeier, J., Sandoval-Soto, L., Kuhn, U., and Berry, J.: A kinetic analysis of leaf uptake of COS and its relation to transpiration, photosynthesis and carbon isotope fractionation, *Biogeosciences*, 7, 333-341, 2010.
- 1035 [Shaw, D. C., Franklin, J. F., Bible, K., Klopatek, J., Freeman, E., Greene, S., and Parker, G. G.: Ecological setting of the Wind River old-growth forest, *Ecosystems*, 7, 427-439, 2004.](#)
- [Smith, B., Knorr, W., Widlowski, J.-L., Pinty, B., and Gobron, N.: Combining remote sensing data with process modelling to monitor boreal conifer forest carbon balances, *Forest Ecology and Management*, 255, 3985-3994, 2008.](#)
- 1040 Spielmann, F., Wohlfahrt, G., Hammerle, A., Kitz, F., Migliavacca, M., Alberti, G., Ibrom, A., El-Madany, T. S., Gerdel, K., and Moreno, G.: Gross primary productivity of four European ecosystems constrained by joint CO₂ and COS flux measurements, *Geophysical research letters*, 46, 5284-5293, 2019.
- Staudt, K., Falge, E., Pyles, R. D., Paw U, K. T., and Foken, T.: Sensitivity and predictive uncertainty of the ACASA model at a spruce forest site, *Biogeosciences*, 7, 3685-3705, 2010.
- 1045 Stimler, K., [Berry, J. A., and Yakir, D.: Effects of carbonyl sulfide and carbonic anhydrase on stomatal conductance, *Plant Physiology*, 158, 524-530, 2012.](#)
- [Stimler, K., Montzka, S. A., Berry, J. A., Rudich, Y., and Yakir, D.: Relationships between carbonyl sulfide \(COS\) and CO₂ during leaf gas exchange, *New Phytologist*, 186, 869-878, 2010.](#)
- Sun, W., Berry, J. A., Yakir, D., and Seibt, U.: Leaf relative uptake of carbonyl sulfide to CO₂ seen through the lens of stomatal conductance–photosynthesis coupling, *New Phytologist*, 235, 1729-1742, 2022.
- 1050 Sun, W., Maseyk, K., Lett, C., and Seibt, U.: A soil diffusion–reaction model for surface COS flux: COSSM v1, *Geoscientific Model Development*, 8, 3055-3070, 2015.
- [Sun, W., Kooijmans, L. M., Maseyk, K., Chen, H., Mammarella, I., Vesala, T., Levula, J., Keskinen, H., and Seibt, U.: Soil fluxes of carbonyl sulfide \(COS\), carbon monoxide, and carbon dioxide in a boreal forest in southern Finland, *Atmospheric Chemistry and Physics*, 18, 1363-1378, 2018.](#)
- 1055 Tang, J. and Zhuang, Q.: A global sensitivity analysis and Bayesian inference framework for improving the parameter estimation and prediction of a process-based Terrestrial Ecosystem Model, *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- Tonkin, M. and Doherty, J.: Calibration-constrained Monte Carlo analysis of highly parameterized models using subspace techniques, *Water Resources Research*, 45, 2009.
- 1060 Vesala, T., Kohonen, K.-M., Kooijmans, L. M., Praplan, A. P., Foltýnová, L., Kolari, P., Kulmala, M., Bäck, J., Nelson, D., and Yakir, D.: Long-term fluxes of carbonyl sulfide and their seasonality and interannual variability in a boreal forest, *Atmospheric Chemistry and Physics*, 22, 2569-2584, 2022.

- 1065 Wang, J., Jiang, F., Wang, H., Qiu, B., Wu, M., He, W., Ju, W., Zhang, Y., Chen, J. M., and Zhou, Y.: Constraining global terrestrial gross primary productivity in a global carbon assimilation system with OCO-2 chlorophyll fluorescence data, *Agricultural and Forest Meteorology*, 304, 108424, 2021.
- [Wang, S., Ibrom, A., Bauer-Gottwein, P., and Garcia, M.: Incorporating diffuse radiation into a light use efficiency and evapotranspiration model: An 11-year study in a high latitude deciduous forest, *Agricultural and Forest Meteorology*, 248, 479-493, 2018.](#)
- 1070 Wehr, R., Commane, R., Munger, J. W., McManus, J. B., Nelson, D. D., Zahniser, M. S., Saleska, S. R., and Wofsy, S. C.: Dynamics of canopy stomatal conductance, transpiration, and evaporation in a temperate deciduous forest, validated by carbonyl sulfide uptake, *Biogeosciences*, 14, 389-401, 2017.
- Whelan, M. E., Hilton, T. W., Berry, J. A., Berkelhammer, M., Desai, A. R., and Campbell, J. E.: Carbonyl sulfide exchange in soils for better estimates of ecosystem carbon uptake, *Atmospheric Chemistry and Physics*, 16, 3711-3726, 2016.
- 1075 Whelan, M. E., Shi, M., Sun, W., Vries, L. K. d., Seibt, U., and Maseyk, K.: Soil carbonyl sulfide (OCS) fluxes in terrestrial ecosystems: an empirical model, *Journal of Geophysical Research: Biogeosciences*, 127, e2022JG006858, 2022.
- Whelan, M. E., Lennartz, S. T., Gimeno, T. E., Wehr, R., Wohlfahrt, G., Wang, Y., Kooijmans, L. M., Hilton, T. W., Belviso, S., and Peylin, P.: Reviews and syntheses: Carbonyl sulfide as a multi-scale tracer for carbon and water cycles, *Biogeosciences*, 15, 3625-3657, 2018.
- 1080 Wohlfahrt, G., Brilli, F., Hörtnagl, L., Xu, X., Bingemer, H., Hansel, A., and Loreto, F.: Carbonyl sulfide (COS) as a tracer for canopy photosynthesis, transpiration and stomatal conductance: potential and limitations, *Plant, cell & environment*, 35, 657-667, 2012.
- Woodward, F. I., Smith, T. M., and Emanuel, W. R.: A global land primary productivity and phytogeography model, *Global biogeochemical cycles*, 9, 471-490, 1995.
- 1085 Wu, M., Ran, Y., Jansson, P.-E., Chen, P., Tan, X., and Zhang, W.: Global parameters sensitivity analysis of modeling water, energy and carbon exchange of an arid agricultural ecosystem, *Agricultural and Forest Meteorology*, 271, 295-306, 2019.
- Wu, M., Tan, X., Wu, J., Huang, J., Jansson, P.-E., and Zhang, W.: Coupled water transport and heat flux in seasonally frozen soils: uncertainties identification in multi-site calibration, *Environmental Earth Sciences*, 79, 524, 2020.
- 1090 Xiao, Z., Liang, S., Wang, J., Xiang, Y., Zhao, X., and Song, J.: Long-time-series global land surface satellite leaf area index product derived from MODIS and AVHRR surface reflectance, *IEEE Transactions on Geoscience and Remote Sensing*, 54, 5301-5318, 2016.
- Xing, X., Wu, M., Zhang, W., Ju, W., Tagesson, T., He, W., Wang, S., Wang, J., Hu, L., and Yuan, S.: Modeling China's terrestrial ecosystem gross primary productivity with BEPS model: Parameter sensitivity analysis and model calibration, *Agricultural and Forest Meteorology*, 343, 109789, 2023.
- 1095 ~~Yang, F., Qubaja, R., Tatarinov, F., Rotenberg, E., and Yakir, D.: Assessing canopy performance using carbonyl sulfide measurements, *Global Change Biology*, 24, 3486-3498, 2018.~~
- Yi, D. H., Kim, D. W., and Park, C. S.: Parameter identifiability in Bayesian inference for building energy models, *Energy and Buildings*, 198, 318-328, 2019.
- Yuan, W., Liu, S., Zhou, G., Zhou, G., Tieszen, L. L., Baldocchi, D., Bernhofer, C., Gholz, H., Goldstein, A. H., and Goulden, M. L.: Deriving a light use efficiency model from eddy covariance flux data for predicting daily gross primary production across biomes, *Agricultural and Forest Meteorology*, 143, 189-207, 2007.
- 1100 Zaehle, S., Sitch, S., Smith, B., and Hatterman, F.: Effects of parameter uncertainties on the modeling of terrestrial biosphere dynamics, *Global Biogeochemical Cycles*, 19, 2005.
- [Zhu, H., Wu, M., Jiang, F., Vossbeck, M., Kaminski, T., Xing, X., Wang, J., Ju, W., and Chen, J. M.: Assimilation of Carbonyl Sulfide \(COS\) fluxes within the adjoint-based data assimilation system–Nanjing University Carbon Assimilation System \(NUCAS v1. 0\), *EGU*sphere, 2023, 1-35, 2023.](#)
- 1105 Zierl, B.: A water balance model to simulate drought in forested ecosystems and its application to the entire forested area in Switzerland, *Journal of Hydrology*, 242, 115-136, 2001.