

General information

Please consider the following information that we intend to add in the submitted manuscript if we are invited to submit a revised version:

- Kirsten Zickfeld was moved from the last to the second position in the list of authors.
- Figure 11b features a revised curve for WETCHIMP emission intensities after double-checking zonally-averaged wetland areas from each model.
- We included new references on the optimal temperature for soil microbial processes and in particular CH₄ production in wetlands: (i) Schipper et al. (2014), *Global Change Biology*, 20, 3578-3586; (ii) Metje and Frenzel (2007), *Environmental Microbiology*, 9, 954-964.
- We modified the framing and boxing of tables in the manuscript and supplements.

In the remainder of the document, reviewer comments are reproduced in plain text and our responses are given in bold text.

Response to reviewer #1

Summary:

The WETMETH model represents some advances in modelling science, although one of its “novel” components (vertical methane production) is also in other models. Even though WETMETH is to be included in an Earth system model, its detailed validation is highly skewed to the middle and high latitudes. More detailed validation needs to be undertaken in tropical regions. The scheme also needs a more thorough justification and calibration of its methane production optimal temperature formulation. The manuscript is generally well-written and constructed, with appropriate figures and tables.

Thanks for the positive and constructive comments on the manuscript. Below we provide our responses to each comment.

Model design, validation and calibration:

The manuscript describes the three pathways by which methane is produced by methanogens in the soil, with the main pathways utilising either acetate or CO₂. Root exudates provide a large source of acetate, but it is unclear from the WETMETH model description that any of the soil carbon pools include root exudates. If root exudates are included in a soil pool, does this pool have a fast enough turnover time appropriate for their decomposition, or is it combined with other compounds which have slower decomposition rates?

WETMETH does not distinguish root exudates from other organic matter compounds in the soil. That is, (1) WETMETH considers the soil carbon pool as a single pool aggregating all sources of organic matter substrates (e.g. litter fall and root exudates), and (2) there is currently no distinction between fast and slow pathways of CH₄ production in our model. We assume that rates of CH₄ production in WETMETH encompass all possible CH₄ production pathways when combined together.

To clarify these points, we will include the following statements in Section 3.1.1:

- **First paragraph (statement to be included after line 187 of the original text) -> We consider C_i to be the aggregate of all sources of soil carbon (i.e. organic matter) such as litter-fall and root exudates.**
- **Second paragraph (statement to be included after line 195 of the original text) -> In this first version of WETMETH, we combine all possible pathways for CH_4 production in wetlands (see Section 2.1) without distinguishing fast and slow pathways.**

An optimal temperature for methane production is included in WETMETH, with the model being particularly sensitive to the value it is assigned. The formulation used is taken from Wu et al 2016, which is calibrated using data from mid and high latitudes bogs and fens. It is also based on the optimal temperature for soil CO_2 production. Methane production via the acetoclastic pathway is not dependent on CO_2 availability however. So is it valid to use the same optimum temperature for methane production? Also is this optimum temperature value used equally valid in the tropics where one might expect soil microbes to be acclimatized to higher temperatures?

Optimum temperature for CH_4 production versus CO_2 production and availability

This is an interesting point linking pathways of CH_4 production in wetlands to soil CO_2 production and CO_2 availability. While it is possible that there exist different optimal temperatures for CH_4 production via the acetoclastic pathway versus other methanogenesis pathways, there is a paucity of literature on these optimal temperatures for the various methanogenesis pathways. For this reason, we use one formulation for the optimal temperature for CH_4 production in our initial development of WETMETH.

Same optimum temperature for CH_4 production across the globe

We were not able to find specific optimal temperatures for CH_4 production in different climate zones. Hence, for our first implementation of WETMETH, we chose to apply a constant optimum temperature for CH_4 production based on published studies, which is mainly based on data from northern high-latitude environments (e.g. Dunfield et al. 1993; Metje and Frenzel 2007). The default optimum temperature for CH_4 production in WETMETH (~27°C) is subject to a sensitivity analysis (See Section 6).

Reference to Wu et al. (2016) for our formulation of the optimal temperature for CH_4 production

Due to this comment, we realize the way we cite Wu et al. (2016) with regard to the Q_{10} coefficient and optimal temperature formulation may not be clear. The form of the equation used for the Q_{10} coefficient in WETMETH is analogous to the one used in Wu et al. (2016), but the coefficients in the equation differ (our mathematical expression for the CH_4 production is $Q_{10}(T_i) = 1.7 + 2.5 \tanh[(0.1(308.15 - T_i))]$, whereas that used in Wu et al. (2016) for soil CO_2 production and respiration is $Q_{10}(T_i) = 1.44 + 0.56 \tanh[0.075(46 - T_i)]$).

To avoid further confusions, we will revise the related statement in Section 3.1.1 of the submitted manuscript (lines 207-208) as follows:

- **“This mathematical formulation is analogous to one used for soil respiration in another study (Wu et al., 2016)”.**

The model produces very different latitudinal patterns in flux per unit area to other estimates (Figure 11b, page 24), which could be due to the optimum temperature. This is especially the case over the tropics where the authors acknowledge that the overestimation

of wetland area may be compensating for the low flux intensity, resulting in an overall flux which is more comparable to other estimates. This optimal temperature parameterization and the value chosen needs more justification, analysis and discussion.

Justification of the optimal temperature parameterization

The existence of an optimum temperature for CH₄ production in wetlands is relatively well established in the literature (e.g. Dunfield et al. 1993; Metje and Frenzel, 2007; Dean et al. 2018). As indicated in Appendix A1, the chosen value for this optimum temperature in WETMETH (~27°C) is within the range of available estimates (i.e. 25-30°C). This default value is subject to a sensitivity analysis (through T_{ref}) in Section 6.

To clarify these points, we will revise the following statement in Section 3.1.1 of the revised manuscript (with new text highlighted in yellow):

- “In order to account for ... the occurrence of an optimal temperature for CH₄ production (Blake et al., 2015; Dean et al., 2018; Dunfield et al., 1993), a temperature-dependent Q_{10} is considered in WETMETH. Its mathematical formulation is $Q_{10}(T_i) = 1.7 + 2.5 \tanh [0.1 (T_{ref} - T_i)]$, where $T_{ref} = 308.15$ K is a reference temperature that is used to define an optimal temperature for CH₄ production (Table 1). This formulation is defined in analogy to a mathematical expression used for soil respiration in another study (Wu et al., 2016), and it enables to account for an optimal temperature for CH₄ production of ~300.15 K (i.e. 27°C) which is consistent with previous studies (Dunfield et al., 1993; Metje and Frenzel, 2007). Additional information on this formulation and its implications for the temperature-dependency of CH₄ production are provided in Appendix A1”.

In the above paragraph, (i) we clarify that the optimal temperature for CH₄ production is defined (or parameterized) through T_{ref} in the Q10 formulation, and (ii) we justify the default value chosen in WETMETH for this optimal temperature.

Analysis with regards to the optimal temperature parameterization

In our model, we parameterize the optimal temperature for CH₄ production through T_{ref} in the Q10 formula described in Section 3.1.1. (third paragraph) as well as Appendix A1. Given that the optimal temperature for CH₄ production is uncertain, we conducted a sensitivity analysis on the default value for the optimal temperature in WETMETH (i.e. ~27°C) which is defined through T_{ref} in the Q10 formula. In Section 6 (see third paragraph), we show that simulated CH₄ emissions from tropical wetlands are more sensitive to changes in T_{ref} (therefore changes in the optimal temperature value for CH₄ production) than CH₄ emissions from northern wetlands. This result suggests that, in the present-day climate, wetland CH₄ emissions in the tropics are more dependent on the optimal temperature for CH₄ production than wetland CH₄ emissions in the boreal and Arctic regions.

Discussion

We plan to include a new bullet point (see below) in Section 7 discussing the optimal temperature for CH₄ production in the context of this manuscript and the modelling of wetland CH₄ emissions.

- “While the existence of an optimal temperature for CH₄ production in wetlands is relatively well established in the literature (Dean et al., 2018), there are currently no estimates of such an optimal temperature for different climate zones across the globe. Previous studies suggest a range of 25-30°C for such an optimal temperature based on measurements of CH₄

production in northern wetlands (Dunfield et al., 1993; Metje and Frenzel, 2007). In WETMETH, we use a global value for this optimal temperature (~27°C) which is assumed to be valid for CH₄ production in both tropical and extra-tropical wetlands (see Section 3.1.1 and Appendix A1). However, our sensitivity analysis suggests that, in the present-day climate, wetland CH₄ emissions in the tropics are more dependent on the optimal temperature for CH₄ production than wetland CH₄ emissions in the boreal and Arctic regions (see Section 6). The optimal temperature for CH₄ production in WETMETH, along with other factors such as areal wetland extents, contributes to inter-model differences in simulated wetland CH₄ intensities in the tropics (see Figure 11b)".

The detailed validation of WETMETH is skewed towards the extra-tropics. There is insufficient validation against available tropical methane measurements. The assessment of tropical fluxes is made by comparison against other models from (climate model) grid to regional scales. This issue is highlighted by the fact that WETMETH produces very different flux rates per unit area in the tropics relative to other models (see above). Adequate validation over the tropics is of particular importance because tropical wetland methane emissions dominate the global wetland budget, and WETMETH is intended for use in Earth system models.

We agree that adequate evaluation in the tropics is needed, however we are limited by the scarcity of large-scale observations across the tropics; a challenge shared by the wide community of scientists working on large-scale CH₄ emissions: from those contributing to observation syntheses and database development (e.g. FLUXNET-CH₄ by Knox et al. 2019) to those putting together CH₄ budget reports (e.g. Saunio et al. 2020). As we state at the beginning of Section 5.2, our model validation focuses on large-scale wetland CH₄ emissions given the coarse grid resolution of the UVic ESCM into which WETMETH has been embedded, and thus we require large-scale observations which, to our knowledge, are not available.

Nevertheless, we show (in Table 2) that tropical wetland CH₄ emissions simulated by WETMETH fall in the range of estimates available at the large spatial scales needed for an appropriate comparison - namely upscaled measurements (Sjörgesten et al. 2014) as well as results from process-based models (Melton et al. 2013) - although we are aware that these estimates are poorly constrained.

Furthermore, we acknowledge the particular importance of tropical wetland CH₄ emissions in the global CH₄ budget and suggest that more observations and improved process-understanding are needed to constrain these CH₄ emissions (see Section 8 - Conclusions, second paragraph).

Finally, we also acknowledge (e.g. in Section 5.1 & Section 8) that tropical wetlands simulated by the UVic ESCM are generally more widespread than predicted by observation-based datasets, implying the need for future work to improve the spatial distribution of these wetlands in the UVic ESCM. In the advent of more reliable data products for large-scale wetland CH₄ emissions in the tropics in the future, along with improved tropical wetlands in the UVic ESCM, we expect that further model development will consider detailed model validation with respect to tropical wetland CH₄ emissions.

Specific comment:

The manuscript states: " : : : particularity of WETMETH among relatively simple models is that the model accounts for : : : : a depth-dependent representation for methane production" (page 27 line 620). The addition of a vertical dependent methane production

is not entirely novel within models of similar complexity (e.g. Comyn-Platt et al 2018).

Our intention with that statement (in the last paragraph on page 27) is to highlight the novelty with respect to calibrating model parameters for depth-dependent CH₄ production against measurements from laboratory incubations. To the best of our knowledge, such a calibration for CH₄ production has never been done in other global wetland CH₄ models.

For better clarity, we intend to rephrase our statement as follows:

- **“The particularity of WETMETH among relatively simple models is that the model accounts for ..., a calibration of depth-dependent CH₄ production rates against potential CH₄ production rates from laboratory incubations, dynamic CH₄ oxidation ...”.**

Response to reviewer #2

The paper describes an intermediate complexity approach to represent methane production and oxidation in an Earth system model. Modeling the two component fluxes of methane from wetlands is challenging and this paper is a good contribution to the existing land surface models engaged in wetland methane emission research. The paper is very well written and is excellently referenced.

I have just minor comments and clarifications:

We thank the reviewer for the positive and constructive comments on the manuscript. Please see below our responses to each of their comments and requests for clarifications.

In the introduction, please mention ‘acetotrophic methanogenesis’ and ‘hydrogenotrophic methanogenesis’ when discussing the three pathways for methane production from microbes.

Thank you for this suggestion. Following your recommendation, we will mention the terminologies associated with the three methanogenesis pathways (acetotrophic methanogenesis, hydrogenotrophic methanogenesis, and methylotrophic methanogenesis) in the revised manuscript.

Fig 1 – suggest adding a CH₄ production figure

The reason for depicting CH₄ concentration rather than production in Figure 1 is that concentration is what is typically measured in the vast majority of field observations. As a simplification for the purpose of this illustration, the CH₄ concentration curve in Figure 1 was assumed to mirror CH₄ production rates at depth within the soil system. The reason for assuming such correspondence (as explained in the text of the paper) is that the main effect on production rates is likely to be % organic content under anaerobic conditions, but under aerobic conditions, the main effect is more likely to be a balance between organic content versus increasing availability of O₂ that could support aerobic decomposer communities that compete for the available organic carbon substrate. Under the aerobic scenario, we assumed CH₄ oxidation would also contribute to lower CH₄ concentrations in the upper soil, but that much of this oxidation would be based on consuming CH₄ produced deeper in the soil that is diffusing upward into and through the non-saturated layer.

For clarification, the following sentence will be added to the figure caption:

- “As an illustrative simplification, the CH₄ concentration profile is assumed to mirror CH₄ production rates at depth within the soil column (see explanatory text in Section 2.4)”.

Eq 1 – is C_i representative of total soil C, or more mobile source of C, like the intermediate pool (if using a CENTURY type model) ?

C_i represents total soil C in a given ground layer *i* (please see the first statement of Section 3.1.1 from the original text as reproduced in our next response).

Eq 1 – specify that S(theta) is for vertical profile, not for the entire grid fraction
Is there any coupling of permafrost sub routines w your CH₄ production scheme? Do they both use same soil temperature – and so if soil temperature is cold, then no CH₄ emissions?

S(theta)

In the first paragraph of Section 3.1.1 of the submitted manuscript (reproduced below), we indicated that CH₄ production in our model (as described by Equation 1) is calculated in individual soil layers whereby S(theta_{*i*}) characterizes soil moisture saturation in a given soil layer *i*, and not for the entire grid fraction.

“For any land location, the rate of CH₄ production in an underlying soil layer *i* (P_{*i*} in kg C m⁻³ s⁻¹) is parameterized as:

$$P_i = S(\theta_i) C_i r Q_{10}^{\frac{T_i - T_0}{10}} \exp\left(-\frac{z_i}{\tau_{\text{prod}}}\right), \quad (1)$$

where S(theta_{*i*}) is the fraction of the soil layer that is saturated with water, and C_{*i*} is the amount of soil carbon (in kg C m⁻³) in the layer”.

Permafrost sub routines & CH₄ production scheme

Our CH₄ production scheme is coupled to the permafrost sub routines in the UVic ESCM and they both use the same soil temperature. In the UVic ESCM-WETMETH coupling, we assume that:

- 1. When a given soil layer *i* is frozen (T_{*i*} < 0), CH₄ production shuts down in that layer;**
- 2. CH₄ production would still occur in unfrozen soil layer (T_{*i*} > 0) contingent to other factors (e.g. soil carbon content and moisture saturation).**

The above assumptions are described in Section 3.1.1 and reflected in Equation 1. A direct implication of these assumptions is that no CH₄ production and emissions occur when the entire soil column is frozen. We acknowledge that any bias in soil temperature (as simulated by the UVic ESCM) will be reflected in CH₄ production and emission rates.

What is the model timestep of CH₄ production and oxidation?

CH₄ production and oxidation in our model (that is, the UVic ESCM-WETMETH coupling) are calculated every 5 days and integrated with a 30-day timestep. This 30-day timestep integration is similar to that of soil respiration and other terrestrial carbon fluxes defined in the dynamic vegetation model (TRIFFID) component of the UVic ESCM.

To clarify this point, we will add a statement (see below) on this model timestep for all terrestrial carbon fluxes in the UVic ESCM through TRIFFID. This statement will be added in Section 3.2 (third paragraph):

- “Through TRIFFID, all terrestrial carbon fluxes in the UVic ESCM are integrated with a 30-day timestep (Meissner et al., 2003)”.

Does soil carbon get reduced to account for carbon lost through CH₄ production?

No, such a reduction of soil carbon is not implemented in the model presented in the manuscript. However, the reduction of soil carbon following CH₄ production is considered in further model developments with the global CH₄ cycle in order to account for the conservation of carbon in the Earth system. This effect is negligible in the historical simulation of wetland CH₄ emissions and the climate system.

Is Zoatz the same as Zoxic, I was a confused here what the difference is.

Zoxic and Zoatz are different: Zoxic represents the oxic zone depth (illustrated with the orange two-sided arrows in Figure 2), whereas Zoatz represents the thickness of the oxic-anoxic transition zone to account for the availability of oxygen in the top layer saturated with water (i.e. top of the anoxic zone). The bottom of Zoatz defines the oxic-anoxic interface (illustrated with the red dashed horizontal lines in Figure 2).

Overall:

1. Zoatz is included in Zoxic;
2. The two terms are only identical when the soil surface is inundated with water (e.g. Figure 2a).

To clarify these points, we will include the following statement in the revised manuscript:

- “When the soil surface is inundated, Z_{oatz} is identical to Z_{oxic} (Fig. 2a). Otherwise, Z_{oatz} is only a fraction of Z_{oxic} (Fig. 2b)”.

Ultimately, the oxidation rate is fixed through space and time, my calculation is that 96% of gross methane production is oxidized, e.g., $1 - \exp(-0.05/0.0146)$ – is this correct?

Why not vary Zoxic to represent a dynamic water table height? I didn’t understand why this is fixed at 0.05 globally and over time.

In our model, CH₄ oxidation does vary with a dynamic Zoxic as illustrated by Figure 2 (see panel a versus panel b). Your calculation is correct for a case when the soil is inundated (i.e. for Zoxic = Zoatz = 0.05 m, 96.7% of CH₄ produced in the soil column gets oxidized in transit to emission). However, when the soil is not inundated, the fraction of produced CH₄ that gets oxidized before release (1) is greater than 97%; and (2) increases as Zoxic deepens.

To clarify this point, we will include the following sentence in the revised manuscript (Section 4.2):

- “This tuned value for τ_{oxid} is listed in Table 1 and implies that ~97% of the CH₄ produced in the soil column gets oxidized when the soil surface is inundated”.

Please note that we assume a static Zoatz (0.05 m) in this first version of WETMETH (as stated in Section 3.1.2, second paragraph). Further model development could explore how to represent Zoatz in a dynamic way.