

## ***Interactive comment on “Capillary rise affecting crop yields under different environmental conditions” by Joop Kroes et al.***

### **Anonymous Referee #1**

Received and published: 5 December 2016

### OVERALL COMMENTS

I read this paper with interest, as it shares some overlap with my own research. In this manuscript, the authors set out to answer two questions stated in the introduction: (quoted from text)

- (i) What is the impact of capillary rise on crop yield and downward leaching of water?
- (ii) What is the impact of soil types on capillary rise?

Understanding soil hydrology in agroecosystems is very important in the global context of sustainably increasing agricultural production while protecting water resources, and there is still uncertainty regarding the sources of water used by crops. However, these questions are rather broad, which I believe contributes to significant issues with the

[Printer-friendly version](#)

[Discussion paper](#)



paper's narrative. Overall, my recommendation is that this manuscript be rejected, for three primary reasons which are elaborated on in more detail below. After significant revisions, the paper may be suitable for a journal with a narrower disciplinary focus such as Agricultural Water Management instead, but as it is currently written there are not wide-reaching implications for this study that justifies a more generalist journal such as HESS.

## MAJOR CONCERNS

### CONCERN 1: Lack of Clarity Regarding Experimental Design

I do not understand the implementation of hydrological boundary condition b (“free drainage with capillary rise”). In my understanding, free drainage is typically implemented by setting the bottom boundary condition to a 0 pressure head gradient, which means that the total hydraulic head gradient is equal to 1 ( $dH/dz=dz/dz$ ). In the SWAP manual, and it states that under free drainage boundary conditions there is downward flow out of the bottom boundary at a rate equal to the hydraulic conductivity of the bottommost soil layer (see equation 2.20 in SWAP manual), which would mean capillary rise through the bottom boundary is not possible.

Therefore, it seems that the authors define capillary rise as any water that moves upward across the depth in the soil column which is defined as the bottom of the root zone (line 180, “We calculated capillary rise as the upward flux at the bottom of the root zone”). Several points of clarification are needed. Is this the same boundary at which the free drainage condition is defined, or is free drainage defined at the bottom of the soil column (the depth of which is not stated) and the capillary rise boundary condition turned on/off at the bottom of the root zone? Could the authors describe in more detail how the Richards Equation is implemented such that water movement is only permitted in a downward direction – are there separate upwards and downwards hydraulic conductivities in the vertical direction, as implied in lines 190-192, or are both upwards and downwards conductivities reduced at the bottom of the root zone? If so,

[Printer-friendly version](#)[Discussion paper](#)

is this physically realistic, or simply a modeling experiment? I note that the authors of the study are the developers of the SWAP model, so perhaps things which seem obvious to them are not quite as clear to the readers (such as myself). Given that the difference between the three boundary conditions is the fundamental concept upon which the entire study is based, it is essential that these boundary conditions are clear and defensible, and they currently are not. They reference a previous study (Kroes and Supit, 2011) which introduces the coupled SWAP-WOFOST model, but this boundary condition is not used in the previous study.

Also, if my above interpretation of the boundary conditions are correct, it seems that the authors' results would be highly sensitive to the depth of the soil column relative to the depth of the rooting zone, as the difference between these two depths provides the "deep soil moisture" that is allowed to move upwards through the bottom boundary of the root zone and therefore constitutes capillary rise in their simulations. How was the total soil column depth selected, and how sensitive are the results to this decision?

#### CONCERN 2: Lack of Narrative Discretion

Overall, the paper's narrative requires significant work. I feel that the authors include too much information. While it is evident that the authors did quite a bit of work, it is also important to distill the results to aid the reader in interpreting and understanding how they relate to the study's research questions; I am reminded of the quote, "If I had more time I would have written a shorter letter" (<http://quoteinvestigator.com/2012/04/28/shorter-letter/>). Their figures and tables seem to consist of a laundry list of all their results, when it is the job of the author to synthesize and condense their results to a coherent message. A more effective paper would synthesize the key messages from these plots to a smaller number of figures targeted closely at their specific research questions, and present the full results in the supplementary material as necessary. For example, figures 4, 6, and 8 only have a few actual observed data points in them; they could easily be condensed into a single scatterplot, color-coded by research site.

[Printer-friendly version](#)[Discussion paper](#)

The other aspect of the narrative, the framing of the study in the Introduction section and situating their results in the literature via the Discussion section, seem incomplete. In the Introduction section, they describe several studies that find that capillary rise from groundwater into the root zone can be an important source of water to crops. The importance of capillary rise to crops in shallow groundwater environments is well known; in addition to the studies cited, many others have found this to be the case (Kang et al., 2001; Sepaskhah et al., 2003; Ghamarnia et al., 2010; Luo and Sophocleous, 2010; Huo et al., 2012; Xu et al., 2013; Talebnejad and Sepaskhah, 2015; Han et al., 2015). Similarly, their comparison of yield difference between cases including groundwater vs those with free drainage seems analogous the concept of groundwater yield subsidy of Zipper et al. (2015). Therefore, it is key for the authors to identify a knowledge gap in the literature that they propose to fill; the closest they come is the statement, “This paper quantifies the effects of capillary rise on crop growth under different conditions of soil hydrology, soil type and weather.” (Lines 94-95). Identifying the relative importance of different drivers (soil, crop choice, weather) to variability in the capillary contribution of groundwater to crop water requirements, but this is not clearly motivated in the introduction, included in the research questions, or answered by the results. (And, as noted in concern #1, I am not convinced their method is appropriate to answer this question). there is no analysis conducted or information given about weather conditions, which is presumably driving much of the interannual variability observed in plots. Nor are any conclusions discussed regarding the impact of weather on capillary rise. Nor are there any cross-site analyses or discussion of variability in soil texture or crop type. Thus, it is hard to say what lessons can be learned from the data presented here. The Discussion is brief (<1 page) and could be enhanced by situating their results within the existing body of literature detailing the importance of capillary rise to crop water requirements; see the studies referenced above. As mentioned above, the questions the authors’ research questions are broad, so expanding the discussion can help make clear what new contribution their results add to the existing body of literature.

CONCERN 3: Issues in Tables/Figures

Figures are the key to any paper and there are many fundamental flaws or oversights here. Figure 2 does not have a legend explaining what the colors mean. I understand that 72 different types of soils is too many to display in a legend, but perhaps there are general categories such as those used in Figure 9? Figures 3-8 do not have axis labels. In Figure 7 (top panel), the observed groundwater levels are provided to the model as a bottom boundary condition, and are therefore not an appropriate validation dataset. In general it takes quite a lot of effort to determine what plots correspond to what scenarios. For example, in the text it states “Observed groundwater levels were used as a lower boundary condition for Borgerswold” (lines 166-167). To determine which plot this was, I then had to go to Table 1 (or Table 3) to determine the case study number, and then look at plots until I found the groundwater levels corresponding to Case Study 5. Perhaps the case studies could be given more descriptive names based on the location and crop (e.g. case study 1 = DM-Grass, case study 2 = DM-Maize, case study 3 = C-Maize, etc.) to aid in interpretation of the figures. In the context of yield, I do not understand what the continuous lines showing simulation results are (e.g. in Figure 8); yield is typically measured at the time of harvest. Perhaps “dry biomass” would be a more accurate term? The axes chosen cut off parts of data in some plots; for example, parts of the boxplots are missing from Figures 9, 11, 12 (upper right panel).

There are also quite a few tables, which could be condensed: Tables 1-3 all have the same first column and can be combined Table 4 has no information which is not already presented in the text In Table 5, I do not think that Nash-Sutcliffe Efficiency is an appropriate metric for evaluating yield, as the authors are typically comparing a single annual observed point to a timeseries of model output; typically, NSE is used for comparing timeseries data (e.g. a hydrograph).

## OTHER, MINOR COMMENTS

While readable, throughout the manuscript there are issues with the writing, grammar, and punctuation; a careful proofreading of the manuscript should be able to fix these

[Printer-friendly version](#)

[Discussion paper](#)



problems.

Lines 45-46: “This is especially the case in delta areas where high population densities occur and agriculture is the predominant land use” – this seems to be mutually exclusive? Are the deltas urbanized or agriculture? Or, are the authors trying to say that there are dense cities surrounded by agriculture and thus potentially competing water users?

#### REFERENCES IN REVIEW:

Ghamarnia, H., M. Golamian, S. Sepehri, and I. Arji. 2010. Shallow groundwater use by Safflower (*Carthamus tinctorius* L.) in a semi-arid region. *Irrig. Sci.* 29(2): 147–156.

Han, M., C. Zhao, J. Šimůnek, and G. Feng. 2015. Evaluating the impact of groundwater on cotton growth and root zone water balance using Hydrus-1D coupled with a crop growth model. *Agric. Water Manag.* 160: 64–75.

Huo, Z., S. Feng, G. Huang, Y. Zheng, Y. Wang, and P. Guo. 2012. Effect of groundwater level depth and irrigation amount on water fluxes at the groundwater table and water use of wheat. *Irrig. Drain.* 61(3): 348–356.

Kang, S., F. Zhang, X. Hu, P. Jerie, and L. Zhang. 2001. Effects of shallow water table on capillary contribution, evapotranspiration, and crop coefficient of maize and winter wheat in a semi-arid region. *Crop Pasture Sci.* 52(3): 317–327.

Kroes, J.G., and I. Supit. 2011. Impact analysis of drought, water excess and salinity on grass production in The Netherlands using historical and future climate data. *Agric. Ecosyst. Environ.* 144(1): 370–381.

Luo, Y., and M. Sophocleous. 2010. Seasonal groundwater contribution to crop-water use assessed with lysimeter observations and model simulations. *J. Hydrol.* 389(3–4): 325–335.

Sepaskhah, A.R., A. Kanooni, and M.M. Ghasemi. 2003. Estimating water table con-

[Printer-friendly version](#)

[Discussion paper](#)



tributions to corn and sorghum water use. *Agric. Water Manag.* 58(1): 67–79.

Talebnejad, R., and A.R. Sepaskhah. 2015. Effect of different saline groundwater depths and irrigation water salinities on yield and water use of quinoa in lysimeter. *Agric. Water Manag.* 148: 177–188.

Xu, X., G. Huang, C. Sun, L.S. Pereira, T.B. Ramos, Q. Huang, and Y. Hao. 2013. Assessing the effects of water table depth on water use, soil salinity and wheat yield: Searching for a target depth for irrigated areas in the upper Yellow River basin. *Agric. Water Manag.* 125: 46–60.

Zipper, S.C., M.E. Soyulu, E.G. Booth, and S.P. Loheide. 2015. Untangling the effects of shallow groundwater and soil texture as drivers of subfield-scale yield variability. *Water Resour. Res.* 51(8): 6338–6358.

---

[Interactive comment on Hydrol. Earth Syst. Sci. Discuss., doi:10.5194/hess-2016-598, 2016.](#)

## HESD

---

[Interactive  
comment](#)

[Printer-friendly version](#)

[Discussion paper](#)

