

Point-by-point reply to review comments:

We would like to thank the reviewers for providing critical feedback to our manuscript. We provided a reply to each comment which is followed by the revisions we made to the manuscript. The text for the reviewer comments are in grey while our replies and revisions are in black.

Reviewer 1:

Specific comments

Overall, the paper is well written but can be improved in what concerns the Methods section organization.

I have two main comments:

1) The first is related to the the choice of the value of the parameter β_c . I think it is a key parameter for the study and should deserve further discussion (or results) on its impact on the results.

We agree with the point raised by the reviewer on the parameter β_c . We adapted the manuscript in Section 5.2 page 11 from line 1. The text now reads:

"Weaknesses: The first aspect that needs further investigation is the selected β_c value for identifying the decoupled soil moisture range. Although the selection in this study was based on trends identified from time series datasets, the methods applied should be tested further using other datasets to confirm the suitability of $\beta_c = 1$ for other depths and soil types. The choice of β_c is crucial as it dictates which soil moisture values are expected to be decoupled. For instance, at the sites where decoupling occurs during dry conditions, a higher β_c value would enlarge the decoupled range. A similar effect would be expected for the site with decoupling during wet conditions. However, a lower β_c value could result to decoupling only during extreme soil moisture conditions (e.g very wet or very dry)."

2) While the authors include different stations characterized by different vegetation cover (grass, corn and forest) its potential effect on the results is not examined in deep, especially for station SM20 which is located in the forested area, this effect could be significant. Can the authors provide further discussion on it?

We agree with the reviewer that the discussion on the controls, such as the effect of vegetation was not very extensive. This is because our main focus for the study was to investigate the methodology applied to quantify coupled and decoupled soil moisture values. However, we do realize that providing further discussion on said topic would improve the quality of the discussion in the paper. We adapted the manuscript in 5.1 in page 9 starting from line 25

"The vegetation type at each site exerts some influence on the soil moisture variability and the resulting (de)coupled values. First, the vegetation type affects how much ground surface is directly exposed to atmospheric conditions. Forested areas and grass fields are almost fully covered by vegetation compared to a corn field where the crops are organized in equidistant rows. Vegetation or canopy cover will determine how atmospheric conditions affect the soil moisture values. For instance, the amount of intercepted precipitation and evaporation are both dependent on vegetation cover. This in turn will have direct impacts to the surface soil moisture dynamics at each of the sites. For comparison, the variability given by the standard deviation bars in fig.4 and variance in fig.5 at for the cornfield (SM09) is higher compared to that of the grass field (SM05) or the forested area (SM20). In addition, the forested area (SM20) has the smallest range of soil moisture values among the four sites. This may be due to the high amounts of intercepted rainfall by the forest canopy. Root water uptake (RWU) is another way by which vegetation affects soil moisture variability. RWU can have significant influence on the subsurface dynamics. The influence of RWU may vary for different vegetation types as it can be exerted over a range of depths, leading to differences in the resulting (de)coupled values."

I have some additional and technical comments that I will list below in order of appearance in the manuscript:

1) Section 3.1: It is not clear to which figure the authors are referring to. Please point to a specific figure when describing graphic features or speak more in general. This section should contain method description and choices made for carrying out analysis.

We adjusted the text in section 3.1 and added a schematic diagram which the text refers to.




Figure 3. Schematic diagram using hypothetical soil moisture values to show vertical variability. Left side of the figure shows the trend with a fitted loess function. The variability can be seen using the standard deviation bars. The right side shows the residuals of the fitted function. Soil moisture variability is more visible from the variance given by the vertical bars and the cumulative variance is given by the black line. A change in variability at an intermediate soil moisture value is marked by a change in the slope of the cumulative variance line, marked by θ_c

2) Section 3.1: Define what vertical variability exactly means.

We added a line in Section 3.1 in page 4 line 6:

"We referred to vertical variability as the unevenness or irregularity in soil moisture distribution within a certain depth of interest along the vertical profile, in this case up to depths of 40 cm"

3) The cumulative variance is presented in Figure 4. Same as before, here the authors speak about slope but it is not clear at all for the reader to what slope they are referring to. Please define what cumulative residual variance means and clarify better all the related concepts.

-We adjusted the text in section 3.1 and added a schematic diagram, which is now figure 3, to refer to the parts of the graph being described in the text. Please see response for technical comment #1

4) Section 3.2.2 Distributed lag model: It is not clear enough how the DLNM model is used with soil moisture observations. I suggest to describe the basic concepts here rather than mathematical details (which could be included in an appendix). Please try to simplify and clarify this section so as it can be more easily interpreted

We realize the difficulty encountered by the reviewer in following the description of methods applied using the DLNM. We completely revised 3.2.2 section of the manuscript. The section now becomes:

"3.2.2 Distributed lag model

We incorporated delayed or lagged effects in evaluating the relation between surface and subsurface values, and eventually in determining (de)coupled values. It should be emphasized that the analysis was primarily focused on examining the trends and relations between surface and subsurface soil moisture. Moreover, it was not intended to contradict or replace other models for estimating soil moisture or examining its patterns.

A distributed lag non-linear model (DLNM) developed by Gasparrini et al. (2010) was applied to the 5 cm and 40 cm time series datasets at the study sites. Briefly, the model is capable of simultaneously representing both functional dependency and delayed response between exposure and response values. We considered surface soil moisture as the exposure values that produced delayed effects to the response values at the subsurface. A non-linear model was selected in order to capture the non-linear dynamics of flow and transport along the soil profile (Mohanty and Skaggs, 2001; Kim and Barros, 2002). Furthermore, DLNM offered enough flexibility to model a variety of dependencies in the time series dataset by selecting a suitable basis function. DLNM could be thought of as equivalent to a linear time series model (e.g. autoregressive model) just as a generalized linear model is equivalent to a linear model.

In assessing lagged dependence, event scale patterns were of interest rather than large scale trends within the time series (Wilson et al., 2004). This required seasonal patterns to be addressed prior to applying the DLNM. This was done by fitting a loess function to the time series and then subtracting it from the original soil moisture values (Cleveland et al., 1990). Removal of seasonality was further justified by the scatterplot results (see Section 4.1). The influence of seasonality on the vertical soil moisture variability is indicated by clustering of observation points occurring within the same months (fig.4). De-seasonalized soil moisture values were used for identifying (de)coupled soil moisture conditions.

For consistency in modeling, the range of surface soil moisture values used was from 0-0.50 cm³ cm⁻³. This was based on the highest surface soil moisture value encountered among the four sites. A lag value of up to 30 days was considered long enough to investigate delayed effects. This period also approximated the recurrence of heavy rainfall within the study sites. A spline function was the basis function chosen to represent the functional dependence as well as delayed effects as it offered flexibility to capture non-linearities. In addition, contributions from daily rainfall data were used to incorporate current and past meteorological conditions. This was applied as a covariate and was represented with an additional basis function. We only considered delayed effects in vertical flow as lateral movement is deemed negligible in flat to slightly sloping terrain (Table.1). The analysis was performed in R software using *dlnm* (Gasparrini, 2011) and *mgcv* (Wood, 2006a) packages.

The following section concisely describes the mathematical formulation of a DLNM. However, the reader may choose to skip this section as the general methods applied have already been described in the text above. For a more detailed explanation, readers are referred to Gasparrini et al. (2010) and Gasparrini et al. (2017).

####Some lines for the mathematical description of DLNM is not included here. The structure of this section was not revised#####

3.3 Evaluating (de)coupled soil moisture values

Application of a DLNM resulted in the estimation of parameter β for each surface soil moisture value. This indicated the strength of dependence between surface and subsurface soil moisture. Higher β values indicated stronger dependence or coupling between the two. Hence, we referred to β as the relative influence of surface soil moisture on subsurface values."

5) Pag. 6 lines 19-24: here the authors seem to anticipate the results of the paper about the value to be assigned to β c . However, I suggest to try to organize the paper in a way that the choice of β c is described in the results section (and supported by analysis).

We agree with the reviewer that this paragraph is more suitable in the results section. This was moved to section 4.3 in page 7 starting from line 24.

6) Pag 8 line 17-19. Clarify this sentence.

We changed the text in page 8 line 17-19. It now reads as:

"For instance, at SM05 and SM09, there is a general increase in β from dry towards wet surface soil moisture values. SM20 also shows increasing β over a limited soil moisture range (0.1 –0.25 cm³ cm⁻³). Outside this range, the estimated β values for SM20 were less than one and have very broad confidence intervals. Recall that the range used for DLNM was only for uniformity among the four study sites. The lack of or very few observations for very dry or very wet soil moisture conditions led to wider confidence intervals not only for SM20 but also for the other three sites. Compared to the three sites, the estimated β values for SM13 show decreasing values towards the wet soil moisture range (> 0.3 cm³ cm⁻³). From the intermediate to dry soil moisture conditions, the values fluctuate around the designated β c."

7) Pag 8 line 11. Replace 40cm with "40 cm"

This was now changed. The whole manuscript was also reviewed for similar errors.

8) Pag 9 . Line 25. T is missing.

A "T" is added in the beginning of the sentence. There was another round of proof reading to check for similar textual and grammatical errors.

Reviewer 2:

General comments: This study examines the coupling between near-surface and sub-surface soil moisture at four sites in the Netherlands. Specifically the authors develop a methodology for determining when the two layers are decoupled, thereby providing an important analysis for surface soil moisture assimilation into models. The manuscript is very well written and the figures are well crafted. The use of a distributed lag non-linear model for quantifying decoupled soil moisture ranges is novel and, as the authors point out, does not suffer from many of the assumptions and limitations of previously implemented methods. I recommend the manuscript for publication given appropriate consideration of the following concern.

C1

Specific comments: The primary concern I have with the manuscript is the conclusion that the decoupled range is not limited to dry conditions. Evidence of this is provided at one of four sites (SM13), at which the authors confirm the presence of burrowing animals. Given the potential data quality issues at site SM13 and the fact that decoupling at the the other three sites was confined to the dry end of the soil moisture range, I believe the strong statements regarding soil moisture decoupling outside of dry conditions (i.e., section 5.1, line 9; section 5.2, lines 26-27) are not adequately supported by the results. Therefore I recommend the authors either soften this conclusion by adequately describing the uncertainty and lack of consistent supporting evidence, or assess why SM13 shows decoupling outside of dry conditions when the other three sites do not.

We acknowledge the reviewer's concern about the occurrence of decoupling at wet soil moisture conditions. Indeed, compared to the other three sites, this result is intriguing. We decided to follow the reviewer's recommendation to asses why SM13 shows decoupling outside conditions. In our view, the occurrence of decoupling during wet conditions is plausible as certain combinations of atmospheric, biological, and pedological conditions can promote this. We elaborated further on whether burrowing animals may have compromised the quality of the dataset at SM13. We argue that the creation of macropores by burrows did not result in lowered data quality as this would result in subsurface soil moisture dynamics that is opposite of what is observed from the time series datasets. We expanded the discussion in Section 5.1 in page 10 starting from line 4.

"Among the four sites, the subsurface trends observed for the 40 cm values at SM13 show consistently high values, which can be more pronounced during winter months. This resulted in decoupling during wet soil moisture conditions fig.8. This trend is different from the other three sites which only show a slight increase in the subsurface values. Further inspection of the time series data at SM13 reveals no sudden disturbance in the signal which could be attributed to errors in the sensor. Field investigation confirmed an increase in silt content at 40 cm compared to the upper layers. The increase in silt content promotes a decrease of hydraulic conductivity over depth that results in a slower vertical flow towards deeper layers. The presence of burrowing and hibernating animals was also observed at the site during winter. These create macropores which eventually alter the hydraulic properties of the soil (Kodešová et al., 2006; Beven and Germann, 2013). We infer that, at the measurement domain of the sensor, these burrows or macropores facilitated faster vertical flow to the subsurface. Alternatively, if the burrows produced voids around the measurement domain, this would result in lowered soil moisture or data gaps due to the loss of sensor to soil media contact. However, there were no gaps observed that coincided with the burrowing animals' period of hibernation. During precipitation events, soil moisture flowing from upper layers arrived more rapidly at 40 cm depths due to the presence of macropores. There it accumulated and flowed more slowly to deeper layers because of the low hydraulic conductivity promoted by the increase in silt content. The overall effect of these factors was the pronounced increase in

soil moisture values at 40 cm compared to those at 5 cm during winter periods as observed from the time series dataset fig.2."

Reviewer 3:

Received and published: 25 January 2018

This is a strong paper which makes a clear contribution. In particular, I appreciate the attempt to inject more rigor into the discussion surrounding vertically-coupling among multi-depth soil moisture measurements. I view the paper's contribution as being mainly methodological; however, some interesting preliminary conclusions regarding the occurrence of vertical de-coupling are presented (specifically, that – at least at one site – de-coupling is not limited to dry soil conditions). The overall presentation of the manuscript is very good and the topic is of sufficient interest for HESS' readership. Therefore, I recommend publication following adequate response to the following minor points:

1) The manuscript would benefit from a more detailed description of exactly how their approach(es) can be used to improve the performance of a land data assimilation system. There are really two issues here. The first is a fundamental observability issue (i.e., what is the upper limit on how effectively surface soil moisture can be used to constrain sub-surface soil moisture given a perfect data assimilation issue). The second issue is the vertical accuracy of the assimilation paper. That is, even if you have high theoretical observability (i.e. high vertical coupling) at a particular point, you can still squander this potential constraint by using an assimilation model that does not properly represent this coupling. This is the model accuracy issue addressed previously by the Kumar et al. 2009 JHM paper (already cited in the original paper). To me, the second point is really the most important; however, addressing it requires the additional step of cross-comparing observed vertical coupling to the vertical coupling predicted by the assimilation model (when run off-line). It sounds like this is the ultimate intention of the authors; however, it is not explicitly spelled out in the current manuscript. So, in summary, I would encourage the authors to be more specific/detailed in describing exactly how these results can be used to improve the performance of a land data assimilation system.

We thank the reviewer for pointing out this aspect. Our intention with this paper was to develop a data driven assumption free method. We agree completely with the cross-comparison of observed vs modeled vertical coupling, but as the focus in this paper is indeed on the methodology itself we discussed to what extent such a comparison should be included in this paper. Nonetheless, based on the suggestion of the reviewer we revised the manuscript largely in Section 5.2 to accommodate the reviewers suggestion in page 11 starting from line 26

"For data assimilation (DA) applications, the applied (de)coupling methods can be used for cross-comparison of the vertical coupling derived from DA model outputs with those observed from long term in situ measurements. This can aid in examining the adequacy of the assumed inherent connection between surface and subsurface values. As Kumar et al. (2009) pointed out, land surface models vary in their representation of the strength of this connection (e.g. weak or strong connection) which contributes the degree in which modeling results are improved. They also suggested that strong coupling is a more robust choice unless independent information suggests that a more decoupled surface-subsurface representation is more realistic. In this aspect, the analysis applied in this study could be a valuable tool in determining which type of surface-subsurface coupling is the more optimal choice. Furthermore, the assumed connection strength is adopted for the whole range of soil moisture values. The results of our analysis show that at any given site, decoupling will occur regardless of degree of soil moisture variability. We suggest that perhaps a variable coupling strength could be adopted based on the soil moisture range where decoupling is likely to occur rather than a single value for the whole range."

2) The discussion in Section 3.2.2 is not very accessible. For example, equation (1) is introduced as describing the time series of "outcomes" Y_t , yet Y_t only appears on the LHS of the equation as $g(E(Y))$. "g" is apparently a "monotonic link function" (??) and E is some kind of an expectation operator (in space, in time, across an ensemble?). So it's hard for me to see how Y_t is actually "described" here. In the next function sentence "s" is introduced as a "basis function" (?) and all this is

before the actual DLNM model is introduced. I suspect that all this terminology is correct and adequate for an applied math audience; however, HESS readership will likely need a bit more help and conceptual background to get through this section. I'd strongly recommend that the authors revise/expand Section 3.2.2 with an eye towards making it more accessible to a general earth science audience. Especially the early part of the section between equations (1) and (2)...I struggled there to follow the authors' approach.

Similar response to reviewer #1, comment #4.

We realize the difficulty encountered by the reviewer in following the description of methods applied using the DLNM. We completely revised 3.2.2 section of the manuscript.

"3.2.2 Distributed lag model

We incorporated delayed or lagged effects in evaluating the relation between surface and subsurface values, and eventually in determining (de)coupled values. It should be emphasized that the analysis was primarily focused on examining the trends and relations between surface and subsurface soil moisture. Moreover, it was not intended to contradict or replace other models for estimating soil moisture or examining its patterns.

A distributed lag non-linear model (DLNM) developed by Gasparrini et al. (2010) was applied to the 5 cm and 40 cm time series datasets at the study sites. Briefly, the model is capable of simultaneously representing both functional dependency and delayed response between exposure and response values. We considered surface soil moisture as the exposure values that produced delayed effects to the response values at the subsurface. A non-linear model was selected in order to capture the non-linear dynamics of flow and transport along the soil profile (Mohanty and Skaggs, 2001; Kim and Barros, 2002). Furthermore, DLNM offered enough flexibility to model a variety of dependencies in the time series dataset by selecting a suitable basis function. DLNM could be thought of as equivalent to a linear time series model (e.g. autoregressive model) just as a generalized linear model is equivalent to a linear model.

In assessing lagged dependence, event scale patterns were of interest rather than large scale trends within the time series (Wilson et al., 2004). This required seasonal patterns to be addressed prior to applying the DLNM. This was done by fitting a loess function to the time series and then subtracting it from the original soil moisture values (Cleveland et al., 1990). Removal of seasonality was further justified by the scatterplot results (see Section 4.1). The influence of seasonality on the vertical soil moisture variability is indicated by clustering of observation points occurring within the same months (fig.4). De-seasonalized soil moisture values were used for identifying (de)coupled soil moisture conditions.

*For consistency in modeling, the range of surface soil moisture values used was from 0-0.50 cm³ cm⁻³. This was based on the highest surface soil moisture value encountered among the four sites. A lag value of up to 30 days was considered long enough to investigate delayed effects. This period also approximated the recurrence of heavy rainfall within the study sites. A spline function was the basis function chosen to represent the functional dependence as well as delayed effects as it offered flexibility to capture non-linearities. In addition, contributions from daily rainfall data were used to incorporate current and past meteorological conditions. This was applied as a covariate and was represented with an additional basis function. We only considered delayed effects in vertical flow as lateral movement is deemed negligible in flat to slightly sloping terrain (Table.1). The analysis was performed in R software using *dlm* (Gasparrini, 2011) and *mgcv* (Wood, 2006a) packages.*

The following section concisely describes the mathematical formulation of a DLNM. However, the reader may choose to skip this section as the general methods applied have already been described in the text above. For a more detailed explanation, readers are referred to Gasparrini et al. (2010) and Gasparrini et al. (2017).

3.3 Evaluating (de)coupled soil moisture values

Application of a DLNM resulted in the estimation of parameter β for each surface soil moisture value. This indicated the strength of dependence between surface and subsurface soil moisture. Higher β values indicated stronger dependence or coupling between the two. Hence, we referred to β as the relative influence of surface soil moisture on subsurface values."

"We incorporated delayed or lagged effects in evaluating the relation between surface and subsurface values, and eventually in determining (de)coupled values. It should be emphasized that the analysis was primarily focused on examining the trends and relations between surface and subsurface soil moisture. Moreover, it was not intended to contradict or replace other models for estimating soil moisture or examining its patterns.

A distributed lag non-linear model (DLNM) developed by Gasparrini et al. (2010) was applied to the 5 cm and 40 cm time series datasets at the study sites. Briefly, the model is capable of simultaneously representing both functional

dependency and delayed response between exposure and response values. We considered surface soil moisture as the exposure values that produced delayed effects to the response values at the subsurface. A non-linear model was selected in order to capture the non-linear dynamics of flow and transport along the soil profile (Mohanty and Skaggs, 2001; Kim and Barros, 2002). Furthermore, DLNM offered enough flexibility to model a variety of dependencies in the time series dataset by selecting a suitable basis function. DLNM could be thought of as equivalent to a linear time series model (e.g. autoregressive model) just as a generalized linear model is equivalent to a linear model.

In assessing lagged dependence, event scale patterns were of interest rather than large scale trends within the time series (Wilson et al., 2004). This required seasonal patterns to be addressed prior to applying the DLNM. This was done by fitting a loess function to the time series and then subtracting it from the original soil moisture values (Cleveland et al., 1990). Removal of seasonality was further justified by the scatterplot results (see Section 4.1). The influence of seasonality on the vertical soil moisture variability is indicated by clustering of observation points occurring within the same months (fig.4). De-seasonalized soil moisture values were used for identifying (de)coupled soil moisture conditions.

For consistency in modeling, the range of surface soil moisture values used was from 0-0.50 cm³ cm⁻³. This was based on the highest surface soil moisture value encountered among the four sites. A lag value of up to 30 days was considered long enough to investigate delayed effects. This period also approximated the recurrence of heavy rainfall within the study sites. A spline function was the basis function chosen to represent the functional dependence as well as delayed effects as it offered flexibility to capture non-linearities. In addition, contributions from daily rainfall data were used to incorporate current and past meteorological conditions. This was applied as a covariate and was represented with an additional basis function. We only considered delayed effects in vertical flow as lateral movement is deemed negligible in flat to slightly sloping terrain (Table.1). The analysis was performed in R software using *dlnm* (Gasparrini, 2011) and *mgcv* (Wood, 2006a) packages.

The following section concisely describes the mathematical formulation of a DLNM. However, the reader may choose to skip this section as the general methods applied have already been described in the text above. For a more detailed explanation, readers are referred to Gasparrini et al. (2010) and Gasparrini et al. (2017).

####Some lines for the mathematical description of DLNM is not included here. The structure of this section was not revised#####

3.3 Evaluating (de)coupled soil moisture values

Application of a DLNM resulted in the estimation of parameter β for each surface soil moisture value. This indicated the strength of dependence between surface and subsurface soil moisture. Higher β values indicated stronger dependence or coupling between the two. Hence, we referred to β as the relative influence of surface soil moisture on subsurface values."

3) The analysis here is based solely on vertically-discrete soil moisture measurements (i.e. soil moisture observed at a depth of 40 cm). However, in remote-sensing, modeling and data assimilation, soil moisture estimates reflect vertically-integrated values (within the measurement depth of the remote sensor or across the vertically-discrete soil layer specified in a land model). Will the transition between vertically-discrete versus vertically-averaged soil moisture values affect the applicability of these results in a modeling or data assimilation context? I would recommend more discussion on this point.

We thank the reviewer for raising this point. We added another paragraph in section 5.2 in page 12 starting from line 4 to tackle this topic.

"Although the study focused on vertically discrete values, the results are also applicable for depth-average values commonly used in remote sensing and DA applications. This requires that the vertically discrete values adequately capture the overall dynamics within zone being investigated. In such a case, we infer that the translation to depth-averaged values would result in (de)coupled values that are close, but not identical, to the values obtained when only comparing two discrete depths. As an illustration, we calculated the depth-average values using all the available measurements at each site (i.e. 5, 10, 20 and 40 cm depth) following the formula from Qiu et al. (2014). Figure 9 (left) reveals highly similar dynamics for both discrete and depth-average values. Therefore, it can be expected that the results from a regression and DLNM analyses using depth-average values would be highly similar to the original results in fig.5 and fig.8. However, if the vertically discrete values insufficiently represent the subsurface dynamics, larger deviations in the resulting decoupled values can be expected."




Figure 9. Subsurface soil moisture dynamics from vertically-discrete (40cm) and depth-average value. Left: Time series of soil moisture at 40 cm and depth-averaged values. The dynamics observed for depth-average values are highly similar to those at 40 cm. The scatterplots on the right further confirms that these two sets of values are highly correlated.

4) While the manuscript is generally well-written, it does contain a large number of minor English usage errors. Additional proof-reading is recommended.

There was another round of proof reading to check for textual and grammatical errors.

List of changes made to the manuscript:

1. Added sentences or paragraphs to methods, results, and discussion sections based the comments from the reviewers.
2. Added figures to methods and discussion based on the reviewer comments
3. Changed some text after re-checking the grammar and spelling of the whole manuscript

Using lagged dependence to identify (de)coupled surface and subsurface soil moisture values

Coleen D.U. Carranza¹, Martine J. van der Ploeg¹, and Paul J.J.F. Torfs²

¹Soil Physics and Land Management Group, Wageningen University, the Netherlands

²Hydrology and Quantitative Water Management Group, Wageningen University, the Netherlands

Correspondence to: Coleen Carranza (coleen.carranza@wur.nl)

Abstract. Recent advances in radar remote sensing popularized [the](#) mapping of surface soil moisture at different spatial scales. Surface soil moisture measurements are used in combination with hydrological models to determine subsurface soil moisture values. However, variability of soil moisture across the soil column is important for estimating depth-integrated values as decoupling between surface and subsurface can occur. In this study, we ~~employed~~ [employ](#) new methods to investigate the occurrence of (de)coupling between surface and subsurface soil moisture. ~~Lagged~~ [Using time series datasets, lagged](#) dependence was incorporated in assessing (de)coupling with the idea that surface soil moisture conditions will be reflected at the subsurface after a certain delay. ~~An exploratory step using residuals from a fitted loess function was performed as a posteriori information to determine (de)coupled values.~~ The main approach ~~was applying~~ [involves the application of](#) a distributed lag non-linear model (DLNM) to simultaneously represent both [the](#) functional relation and ~~lag structure~~ [the lag structure in the](#) [time series. The results of an exploratory analysis using residuals from a fitted loess function serve as a posteriori information to determine \(de\)coupled values.](#) Both methods allow for a range of (de)coupled soil moisture values to be quantified. Results provide new insights on the decoupled range as its occurrence [among the sites investigated](#) is not limited to dry conditions.

1 Introduction

Although recent decades have seen great advances in remote sensing applications for mapping surface soil moisture (Jackson, 1993; Njoku et al., 2003; Mohanty et al., 2017), most hydrological studies that make use of soil moisture data require integrated values over a certain soil depth (Brocca et al., 2017). Extrapolation of surface soil moisture from remote sensing techniques to depths beyond the sensor's capacity (up to ~~5cm~~ [5 cm](#)) is not a trivial task given the spatio-temporal variability of soil moisture. The vertical distribution of soil moisture, which determines integrated soil moisture content over a soil column, is rarely uniform as more pronounced dynamics are expected closer to the surface compared to deeper in the soil (Hupet and Vanclooster, 2002). Currently, information derived from remote sensing are assimilated into hydrological models to obtain integrated soil moisture values (Houser et al., 1998; Das et al., 2008). However, Kumar et al. (2009) stressed that it is important to assess vertical variability, especially the strength of coupling between surface and subsurface soil moisture ~~,~~ for improvement of data assimilation results. [Analyses of vertical soil moisture distributions also have important implications for modeling studies, as they could be used for calibration or validation of model parameters \(De Lannoy et al., 2006\).](#)

The amount of soil moisture at any given time is controlled by factors operating a different time scales. While prevailing atmospheric conditions directly affect surface layers and control the temporal dynamics of soil moisture (Albertson and Montaldo, 2003; Koster et al., 2004), it is the downward movement of water from the surface that dictates the amount of subsurface soil moisture at a given time (Belmans et al., 1983; Rodriguez-Iturbe et al., 1999). Flow rates to the subsurface are driven by hydraulic properties, which are in turn controlled by physical soil characteristics such as texture, bulk density, and structure. Relative to changes in atmospheric conditions, soil physical properties change over longer timescales. Vegetation further modifies vertical soil moisture distribution by root water uptake (Yu et al., 2007) and by changing soil structure (Angers and Caron, 1998). ~~Analyses of vertical soil moisture distributions also have important implications for modeling studies, as they could be used for calibration or validation of model parameters (De Lannoy et al., 2006).~~

Given the variability along ~~the a~~ soil column, during which conditions ~~does do~~ surface soil moisture reflect subsurface soil moisture? Several studies have investigated this relation to address the correspondence between surface and subsurface soil moisture content. One of the earliest studies is by Capehart and Carlson (1997) ~~using modeling outputs for comparison wherein they compared modeling outputs~~ with remote sensing measurements. Using very shallow depths of ~~5mm and 5cm, they observed deviations from linear correlation due to differences in drying rates which they referred to as “decoupling 10005 mm~~ and 5 cm, they referred to decoupling as the deviation from a linear correlation between these depths due to variable drying rates. Further assessment of decoupling from model-generated time series soil moisture data have been investigated using cross-correlation values (Martinez et al., 2008; Mahmood et al., 2012; Ford et al., 2014). High correlation to the subsurface was obtained using lagged values of surface soil moisture. However, cross-correlation is limited to providing a single value throughout the range of soil moisture encountered per lag. Furthermore, cross-correlation generally aims to evaluate the strength of lagged linear dependence between two variables (Shumway and Stoffer, 2010). However, lagged dependence between surface and subsurface soil moisture may not be linear given that non-linear processes determine water flow along the soil profile. Using in situ field measurements, Wilson et al. (2003) investigated spatial surface (~~0-6cm~~0-6 cm) and subsurface (~~0-30cm~~0-30 cm) soil moisture distribution by calculating statistical metrics and by means of a variogram. Decoupling between the two depths was observed which they suggested to be influenced by vegetation, especially root density at surface soil. Their results were also affected by the dry soil moisture range and emphasized the importance of distinguishing between surface and total soil moisture for future applications of remote sensing to atmospheric studies.

Based on previous studies, the term decoupling refers to a weak dependence between soil moisture contents at the surface and subsurface. Recognition of decoupling is important, however most studies have been limited to providing qualitative characterization of conditions when decoupling occurs (e.g. dry period). Only Capehart and Carlson (1997) identified a mid-range soil moisture ($\sim 0.3 \text{ cm}^3 \text{ cm}^{-3}$) when the surface and very near surface begin to decouple. Their results, however, are limited to a thin layer of the soil column. In this paper, our main objective is to quantitatively identify a range of surface soil moisture values that is decoupled from the subsurface. Furthermore, we consider depths greater than those investigated by Capehart and Carlson (1997). The ability to quantify (de)coupled surface and subsurface soil moisture contents will contribute to more effective estimation of depth-integrated soil moisture data using remote sensing methods and improved data assimilation results in hydrological models.

We ~~utilize~~utilized in situ time series datasets at depths of ~~5cm and 40cm~~5 cm and 40 cm to represent surface and sub-surface, respectively. Values outside the decoupled range are considered coupled since soil moisture is inherently bounded up a maximum value equal to soil's porosity. ~~Investigation~~The investigation of (de)coupling is based on the idea that surface conditions will be reflected at the subsurface after a certain delay indicating strong coupling between the two zones, and vice versa. More focus is given to the decoupled soil moisture range since it has greater implications for extrapolation of surface soil moisture values to deeper soil layers. We applied statistical methods to identify conditions of decoupling with no prior assumptions on the type of functional relation between surface and subsurface. As an exploratory step, we first assessed the dependence without considering lags using regression and residuals analysis. The main approach for assessing decoupling was application of distributed lag non-linear models (Gasparri et al., 2010) to incorporate both the lag structure and the functional relation between surface and subsurface soil moisture. Applications of distributed lag models to econometrics and environmental epidemiology have been well documented (Almon, 1965; Zanobetti et al., 2002; Bhaskaran et al., 2013; Wu et al., 2013). However, their application to hydrological studies have rarely been explored.

INSERT Fig.1 here

2 Description of datasets and study sites

Four time series datasets from the Twente soil moisture and temperature monitoring network (Dente et al., 2011) were used in this study (fig.1). Datasets from 2014-2016 are available with only short periods of missing data. The stations are located in agricultural fields with sensors installed at ~~5cm, 10cm, 20cm, and 40cm~~5 cm, 10 cm, 20 cm, and 40 cm depths. To investigate decoupling, only the ~~5cm and 40cm~~5 cm and 40 cm depths were considered because the largest possible distance was desired. Each station consists of EC-TM ECH2O capacitance probes (Decagon Devices, Inc., USA) that logged soil moisture data every 15 minutes. A calibration procedure using gravimetric measurements was applied prior to analysis (Dente et al., 2011).

Land cover in the area varies from corn in one field (SM05), to grass in two fields (SM05 and SM13), to a forest area (SM20). Values at ~~40cm~~40 cm capture the root zone of vegetation for each site. In reality, rooting depths vary and depend on species composition, climate, and plant growth rate. However, the depth considered would still allow for approximation of root zone conditions. The landscape is characterized by flat to slightly sloping terrain. It is important to note that SM20 is located at the eastern foot of a small hilly terrain. Throughout the study period, either land cover remained unchanged or the same crop was planted. The soil types for the stations range from coarse sandy soils to weakly silty soils (Wosten et al., 2013). A summary of the land cover and relevant characteristics of the stations are summarized in Table 1.

Soil moisture values were averaged into daily values to match the available daily rainfall data from the Dutch national weather service (KNMI). For SM13 and SM20, there are some missing data from the beginning of 2014. The datasets from SM13 begins on April 25, 2014 while SM20 begins on May 2, 2015 (fig.2).

INSERT Fig.2 here

INSERT Table 1 here

3 Methods

3.1 Regression and residuals analysis

As an exploratory step, the dependence between surface and subsurface soil moisture was initially visualized using scatterplots. Conditional means for every $0.01 \text{ cm}^3\text{cm}^{-3}$ interval and vertical bars representing \pm standard deviation were added to show changes in vertical variability across the soil moisture range. Longer standard deviation bars indicate higher vertical variability (fig.3, left). We referred to vertical variability as the unevenness or irregularity in the soil moisture distribution within a certain depth of interest along the vertical profile, in this case up to depths of 40 cm. For the rest of this paper, variability will refer to vertical soil moisture variability, unless otherwise stated. Points-The plotted points were colored per month to show any impacts of seasonality. The effect of rainfall was included by adjusting the sizes of the points proportional to rainfall intensity measured from the nearest KNMI stations. For the overall measure of dependence, Spearman's rank correlation coefficient R_s was computed for every pair of ranked values in the time series. This was chosen as the assumption of linear dependence is was not made.

A flexible non-parametric locally weighted regression function (commonly called a loess function, Cleveland and Devlin (1988)) was fitted along the soil moisture range. This was used to explore and identify trends across the soil moisture range. A linear regression was also fitted only for comparison. Residuals were analyzed further for variability not captured by the fitted function -

~~The cumulative residual variance, which more clearly shows the changes in (fig.3, right). The residuals variance for every $0.1 \text{ cm}^3\text{cm}^{-3}$ interval as well as the resulting cumulative residuals variance were analyzed to examine variability across the range, was further analyzed. A. The degree of variability was related to the slope of the cumulative variance line, with steep slopes indicating high variability. In addition, a significant change in slope between two neighboring points is identified along the cumulative residual variance line. This change occurs at an intermediate variance between two points was indicated by a significant change in the slope of the line. The soil moisture value , where a significant change in slope occurred was marked by θ_c , that this divides the soil moisture range into two groups. The group with the steeper slope is a steeper slope was interpreted as the decoupled range, and vice versa. Since the measured variance is sensitive to sample size, a correlation coefficient was calculated to determine if there was significant dependence between the two variables. Residuals variance were first normalized from 0-1 because of the varying soil moisture range encountered at each station.~~

Results of the exploratory methods are were considered *a posteriori* knowledge for analysis of lagged dependence and interpretation of results.

INSERT Fig.3 here

3.2 Analysis of Lagged Dependence

3.2.1 Cross correlation

Since decoupling is based on the strength of lagged dependence, the existence of lag between surface and subsurface soil moisture values was first determined. Cross-correlation is known to be a quick and easy method to apply for this objective.

- 5 Lagged values of surface soil moisture ~~are~~were correlated with instantaneous values at the subsurface. ~~Maximum~~A maximum cross-correlation at negative lags ~~would indicate~~indicated that surface soil moisture is leading subsurface soil moisture, and vice versa (Shumway and Stoffer, 2010). A 10-day lag was deemed long enough to show the presence of lag-lead relations in the time series since the maximum correlation ~~occurs~~occurred within this period.

3.2.2 Distributed lag non-linear model

- 10 ~~The main approach for investigating decoupling is based on lagged dependence~~We incorporated delayed or lagged effects in evaluating the relation between surface and subsurface values. ~~Decoupling is inferred when subsurface values show weak lagged dependence to surface soil moisture values. For the analysis, we only considered lag in vertical flow as lateral movement is deemed negligible in flat to slightly sloping terrain (Table.1). The strength of lagged dependence is determined using a , and eventually in determining the (de)coupled values. It should be emphasized that the analysis was primarily focused on examining~~
15 the trends and relation between surface and subsurface soil moisture. Moreover, it was not intended to replace other existing models for estimating soil moisture or examining its patterns.

- A distributed lag non-linear model (DLNM, (Gasparrini et al., 2010)). ~~DLNM simultaneously represents both exposure-response dependencies and delayed effects in the time series~~ developed by Gasparrini et al. (2010) was applied to the 5 cm and 40 cm time series datasets at the study sites. Briefly, the model is capable of simultaneously representing both functional the dependence
20 and delayed response between exposure and response values. We considered surface soil moisture as the exposure ~~which produces a delayed response in values that produced delayed effects to the response values at~~ the subsurface. A non-linear model ~~is used~~was selected in order to capture the non-linear dynamics of flow and transport along the soil profile (Mohanty and Skaggs, 2001; Kim and Barros, 2002). ~~In the following paragraphs, we provide a concise description of the DLNM concept. The full mathematical explanation of distributed lag non-linear models is described in Gasparrini et al. (2010) and~~
25 (Gasparrini et al., 2017) Furthermore, DLNM offered enough flexibility to model a variety of dependencies in the time series dataset by selecting a suitable basis function. As an analogy, a DLNM is to a linear time series model (e.g. autoregressive model) just as a generalized linear model is to a linear model, as can be seen in Eq. 1.

- In assessing lagged dependence, event scale patterns were of interest rather than large scale trends within the time series (Wilson et al., 2004). This required seasonal patterns to be addressed prior to applying the DLNM. This was done by fitting a
30 loess function to the time series and then subtracting it from the original soil moisture values (Cleveland et al., 1990). Removal of seasonality was further justified by the scatterplot results (see Section 4.1). The influence of seasonality on the vertical soil moisture variability is indicated by clustering of observation points occurring within the same months (fig.4). De-seasonalized soil moisture values were used for identifying (de)coupled soil moisture conditions.

For consistency in modeling, the range of surface soil moisture values used was from 0-0.50 cm³cm⁻³. This was based on the highest surface soil moisture value encountered among the four sites. A lag value of up to 30 days was considered long enough to investigate delayed effects. This period also approximated the recurrence of heavy rainfall within the study sites. A spline function was the basis function chosen to represent the functional dependence and delayed effects as it offered flexibility to capture non-linearities. In addition, contributions from daily rainfall data were used to incorporate current and past meteorological conditions. This was applied as a covariate and was represented with an additional basis function. We only considered delayed effects in vertical flow as lateral movement is deemed negligible in a flat to slightly sloping terrain (Table.1). The analysis was performed in R software using *dlnm* (Gasparrini, 2011) and *mgcv* (Wood, 2006a) packages.

~~From the general model of a time series,~~ The following section concisely describes the mathematical formulation of a DLNM. However, the reader may choose to skip this section as the general description of the methods applied have already been given in the text above. For a more detailed explanation, readers are referred to Gasparrini et al. (2010) and Gasparrini et al. (2017).

To more formally describe a DLNM, let us first consider a general time series model, where outcomes Y_t with $t = 1, \dots, n$ are can be described by:

$$g(\mu_t) = \alpha + \sum_{j=1}^J s_j(x_{tj}; \beta_j) + \sum_{k=1}^K \gamma_k u_{tk} \quad (1)$$

where $\mu \equiv E(Y)$, which is assumed to be derived from a Poisson distribution, and g is a monotonic link function. The functions s_j denote relationships between the variables x_j and vector parameters β_j . Other u_k variables with predictors are included in coefficients γ_k specifying to specify their related effects. The relation between x and $g(\mu)$ is represented by $s(x)$ through a basis function. The complexity of this estimated relationship depends on the type basis function chosen and its dimensiondimensions. In the presence of delayed effects, the outcome Y at any time t is explained by the past exposures x_{t-l} with l as the lag representing the elapsed time between exposure and response. The final goal of a DLNM is to simultaneously describe the dependency along both the predictor space and lag dimension. This is achieved by selecting two sets of basis functions that are combined to obtain the cross-basis functions (Gasparrini et al., 2010).

Within the DLNM framework, a response Y_t at time $t = 1$ is based on lagged occurrences of predictor x_t , which is represented by vector $q_t = [x_{t-l_0}; \dots; x_{t-L}]^T$. The minimum and maximum lags are given by l_0 and L_T , respectively. The function represents dependence through:

$$s(q, t) = s(x_{t,t-l_0}, \dots, x_{t-L}) = \sum_{l=l_0}^L f \cdot w(x_{t-L}, l) \quad (2)$$

where $f \cdot w(x_{t-L}, l)$ represents the exposure-lag-response function, which is composed of two marginal functions: the exposure-response function $f(x)$ and lag-response function $w(l)$ in the space of the lag. Parameterization of f and w is achieved

by application of the known basis functions to the vectors q_t and l . The result can be expressed as matrices \mathbf{R} and \mathbf{C} with dimensions $(L - l_0 + 1) \times v_x$ and $(L - l_0 + 1) \times v_l$, respectively.

The cross basis function s and parameterized coefficients $\boldsymbol{\eta}$ are given by:

$$s(x_{t,t-l_0}, \dots, x_{t-L}; \boldsymbol{\eta}) = (\mathbf{1}_{L-l_0+1}^T \mathbf{A}_t) \boldsymbol{\eta} = \mathbf{w}_t^T \boldsymbol{\eta} \quad (3)$$

- 5 The values of \mathbf{w} are derived from \mathbf{A}_t , which is computed from the row-wise Kronecker product between matrices \mathbf{R} and \mathbf{C} . The dependence is expressed through \mathbf{w} and parameters $\boldsymbol{\eta}$. The cross-basis function represents the integral of $s(x, t)$ over the interval $[l_0, L]$, summing the contributions from the exposure history. ~~Estimated~~ The estimated dependence to specific exposure values is determined by prediction of $\hat{\beta}$, called lag coefficients. The estimated $\hat{\beta}$ and covariance matrix $V(\hat{\beta})$ is given by:

$$\hat{\beta} = \mathbf{A}_x \hat{\boldsymbol{\eta}} \quad (4)$$

10 $V(\hat{\beta}) = \mathbf{A}_x V(\hat{\boldsymbol{\eta}}) \mathbf{A}_x^T \quad (5)$

~~The values of $\hat{\beta}$ indicate the strength of dependence between surface and subsurface soil moisture. Higher $\hat{\beta}$ values indicate stronger dependence or coupling between the two. Hence, we refer to $\hat{\beta}$ as the relative influence of surface soil moisture on subsurface values.~~ A further extension to DLNM is the application of penalties for smoothness of the lag structure and shrinkage of lag coefficients to null at very high lags. These penalties ~~are applied~~ were applied in the analysis using a second-
 15 order difference (Wood, 2006b) and varying ridge penalties (Obermeier et al., 2015; Gasparrini et al., 2017), respectively. Application of penalties ~~is was~~ based on the assumption that, at higher lags, the lag coefficients become smaller and approach the null value.

~~In order to identify a range that is decoupled, a threshold value ($\hat{\beta}_c$) must be specified. This value is comparable to the intermediate soil moisture θ_c identified from Section 4.1. The values of θ_c provided a suitable guide for identifying a threshold common to all four sites. The corresponding $\hat{\beta}$ values obtained at θ_c were very close to 1, therefore, setting the threshold $\hat{\beta}_c = 1$ seemed a reasonable choice. This was preferred over the exact~~

20

3.3 Evaluating (de)coupled soil moisture values

Application of a DLNM resulted in the estimation of parameter $\hat{\beta}$ at each θ_c since the latter was defined using exploratory methods at $lag = 0$. Using the chosen $\hat{\beta}_c = 1$, for each surface soil moisture values with $\hat{\beta} < 1$ are considered decoupled while
 25 those with $\hat{\beta} \geq 1$ are coupled.

~~In assessing lagged dependence, event scale analysis of decoupling is of interest rather than large scale patterns within the time series (Wilson et al., 2004). This requires seasonal patterns to be addressed, which was done by fitting a loess function and then subtracting it from the time series data (Cleveland et al., 1990). Removal of seasonality was further justified by scatterplot results (see Section 4.1). The influence of seasonality on vertical soil moisture variability is indicated by clustering~~

of observation points occurring within the same months (fig.4). De-seasonalized soil moisture values were used for determining decoupling from the time series datasets.

For consistency in modeling, value (Eq. 4 and 5). This indicated the range strength of dependence between surface and subsurface soil moisture. Higher $\hat{\beta}$ values indicated stronger dependence or coupling between the two. Hence, we referred to $\hat{\beta}$ as the relative influence of surface soil moisture values used was from 0-0.50 cm³ cm⁻³. This is based on the highest value encountered for surface soil moisture among the four sites. A lag value of up to 30 days was considered long enough to investigate delayed effects. This period also approximates the recurrence of heavy rainfall within the study sites. Spline function was the basis function chosen for both exposure-response and lag-response functions as it offers flexibility to capture non-linearities. In addition, contributions from daily rainfall data were used to incorporate current and past meteorological conditions. This was used as a covariate that was represented with an additional basis function. It is a covariate that is represented with an additional basis function. The analysis was performed in R software using *dlnm* (Gasparrini, 2014) and *mgev* (Wood, 2006a) packages on subsurface values.

4 Results

4.1 Regression and Residuals analysis

The overall dependence between surface and subsurface given by the Spearman's rank coefficient (R_s) range from 0.746 to 0.866 (fig.4). However, even with a high overall dependence, variability is not uniform across the soil moisture range (fig.4). Except for SM13, increased variability is observed towards drier soil moisture values. Furthermore, the degree of variability also differs among the four sites. Most The most pronounced variability is observed at SM13 and the least at SM05. Clustering of observation points occurring within the same months indicate that seasonality dictates soil moisture values and impacts soil moisture variability. Rainfall events measured on the same day do not show a clear effect on surface and subsurface soil moisture dependence. Observations with higher rainfall intensities appear scattered in the plots (fig.4). In addition, the said observation points do not necessarily fall along the fitted functions or at the wet soil moisture region of the scatterplots. As lag is not considered, the impact of rainfall on variability is not fully captured from in the scatterplots alone.

INSERT Fig.4 here

Assessment of the regression fit quality was performed by comparison using residual standard errors (RSE). The results for both linear and loess functions show highly similar values (fig.4). This indicates that, in this case, a linear function captures the relation between surface and subsurface values. Nevertheless, the more flexible loess function was preferred for further residuals analysis because of its slightly better model fit and, using only visual inspection of fig.4, it more closely approximates the calculated conditional mean.

Figure 5 shows the residual plots with lines of the cumulative residual-residuals variance. The change in slope of the line is a feature consistent for all sites regardless of the magnitude of residual variance. Furthermore, The changes in variability are more clearly observed from the residuals than from the standard deviation bars in the scatterplots. The location of the change

in slope β at θ_c is highlighted by the vertical dashed line. The Decoupled soil moisture range corresponds to the section of cumulative residuals variance line with a steeper slope.

Specifically, the range of decoupled surface soil moisture values (in $\text{cm}^3\text{cm}^{-3}$) is 0.08-0.21 for SM05, 0.12-0.27 for SM09, 0.30-0.39 for SM13, and 0.08-0.12 for SM20. Except for SM13, the decoupled values are within the dry to intermediate soil moisture range. The cumulative residuals variance line for SM13 appears to increase exponentially with increasing surface soil moisture. This differs from the other three sites which show a distinct decrease in slope at increasing soil moisture values. For SM20, a second point is identified with a change in slope. The flat line starting from 0.24-0.28 $\text{cm}^3\text{cm}^{-3}$ indicates there is still lowered variance at the very wet soil moisture range.

The correlation between normalized variance and sample size yielded a value of -0.24 (fig.6). This low correlation magnitude confirms that the variance obtained for the soil surface moisture intervals was not strongly influenced by the sample size used.

[INSERT Fig.6 here](#)

[INSERT Fig.7 here](#)

4.2 Cross-correlation

Figure 7 shows cross-correlation values at the four sites. Maximum correlation occurs at -1 to -2 days lag, except at SM20. This translates to a 1-2 day lead of surface soil moisture values. For SM20, the maximum correlation occurs at positive lags. Correlation values from lag = 0 to lag = 10 are almost equal at SM20. Although this indicates leading subsurface values, it does not eliminate the possibility of having a lag between surface and subsurface values (see Section 5.2). Other factors may play a role in having leading subsurface values in the cross-correlation plots. Hence, SM20 was still analyzed for decoupling using DLNM.

4.3 Distributed lag model

Figure 8 shows the overall $\hat{\beta}$ for each surface soil moisture value with 5% and 95% confidence intervals in shaded gray regions. In order to identify a range that is decoupled, a threshold value ($\hat{\beta}_c$) must be specified. This value is comparable to the intermediate soil moisture θ_c identified from Section 4.1. The values of θ_c provided a suitable guide for identifying a threshold common to all four sites (Table 2). The corresponding $\hat{\beta}$ values obtained at θ_c were very close to 1, therefore, setting the threshold $\hat{\beta}_c = 1$ seemed a reasonable choice. This was preferred over the exact $\hat{\beta}$ at each θ_c since the latter was defined using exploratory methods at lag = 0. Using the chosen $\hat{\beta}_c = 1$, surface soil moisture values with $\hat{\beta} < 1$ are considered decoupled while those with $\hat{\beta} \geq 1$ are coupled.

Based on $\hat{\beta}_c$, the identified decoupled values are generally in the dry to intermediate soil moisture range (fig.8), except for SM13 where decouple values are at the wet range. Table 2 shows the decoupled values identified based on the selected $\hat{\beta}_c$. Behavior-The behavior and trends of $\hat{\beta}$ also differ for each station. For instance, at SM05 and SM09, there is a general increase in $\hat{\beta}$ from dry towards wet surface soil moisture values. However, the predicted $\hat{\beta}$ in the very dry soil moisture range ($<0.10\text{cm}^3\text{cm}^{-3}$) are different for the two sites as decoupling is predicted for only SM05. At SM13, $\hat{\beta}$ values mostly fluctuate

around $\hat{\beta}_c$ and then proceed to decrease toward the decoupled wet range. The behavior of $\hat{\beta}$ for SM20 has an increasing trend also shows increasing $\hat{\beta}$ over a limited range. Among the four sites, SM20 has the smallest soil moisture range, only reaching a maximum of $0.28 \text{ cm}^3 \text{ cm}^{-3}$. The limited soil moisture range in SM20 leads to very high uncertainties in the predicted ($0.1 - 0.25 \text{ cm}^3 \text{ cm}^{-3}$). Outside this range, the estimated $\hat{\beta}$ for wet soil moisture conditions ($> 0.30 \text{ cm}^3 / \text{cm}^3$) values for SM20 were less than one and have very broad confidence intervals. Recall that the range used for DLNM was only for uniformity among the four study sites. The lack of or very few observations for very dry or very wet soil moisture conditions led to wider confidence intervals not only for SM20 but also for the other three sites. Compared to the three sites, the estimated $\hat{\beta}$ values for SM13 show decreasing values towards the wet soil moisture range ($> 0.3 \text{ cm}^3 \text{ cm}^{-3}$). From the intermediate to dry soil moisture conditions, the values fluctuate around the designated $\hat{\beta}_c$.

INSERT Fig.8 here

INSERT Table 2 here

5 Discussion

5.1 Decoupled soil moisture values

Regression and residuals analysis analyses show that there is an inherent vertical variability between surface and subsurface soil moisture values based on the lack of 1:1 correspondence between the two (fig.4). This inherent variability is also not uniform as higher variability is observed in at certain soil moisture ranges. The cumulative residual variance plots (fig.5) clearly indicate the soil moisture values where vertical variability starts to become consistently larger. The increase in variability further translates to weak lagged dependence which we observe as low $\hat{\beta}$ values from DLNM. The increase in vertical variability and weakening of lagged dependence is what we consider to be considered as decoupling between the surface and subsurface soil moisture.

Both residuals analysis and DLNM were successful in identifying a decoupled soil moisture range and there is good agreement between the results from both. Three out of four sites show decoupled values in the dry to intermediate soil moisture range (fig.5 and Table 2). These results agree with the known range where decoupling is expected (Capehart and Carlson, 1997; Hirschi et al., 2014; Wilson et al., 2003). For SM05 and SM09, the intermediate soil moisture value, θ_c that marks when decoupling begins (Table 2) is close to that identified by Capehart and Carlson (1997). They obtained a value of $0.3 \text{ cm}^3 \text{ cm}^{-3}$ as the point below which decoupling begins. However, results for SM13 do not conform to the traditional concept of decoupling. This result is significant as it implies that decoupling can may occur at any value and is not confined to dry soil moisture range.

The physical characteristics at SM13 allow some insights into the potential controls for decoupling vegetation type at each site exerts some influence on the soil moisture variability and the resulting (de)coupled values. First, the vegetation type affects how much ground surface is directly exposed to atmospheric conditions. Forested areas and grass fields are almost fully covered by vegetation compared to a corn field where the crops are organized in equidistant rows. Vegetation or canopy cover will determine how atmospheric conditions affect the soil moisture values. For instance, further inspection of the soil profile

showed a slight increase in silt content at 40cm compared to 5cm. The sensor is also located close to a small shallow ditch (Table 1). These factors contribute to increased water retention and soil water content at the 40cm depth which may affect decoupling at the site. The identified decoupled values occur during colder winter periods (fig.4). One factor that could potentially induce changes to soil physical characteristics during winter periods is the presence of burrowing animals that are in hibernation. Site inspection confirmed the presence of burrows at SM13. These burrows the amount of intercepted precipitation and evaporation are both dependent on vegetation cover. This in turn will have direct impacts to the surface soil moisture dynamics at each of the sites. For comparison, the variability given by the standard deviation bars in fig.4 and variance in fig.5 at the cornfield (SM09) is higher compared to that of the grass field (SM05) or the forested area (SM20). In addition, the forested area (SM20) has the smallest range of soil moisture values among the four sites. This may be due to the large intercepted rainfall by the forest canopy. Root water uptake (RWU) is another way by which vegetation affects soil moisture variability. RWU can have significant influence on the subsurface dynamics. The influence of RWU may vary for different vegetation types as it can be exerted over a range of depths, leading to differences in the resulting (de)coupled values.

Among the four sites, the subsurface trends observed for the 40 cm values at SM13 show consistently high values, which can be more pronounced during winter months. This resulted in decoupling during wet soil moisture conditions fig.8. This trend is different from the other three sites which only show a slight increase in the subsurface values. Further inspection of the time series data at SM13 reveals no sudden disturbance in the signal which could be attributed to errors in the sensor. Field investigation confirmed an increase in silt content at 40 cm compared to the upper layers. The increase in silt content promotes a decrease of hydraulic conductivity over depth that results in a slower vertical flow towards deeper layers. The presence of burrowing and hibernating animals was also observed at the site during winter. These create macropores which eventually alter the hydraulic properties of the soil (Kodešová et al., 2006; Beven and Germann, 2013). We infer that, at the measurement domain of the sensor, these burrows or macropores facilitated faster vertical flow to the subsurface. Alternatively, if the burrows produced voids around the measurement domain, this would result in lowered soil moisture or data gaps due to the loss of sensor to soil media contact. However, there were no gaps observed that coincided with the burrowing animals' period of hibernation. During precipitation events, soil moisture flowing from upper layers arrived more rapidly at 40 cm depths due to the presence of macropores. There it accumulated and flowed more slowly to deeper layers because of the low hydraulic conductivity promoted by the increase in silt content. The overall effect of these factors was the pronounced increase in soil moisture values at 40 cm compared to those at 5 cm during winter periods as observed from the time series dataset fig.2.

Site-specific characteristics at each station control the magnitude of variability as well as the range where-at which decoupling is observed. However, the occurrence of decoupling is independent of the magnitude of variability since it was observed from SM05 where variability is least up to SM13 where it is greatest. The methods applied in this study only identify conditions when decoupling occurs but do not explicitly determine its controls. Identification of controls for decoupling requires a separate analysis where mechanistic models or statistical approaches can be applied.

5.2 Assessing the use of lagged dependence for ~~identification of decoupling~~identifying decoupled conditions

To assess the applicability of the methods applied, we further discuss their strengths and weaknesses. We also present opportunities for further studies as well as foreseen limitations for other sites.

Strengths: ~~he~~The residuals analysis and DLNM methods allow quantification of a range of soil moisture values where decoupling occurs. This provides further extension to previous studies where decoupling is only described qualitatively. As seen from the results at the four sites, decoupling can occur at any soil moisture value, and is not confined to dry periods or ranges. Furthermore, by making no initial assumptions on data distributions and the type of functional relation and lag structure, the methods applied were considered robust. Non-linear functions were applied as they conform to the nonlinearity of water flow in the unsaturated zone. They can also handle a variety of bivariate dependence, even in cases where the relation is linear, as shown by the highly similar fit of the loess and linear functions in Section.4.1.

Weaknesses: The first aspect that needs to be further investigated is the selected ~~$\hat{\beta}$ value used~~ $\hat{\beta}_c$ value for identifying the decoupled soil moisture range. Although the selection in this study was based on trends identified from time series datasets, the methods applied should be tested further using other datasets to confirm the suitability of $\hat{\beta}_c = 1$ for other depths and soil types. ~~Another aspect~~The choice of β_c is crucial as it dictates which soil moisture values are expected to be decoupled. For instance, at the sites where decoupling occurs during dry conditions, a higher β_c value would enlarge the decoupled range. A similar effect would be expected for the site with decoupling during wet conditions. However, a lower β_c value could result to decoupling only during extreme soil moisture conditions (e.g very wet or very dry).

Another aspect to further examine is the use of cross-correlation for confirming the presence of leading surface soil moisture values. Results from SM20 show maximum correlation at positive lags which indicate leading subsurface values (fig.7). The weakness of using cross-correlation as a test for the presence of lag can be two-fold. First, cross-correlation can also capture the effect of subsurface dynamics such as groundwater influence and lateral flow. We infer that in SM20, subsurface dynamics dominates and masks the lag relation sought. An additional covariate representing subsurface dynamics was not included in the DLNM analysis since a dominant downward vertical flow was assumed. This assumption was based on the flat slopes encountered at SM20 (Table 1). Therefore, the occurrence of subsurface lateral flow or groundwater influence pose limitations to the applicability of DLNM for assessing decoupling. Second, cross-correlation is limited to evaluating linear lagged dependence. ~~Incorporating and in incorporating~~non-linear lagged dependence can make the test more robust. Equivalent methods exist (e.g. mutual information content (Qiu et al., 2014)) but they are much more computationally demanding when the goal is simply to check for the existence of lag-lead relation.

Opportunities: In relation to utilizing remote sensing techniques, our results imply that the accuracy of estimating subsurface values from surface soil moisture can be greatly affected by vertical coupling. Lower variability and hence lower uncertainties are expected in the coupled soil moisture range. Assessment of decoupling can be used in combination with modeling studies as a preliminary method to determine the range where variability is expected to be higher. Furthermore, it can be helpful in assessing whether simulation results capture the variabilities observed in both the coupled and decoupled ranges. Taking

decoupling into account can also assist in evaluating the necessity of complex models for simulating vertical soil moisture content.

For data assimilation (DA) applications, (de)coupling methods can be used for cross-comparison of the vertical coupling derived from DA model outputs with those observed from long term in situ measurements. This can aid in examining the adequacy of the assumed inherent connection between surface and subsurface values. As Kumar et al. (2009) pointed out, land surface models vary in their representation of the strength of this connection (e.g. weak or strong connection) which contributes the degree in which modeling results are improved. They also suggested that strong coupling is a more robust choice unless independent information suggests that a more decoupled surface-subsurface representation is more realistic. In this aspect, the analysis applied in this study could be a valuable tool in determining which type of surface-subsurface coupling is the more optimal choice. Furthermore, the assumed connection strength is adopted for the whole range of soil moisture values. The results of our analysis show that at any given site, decoupling will occur regardless of degree of soil moisture variability. A variable coupling strength could be adopted based on the soil moisture range where decoupling is likely to occur as an alternative to the single value for the whole range.

INSERT Fig.9 here

Although the study focused on vertically discrete values, the results are also applicable for depth-average values commonly used in remote sensing and DA applications. This requires that the vertically discrete values adequately capture the overall dynamics within zone being investigated. In such a case, we infer that the translation to depth-averaged values would result in (de)coupled values that are close, but not identical, to the values obtained when only comparing two discrete depths. As an illustration, we calculated the depth-average values using all the available measurements at each site (i.e. 5, 10, 20 and 40 cm depth) following the formula from Qiu et al. (2014). Figure 9 (left) reveals highly similar dynamics for both discrete and depth-average values. Therefore, it can be expected that the results from a regression and DLNM analyses using depth-average values would be highly similar to the original results in fig.5 and fig.8. However, if the vertically discrete values insufficiently represent the subsurface dynamics, larger deviations in the resulting decoupled values can be expected.

Limitations: In this study, only meteorological factors were incorporated in the DLNM analysis since vertical movement was assumed to be the dominant flow mechanism. However, the subsurface can also be influenced by lateral movement or groundwater by capillary rise. In such scenarios, decoupling will not be limited to changes in surface conditions. For this, SM20 provides an excellent example. This station is located at the foot of a small hill (fig.2) where the occurrence of lateral subsurface movement is highly probable. This shows that although the analysis would be limited to smaller scales, or even a single point, recognition of regional setting is important for interpretation of results. In addition, subsurface dynamics can also be affected by capillary rise in areas with shallow groundwater. For future applications, the effect of both capillary rise and lateral movements to subsurface dynamics should be assessed and included in the DLNM analysis, but caution should be exercised when interpreting the results. Assessment of decoupling with DLNM is deemed more applicable to areas where the subsurface has insignificant groundwater influence and where vertical downward movement is the dominant flow mechanism.

6 Conclusion

The methods applied in this study allow for investigation of vertical soil moisture variability. More importantly, application of DLNM allowed for decoupled soil moisture range to be quantitatively identified. The results also reveal that decoupling is not confined to dry soil moisture range as implied by previous studies. The reasons for decoupling are manifold and controls for the dry soil moisture range may differ from those for the wet range. The results of this study have implications for remote sensing and data assimilation methods, especially for uncertainties related to the use of surface soil moisture to obtain integrated soil moisture values.

Data availability. The datasets for soil moisture were obtained from the Water Resource Department of ITC-Twente University. At the moment, the datasets are not publicly available. Access to the datasets may be granted upon request from the institute thru Prof. Rogier van der Velde, PhD (r.vandervelde@utwente.nl).

Author contributions. Coleen Carranza and Martine van der Ploeg initially conceptualized the idea for investigating the relation between surface and subsurface soil moisture values. Paul Torfs provided significant contributions to the statistical analysis applied. All three authors contributed to writing and editing of the manuscript.

Competing interests. No competing interest present

Acknowledgements. The authors are grateful for the Water Resources Department of ITC-Twente University, the Netherlands for sharing the datasets from their network. [We thank the three anonymous reviews for providing critical insights that improved the manuscript.](#) This work is part of the research programme Optimizing Water Availability through Sentinel-1 Satellites (OWAS1S) with project number 13871 which is (partly) financed by the Netherlands Organisation for Scientific Research (NWO). ~~We also thank Demie Moore for reviewing the manuscript.~~

References

- Albertson, J. D. and Montaldo, N.: Temporal dynamics of soil moisture variability: 1. Theoretical basis, *Water Resources Research*, 39, 2003.
- Almon, S.: The distributed lag between capital appropriations and expenditures, *Econometrica: Journal of the Econometric Society*, pp. 178–196, 1965.
- 5 Angers, D. A. and Caron, J.: Plant-induced changes in soil structure: processes and feedbacks, in: *Plant-induced soil changes: processes and feedbacks*, pp. 55–72, Springer, 1998.
- Belmans, C., Wesseling, J., and Feddes, R. A.: Simulation model of the water balance of a cropped soil: SWATRE, *Journal of hydrology*, 63, 271–286, 1983.
- Beven, K. and Germann, P.: Macropores and water flow in soils revisited, *Water Resources Research*, 49, 3071–3092, 2013.
- 10 Bhaskaran, K., Gasparrini, A., Hajat, S., Smeeth, L., and Armstrong, B.: Time series regression studies in environmental epidemiology, *International journal of epidemiology*, 42, 1187–1195, 2013.
- Brocca, L., Ciabatta, L., Massari, C., Camici, S., and Tarpanelli, A.: Soil Moisture for Hydrological Applications: Open Questions and New Opportunities, *Water*, 9, 140, 2017.
- Capenhart, W. J. and Carlson, T. N.: Decoupling of surface and near-surface soil water content: A remote sensing perspective, *Water Resources*
- 15 *Research*, 33, 1383–1395, 1997.
- Cleveland, R. B., Cleveland, W. S., and Terpenning, I.: STL: A seasonal-trend decomposition procedure based on loess, *Journal of Official Statistics*, 6, 3, 1990.
- Cleveland, W. S. and Devlin, S. J.: Locally weighted regression: an approach to regression analysis by local fitting, *Journal of the American statistical association*, 83, 596–610, 1988.
- 20 Das, N., Mohanty, B., Cosh, M., and Jackson, T.: Modeling and assimilation of root zone soil moisture using remote sensing observations in Walnut Gulch Watershed during SMEX04, *Remote Sensing of Environment*, 112, 415–429, 2008.
- De Lannoy, G. J., Verhoest, N. E., Houser, P. R., Gish, T. J., and Van Meirvenne, M.: Spatial and temporal characteristics of soil moisture in an intensively monitored agricultural field (OPE 3), *Journal of Hydrology*, 331, 719–730, 2006.
- Dente, L., Verkedy, Z., Su, Z., and Ucer, M.: Twente Soil Moisture and Soil Temperature Monitoring Network, Tech. rep., University of Twente—ITC: Enschede, The Netherlands, http://www.itc.nl/library/papers_2011/scie/dente_twe.pdf, 2011.
- 25 Ford, T., Harris, E., and Quiring, S.: Estimating root zone soil moisture using near-surface observations from SMOS, *Hydrology and Earth System Sciences*, 18, 139–154, 2014.
- Gasparrini, A.: Distributed lag linear and non-linear models in R: the package dlnm, *Journal of statistical software*, 43, 1, 2011.
- Gasparrini, A., Armstrong, B., and Kenward, M.: Distributed lag non-linear models, *Statistics in medicine*, 29, 2224–2234, 2010.
- 30 Gasparrini, A., Scheipl, F., Armstrong, B., and Kenward, M.: A penalized framework for distributed lag non-linear models, *Biometrics*, 2017.
- Hirschi, M., Mueller, B., Dorigo, W., and Seneviratne, S.: Using remotely sensed soil moisture for land–atmosphere coupling diagnostics: The role of surface vs. root-zone soil moisture variability, *Remote sensing of environment*, 154, 246–252, 2014.
- Houser, P. R., Shuttleworth, W. J., Famiglietti, J. S., Gupta, H. V., Syed, K. H., and Goodrich, D. C.: Integration of soil moisture remote sensing and hydrologic modeling using data assimilation, *Water Resources Research*, 34, 3405–3420, 1998.
- 35 Hupet, F. and Vanclooster, M.: Intraseasonal dynamics of soil moisture variability within a small agricultural maize cropped field, *Journal of Hydrology*, 261, 86–101, 2002.
- Jackson, T. J.: III. Measuring surface soil moisture using passive microwave remote sensing, *Hydrological processes*, 7, 139–152, 1993.

- Kim, G. and Barros, A. P.: Space–time characterization of soil moisture from passive microwave remotely sensed imagery and ancillary data, *Remote sensing of environment*, 81, 393–403, 2002.
- Kodešová, R., Kodeš, V., Žigová, A., and Šimnek, J.: Impact of plant roots and soil organisms on soil micromorphology and hydraulic properties, *Biologia*, 61, S339–S343, 2006.
- 5 Koster, R. D., Dirmeyer, P. A., Guo, Z., Bonan, G., Chan, E., Cox, P., Gordon, C., Kanae, S., Kowalczyk, E., Lawrence, D., et al.: Regions of strong coupling between soil moisture and precipitation, *Science*, 305, 1138–1140, 2004.
- Kumar, S. V., Reichle, R. H., Koster, R. D., Crow, W. T., and Peters-Lidard, C. D.: Role of subsurface physics in the assimilation of surface soil moisture observations, *Journal of hydrometeorology*, 10, 1534–1547, 2009.
- Mahmood, R., Littell, A., Hubbard, K. G., and You, J.: Observed data-based assessment of relationships among soil moisture at various
10 depths, precipitation, and temperature, *Applied Geography*, 34, 255–264, 2012.
- Martinez, C., Hancock, G., Kalma, J., and Wells, T.: Spatio-temporal distribution of near-surface and root zone soil moisture at the catchment scale, *Hydrological Processes*, 22, 2699–2714, 2008.
- Mohanty, B. and Skaggs, T.: Spatio-temporal evolution and time-stable characteristics of soil moisture within remote sensing footprints with varying soil, slope, and vegetation, *Advances in water resources*, 24, 1051–1067, 2001.
- 15 Mohanty, B. P., Cosh, M. H., Lakshmi, V., and Montzka, C.: *Soil Moisture Remote Sensing: State-of-the-Science*, 16, <https://doi.org/10.2136/vzj2016.10.0105>, 2017.
- Njoku, E. G., Jackson, T. J., Lakshmi, V., Chan, T. K., and Nghiem, S. V.: Soil moisture retrieval from AMSR-E, *IEEE transactions on Geoscience and remote sensing*, 41, 215–229, 2003.
- Obermeier, V., Scheipl, F., Heumann, C., Wassermann, J., and Küchenhoff, H.: Flexible distributed lags for modelling earthquake data,
20 *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 64, 395–412, 2015.
- Qiu, J., Crow, W. T., Nearing, G. S., Mo, X., and Liu, S.: The impact of vertical measurement depth on the information content of soil moisture times series data, *Geophysical Research Letters*, 41, 4997–5004, 2014.
- Rodriguez-Iturbe, I., Porporato, A., Ridolfi, L., Isham, V., and Cox, D.: Probabilistic modelling of water balance at a point: the role of climate, soil and vegetation, *Proceedings of the Royal Society of London Series A*, 455, 3789, 1999.
- 25 Shumway, R. H. and Stoffer, D. S.: *Time series analysis and its applications: with R examples*, Springer Science & Business Media, 2010.
- Wilson, D. J., Western, A. W., Grayson, R. B., Berg, A. A., Lear, M. S., Rodell, M., Famiglietti, J. S., Woods, R. A., and McMahon, T. A.: Spatial distribution of soil moisture over 6 and 30cm depth, Mahurangi river catchment, New Zealand, *Journal of Hydrology*, 276, 254–274, 2003.
- Wilson, D. J., Western, A. W., and Grayson, R. B.: Identifying and quantifying sources of variability in temporal and spatial soil moisture
30 observations, *Water Resources Research*, 40, 2004.
- Wood, S.: *Generalized Additive Models: An Introduction with R*, CRC Press, 2006a.
- Wood, S. N.: Low-Rank Scale-Invariant Tensor Product Smooths for Generalized Additive Mixed Models, *Biometrics*, 62, 1025–1036, <https://doi.org/10.1111/j.1541-0420.2006.00574.x>, 2006b.
- Wosten, J., de Vries, F., Hoogland, T., Massop, H., Veldhuizen, A., Vroon, H., Wesseling, J., Heijkers, J., and Bolman, A.: BOFEK2012, de
35 nieuwe bodemfysische schematisatie van Nederland, Tech. rep., Alterra, 2013.
- Wu, W., Xiao, Y., Li, G., Zeng, W., Lin, H., Rutherford, S., Xu, Y., Luo, Y., Xu, X., Chu, C., et al.: Temperature–mortality relationship in four subtropical Chinese cities: a time-series study using a distributed lag non-linear model, *Science of the Total Environment*, 449, 355–362, 2013.

Yu, G.-R., Zhuang, J., Nakayama, K., and Jin, Y.: Root water uptake and profile soil water as affected by vertical root distribution, *Plant Ecology*, 189, 15–30, <https://doi.org/10.1007/s11258-006-9163-y>, <https://doi.org/10.1007/s11258-006-9163-y>, 2007.

Zanobetti, A., Schwartz, J., Samoli, E., Gryparis, A., Touloumi, G., Atkinson, R., Le Tertre, A., Bobros, J., Celko, M., Goren, A., et al.: The temporal pattern of mortality responses to air pollution: a multicity assessment of mortality displacement, *Epidemiology*, 13, 87–93, 2002.




Figure 1. Location of study site in the eastern part of the Netherlands (inset). Triangles represent stations in-used within the Twente soil moisture and temperature monitoring network (Dente et al., 2011). Squares represent meteorological stations. Symbols with similar colors indicate the pair of measurements used for the analysis.




Figure 2. Time series plots of surface (5cm-5 cm in light blue) and subsurface (40cm-40 cm in dark blue) soil moisture. Vertical black bars at the top show daily precipitation data from the nearest KNMI station.




Figure 3. Schematic diagram using hypothetical soil moisture values to show vertical variability. Left: Scatterplot showing the trend with a fitted loess function. The variability can be seen using the standard deviation bars. Right: Scatterplot of the residuals from the fitted function. Soil moisture variability is visible from the variance given by the vertical bars and the cumulative residuals variance given by the black line. A change in variability at an intermediate soil moisture value is marked by a change in the slope of the cumulative variance line, indicated by θ_c .




Figure 4. Scatter plots of $5\text{cm}-5\text{ cm}$ vs $40\text{cm}-40\text{ cm}$ soil moisture values at $\text{lag}=0$. Colors correspond to the months in a year and sizes of points are proportional to rainfall intensity. Trends along the soil moisture range shown with the fitted loess function (black line). Vertical variability across the soil moisture range is indicated by the lengths of standard deviation bars. A linear function (red line) is also fitted for comparison. The overall dependence using Spearman's rank correlation R_s is given in the upper left corner each plot. Residual standard errors (RSE) for loess (lo) and linear (lm) fits are also shown.




Figure 5. Residual variance plots from the fitted loess function. Vertical bars at the bottom of each plot represent the variance for every $0.1 \text{ cm}^3 \text{ cm}^{-3}$ interval. Cumulative The cumulative residual variance (gray line) shows a change in slope at an intermediate soil moisture value (referred to as θ_c , indicated by the vertical black dashed line). The change in slope This separates the soil moisture values into a range with higher variance (steeper slope) and another range with lower variance (gentler slopes). The range with higher variance is considered decoupled, and vice versa.

Table 1. Summary of land cover descriptions at each station covering the period of 2014–2016. Soil descriptions and codes are based on BOFEK 2012 (Wosten et al., 2013). Slope Both the slope and distance to nearest ditch were determined from 5m resolution DEM. Datasets are from 2016 and were obtained from the publicly available national topographic database of the Netherlands (TOPNL)

Station No.	Land cover	BOFEK Soil description	Slope (degrees)	Aerial distance to nearest ditch (m)
SM05	Grass	Loamy sandy soils with a thick cultivated layer(317)	2.22	18.97
SM09	Corn	Weakly loamy sand soils with a thick cultivated layer(311)	2.70	1.41
SM13	Grass	Weak silty soils (podsols)(304)	1.0	17.09
SM20	Forest	Coarse sand (podsols)(320)	2.30	875.26




Figure 6. Scatterplot of sample size vs. normalized residual variance calculated for each $0.01 \text{ cm}^3 \text{ cm}^{-3}$ interval. Colors indicate soil moisture conditions at each point. The plot of points indicate very weak linear dependence quantified, which is further confirmed by a -0.24 correlation coefficient.




Figure 7. Cross-correlation plots of 5cm vs 40cm soil moisture values. Lagged 5cm The lagged surface soil moisture values at 5 cm are correlated with 40cm subsurface values at 40 cm. A 1-2 day lead of surface soil moisture is observed, except for SM20. This is indicated by having maximum the correlation values at lag-lags of -1 to -2 days. At SM20, the maximum correlation occurs at positive lags.




Figure 8. The relative influence of surface soil moisture on subsurface values obtained by summing the predicted $\hat{\beta}$ along the 30-day lag. The threshold value ($\hat{\beta}_c$) used to identify the decoupled range is indicated by the horizontal line. Surface soil moisture values below $\hat{\beta}_c$ are considered decoupled. The 5% and 95% confidence intervals of the predicted values are shown as shaded regions.

Table 2. List of surface soil moisture values (SSM in $\text{cm}^3\text{cm}^{-3}$) obtained from fig.5 and fig.8. SSM at θ_c in fig.5 were used to determine $\hat{\beta}$ in fig.8. A common threshold $\hat{\beta}_c$ was used for all sites since $\hat{\beta}$ are all close to 1. The resulting decoupled SSM values are shown in the fourth column.

Station No.	SSM at θ_c	$\hat{\beta}$ at θ_c	Decoupled values using threshold $\hat{\beta}_c = 1$
SM05	0.21	0.90	<0.24
SM09	0.27	0.97	<0.28
SM13	0.30	1.18	>0.34
SM20	0.12	0.94	0.16>SSM>0.23




Figure 9. Subsurface soil moisture dynamics for vertically-discrete (40cm) and depth-average value. Left: Time series of soil moisture at 40 cm and depth-averaged values. The dynamics observed for depth-average values are highly similar to those at 40 cm. Right: Scatterplot showing that these two sets of values are highly correlated.