

# Second International Guidelines for the Diagnosis and Management of Hereditary Hemorrhagic Telangiectasia

Marie E. Faughnan, MD, MSc; Johannes J. Mager, MD, PhD; Steven W. Hetts, MD; Valerie A. Palda, MD, MSc; Kelly Lang-Robertson; Elisabetta Buscarini, MD; Erik Deslandres, MD; Raj S. Kasthuri, MD; Andrea Lausman, MD; David Poetker, MD, MA; Felix Ratjen, MD; Mark S. Chesnutt, MD; Marianne Clancy, RDH, MPA; Kevin J. Whitehead, MD; Hanny Al-Samkari, MD; Murali Chakinala, MD; Miles Conrad, MD; Daniel Cortes, BScPhm; Claudia Crocione; Jama Darling, MD; Els de Gussem, MD; Carol Derksen; Sophie Dupuis-Girod, MD, PhD; Patrick Foy, MD; Urban Geisthoff, MD; James R. Gossage, MD; Adrienne Hammill, MD; Ketil Heimdal, MD; Katharine Henderson, MS, CGC; Vivek N. Iyer, MD, MPH; Anette D. Kjeldsen, MD; Masaki Komiyama, MD; Kevin Korenblatt, MD; Jamie McDonald, MS, CGC; Jack McMahon; Justin McWilliams, MD; Mary E. Meek, MD; Meir Mei-Zahav, MD; Scott Olitsky, MD, MBA; Sara Palmer, PhD; Rose Pantalone, RN; Jay F. Piccirillo, MD; Beth Plahn, RN, MHA; Mary E.M. Porteous, MD; Marco C. Post, MD, PhD; Ivan Radovanovic, MD; Paul J. Rochon, MD; Josanna Rodriguez-Lopez, MD; Carlo Sabba, MD; Marcelo Serra, MD; Claire Shovlin, PhD, MA; Dennis Sprecher, MD; Andrew J. White, MD; Ingrid Winship, MBChB, MD; and Roberto Zarrabeitia, MD

**Description:** Hereditary hemorrhagic telangiectasia (HHT) is an autosomal dominant disease with an estimated prevalence of 1 in 5000 that is characterized by the presence of vascular malformations (VMs). These result in chronic bleeding, acute hemorrhage, and complications from shunting through VMs. The goal of the Second International HHT Guidelines process was to develop evidence-based consensus guidelines for the management and prevention of HHT-related symptoms and complications.

**Methods:** The guidelines were developed using the AGREE II (Appraisal of Guidelines for Research and Evaluation II) framework and GRADE (Grading of Recommendations Assessment, Development and Evaluation) methodology. The guidelines expert panel included expert physicians (clinical and genetic) in HHT from 15 countries, guidelines methodologists, health care workers, health care administrators, patient advocacy representatives, and persons with HHT. During the preconference process, the expert panel generated clinically relevant questions in 6 priority topic areas. A systematic literature search was done in

June 2019, and articles meeting a priori criteria were included to generate evidence tables, which were used as the basis for recommendation development. The expert panel subsequently convened during a guidelines conference to conduct a structured consensus process, during which recommendations reaching at least 80% consensus were discussed and approved.

**Recommendations:** The expert panel generated and approved 6 new recommendations for each of the following 6 priority topic areas: epistaxis, gastrointestinal bleeding, anemia and iron deficiency, liver VMs, pediatric care, and pregnancy and delivery (36 total). The recommendations highlight new evidence in existing topics from the first International HHT Guidelines and provide guidance in 3 new areas: anemia, pediatrics, and pregnancy and delivery. These recommendations should facilitate implementation of key components of HHT care into clinical practice.

*Ann Intern Med.* doi:10.7326/M20-1443

Annals.org

For author, article, and disclosure information, see end of text.

This article was published at Annals.org on 8 September 2020.

**H**ereditary hemorrhagic telangiectasia (HHT) is an autosomal dominant disease with an estimated prevalence of approximately 1 in 5000 (1). It is characterized by clinically significant vascular malformations (VMs) of skin and mucous membranes of the nose and gastrointestinal (GI) tract as well as the brain, lung, and liver. It is underdiagnosed, and a long diagnostic delay is common (2). A diagnosis of HHT allows appropriate screening and preventive treatment to be undertaken in a patient and their affected family members. The most common symptom of HHT, epistaxis, has an age-related expression, as does the appearance of the typical telangiectasia (3). Consensus clinical diagnostic criteria known as the Curaçao criteria were published in 2000 (4) (Table 1 of Supplement 1, available at Annals.org) and upheld in the first International HHT Guidelines (5). The first guidelines also recommended genetic testing for HHT diagnosis, primarily for asymptomatic persons from a family with known HHT, as detailed in the Table. In 97% of patients with a definite clinical diagnosis of HHT, a causative mutation is identified in one of these genes: endoglin (*ENG*, HHT type 1), activin receptor-like kinase-1 (*ACVRL1*, HHT type 2), and Mothers against decapentaplegic homolog 4 (*SMAD4*, juvenile polyposis-HHT overlap) (6).

The goal of this Second International HHT Guidelines process was to develop evidence-informed consensus guidelines regarding the diagnosis of HHT, prevention of HHT-related complications, and treatment of symptomatic disease in areas not previously addressed by guidelines and those where significant new literature had been published. Several other recommendations from the first International HHT Guidelines were not reassessed during this process and remain currently recommended (Table).

## METHODS

The Second International HHT Guidelines were developed using the AGREE II (Appraisal of Guidelines for

### See also:

Web-Only  
Supplements

**Table.** Clinical Recommendations From the Second International HHT Guidelines and Currently Recommended Clinical Recommendations From the First International HHT Guidelines\*

Listed here are the clinical recommendations with  $\geq 80\%$  consensus at the Second International HHT Guidelines conference. These are followed by the clinical recommendations from the first International Guidelines that had  $\geq 80\%$  consensus and were not reassessed by the 2019 International Guidelines Working Group. The expert panel is aware of new evidence and insights regarding some of these existing guidelines, and they have been prioritized for updating in the next guidelines process (Part 4 of Supplement 2, available at [Annals.org](https://annals.org)).

#### Epistaxis management

- A1:** The expert panel recommends that patients with HHT-related epistaxis use moisturizing topical therapies that humidify the nasal mucosa to reduce epistaxis.\*  
Quality of evidence: moderate (agreement, 98%)  
Strength of recommendation: strong (agreement, 100%)
- A2:** The expert panel recommends that clinicians consider the use of oral tranexamic acid for the management of epistaxis that does not respond to moisturizing topical therapies.\*  
Quality of evidence: high (agreement, 92%)  
Strength of recommendation: strong (agreement, 94%)
- A3:** The expert panel recommends that clinicians should consider ablative therapies for nasal telangiectasias, including laser treatment, radiofrequency ablation, electrosurgery, and sclerotherapy, in patients that have failed to respond to moisturizing topical therapies.\*  
Quality of evidence: moderate (agreement, 83%)  
Strength of recommendation: weak (agreement, 94%)
- A4:** The expert panel recommends that clinicians consider the use of systemic antiangiogenic agents for the management of epistaxis that has failed to respond to moisturizing topical therapies, ablative therapies, and/or tranexamic acid.\*  
Quality of evidence: moderate (agreement, 92%)  
Strength of recommendation: strong (agreement, 82%)
- A5:** The expert panel recommends that clinicians consider a septodermoplasty for patients whose epistaxis has failed to respond sufficiently to moisturizing topical therapies, ablative therapies, and/or tranexamic acid.\*  
Quality of evidence: low (agreement, 92%)  
Strength of recommendation: weak (agreement, 88%)
- A6:** The expert panel recommends that clinicians consider a nasal closure for patients whose epistaxis has failed to respond sufficiently to moisturizing topical therapies, ablative therapies, and/or tranexamic acid.\*  
Quality of evidence: moderate (agreement, 86%)  
Strength of recommendation: strong (agreement, 82%)
- A7:** The expert panel recommends that physicians advise patients with HHT-related epistaxis to use agents that humidify the nasal mucosa to prevent epistaxis. (Agreement, 94%)†  
Level of evidence: III  
Strength of recommendation: weak
- A8:** The expert panel recommends that clinicians refer HHT patients with epistaxis and who desire treatment to otorhinolaryngologists with HHT expertise for evaluation and treatment. (Agreement, 87%)†  
Level of evidence: III  
Strength of recommendation: weak
- A9:** The expert panel recommends that when considering nasal surgery for reasons other than epistaxis, the patient and clinician obtain consultation from an otorhinolaryngologist with expertise in HHT-related epistaxis. (Agreement, 100%)†  
Level of evidence: III  
Strength of recommendation: weak
- A10:** The expert panel recommends that the treatment for acute epistaxis requiring intervention include packing with material or products that have a low likelihood of causing rebleeding with removal (e.g., lubricated low-pressure pneumatic packing). (Agreement, 93%)†  
Level of evidence: III  
Strength of recommendation: weak

**Table—Continued**

#### GI bleeding management\*

- B1:** The expert panel recommends esophagogastroduodenoscopy as the first-line diagnostic test for suspected HHT-related bleeding. Patients who meet colorectal cancer screening criteria and patients with SMAD4-HHT (genetically proven or suspected) should also undergo colonoscopy.  
Quality of evidence: low (agreement, 82%)  
Strength of recommendation: strong (agreement, 94%)
- B2:** The expert panel recommends considering capsule endoscopy for suspected HHT-related bleeding when esophagogastroduodenoscopy does not reveal significant HHT-related telangiectasia.  
Quality of evidence: low (agreement, 92%)  
Strength of recommendation: strong (agreement, 88%)
- B3:** The expert panel recommends that clinicians grade the severity of HHT-related GI bleeding and proposes the following framework:  
• Mild HHT-related GI bleeding: patient who meets their hemoglobin goals‡ with oral iron replacement  
• Moderate HHT-related GI bleeding: patient who meets their hemoglobin goals‡ with intravenous iron treatment  
• Severe HHT-related GI bleeding: patient who does not meet their hemoglobin goals‡ despite adequate iron replacement or requires blood transfusions  
Quality of evidence: low (expert consensus) (agreement, 96%)  
Strength of recommendation: strong (agreement, 96%)
- B4:** The expert panel recommends that endoscopic argon plasma coagulation be only used sparingly during endoscopy.  
Quality of evidence: low (expert consensus) (agreement, 88%)  
Strength of recommendation: weak (agreement, 81%)
- B5:** The expert panel recommends that clinicians consider treatment of mild HHT-related GI bleeding with oral antifibrinolytics.  
Quality of evidence: low (agreement, 94%)  
Strength of recommendation: weak (agreement, 90%)
- B6:** The expert panel recommends that clinicians consider treatment of moderate to severe HHT-related GI bleeding with intravenous bevacizumab or other systemic antiangiogenic therapy.  
Quality of evidence: moderate (agreement, 94%)  
Strength of recommendation: strong (agreement, 98%)
- #### Anemia and anticoagulation\*
- C1:** The expert panel recommends that the following HHT patients be tested for iron deficiency and anemia:  
• All adults, regardless of symptoms  
• All children with recurrent bleeding and/or symptoms of anemia  
Quality of evidence: high (agreement, 98%)  
Strength of recommendation: strong (agreement, 96%)
- C2:** The expert panel recommends iron replacement for treatment of iron deficiency and anemia as follows:  
• Initial therapy with oral iron  
• Intravenous iron replacement for patients in whom oral is not effective, not absorbed, or not tolerated, or who are presenting with severe anemia  
Quality of evidence: moderate (agreement, 88%)  
Strength of recommendation: strong (agreement, 100%)
- C3:** The expert panel recommends red blood cell transfusions in the following settings:  
• Hemodynamic instability/shock  
• Comorbidities that require a higher hemoglobin target  
• Need to increase the hemoglobin acutely, such as prior to surgery or during pregnancy  
• Inability to maintain an adequate hemoglobin despite frequent iron infusions  
Quality of evidence: low (agreement, 92%)  
Strength of recommendation: strong (agreement, 96%)
- C4:** The expert panel recommends considering evaluation for additional causes of anemia in the setting of an inadequate response to iron replacement.  
Quality of evidence: low (agreement, 100%)  
Strength of recommendation: strong (agreement, 100%)
- C5:** The expert panel recommends that HHT patients receive anticoagulation (prophylactic or therapeutic) or antiplatelet therapy when there is an indication, with consideration of their individualized bleeding risks; bleeding in HHT is not an absolute contraindication for these therapies.

Continued on following page

## Table—Continued

Quality of evidence: low (agreement, 98%)  
 Strength of recommendation: strong (agreement, 98%)  
 C6: The panel recommends avoiding the use of dual antiplatelet therapy and/or combination of antiplatelet therapy and anticoagulation, where possible, in patients with HHT.  
 Quality of evidence: low (expert consensus) (agreement, 83%)  
 Strength of recommendation: weak (agreement, 92%)

**Liver VMs in HHT**

- D1: The expert panel recommends that screening for liver VMs be offered to adults with definite or suspected HHT.\*  
 Quality of evidence: low (agreement, 84%)  
 Strength of recommendation: weak (agreement, 93%)  
 D2: The expert panel recommends diagnostic testing for liver VMs in HHT patients with symptoms and/or signs suggestive of complicated liver VMs (including heart failure, pulmonary hypertension, abnormal cardiac biomarkers, abnormal liver function tests, abdominal pain, portal hypertension, or encephalopathy), using Doppler ultrasound, multiphase contrast CT scan, or contrast abdominal MRI for diagnostic assessment of liver VMs.\*  
 Quality of evidence: high (agreement, 98%)  
 Strength of recommendation: strong (agreement, 100%)  
 D3: The expert panel recommends an intensive first-line management only for patients with complicated and/or symptomatic liver VMs, tailored to the type of liver VM complication(s).  
 The expert panel recommends that HHT patients with high-output cardiac failure and pulmonary hypertension be managed by the HHT Center of Excellence and an HHT cardiologist or a pulmonary hypertension specialty clinic.\*  
 Quality of evidence: moderate (agreement, 88%)  
 Strength of recommendation: strong (agreement, 88%)  
 D4: The expert panel recommends that clinicians estimate prognosis of liver VMs using available predictors, to identify patients in need of closer monitoring.\*  
 Quality of evidence: moderate (agreement, 89%)  
 Strength of recommendation: strong (agreement, 82%)  
 D5: The expert panel recommends considering intravenous bevacizumab for patients with symptomatic high-output cardiac failure due to liver VMs who have failed to respond sufficiently to first-line management.\*  
 Quality of evidence: moderate (agreement, 98%)  
 Strength of recommendation: strong (agreement, 98%)  
 D6: The expert panel recommends referral for consideration of liver transplantation for patients with symptomatic complications of liver VMs, specifically refractory high-output cardiac failure, biliary ischemia, or complicated portal hypertension.\*  
 Quality of evidence: moderate (agreement, 83%)  
 Strength of recommendation: strong (agreement, 92%)  
 D7: The expert panel recommends that liver biopsy be avoided in any patient with proven or suspected HHT. (Agreement, 97%)†  
 Level of evidence: III  
 Strength of recommendation: strong  
 D8: The expert panel recommends that hepatic artery embolization be avoided in patients with liver VMs as it is only a temporizing procedure associated with significant morbidity and mortality. (Agreement, 94%)†  
 Level of evidence: III  
 Strength of recommendation: strong

**Pediatric care\***

- E1: The expert panel advises that diagnostic genetic testing be offered for asymptomatic children of a parent with HHT.  
 Quality of evidence: high (agreement, 96%)  
 Strength of recommendation: strong (agreement, 94%)  
 E2: The expert panel recommends screening for pulmonary AVMs in asymptomatic children with HHT or at risk for HHT at the time of presentation/diagnosis.  
 Quality of evidence: moderate (agreement, 94%)  
 Strength of recommendation: strong (agreement, 94%)  
 E3: The expert panel recommends that large pulmonary AVMs and pulmonary AVMs associated with reduced oxygen saturation be treated in children to avoid serious complications.  
 Quality of evidence: moderate (agreement, 98%)  
 Strength of recommendation: strong (agreement, 98%)

## Table—Continued

- E4: The expert panel recommends repeating pulmonary AVM screening in asymptomatic children with HHT or at risk for HHT, typically at 5-year intervals.  
 Quality of evidence: low (agreement, 92%)  
 Strength of recommendation: strong (agreement, 86%)  
 E5: The expert panel recommends screening for brain VM in asymptomatic children with HHT or at risk for HHT, at the time of presentation/diagnosis.  
 Quality of evidence: low (agreement, 86%)  
 Strength of recommendation: strong (agreement, 86%)  
 E6: The expert panel recommends that brain VMs with high-risk features be treated.  
 Quality of evidence: low (agreement, 100%)  
 Strength of recommendation: strong (agreement, 98%)

**Pregnancy and delivery\***

- F1: The expert panel recommends that clinicians discuss preconception and prenatal diagnostic options, including preimplantation genetic diagnosis, with HHT-affected individuals.  
 Quality of evidence: very low (agreement, 86%)  
 Strength of recommendation: strong (agreement, 83%)  
 F2: The expert panel recommends testing with unenhanced MRI in pregnant women with symptoms suggestive of brain VMs.  
 Quality of evidence: very low (agreement, 98%)  
 Strength of recommendation: strong (agreement, 92%)  
 F3: The expert panel recommends that pregnant women with HHT who have not been recently screened and/or treated for pulmonary AVM should be approached as follows:  
  - In asymptomatic patients, initial pulmonary AVM screening should be performed using either agitated saline transthoracic contrast echocardiography (TTCE) or low-dose noncontrast chest CT, depending on local expertise. Chest CT, when performed, should be done early in the second trimester.
  - In patients with symptoms suggestive of pulmonary AVM, diagnostic testing should be performed using low-dose noncontrast chest CT. This testing can be performed at any gestational age, as clinically indicated.
  - Pulmonary AVMs should be treated starting in the second trimester unless otherwise clinically indicated.
 Quality of evidence: moderate (agreement, 88%)  
 Strength of recommendation: strong (agreement, 83%)  
 F4: The expert panel recommends that pregnant women with HHT be managed at a tertiary care center by a multidisciplinary team if they have untreated pulmonary AVMs and/or brain VMs or have not been recently screened for pulmonary AVMs.  
 Quality of evidence: very low (agreement, 94%)  
 Strength of recommendation: strong (agreement, 85%)  
 F5: The expert panel recommends not withholding an epidural because of a diagnosis of HHT, and that screening for spinal vascular malformations is not required.  
 Quality of evidence: low (agreement, 98%)  
 Strength of the recommendation: strong (agreement, 92%)  
 F6: The expert panel recommends that women with known, non-high-risk brain VMs can labor and proceed with vaginal delivery. Patients may require an assisted second stage on a case-by-case basis.  
 Quality of evidence: moderate (agreement, 94%)  
 Strength of the recommendation: strong (agreement, 94%)

**Diagnosis of HHT†**

- G1: The expert panel recommends that clinicians diagnose HHT using the Curaçao criteria or by identification of a causative mutation. (Agreement, 82%)  
 Level of evidence: III  
 Strength of recommendation: weak  
 G2: The expert panel recommends that clinicians consider the diagnosis of HHT in patients with one or more Curaçao criteria. (Agreement, 91%)  
 Level of evidence: III  
 Strength of recommendation: weak  
 G3: The expert panel recommends that asymptomatic children of a parent with HHT be considered to have possible HHT, unless excluded by genetic testing. (Agreement, 87%)  
 Level of evidence: III  
 Strength of recommendation: weak

Continued on following page

## Table—Continued

G4: The expert panel recommends that clinicians refer patients for diagnostic genetic testing for HHT (Agreement, 80%)

1. to identify the causative mutation in a family with clinically confirmed HHT;
2. to establish a diagnosis in relatives of a person with a known causative mutation, including
  - a. individuals who are asymptomatic or minimally symptomatic and
  - b. individuals who desire prenatal testing; and
3. to assist in establishing a diagnosis of HHT in individuals who do not meet clinical diagnostic criteria.

Level of evidence: III  
Strength of recommendation: weak

G5: The expert panel recommends that for individuals who test negative for ENG and ACVRL1 coding sequence mutations, SMAD4 testing should be considered to identify the causative mutation. (Agreement, 93%)

Level of evidence: III  
Strength of recommendation: weak

## Brain VMs†

H1: The expert panel recommends the use of MRI for brain VM screening in adults with possible or definite HHT using a protocol with and without contrast administration and using sequences that detect blood products, to maximize sensitivity. (Agreement, 100%)

Level of evidence: III  
Strength of recommendation: weak

H2: The expert panel recommends that adults presenting with an acute hemorrhage secondary to a brain VM be considered for definitive treatment in a center with neurovascular expertise. (Agreement, 94%)

Level of evidence: III  
Strength of recommendation: strong

H3: The expert panel recommends that all other adults with brain VMs be referred to a center with neurovascular expertise to be considered for invasive testing and individualized management. (Agreement, 84%)

Level of evidence: III  
Strength of recommendation: strong

H4: The expert panel recommends that pregnant women with suspected or confirmed HHT harboring an asymptomatic brain VM during pregnancy have definitive treatment of their brain VM deferred until after delivery of their fetus. The expert panel recommends that the delivery of the fetus follow obstetrical principles. (Agreement, 80%)

Level of evidence: III  
Strength of recommendation: weak

## Pulmonary AVMs†

I1: The expert panel recommends that clinicians screen all patients with possible or confirmed HHT for pulmonary AVMs. (Agreement, 96%)

Level of evidence: III  
Strength of recommendation: strong

I2: The expert panel recommends that clinicians use transthoracic contrast echocardiography as the initial screening test for pulmonary AVM. (Agreement, 96%)

Level of evidence: II  
Strength of recommendation: weak

I3: The expert panel recommends that clinicians treat pulmonary AVMs with transcatheter embolotherapy. (Agreement, 96%)

Level of evidence: II  
Strength of recommendation: strong

I4: The expert panel recommends that clinicians provide the following long-term advice to patients with documented pulmonary AVMs (treated or untreated):

1. Antibiotic prophylaxis for procedures with risk of bacteremia
2. When IV access is in place, take extra care to avoid IV air
3. Avoidance of SCUBA diving (Agreement, 87%)

Level of evidence: III  
Strength of recommendation: weak

I5: The expert panel recommends that clinicians provide long-term follow-up for patients who have pulmonary AVMs, in order to detect growth of untreated pulmonary AVMs and also reperfusion of treated AVMs. (Agreement, 100%)

## Table—Continued

Level of evidence: II  
Strength of recommendation: strong

AVM = arteriovenous malformation; CT = computed tomography; EGD = esophagogastroduodenoscopy; GI = gastrointestinal; HHT = hereditary hemorrhagic telangiectasia; IV = intravenous; MRI = magnetic resonance imaging; VM = vascular malformation.

\* Second HHT Guidelines.

† First HHT Guidelines.

‡ Hemoglobin goals should reflect age, gender, symptoms, and comorbidities.

Research and Evaluation II) framework and GRADE (Grading of Recommendations Assessment, Development and Evaluation) methodology. The international HHT community provided priority topics to be updated on the basis of new evidence or added because not previously addressed. The Table details recommendations that were not revisited but are still currently recommended. Topic groups were appointed for each of the 6 areas selected for update or new review. They identified key questions to guide the systematic search strategy of the literature. A medical librarian (K.L.R.) developed and executed 6 sets of search strategies between May and June 2019 in Ovid MEDLINE with input from the chair of the Guidelines Working Group (GWG). Through a series of predetermined steps illustrated in Part 1 of Supplement 2 (available at [Annals.org](http://Annals.org)), including double review of both abstracts and full-text articles, 221 articles were summarized into evidence tables. The quality of included randomized controlled trials (RCTs) was assessed (Part 2 of Supplement 2) using the structured framework of the Cochrane Risk of Bias Tool (7). In the months preceding the conference, the 6 topic groups generated draft recommendations that were based on key questions and the evidence tables and were consistent with GRADE (8) formatting for levels of evidence and strength of recommendation. Draft recommendations were distributed to all panel members 2 weeks before the consensus meeting.

The GWG convened at the guidelines conference in November 2019 in Toronto, Canada, to partake in a structured consensus process. The GWG included clinical and genetic experts in all aspects of HHT from 15 countries, guidelines methodologists, health care workers, health care administrators, HHT clinic staff, medical trainees, patient advocacy representatives, and patients with HHT. The GWG completed individual conflict-of-interest disclosures, and the chair reviewed potential conflicts. The GWG was presented draft recommendations with supporting quality of evidence, voted anonymously on the wording and quality of evidence, was presented the draft strength of recommendation with justification by GRADE methodology, and then voted on the strength of recommendation. Consensus of 80% was required for the recommendation to be included in the guidelines. A structured process was used to identify sources of disagreement for votes with less than 80% agreement (Supplement 1). The recommendations were sent for external review to HHT experts and organizations; their comments were collated and ad-



dressed (Part 3 of **Supplement 2**). The funding sources had no role in the design, conduct, or reporting of the guidelines or in the decision to submit for publication. Although the funding sources were not directly involved in the generation of the recommendations, some participants in the guidelines process were also board members, officers, or committee members of Cure HHT.

## RECOMMENDATIONS

A summary of all the recommendations is available in the **Table**.

### Epistaxis Management

*Recommendation A1: The expert panel recommends that patients with HHT-related epistaxis use moisturizing topical therapies that humidify the nasal mucosa to reduce epistaxis.* (Quality of evidence: moderate [agreement, 98%])

Topical saline has been shown to reduce epistaxis severity score compared with baseline in an RCT of topical therapies (9) (**Table 2 of Supplement 1**). (Strength of recommendation: strong [agreement, 100%])

*Clinical considerations:* Topical saline (spray or gel) is typically used twice daily.

*Recommendation A2: The expert panel recommends that clinicians consider the use of oral tranexamic acid for the management of epistaxis that does not respond to moisturizing topical therapies.* (Quality of evidence: high [agreement, 92%])

Two RCTs of oral tranexamic acid showed a significant decrease in epistaxis severity (10, 11) with minimal adverse events (**Table 2 of Supplement 1**). (Strength of recommendation: strong [agreement, 94%])

*Clinical considerations:* **Table 4 of Supplement 1** gives prescribing and safety monitoring guidance for oral tranexamic acid.

*Recommendation A3: The expert panel recommends that clinicians should consider ablative therapies for nasal telangiectasias, including laser treatment, radiofrequency ablation, electrosurgery, and sclerotherapy, in patients that have failed to respond to moisturizing topical therapies.* (Quality of evidence: moderate [agreement, 83%])

One RCT showed reduced epistaxis severity score with sclerotherapy (12). Multiple uncontrolled series demonstrated that various ablative therapies temporarily reduced epistaxis (13–15) (**Tables 2 and 3 of Supplement 1**). (Strength of recommendation: weak [agreement, 94%])

*Clinical considerations:* Clinicians and patients should choose a specific ablative therapy on the basis of local expertise, understanding that ablative therapy is a temporizing treatment of epistaxis and that perforation of the nasal septum is a known complication of all techniques.

*Recommendation A4: The expert panel recommends that clinicians consider the use of systemic antiangiogenic agents for the management of epistaxis that has failed to respond to moisturizing topical therapies,*

*ablative therapies, and/or tranexamic acid.* (Quality of evidence: moderate [agreement, 92%])

Multiple uncontrolled series demonstrated that intravenous (IV) bevacizumab reduced epistaxis, improved anemia, reduced transfusion requirements, or improved quality of life (QOL) (16–23) (**Table 3 of Supplement 1**). (Strength of recommendation: strong [agreement, 82%])

*Clinical considerations:* **Table 4 of Supplement 1** gives prescribing and safety monitoring guidance for IV bevacizumab.

*Recommendation A5: The expert panel recommends that clinicians consider a septodermoplasty for patients whose epistaxis has failed to respond sufficiently to moisturizing topical therapies, ablative therapies, and/or tranexamic acid.* (Quality of evidence: low [agreement, 92%])

Multiple uncontrolled series demonstrated that septodermoplasty reduced epistaxis, improved anemia, reduced surgical reintervention, or improved QOL (24–29) (**Table 3 of Supplement 1**). (Strength of recommendation: weak [agreement, 88%])

*Clinical considerations:* Clinicians and patients should consider septodermoplasty when epistaxis affects QOL or is life-threatening; they should consider the risks and benefits, as well as alternatives, such as nasal closure and antiangiogenic medications.

*Recommendation A6: The expert panel recommends that clinicians consider a nasal closure for patients whose epistaxis has failed to respond sufficiently to moisturizing topical therapies, ablative therapies, and/or tranexamic acid.* (Quality of evidence: moderate [agreement, 86%])

Multiple uncontrolled series demonstrated that nasal closure reduced epistaxis (26, 28) (**Table 3 of Supplement 1**). (Strength of recommendation: strong [agreement, 82%])

*Clinical considerations:* Clinicians and patients should consider nasal closure when epistaxis affects QOL or is life-threatening; they should consider the risks and benefits, as well as alternatives, such as septodermoplasty and antiangiogenic medications.

### GI Bleeding Management

*Recommendation B1: The expert panel recommends esophagogastroduodenoscopy as the first-line diagnostic test for suspected HHT-related bleeding. Patients who meet colorectal cancer screening criteria and patients with SMAD4-HHT (genetically proven or suspected) should also undergo colonoscopy.* (Quality of evidence: low [agreement, 82%])

Several cross-sectional studies of diagnostic yield demonstrated a high yield from esophagogastroduodenoscopy (EGD) for upper GI telangiectasias in patients with HHT and suspected GI bleeding (30–32) (**Table 5 of Supplement 1**). (Strength of recommendation: strong [agreement, 94%])

*Clinical considerations:* Clinicians should consider performing EGD in an experienced center given potential unusual complications during the procedure (such as massive epistaxis) and should also be aware of the

precautions required for patients with HHT and pulmonary arteriovenous malformations (AVMs) (Table).

In suspected or proven *SMAD4*-HHT, screening colonoscopy is recommended starting at age 15 years and repeated every 3 years if no polyps are found or every year along with EGD if colonic polyps are found. Other patients with HHT (non-*SMAD4*) should be screened for colon cancer following general population guidelines.

*Recommendation B2: The expert panel recommends considering capsule endoscopy for suspected HHT-related bleeding when esophagogastroduodenoscopy does not reveal significant HHT-related telangiectasia.* (Quality of evidence: low [agreement, 92%])

Several cross-sectional studies of diagnostic yield demonstrated a high yield from capsule endoscopy, with an excellent safety profile, for small-bowel GI telangiectases in patients with HHT and suspected GI bleeding (30–34) (Table 5 of Supplement 1). (Strength of recommendation: strong [agreement, 88%])

*Clinical considerations:* Capsule endoscopy remains complementary to EGD when anemia is unexplained by the severity of epistaxis and gastric involvement or when the EGD findings are negative.

*Recommendation B3: The expert panel recommends that clinicians grade the severity of HHT-related GI bleeding and proposes the following framework:*

- *Mild HHT-related GI bleeding: patient who meets their hemoglobin goals\* with oral iron replacement*
- *Moderate HHT-related GI bleeding: patient who meets their hemoglobin goals\* with intravenous iron treatment*
- *Severe HHT-related GI bleeding: patient who does not meet their hemoglobin goals\* despite adequate iron replacement or requires blood transfusions*

\* Hemoglobin goals should reflect age, gender, symptoms, and comorbidities. (Quality of evidence: low [expert consensus] [agreement, 96%])

Case series describe a severity range for HHT-related GI bleeding, with secondary anemia, reduced QOL, blood transfusion requirements, hospitalization, morbidity, and mortality (32, 35–41). (Strength of recommendation: strong [agreement, 96%])

*Clinical considerations:* Hemoglobin goals (not levels) are specified to reflect the patient's individual physiologic needs. This classification applies to patients who have had at least 3 months of iron therapy.

*Recommendation B4: The expert panel recommends that endoscopic argon plasma coagulation be only used sparingly during endoscopy.* (Quality of evidence: low [expert consensus] [agreement, 88%])

Expert consensus in HHT and case series in patients without HHT show some benefit from endoscopic argon plasma coagulation (42, 43). (Strength of recommendation: weak [agreement, 81%])

*Clinical considerations:* Argon plasma coagulation is best administered concurrent with the initial endoscopic evaluation for bleeding lesions and significant

(1- to 3-mm) nonbleeding lesions. Repeated sessions are discouraged to avoid repeated iatrogenic injury to the intestinal mucosa.

*Recommendation B5: The expert panel recommends that clinicians consider treatment of mild HHT-related GI bleeding with oral antifibrinolytics.* (Quality of evidence: low [agreement, 94%])

One case series reported reduced need for endoscopic management in patients treated with oral tranexamic acid (44) (Table 6 of Supplement 1) with a good safety profile. (Strength of recommendation: weak [agreement, 90%])

*Clinical considerations:* Table 4 of Supplement 1 gives prescribing and safety monitoring guidance for oral tranexamic acid.

*Recommendation B6: The expert panel recommends that clinicians consider treatment of moderate to severe HHT-related GI bleeding with intravenous bevacizumab or other systemic antiangiogenic therapy.* (Quality of evidence: moderate [agreement, 94%])

Small uncontrolled series showed that systemic antiangiogenic therapies alleviated anemia, reduced transfusion requirements, or improved QOL (19, 21, 45) (Table 6 of Supplement 1). (Strength of recommendation: strong [agreement, 98%])

*Clinical considerations:* Table 4 of Supplement 1 gives prescribing and safety monitoring guidance for IV bevacizumab.

## Anemia and Anticoagulation

*Recommendation C1: The expert panel recommends that the following HHT patients be tested for iron deficiency and anemia:*

- *All adults, regardless of symptoms*
- *All children with recurrent bleeding and/or symptoms of anemia* (Quality of evidence: high [agreement, 98%])

Three case series reported iron deficiency anemia as a common complication of HHT, typically in adults (36, 46, 47). (Strength of recommendation: strong [agreement, 96%])

*Clinical considerations:* Testing typically includes complete blood count and ferritin measurement. If a patient is anemic but ferritin is not reduced, serum iron, total iron-binding capacity, and transferrin saturation should be measured and a hematology consultation should be considered.

*Recommendation C2: The expert panel recommends iron replacement for treatment of iron deficiency and anemia as follows:*

- *Initial therapy with oral iron*
- *Intravenous iron replacement for patients in whom oral is not effective, not absorbed, or not tolerated, or who are presenting with severe anemia* (Quality of evidence: moderate [agreement, 88%])

Evidence for iron replacement and initial dosing are based on case series in HHT and non-HHT iron deficiency anemia (48–53). (Strength of recommendation: strong [agreement, 100%])

*Clinical considerations:* Iron replacement typically starts with once-daily oral dosing of 35 to 65 mg of elemental iron, 2 hours before or 1 hour after meals. An increase in hemoglobin of less than 10 g/L is considered inadequate in anemic patients, and every-other-day dosing or an alternate preparation of oral iron should be attempted. In refractory anemia or severe chronic bleeding, regularly scheduled iron infusions may be required. Initial IV iron dosing can be calculated (54), or a total initial dose of 1 g of IV iron can be provided, as a single infusion or divided doses. **Supplement 1** details additional safety and prescribing information.

*Recommendation C3:* The expert panel recommends red blood cell transfusions in the following settings:

- Hemodynamic instability/shock
- Comorbidities that require a higher hemoglobin target
- Need to increase the hemoglobin acutely, such as prior to surgery or during pregnancy
- Inability to maintain an adequate hemoglobin despite frequent iron infusions (Quality of evidence: low [agreement, 92%])

Expert consensus in HHT. (Strength of recommendation: strong [agreement, 96%])

*Clinical considerations:* Hemoglobin targets and thresholds for red blood cell transfusion should be individualized in HHT, depending on patient symptoms, severity of ongoing HHT-related bleeding, response to other therapies and iron supplementation, presence of comorbid conditions, and acuity.

*Recommendation C4:* The expert panel recommends considering evaluation for additional causes of anemia in the setting of an inadequate response to iron replacement. (Quality of evidence: low [agreement, 100%])

One case series reported folate deficiency and hemolysis as additional causes of anemia in patients with HHT (55). (Strength of recommendation: strong [agreement, 100%])

*Clinical considerations:* Evaluation should include measurement of folate, vitamin B<sub>12</sub>, mean corpuscular volume, and thyroid-stimulating hormone; smear; reticulocyte counts; and work-up for hemolysis, with referral to hematology in unresolved cases.

*Recommendation C5:* The expert panel recommends that HHT patients receive anticoagulation (prophylactic or therapeutic) or antiplatelet therapy when there is an indication, with consideration of their individualized bleeding risks; bleeding in HHT is not an absolute contraindication for these therapies. (Quality of evidence: low [agreement, 98%])

Expert consensus in HHT and 2 case series demonstrated that anticoagulation or antiplatelet therapy is well tolerated by most patients with HHT (56, 57). (Strength of recommendation: strong [agreement, 98%])

*Clinical considerations:* When anticoagulation is pursued, unfractionated heparin, low-molecular-weight heparin, and vitamin K antagonists are preferred over direct-acting oral anticoagulants, which are less well

tolerated in HHT (58). In cases of atrial fibrillation, if anticoagulation is not tolerated, alternate approaches can be considered, such as left atrial appendage closure (59).

*Recommendation C6:* The panel recommends avoiding the use of dual antiplatelet therapy and/or combination of antiplatelet therapy and anticoagulation, where possible, in patients with HHT. (Quality of evidence: low (expert consensus) [agreement, 83%])

Expert consensus in HHT. (Strength of recommendation: weak [agreement, 92%])

*Clinical considerations:* If dual or combination therapies are required, duration of therapy should be minimized and patients should be monitored closely.

### Liver VMs in HHT

*Recommendation D1:* The expert panel recommends that screening for liver VMs be offered to adults with definite or suspected HHT. (Quality of evidence: low [agreement, 84%])

Several cross-sectional diagnostic studies demonstrated high yield and accuracy of Doppler ultrasonography, multiphase contrast computed tomography (CT), and magnetic resonance imaging (MRI) for detection of liver VMs (5, 60–68) (Table 7 of Supplement 1); Doppler ultrasonography severity grading was predictive of outcomes (69). Anicteric cholestasis, reported in one third of patients with liver VMs, correlated with severity of liver VMs and complications (69–71). (Strength of recommendation: weak [agreement, 93%])

*Clinical considerations:* The rationale for screening is that awareness of liver VMs could improve subsequent patient management or help confirm the diagnosis of HHT. The imaging test of choice is Doppler ultrasonography because of its accuracy, safety, tolerability, low costs, and operating characteristics. However, depending on local availability of and expertise in Doppler ultrasonography, as well as patient preference, patients may be screened clinically (history, physical examination, and blood work) or alternative imaging may be considered, including multiphase contrast CT or MRI.

*Recommendation D2:* The expert panel recommends diagnostic testing for liver VMs in HHT patients with symptoms and/or signs suggestive of complicated liver VMs (including heart failure, pulmonary hypertension, abnormal cardiac biomarkers, abnormal liver function tests, abdominal pain, portal hypertension, or encephalopathy), using Doppler ultrasound, multiphase contrast CT scan, or contrast abdominal MRI for diagnostic assessment of liver VMs. (Quality of evidence: high [agreement, 98%])

Several cross-sectional diagnostic studies demonstrated high yield and accuracy of Doppler ultrasonography, multiphase contrast CT, and MRI for diagnosis of liver VMs (5, 60–68) (Table 7 of Supplement 1). (Strength of recommendation: strong [agreement, 100%])

*Clinical considerations:* The choice of imaging method should be informed by the risk-benefit balance, local expertise, and availability or cost. Contrast studies (CT and MRI) should be avoided if kidney dys-



function is present. Echocardiography provides additional information about the hemodynamic effect of liver VMs. These tests will be most informative when done in the context of a clinical assessment at an HHT center of excellence.

*Recommendation D3: The expert panel recommends an intensive first-line management only for patients with complicated and/or symptomatic liver VMs, tailored to the type of liver VM complication(s).*

*The expert panel recommends that HHT patients with high-output cardiac failure and pulmonary hypertension be comanaged by the HHT Center of Excellence and an HHT cardiologist or a pulmonary hypertension specialty clinic. (Quality of evidence: moderate [agreement, 88%])*

One large series showed moderate response to first-line therapy tailored to liver VM complication (69). Expert consensus supported the recommendation for specialized center management. (Strength of recommendation: strong [agreement, 88%])

*Clinical considerations: Supplement 1 describes first-line therapies, by specific liver VM complication. Typically, patients with symptomatic liver VMs are managed by an expert team at an HHT center of excellence, with at least annual follow-up.*

*Recommendation D4: The expert panel recommends that clinicians estimate prognosis of liver VMs using available predictors, to identify patients in need of closer monitoring. (Quality of evidence: moderate [agreement, 89%])*

Three observational studies identified clinical predictors of complications from liver VMs (69, 70, 72). (Strength of recommendation: strong [agreement, 82%])

*Clinical considerations: Clinicians should plan monitoring for patients with liver VMs on the basis of estimated prognosis.*

*Recommendation D5: The expert panel recommends considering intravenous bevacizumab for patients with symptomatic high-output cardiac failure due to liver VMs who have failed to respond sufficiently to first-line management. (Quality of evidence: moderate [agreement, 98%])*

Small uncontrolled series showed that IV bevacizumab improved cardiac output or clinical symptoms in 80% of patients with severe liver VMs, primarily in those with high-output cardiac failure (16) (Table 8 of Supplement 1). The adverse event rate was reported at 50 per 100 person-years, including 1 fatal event probably related to bevacizumab (73). (Strength of recommendation: strong [agreement, 98%])

Table 4 of Supplement 1 gives prescribing and safety monitoring guidance for IV bevacizumab.

*Recommendation D6: The expert panel recommends referral for consideration of liver transplantation for patients with symptomatic complications of liver VMs, specifically refractory high-output cardiac failure, biliary ischemia, or complicated portal hypertension. (Quality of evidence: moderate [agreement, 83%])*

Small uncontrolled series of orthotopic liver transplantation for liver VMs in HHT demonstrated excellent 5- to 10-year survival (82% to 92%) (74, 75) with asymptomatic rare and late recurrence of liver VMs after liver

transplantation (76). (Strength of recommendation: strong [agreement, 92%])

*Clinical considerations: Timing for listing a symptomatic patient for orthotopic liver transplantation should be based on prognostic predictors and the severity of liver VM complications, including pulmonary hypertension. Liver transplantation can be undertaken in the presence of pulmonary hypertension if pulmonary vascular resistance, estimated by right heart catheterization, is less than 3 Woods units.*

## Pediatric Care

*Recommendation E1: The expert panel advises that diagnostic genetic testing be offered for asymptomatic children of a parent with HHT. (Quality of evidence: high [agreement, 96%])*

Two cross-sectional diagnostic studies demonstrated that genetic testing can identify subclinical or presymptomatic disease in children of HHT families with known mutation (77–79). (Strength of recommendation: strong [agreement, 94%])

*Clinical considerations: An affected family member should be tested first to determine the causative mutation before testing an asymptomatic child who does not meet the clinical diagnostic criteria for HHT (Curaçao criteria) (4). The benefits of testing, alternatives, pros, and cons should be discussed with children or—as appropriate—their parents.*

*Recommendation E2: The expert panel recommends screening for pulmonary AVMs in asymptomatic children with HHT or at risk for HHT at the time of presentation/diagnosis. (Quality of evidence: moderate [agreement, 94%])*

Several pediatric case series demonstrated a prevalence of pulmonary AVMs similar to that in adults and a risk for life-threatening complications with good outcomes from embolization (80–85). Several series have reported 2 sensitive screening protocols in children (86–90) (Table 9 of Supplement 1). (Strength of recommendation: strong [agreement, 94%])

*Clinical considerations: Screening may be performed with either chest radiography and pulse oximetry or trans-thoracic contrast echocardiography. Screening with CT is not recommended, although chest CT remains the confirmatory diagnostic test when screening tests have positive findings.*

*Recommendation E3: The expert panel recommends that large pulmonary AVMs and pulmonary AVMs associated with reduced oxygen saturation be treated in children to avoid serious complications. (Quality of evidence: moderate [agreement, 98%])*

Case series demonstrated that children are at risk for serious complications from large pulmonary AVMs (or AVMs causing hypoxemia) (82, 83, 85) and that embolization is safe and effective (85) (Table 10 of Supplement 1). (Strength of recommendation: strong [agreement, 98%])

*Clinical considerations: Pulmonary AVMs with feeding arteries at least 3 mm in diameter are suitable for embolotherapy. Follow-up is indicated to detect recanalization and reperfusion of treated AVMs and growth*



of small untreated AVMs. Specific protocols vary among centers (CT, oximetry, or transthoracic contrast echocardiography), as do intervals.

*Recommendation E4: The expert panel recommends repeating pulmonary AVM screening in asymptomatic children with HHT or at risk for HHT, typically at 5-year intervals.* (Quality of evidence: low [agreement, 92%])

One case series showed growth of pulmonary AVMs during childhood (91). (Strength of recommendation: strong [agreement, 86%])

*Clinical considerations:* Screening is typically repeated every 5 years after negative findings. In children with indeterminate or borderline screening results based on either imaging or oximetry, screening should be repeated sooner.

*Recommendation E5: The expert panel recommends screening for brain VM in asymptomatic children with HHT or at risk for HHT, at the time of presentation/diagnosis.* (Quality of evidence: low [agreement, 86%])

Case series demonstrated risk for intracranial hemorrhage from brain VMs (92-95) in children; MRI as a sensitive screening test (96-98); and benefits of surgical and endovascular management (99, 100), also with significant risk. (Strength of recommendation: strong [agreement, 86%])

*Clinical considerations:* First-line screening is MRI (contrast-enhanced is more sensitive) to identify brain VMs and determine subtype and risk factors for hemorrhage. This typically requires sedation or anesthesia in young children. The decision to treat versus observe is based on risk of treatment versus risk for hemorrhage. As such, the decision to screen the child should be shared among clinicians, caregivers, and the child (where possible). Clinical practice differs across countries, from screening asymptomatic children with MRI in infancy to no routine screening of asymptomatic children for brain VMs. Patient representatives felt strongly that children should be screened for brain VMs and cited anecdotal evidence of disastrous outcomes in unscreened patients.

*Recommendation E6: The expert panel recommends that brain VMs with high-risk features be treated.* (Quality of evidence: low [agreement, 100%])

Case series demonstrated risk for intracranial hemorrhage from brain VMs (92-95); identified high-risk features (95, 101, 102); and showed the benefits of surgical and endovascular management (99, 100), also with significant risk. (Strength of recommendation: strong [agreement, 98%])

*Clinical considerations:* Given the need to balance natural history risk with treatment risk, children with HHT who have brain VMs should be referred to a center with multidisciplinary expertise in neurovascular disease management. Treated brain VMs require close follow-up; the follow-up for small (untreated) brain VMs is not well defined.

## Pregnancy and Delivery

*Recommendation F1: The expert panel recommends that clinicians discuss preconception and prena-*

*tal diagnostic options, including preimplantation genetic diagnosis, with HHT-affected individuals.* (Quality of evidence: very low [agreement, 86%])

Expert consensus in HHT. (Strength of recommendation: strong [agreement, 83%])

*Clinical considerations:* Once the causative familial mutation is identified in an affected parent, it can be screened for in future offspring. Available options, including preimplantation, postconception, and postdelivery testing (Supplement 1), vary internationally. The discussion will be influenced by local legislation pertaining to preimplantation diagnosis and termination of pregnancy.

*Recommendation F2: The expert panel recommends testing with unenhanced MRI in pregnant women with symptoms suggestive of brain VMs.* (Quality of evidence: very low [agreement, 98%])

Expert consensus in HHT. (Strength of recommendation: strong [agreement, 92%])

*Clinical considerations:* For symptomatic patients, including those with previous cerebral hemorrhage, MRI without gadolinium should be planned in the second trimester. Asymptomatic patients do not require routine screening during pregnancy.

*Recommendation F3: The expert panel recommends that pregnant women with HHT who have not been recently screened and/or treated for pulmonary AVM should be approached as follows:*

- *In asymptomatic patients, initial pulmonary AVM screening should be performed using either agitated saline transthoracic contrast echocardiography (TTCE) or low-dose noncontrast chest CT, depending on local expertise. Chest CT, when performed, should be done early in the second trimester.*
- *In patients with symptoms suggestive of pulmonary AVM, diagnostic testing should be performed using low-dose noncontrast chest CT. This testing can be performed at any gestational age, as clinically indicated.*
- *Pulmonary AVMs should be treated starting in the second trimester unless otherwise clinically indicated.* (Quality of evidence: moderate [agreement, 88%])

Case series demonstrated elevated risk for complications from pulmonary AVMs during pregnancy (103-105) and low risk of imaging and embolization in the second trimester (106). (Strength of recommendation: strong [agreement, 83%])

*Clinical considerations:* Technique for embolization in pregnant patients should include measures to reduce fetal radiation exposure, including avoidance of fluoroscopy over the abdomen and pelvis, use of pulsed or low-dose fluoroscopy, minimization of angiography runs, and use of tight collimation. For both CT and angiography, abdominal shielding is not helpful and may in fact increase scattered radiation to the fetus.

*Recommendation F4: The expert panel recommends that pregnant women with HHT be managed at a*

tertiary care center by a multidisciplinary team if they have untreated pulmonary AVMs and/or brain VMs or have not been recently screened for pulmonary AVMs. (Quality of evidence: very low [agreement, 94%])

Expert consensus in HHT. (Strength of recommendation: strong [agreement, 85%])

**Clinical considerations:** Pregnant women with untreated pulmonary AVMs or brain VMs, as well as those who have not been screened, should be considered high-risk for hemorrhagic and neurologic complications and be managed accordingly by a high-risk team with HHT expertise.

**Recommendation F5:** The expert panel recommends not withholding an epidural because of a diagnosis of HHT, and that screening for spinal vascular malformations is not required. (Quality of evidence: low [agreement, 98%])

Two case series showed no evidence of hemorrhagic complications from epidural or spinal anesthesia (103, 107). (Strength of recommendation: strong [agreement, 92%])

**Clinical considerations:** Patients should meet with an anesthesiologist during the early third trimester to discuss anesthesia options. The risks for complications from spinal VMs during epidural anesthesia are unsubstantiated and only theoretical.

**Recommendation F6:** The expert panel recommends that women with known, non-high-risk brain VMs can labor and proceed with vaginal delivery. Patients may require an assisted second stage on a case-by-case basis. (Quality of evidence: moderate [agreement, 94%])

Two case series showed no intracranial hemorrhage during delivery from brain VMs in patients with HHT (103, 107). (Strength of recommendation: strong [agreement, 94%])

**Clinical considerations:** If a brain VM has not previously ruptured, patients may proceed with method of delivery based on obstetric indications and discussion with their obstetric care provider. Vaginal delivery is not contraindicated. Patients with “high-risk” brain VMs should be considered for cesarean section or epidural to allow passive descent of the presenting part, with consideration of an assisted second stage. Diligent management of blood pressure is imperative in these higher-risk cases, and obtaining the opinion of a multidisciplinary neurovascular team is prudent.

From St. Michael's Hospital, Li Ka Shing Knowledge Institute, and University of Toronto, Toronto, Ontario, Canada (M.E.F.); St. Antonius Hospital, Nieuwegein, the Netherlands (J.J.M.); University of California, San Francisco, San Francisco, California (S.W.H., M.C.); University of Toronto, Toronto, Ontario, Canada (V.A.P.); Centre for Effective Practice, Toronto, Ontario, Canada (K.L.); HHT Reference Center ERN, Ospedale Maggiore, ASST Crema, Crema, Italy (E.B.); Centre Hospitalier de l'Université de Montréal, Hôtel-Dieu, Montreal, Quebec, Canada (E.D.); University of North Carolina at Chapel Hill, Chapel Hill, North Carolina (R.S.K., J.D.); St. Michael's Hospital and University of Toronto, Toronto, Ontario, Canada (A.L., R.P.); Froedtert and Medical College of Wisconsin, Milwaukee,

Wisconsin (D.P., P.F.); The Hospital for Sick Children and University of Toronto, Toronto, Ontario, Canada (F.R.); VA Portland Health Care System and Oregon Health & Science University, Portland, Oregon (M.S.C.); Cure HHT, Monkton, Maryland (M.C., S.O.); University of Utah Medical Center, Salt Lake City, Utah (K.J.W., J.M.); Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts (H.A.); Washington University School of Medicine, St. Louis, Missouri (M.C., K.K., J.F.P., A.J.W.); St. Michael's Hospital and Unity Health Toronto, Toronto, Canada (D.C.); HHT Europe, Rome, Italy (C.C.); Grace Hospital, Winnipeg, Manitoba, Canada (E.D.); HHT Canada, Spruce Grove, Alberta, Canada (C.D.); Hospices Civils de Lyon, Femme-Mère-Enfant, Lyon, France (S.D.); University Hospital of Marburg and Phillips University Marburg, Marburg, Germany (U.G.); Augusta University, Augusta, Georgia (J.R.G.); Cincinnati Children's Hospital and University of Cincinnati, Cincinnati, Ohio (A.H.); Oslo University Hospital, Rikshospitalet, Oslo, Norway (K.H.); Yale University School of Medicine, New Haven, Connecticut (K.H.); Mayo Clinic, Rochester, Minnesota (V.N.I.); Odense University Hospital, Odense, Denmark (A.D.K.); Osaka City General Hospital, Osaka, Japan (M.K.); Chester, New Jersey (J.M.); University of California, Los Angeles, Los Angeles, California (J.M.); University of Arkansas for Medical Sciences, Little Rock, Arkansas (M.E.M.); Schneider Children's Medical Center of Israel and Sackler School of Medicine of Tel Aviv University, Tel Aviv, Israel (M.M.); Baltimore, Maryland (S.P.); Sioux Falls, South Dakota (B.P.); University of Edinburgh, Edinburgh, Scotland (M.E.P.); St. Antonius Hospital, Nieuwegein, and University Medical Center Utrecht, Utrecht, the Netherlands (M.C.P.); Toronto Western Hospital and University of Toronto, Toronto, Ontario, Canada (I.R.); University of Colorado Hospital, Aurora, Colorado (P.J.R.); Massachusetts General Hospital, Boston, Massachusetts (J.R.); University of Bari, Bari, Italy (C.S.); Hospital Italiano de Buenos Aires, Buenos Aires, Argentina (M.S.); Hammersmith Hospital, London, England (C.S.); Blue Bell, Pennsylvania (D.S.); Royal Melbourne Hospital and University of Melbourne, Melbourne, Victoria, Australia (I.W.); and Hospital Sierrallana (Servicio Cántabro de Salud), Torrelavega, Spain (R.Z.).

**Note:** Centers with recognized expertise in the diagnosis and management of HHT can be located at <https://curehht.org>, the website for Cure HHT, and <http://vascern.eu>, the website for the European Reference Network for Rare Vascular Diseases.

**Acknowledgment:** The authors thank Nicole Schaefer and Sandy Medeiros for their contributions to guideline development and the guidelines conference.

**Financial Support:** By the Christopher McMahon Family and Cure HHT. Dr. Faughnan is supported by the Nelson Arthur Hyland Foundation and the Li Ka Shing Knowledge Institute of St Michael's Hospital.

**Disclosures:** Disclosures can be viewed at [www.acponline.org/authors/icmje/ConflictOfInterestForms.do?msNum=M20-1443](http://www.acponline.org/authors/icmje/ConflictOfInterestForms.do?msNum=M20-1443).

**Corresponding Author:** Marie E. Faughnan, MD, MSc, St. Michael's Hospital, University of Toronto, 30 Bond Street, Toronto, ON M5B 1W8, Canada; e-mail, [marie.faughnan@unityhealth.to](mailto:marie.faughnan@unityhealth.to).

Current author addresses and author contributions are available at [Annals.org](http://Annals.org).

## References

1. Dakeishi M, Shioya T, Wada Y, et al. Genetic epidemiology of hereditary hemorrhagic telangiectasia in a local community in the northern part of Japan. *Hum Mutat*. 2002;19:140-8. [PMID: 11793473]
2. Pierucci P, Lenato GM, Suppressa P, et al. A long diagnostic delay in patients with hereditary haemorrhagic telangiectasia: a questionnaire-based retrospective study. *Orphanet J Rare Dis*. 2012; 7:33. [PMID: 22676497]
3. Porteous ME, Burn J, Proctor SJ. Hereditary haemorrhagic telangiectasia: a clinical analysis. *J Med Genet*. 1992;29:527-30. [PMID: 1518020]
4. Shovlin CL, Guttmacher AE, Buscarini E, et al. Diagnostic criteria for hereditary hemorrhagic telangiectasia (Rendu-Osler-Weber syndrome). *Am J Med Genet*. 2000;91:66-7. [PMID: 10751092]
5. Faughnan ME, Palda VA, Garcia-Tsao G, et al; HHT Foundation International Guidelines Working Group. International guidelines for the diagnosis and management of hereditary haemorrhagic telangiectasia. *J Med Genet*. 2011;48:73-87. [PMID: 19553198] doi:10.1136/jmg.2009.069013
6. McDonald J, Wooderchak-Donahue W, VanSant Webb C, et al. Hereditary hemorrhagic telangiectasia: genetics and molecular diagnostics in a new era. *Front Genet*. 2015;6:1. [PMID: 25674101] doi: 10.3389/fgene.2015.00001
7. Chandler J, McKenzie J, Boutron I, et al, eds. *Cochrane Methods*. *Cochrane Database Syst Rev*. 2016;10. doi:10.1002/14651858.CD201601
8. Balslem H, Helfand M, Schünemann HJ, et al. GRADE guidelines: 3. rating the quality of evidence. *J Clin Epidemiol*. 2011;64:401-6. [PMID: 21208779] doi:10.1016/j.jclinepi.2010.07.015
9. Whitehead KJ, Sautter NB, McWilliams JP, et al. Effect of topical intranasal therapy on epistaxis frequency in patients with hereditary hemorrhagic telangiectasia: a randomized clinical trial. *JAMA*. 2016; 316:943-51. [PMID: 27599329] doi:10.1001/jama.2016.11724
10. Gaillard S, Dupuis-Girod S, Boutitie F, et al; ATERO Study Group. Tranexamic acid for epistaxis in hereditary hemorrhagic telangiectasia patients: a European cross-over controlled trial in a rare disease. *J Thromb Haemost*. 2014;12:1494-502. [PMID: 25040799] doi:10.1111/jth.12654
11. Geisthoff UW, Seyfert UT, Kübler M, et al. Treatment of epistaxis in hereditary hemorrhagic telangiectasia with tranexamic acid - a double-blind placebo-controlled cross-over phase IIIB study. *Thromb Res*. 2014;134:565-71. [PMID: 25005464] doi:10.1016/j.thromres.2014.06.012
12. Boyer H, Fernandes P, Le C, et al. Prospective randomized trial of sclerotherapy vs standard treatment for epistaxis due to hereditary hemorrhagic telangiectasia. *Int Forum Allergy Rhinol*. 2015;5:435-40. [PMID: 25643928] doi:10.1002/alr.21484
13. Boyer H, Fernandes P, Duran O, et al. Office-based sclerotherapy for recurrent epistaxis due to hereditary hemorrhagic telangiectasia: a pilot study. *Int Forum Allergy Rhinol*. 2011;1:319-23. [PMID: 22287439] doi:10.1002/alr.20053
14. Rothenberg B, Noyek S, Chin CJ. Radiofrequency ablation for treatment of hereditary hemorrhagic telangiectasia lesions: "how I do it". *Am J Rhinol Allergy*. 2015;29:226-7. [PMID: 25975256] doi:10.2500/ajra.2015.29.4163
15. Kuan EC, Peng KA, Thompson CF, et al. Sinonasal quality of life outcomes following laser treatment of epistaxis related to hereditary hemorrhagic telangiectasia. *Lasers Med Sci*. 2017;32:527-531. [PMID: 28116537] doi:10.1007/s10103-017-2144-7
16. Dupuis-Girod S, Ginon I, Saurin JC, et al. Bevacizumab in patients with hereditary hemorrhagic telangiectasia and severe hepatic vascular malformations and high cardiac output. *JAMA*. 2012;307:948-55. [PMID: 22396517] doi:10.1001/jama.2012.250
17. Thompson AB, Ross DA, Berard P, et al. Very low dose bevacizumab for the treatment of epistaxis in patients with hereditary hemorrhagic telangiectasia. *Allergy Rhinol (Providence)*. 2014;5:91-5. [PMID: 25199101] doi:10.2500/ar.2014.5.0091
18. Chavan A, Schumann-Binarsch S, Schmuck B, et al. Emerging role of bevacizumab in management of patients with symptomatic hepatic involvement in hereditary hemorrhagic telangiectasia [Letter]. *Am J Hematol*. 2017;92:E641-E644. [PMID: 28776732] doi:10.1002/ajh.24878
19. Iyer VN, Apala DR, Pannu BS, et al. Intravenous bevacizumab for refractory hereditary hemorrhagic telangiectasia-related epistaxis and gastrointestinal bleeding. *Mayo Clin Proc*. 2018;93:155-166. [PMID: 29395350] doi:10.1016/j.mayocp.2017.11.013
20. Epperla N, Kapke JT, Karafin M, et al. Effect of systemic bevacizumab in severe hereditary hemorrhagic telangiectasia associated with bleeding [Letter]. *Am J Hematol*. 2016;91:E313-4. [PMID: 26994402] doi:10.1002/ajh.24367
21. Al-Samkari H, Kritharis A, Rodriguez-Lopez JM, et al. Systemic bevacizumab for the treatment of chronic bleeding in hereditary haemorrhagic telangiectasia. *J Intern Med*. 2019;285:223-231. [PMID: 30191646] doi:10.1111/joim.12832
22. Rosenberg T, Fialla AD, Kjeldsen J, et al. Does severe bleeding in HHT patients respond to intravenous bevacizumab? Review of the literature and case series. *Rhinology*. 2019;57:242-251. [PMID: 30907391] doi:10.4193/Rhin18.289
23. Guilhem A, Fargeton AE, Simon AC, et al. Intra-venous bevacizumab in hereditary hemorrhagic telangiectasia (HHT): a retrospective study of 46 patients. *PLoS One*. 2017;12:e0188943. [PMID: 29190827] doi:10.1371/journal.pone.0188943
24. Levine CG, Ross DA, Henderson KJ, et al. Long-term complications of septal dermoplasty in patients with hereditary hemorrhagic telangiectasia. *Otolaryngol Head Neck Surg*. 2008;138:721-4. [PMID: 18503842] doi:10.1016/j.otohns.2008.01.005
25. Lesnik GT, Ross DA, Henderson KJ, et al. Septectomy and septal dermoplasty for the treatment of severe transfusion-dependent epistaxis in patients with hereditary hemorrhagic telangiectasia and septal perforation. *Am J Rhinol*. 2007;21:312-5. [PMID: 17621815]
26. Richer SL, Geisthoff UW, Livada N, et al. The Young's procedure for severe epistaxis from hereditary hemorrhagic telangiectasia. *Am J Rhinol Allergy*. 2012;26:401-4. [PMID: 23168156] doi:10.2500/ajra.2012.26.3809
27. Ulsø C, Vase P, Stoksted P. Long-term results of dermatoplasty in the treatment of hereditary haemorrhagic telangiectasia. *J Laryngol Otol*. 1983;97:223-6. [PMID: 6339655]
28. Lund VJ, Darby Y, Rimmer J, et al. Nasal closure for severe hereditary haemorrhagic telangiectasia in 100 patients. The Lund modification of the Young's procedure: a 22-year experience. *Rhinology*. 2017;55:135-141. [PMID: 28064338] doi:10.4193/Rhin16.315
29. Rimmer J, Lund VJ. A modified technique for septodermoplasty in hereditary hemorrhagic telangiectasia. *Laryngoscope*. 2014;124: 67-9. [PMID: 23832829] doi:10.1002/lary.24303
30. van Tuyl SA, Letteboer TG, Rogge-Wolf C, et al. Assessment of intestinal vascular malformations in patients with hereditary hemorrhagic telangiectasia and anemia. *Eur J Gastroenterol Hepatol*. 2007;19:153-8. [PMID: 17273001]
31. Canzonieri C, Centenara L, Orati F, et al. Endoscopic evaluation of gastrointestinal tract in patients with hereditary hemorrhagic telangiectasia and correlation with their genotypes. *Genet Med*. 2014; 16:3-10. [PMID: 23722869] doi:10.1038/gim.2013.62
32. Ingrosso M, Sabbà C, Pisani A, et al. Evidence of small-bowel involvement in hereditary hemorrhagic telangiectasia: a capsule-endoscopic study. *Endoscopy*. 2004;36:1074-9. [PMID: 15578297]
33. Grève E, Moussata D, Gaudin JL, et al. High diagnostic and clinical impact of small-bowel capsule endoscopy in patients with hereditary hemorrhagic telangiectasia with overt digestive bleeding and/or severe anemia. *Gastrointest Endosc*. 2010;71:760-7. [PMID: 20170910] doi:10.1016/j.gie.2009.11.004
34. Nam SJ, Lee HS, Lim YJ. Evaluation of gastric disease with capsule endoscopy. *Clin Endosc*. 2018;51:323-328. [PMID: 30078305] doi:10.5946/ce.2018.092



35. Kjeldsen AD, Vase P, Green A. Hereditary haemorrhagic telangiectasia: a population-based study of prevalence and mortality in Danish patients. *J Intern Med*. 1999;245:31-9. [PMID: 10095814]
36. Kasthuri RS, Montifar M, Nelson J, et al; Brain Vascular Malformation Consortium HHT Investigator Group. Prevalence and predictors of anemia in hereditary hemorrhagic telangiectasia [Letter]. *Am J Hematol*. 2017. [PMID: 28639385] doi:10.1002/ajh.24832
37. Pasculli G, Resta F, Guastamacchia E, et al. Health-related quality of life in a rare disease: hereditary hemorrhagic telangiectasia (HHT) or Rendu-Osler-Weber disease. *Qual Life Res*. 2004;13:1715-23. [PMID: 15651542]
38. Geisthoff UW, Heckmann K, D'Amelio R, et al. Health-related quality of life in hereditary hemorrhagic telangiectasia. *Otolaryngol Head Neck Surg*. 2007;136:726-33; discussion 734-5. [PMID: 17478205]
39. Donaldson JW, McKeever TM, Hall IP, et al. Complications and mortality in hereditary hemorrhagic telangiectasia: a population-based study. *Neurology*. 2015;84:1886-93. [PMID: 25862798] doi:10.1212/WNL.0000000000001538
40. Iyer VN, Brinjikji W, Apala D, et al. Impact of age on outcomes in hospitalized patients with hereditary hemorrhagic telangiectasia. *Adv Hematol*. 2018;2018:4798425. [PMID: 29610574] doi:10.1155/2018/4798425
41. Brinjikji W, Wood CP, Lanzino G, et al. High rates of bleeding complications among hospitalized patients with hereditary hemorrhagic telangiectasia in the United States. *Ann Am Thorac Soc*. 2016;13:1505-11. [PMID: 27387823] doi:10.1513/AnnalsATS.201603-200OC
42. Becq A, Rahmi G, Perrod G, et al. Hemorrhagic angiodysplasia of the digestive tract: pathogenesis, diagnosis, and management. *Gastrointest Endosc*. 2017;86:792-806. [PMID: 28554655] doi:10.1016/j.gie.2017.05.018
43. Kwan V, Bourke MJ, Williams SJ, et al. Argon plasma coagulation in the management of symptomatic gastrointestinal vascular lesions: experience in 100 consecutive patients with long-term follow-up. *Am J Gastroenterol*. 2006;101:58-63. [PMID: 16405534]
44. Zaffar N, Ravichakaravathy T, Faughnan ME, et al. The use of anti-fibrinolytic agents in patients with HHT: a retrospective survey. *Ann Hematol*. 2015;94:145-52. [PMID: 25064693] doi:10.1007/s00277-014-2169-y
45. Faughnan ME, Gossage JR, Chakinala MM, et al. Pazopanib may reduce bleeding in hereditary hemorrhagic telangiectasia. *Angiogenesis*. 2019;22:145-155. [PMID: 30191360] doi:10.1007/s10456-018-9646-1
46. Shovlin CL, Awan I, Cahilog Z, et al. Reported cardiac phenotypes in hereditary hemorrhagic telangiectasia emphasize burdens from arrhythmias, anemia and its treatments, but suggest reduced rates of myocardial infarction. *Int J Cardiol*. 2016;215:179-85. [PMID: 27116331] doi:10.1016/j.ijcard.2016.04.006
47. Chaturvedi S, Kohli R, Schaefer N, et al. Characteristics and outcomes of venous thromboembolism in patients with hereditary hemorrhagic telangiectasia [Letter]. *Thromb Res*. 2018;169:41-43. [PMID: 30015226] doi:10.1016/j.thromres.2018.07.004
48. Hallberg L, Ryttinger L, Sölvell L. Side-effects of oral iron therapy. A double-blind study of different iron compounds in tablet form. *Acta Med Scand Suppl*. 1966;459:3-10. [PMID: 5957969]
49. Finnamore H, Le Couteur J, Hickson M, et al. Hemorrhage-adjusted iron requirements, hematinics and hepcidin define hereditary hemorrhagic telangiectasia as a model of hemorrhagic iron deficiency. *PLoS One*. 2013;8:e76516. [PMID: 24146883] doi:10.1371/journal.pone.0076516
50. Peyrin-Biroulet L, Williet N, Cacoub P. Guidelines on the diagnosis and treatment of iron deficiency across indications: a systematic review. *Am J Clin Nutr*. 2015;102:1585-94. [PMID: 26561626] doi:10.3945/ajcn.114.103366
51. Moretti D, Goede JS, Zeder C, et al. Oral iron supplements increase hepcidin and decrease iron absorption from daily or twice-daily doses in iron-depleted young women. *Blood*. 2015;126:1981-9. [PMID: 26289639] doi:10.1182/blood-2015-05-642223
52. Stoffel NU, Cercamondi CI, Brittenham G, et al. Iron absorption from oral iron supplements given on consecutive versus alternate days and as single morning doses versus twice-daily split dosing in iron-depleted women: two open-label, randomised controlled trials. *Lancet Haematol*. 2017;4:e524-e533. [PMID: 29032957] doi:10.1016/S2352-3026(17)30182-5
53. DeLoughery TG. Safety of oral and intravenous iron. *Acta Haematol*. 2019;142:8-12. [PMID: 30970354] doi:10.1159/000496966
54. Ganzoni AM. [Intravenous iron-dextran: therapeutic and experimental possibilities]. *Schweiz Med Wochenschr*. 1970;100:301-3. [PMID: 5413918]
55. Thielemans L, Layton DM, Shovlin CL. Low serum haptoglobin and blood films suggest intravascular hemolysis contributes to severe anemia in hereditary hemorrhagic telangiectasia [Letter]. *Haematologica*. 2019;104:e127-e130. [PMID: 30337360] doi:10.3324/haematol.2018.205682
56. Edwards CP, Shehata N, Faughnan ME. Hereditary hemorrhagic telangiectasia patients can tolerate anticoagulation. *Ann Hematol*. 2012;91:1959-68. [PMID: 23053175] doi:10.1007/s00277-012-1553-8
57. Devlin HL, Hosman AE, Shovlin CL. Antiplatelet and anticoagulant agents in hereditary hemorrhagic telangiectasia [Letter]. *N Engl J Med*. 2013;368:876-8. [PMID: 23445111] doi:10.1056/NEJMc1213554
58. Shovlin CL, Millar CM, Droegge F, et al; VASCERN-HHT. Safety of direct oral anticoagulants in patients with hereditary hemorrhagic telangiectasia. *Orphanet J Rare Dis*. 2019;14:210. [PMID: 31462308] doi:10.1186/s13023-019-1179-1
59. Vorselaars VM, Velthuis S, Swaans MJ, et al. Percutaneous left atrial appendage closure—an alternative strategy for anticoagulation in atrial fibrillation and hereditary hemorrhagic telangiectasia? *Cardiovasc Diagn Ther*. 2015;5:49-53. [PMID: 25774347] doi:10.3978/j.issn.2223-3652.2015.01.02
60. European Association for the Study of the Liver. EASL clinical practice guidelines: vascular diseases of the liver. *J Hepatol*. 2016;64:179-202. [PMID: 26516032] doi:10.1016/j.jhep.2015.07.040
61. Buscarini E, Danesino C, Olivieri C, et al. Doppler ultrasonographic grading of hepatic vascular malformations in hereditary hemorrhagic telangiectasia—results of extensive screening. *Ultraschall Med*. 2004;25:348-55. [PMID: 15368138]
62. Caselitz M, Bahr MJ, Bleck JS, et al. Sonographic criteria for the diagnosis of hepatic involvement in hereditary hemorrhagic telangiectasia (HHT). *Hepatology*. 2003;37:1139-46. [PMID: 12717395]
63. Schelker RC, Barreiros AP, Hart C, et al. Macro- and microcirculation patterns of intrahepatic blood flow changes in patients with hereditary hemorrhagic telangiectasia. *World J Gastroenterol*. 2017;23:486-495. [PMID: 28210085] doi:10.3748/wjg.v23.i3.486
64. Buonamico P, Suppressa P, Lenato GM, et al. Liver involvement in a large cohort of patients with hereditary hemorrhagic telangiectasia: echo-color-Doppler vs multislice computed tomography study. *J Hepatol*. 2008;48:811-20. [PMID: 18321607] doi:10.1016/j.jhep.2007.12.022
65. Buscarini E, Gebel M, Ocran K, et al. Interobserver agreement in diagnosing liver involvement in hereditary hemorrhagic telangiectasia by Doppler ultrasound. *Ultrasound Med Biol*. 2008;34:718-25. [PMID: 18207308] doi:10.1016/j.ultrasmedbio.2007.11.007
66. Wu JS, Saluja S, Garcia-Tsao G, et al. Liver involvement in hereditary hemorrhagic telangiectasia: CT and clinical findings do not correlate in symptomatic patients. *AJR Am J Roentgenol*. 2006;187:W399-405. [PMID: 16985112]
67. Scardapane A, Stabile Ianora A, Sabbà C, et al. Dynamic 4D MR angiography versus multislice CT angiography in the evaluation of vascular hepatic involvement in hereditary haemorrhagic telangiectasia. *Radiol Med*. 2012;117:29-45. [PMID: 21643641] doi:10.1007/s11547-011-0688-3
68. Milot L, Kamaoui I, Gautier G, et al. Hereditary-hemorrhagic telangiectasia: one-step magnetic resonance examination in evaluation of liver involvement. *Gastroenterol Clin Biol*. 2008;32:677-85. [PMID: 18757148] doi:10.1016/j.gcb.2008.04.033
69. Buscarini E, Leandro G, Conte D, et al. Natural history and outcome of hepatic vascular malformations in a large cohort of patients



- with hereditary hemorrhagic telangiectasia. *Dig Dis Sci*. 2011;56:2166-78. [PMID: 21290179] doi:10.1007/s10620-011-1585-2
70. Singh S, Swanson KL, Hathcock MA, et al. Identifying the presence of clinically significant hepatic involvement in hereditary haemorrhagic telangiectasia using a simple clinical scoring index. *J Hepatol*. 2014;61:124-31. [PMID: 24607625] doi:10.1016/j.jhep.2014.02.028
71. Buscarini E, Buscarini L, Danesino C, et al. Hepatic vascular malformations in hereditary hemorrhagic telangiectasia: Doppler sonographic screening in a large family. *J Hepatol*. 1997;26:111-8. [PMID: 9148001]
72. Young LH, Henderson K, Pollak JS, et al. Predictors of death in patients with HHT, liver vascular malformations (LVMs) and symptomatic heart failure (HF) [Abstract]. In: Bernabeu C, Botella LM, Brady A, et al, eds. 10th International Hereditary Hemorrhagic Telangiectasia Scientific Conference, Cork, Ireland, 12-15 June 2013. *Hematol Rep*. 2013;5. Abstract no. C-008.
73. Buscarini E, Botella LM, Geisthoff U, et al; VASCERN-HHT. Safety of thalidomide and bevacizumab in patients with hereditary hemorrhagic telangiectasia. *Orphanet J Rare Dis*. 2019;14:28. [PMID: 30717761] doi:10.1186/s13023-018-0982-4
74. Lerut J, Orlando G, Adam R, et al; European Liver Transplant Association. Liver transplantation for hereditary hemorrhagic telangiectasia: report of the European Liver Transplant Registry. *Ann Surg*. 2006;244:854-62; discussion 862-4. [PMID: 17122610]
75. Dupuis-Girod S, Chesnais AL, Ginon I, et al. Long-term outcome of patients with hereditary hemorrhagic telangiectasia and severe hepatic involvement after orthotopic liver transplantation: a single-center study. *Liver Transpl*. 2010;16:340-7. [PMID: 20209594] doi:10.1002/lt.21990
76. Dumortier J, Dupuis-Girod S, Valette PJ, et al. Recurrence of hereditary hemorrhagic telangiectasia after liver transplantation: clinical implications and physiopathological insights. *Hepatology*. 2019;69:2232-2240. [PMID: 30549294] doi:10.1002/hep.30424
77. Cymerman U, Vera S, Pece-Barbara N, et al. Identification of hereditary hemorrhagic telangiectasia type 1 in newborns by protein expression and mutation analysis of endoglin. *Pediatr Res*. 2000;47:24-35. [PMID: 10625079]
78. Giordano P, Nigro A, Lenato GM, et al. Screening for children from families with Rendu-Osler-Weber disease: from geneticist to clinician. *J Thromb Haemost*. 2006;4:1237-45. [PMID: 16706966]
79. Bossler AD, Richards J, George C, et al. Novel mutations in ENG and ACVRL1 identified in a series of 200 individuals undergoing clinical genetic testing for hereditary hemorrhagic telangiectasia (HHT): correlation of genotype with phenotype. *Hum Mutat*. 2006;27:667-75. [PMID: 16752392]
80. Al-Saleh S, Mei-Zahav M, Faughnan ME, et al. Screening for pulmonary and cerebral arteriovenous malformations in children with hereditary haemorrhagic telangiectasia. *Eur Respir J*. 2009;34:875-81. [PMID: 19386691] doi:10.1183/09031936.00030009
81. Curie A, Lesca G, Cottin V, et al. Long-term follow-up in 12 children with pulmonary arteriovenous malformations: confirmation of hereditary hemorrhagic telangiectasia in all cases. *J Pediatr*. 2007;151:299-306. [PMID: 17719943]
82. Giordano P, Lenato GM, Suppressa P, et al. Hereditary hemorrhagic telangiectasia: arteriovenous malformations in children. *J Pediatr*. 2013;163:179-86.e1-3. [PMID: 23535011] doi:10.1016/j.jpeds.2013.02.009
83. Gefen AM, White AJ. Asymptomatic pulmonary arteriovenous malformations in children with hereditary hemorrhagic telangiectasia. *Pediatr Pulmonol*. 2017;52:1194-1197. [PMID: 28608633] doi:10.1002/ppul.23686
84. Latino GA, Al-Saleh S, Alharbi N, et al. Prevalence of pulmonary arteriovenous malformations in children versus adults with hereditary hemorrhagic telangiectasia. *J Pediatr*. 2013;163:282-4. [PMID: 23522378] doi:10.1016/j.jpeds.2013.02.021
85. Faughnan ME, Thabet A, Mei-Zahav M, et al. Pulmonary arteriovenous malformations in children: outcomes of transcatheter embolotherapy. *J Pediatr*. 2004;145:826-31. [PMID: 15580209]
86. Al-Saleh S, Dragulescu A, Manson D, et al. Utility of contrast echocardiography for pulmonary arteriovenous malformation screening in pediatric hereditary hemorrhagic telangiectasia. *J Pediatr*. 2012;160:1039-43.e1. [PMID: 22196589] doi:10.1016/j.jpeds.2011.11.038
87. Karam C, Sellier J, Mansencal N, et al. Reliability of contrast echocardiography to rule out pulmonary arteriovenous malformations and avoid CT irradiation in pediatric patients with hereditary hemorrhagic telangiectasia. *Echocardiography*. 2015;32:42-8. [PMID: 24813063] doi:10.1111/echo.12615
88. Fernandopulle N, Mertens L, Klingel M, et al. Echocardiography grading for pulmonary arteriovenous malformation screening in children with hereditary hemorrhagic telangiectasia. *J Pediatr*. 2018;195:288-291.e1. [PMID: 29274888] doi:10.1016/j.jpeds.2017.11.047
89. Westermann CJ, Rosina AF, De Vries V, et al. The prevalence and manifestations of hereditary hemorrhagic telangiectasia in the Afro-Caribbean population of the Netherlands Antilles: a family screening. *Am J Med Genet A*. 2003;116A:324-8. [PMID: 12522784]
90. Hosman AE, de Gussem EM, Balemans WAF, et al. Screening children for pulmonary arteriovenous malformations: evaluation of 18 years of experience. *Pediatr Pulmonol*. 2017;52:1206-1211. [PMID: 28407366] doi:10.1002/ppul.23704
91. Ratjen A, Au J, Carpenter S, et al. Growth of pulmonary arteriovenous malformations in pediatric patients with hereditary hemorrhagic telangiectasia. *J Pediatr*. 2019;208:279-281. [PMID: 30853205] doi:10.1016/j.jpeds.2018.12.069
92. Krings T, Ozanne A, Chng SM, et al. Neurovascular phenotypes in hereditary haemorrhagic telangiectasia patients according to age. Review of 50 consecutive patients aged 1 day-60 years. *Neuroradiology*. 2005;47:711-20. [PMID: 16136265]
93. Morgan T, McDonald J, Anderson C, et al. Intracranial hemorrhage in infants and children with hereditary hemorrhagic telangiectasia (Osler-Weber-Rendu syndrome). *Pediatrics*. 2002;109:E12. [PMID: 11773580]
94. Saleh M, Carter MT, Latino GA, et al. Brain arteriovenous malformations in patients with hereditary hemorrhagic telangiectasia: clinical presentation and anatomical distribution. *Pediatr Neurol*. 2013;49:445-50. [PMID: 24080277] doi:10.1016/j.pediatrneurol.2013.07.021
95. Hetts SW, Keenan K, Fullerton HJ, et al. Pediatric intracranial nongalenic pial arteriovenous fistulas: clinical features, angioarchitecture, and outcomes. *AJNR Am J Neuroradiol*. 2012;33:1710-9. [PMID: 22766672] doi:10.3174/ajnr.A3194
96. Mori H, Aoki S, Okubo T, et al. Two-dimensional thick-slice MR digital subtraction angiography in the assessment of small to medium-size intracranial arteriovenous malformations. *Neuroradiology*. 2003;45:27-33. [PMID: 12525951]
97. Mukherji SK, Quisling RG, Kubilis PS, et al. Intracranial arteriovenous malformations: quantitative analysis of magnitude contrast MR angiography versus gradient-echo MR imaging versus conventional angiography. *Radiology*. 1995;196:187-93. [PMID: 7784565]
98. Gauvrit JY, Oppenheim C, Nataf F, et al. Three-dimensional dynamic magnetic resonance angiography for the evaluation of radiosurgically treated cerebral arteriovenous malformations. *Eur Radiol*. 2006;16:583-91. [PMID: 16220210]
99. Krings T, Chng SM, Ozanne A, et al. Hereditary hemorrhagic telangiectasia in children: endovascular treatment of neurovascular malformations: results in 31 patients. *Neuroradiology*. 2005;47:946-54. [PMID: 16163493]
100. Meybodi AT, Kim H, Nelson J, et al. Surgical treatment vs non-surgical treatment for brain arteriovenous malformations in patients with hereditary hemorrhagic telangiectasia: a retrospective multicenter consortium study. *Neurosurgery*. 2018;82:35-47. [PMID: 289734260] doi:10.1093/neuros/nyx168
101. Hetts SW, Cooke DL, Nelson J, et al. Influence of patient age on angioarchitecture of brain arteriovenous malformations. *AJNR Am J Neuroradiol*. 2014;35:1376-80. [PMID: 24627452] doi:10.3174/ajnr.A3886
102. Guo Y, Saunders T, Su H, et al; University of California, San Francisco Brain Arteriovenous Malformation (UCSF bAVM) Study

**Project.** Silent intralesional microhemorrhage as a risk factor for brain arteriovenous malformation rupture. *Stroke*. 2012;43:1240-6. [PMID: 22308253] doi:10.1161/STROKEAHA.111.647263

103. **Shovlin CL, Sodhi V, McCarthy A, et al.** Estimates of maternal risks of pregnancy for women with hereditary haemorrhagic telangiectasia (Osler-Weber-Rendu syndrome): suggested approach for obstetric services. *BJOG*. 2008;115:1108-15. [PMID: 18518871] doi:10.1111/j.1471-0528.2008.01786.x

104. **Shovlin CL, Winstock AR, Peters AM, et al.** Medical complications of pregnancy in hereditary haemorrhagic telangiectasia. *QJM*. 1995;88:879-87. [PMID: 8593547]

105. **Ference BA, Shannon TM, White RI Jr, et al.** Life-threatening pulmonary hemorrhage with pulmonary arteriovenous malformations and hereditary hemorrhagic telangiectasia. *Chest*. 1994;106:1387-90. [PMID: 7956388]

106. **Gershon AS, Faughnan ME, Chon KS, et al.** Transcatheter embolotherapy of maternal pulmonary arteriovenous malformations during pregnancy. *Chest*. 2001;119:470-7. [PMID: 11171725]

107. **de Gussem EM, Lausman AY, Beder AJ, et al.** Outcomes of pregnancy in women with hereditary hemorrhagic telangiectasia. *Obstet Gynecol*. 2014;123:514-20. [PMID: 24499751] doi:10.1097/AOG.0000000000000120

**Current Author Addresses:** Drs. Faughnan, Palda, and Lausman; Mr. Cortes; and Ms. Pantalone: St. Michael's Hospital, University of Toronto, 30 Bond Street, Toronto, ON M5B 1W8, Canada.

Drs. Mager and Post: St. Antonius Hospital, Koekoekslaan 1, Nieuwegein 3435 CM, the Netherlands.

Dr. Hetts: Department of Neurointerventional Radiology, University of California, San Francisco, 505 Parnassus Avenue, L-351, San Francisco, CA 94143.

Ms. Lang-Robertson: Centre for Effective Practice, 400 University Avenue, Suite 2100, Toronto, ON M5G 1S5, Canada.

Dr. Buscarini: UOC Gastroenterologia ed Endoscopia Digestiva, Ospedale Maggiore, ASST Crema, Largo Dossena 2, Crema, 26013 Italy.

Dr. Deslandres: Department of Gastroenterology, CHUM, Hotel Dieu, 1051 Rue Sanguinet, C.02.7013, Montreal, QC H2X 0C1, Canada.

Dr. Kasthuri: Division of Hematology/Oncology, University of North Carolina, Chapel Hill, 116 Manning Drive, CB7035, 8206B Mary Ellen Jones, Chapel Hill, NC 27599.

Dr. Poetker: Department of Otolaryngology, Froedtert and Medical College of Wisconsin, 8701 Watertown Plank Road, Milwaukee, WI 53226.

Dr. Ratjen: Department of Pediatrics, The Hospital for Sick Children, 555 University Avenue, Toronto, ON M5G 1X8, Canada.

Dr. Chesnutt: VA Portland Health Care System, HHT Center of Excellence, 3710 SW US Veterans Hospital Road, P2IES, Portland, OR 97221.

Ms. Clancy and Dr. Olitsky: Cure HHT, PO Box 329, Monkton, MD 21111.

Dr. Whitehead: Department of Cardiovascular Medicine and Pediatric Cardiology, University of Utah Medical Center, 30 North 1900 South, Room 4A100, Salt Lake City, UT 84132.

Dr. Al-Samkari: Division of Hematology, Massachusetts General Hospital, Harvard Medical School, Zero Emerson Place Suite 118, Office 112, Boston, MA 02114.

Dr. Chakinala: Department of Pulmonology and Critical Care, Washington University School of Medicine, 660 South Euclid Avenue, Campus Box 8052, St. Louis, MO 63110.

Dr. Conrad: Zuckerberg San Francisco General Hospital, 1001 Potrero Avenue, San Francisco, CA 94114.

Ms. Crocione: HHT Europe, Via Europa 63, Rome, 05022 Italy.

Dr. Darling: Department of Hepatology, University of North Carolina, Chapel Hill, 8014 Burnett Womack Building, CB 7584, Chapel Hill, NC 27599.

Dr. de Gussem: Department of Medicine, Section of Respiriology, Grace Hospital, 300 Booth Drive, Winnipeg, MB R3J 3M7, Canada.

Dr. Dupuis-Girod: Hospices Civils de Lyon, Femme-Mère-Enfant, 59 Boulevard Pinel, 69677 Bron, France.

Dr. Foy: Department of Hematology, Froedtert and Medical College of Wisconsin, 9200 West Wisconsin Avenue, Milwaukee, WI 53226.

Dr. Geisthoff: Department of Otorhinolaryngology, Head and Neck Surgery, University Hospital of Marburg, Phillips University Marburg, Baldingerstraße, 35043 Marburg, Germany.

Dr. Gossage: Augusta University, 1120 Fifteenth Street, BBR-5513, Augusta, GA 30912.

Dr. Hammill: Division of Hematology, Cancer and Blood Diseases Institute, Cincinnati Children's Hospital, and Department of Pediatrics, University of Cincinnati, 3333 Burnet Avenue, MLC 7015, Cincinnati, OH 45229.

Dr. Heimdal: Department of Genetics, Oslo University Hospital, Rikshospitalet, PO Box 4950 Nydalen, 4950 Oslo, Norway. Ms. Henderson: Yale University School of Medicine, PO Box 208042, New Haven, CT 06520.

Dr. Iyer: Division of Pulmonary and Critical Care Medicine, Mayo Clinic, Rochester, MN 55905.

Dr. Kjeldsen: Department of Otorhinolaryngology Head and Neck Surgery, HHT-center OUH, J.B.Winsløvs Vej 4, 5000 Odense, Denmark.

Dr. Komiyama: Department of Neurointervention, Osaka City General Hospital, 2-13-22, Miyakojima-Hondori, Miyakojima, Osaka 534-0021, Japan.

Dr. Korenblatt: Department of Hepatology, Washington University School of Medicine, 660 South Euclid Avenue, Campus Box 8124, St. Louis, MO 63110.

Ms. McDonald: Department of Pathology and Radiology, University of Utah Medical Center, 50 North Medical Drive 4A100, Salt Lake City, UT 84132.

Dr. McWilliams: Department of Interventional Radiology, University of California, Los Angeles, 757 Westwood Plaza, Suite 2125C, Los Angeles, CA 90095.

Dr. Meek: Department of Interventional Radiology, University of Arkansas for Medical Sciences, 4301 West Markham Street, Slot 556, Little Rock, AR 72205.

Dr. Mei-Zahav: Pulmonary Institute, Schneider CMCI, 14 Kaplan Street, Petach Tikva, 49202, Israel.

Dr. Piccirillo: Department of Otolaryngology-Head & Neck Surgery, Washington University School of Medicine, 660 South Euclid Avenue, Campus Box 8115, St. Louis, MO 63110.

Dr. Porteous: Department of Genetics University of Edinburgh, Center of Molecular Medicine, Crewe Road, Edinburgh EH4 2XU, Scotland.

Dr. Radovanovic: Department of Neurosurgery, University Health Network, Toronto Western Hospital, 399 Bathurst Street, Toronto, ON M5T 2S8, Canada.

Dr. Rochon: University of Colorado Hospital, 12631 East 17th Avenue, Mailstop 8200, Aurora, CO 80048.

Dr. Rodriguez-Lopez: Department of Pulmonology, Massachusetts General Hospital, 55 Fruit Street, Bulfinch 148, Boston, MA 02114.

Dr. Sabba: Department of Internal Medicine, University of Bari, Policlinico, Piazza Giulio Cesare, Bari, 70124 Italy.

Dr. Serra: Department of Internal Medicine, Hospital Italiano de Buenos Aires, Araoz 2067, Buenos Aires 1425, Argentina.

Dr. Shovlin: Department of Pulmonology, Hammersmith Hospital, Du Cane Road, London W12 0NN, England.

Dr. White: Division of Pediatric Immunology and Rheumatology, Washington University School of Medicine, One Children's Place, St. Louis, MO 63110.

Dr. Winship: Genomic Medicine, Royal Melbourne Hospital and University of Melbourne, Grattan Street, Melbourne, VIC 3002, Australia.

Dr. Zarrabeitia: Hospital Sierrallana, Barrio Ganzo s/n, 39300 Torrelavega, Cantabria, Spain.

**Author Contributions:** Conception and design: M.E. Faughnan, J.J. Mager, S.W. Hetts, V.A. Palda, E. Buscarini, E. Deslandres, R.S. Kasthuri, F. Ratjen, M.S. Chesnutt, E. de Gussem, A. Hammill, A.D. Kjeldsen, J. McDonald, M. Mei-Zahav, S. Olitsky, B. Plahn, M.C. Post, C. Shovlin, A.J. White, I. Winship.

Analysis and interpretation of the data: M.E. Faughnan, J.J. Mager, S.W. Hetts, V.A. Palda, E. Buscarini, E. Deslandres, R.S. Kasthuri, D. Poetker, F. Ratjen, M.S. Chesnutt, M. Clancy, K.J.

Whitehead, H. Al-Samkari, M. Conrad, C. Crocione, J. Darling, E. de Gussem, P. Foy, U. Geisthoff, J.R. Gossage, K. Heimdal, K. Henderson, V.N. Iyer, A.D. Kjeldsen, M. Komiyama, J. McDonald, J. McMahon, J. McWilliams, M.E. Meek, M. Mei-Zahav, S. Olitsky, S. Palmer, R. Pantalone, J.F. Piccirillo, M.E.M. Porteous, I. Radovanovic, P.J. Rochon, C. Sabba, M. Serra, C. Shovlin, D. Sprecher, A.J. White, I. Winship, R. Zarrabeitia.

Drafting of the article: M.E. Faughnan, J.J. Mager, S.W. Hetts, V.A. Palda, K. Lang-Robertson, E. Buscarini, E. Deslandres, R.S. Kasthuri, A. Lausman, D. Poetker, F. Ratjen, M.S. Chesnutt, M. Clancy, K.J. Whitehead, M. Chakinala, E. de Gussem, J.R. Gossage, K. Henderson, V.N. Iyer, M. Mei-Zahav, R. Pantalone, B. Plahn, M.C. Post, P.J. Rochon, M. Serra, R. Zarrabeitia.

Critical revision of the article for important intellectual content: M.E. Faughnan, J.J. Mager, S.W. Hetts, V.A. Palda, E. Buscarini, E. Deslandres, R.S. Kasthuri, A. Lausman, D. Poetker, F. Ratjen, M.S. Chesnutt, K.J. Whitehead, H. Al-Samkari, M. Chakinala, M. Conrad, J. Darling, E. de Gussem, S. Dupuis-Girod, P. Foy, U. Geisthoff, J.R. Gossage, A. Hammill, K. Heimdal, K. Henderson, V.N. Iyer, A.D. Kjeldsen, K. Korenblatt, J. McWilliams, M. Mei-Zahav, R. Pantalone, J.F. Piccirillo, M.E.M. Porteous, M.C. Post, I. Radovanovic, P.J. Rochon, J. Rodriguez-Lopez, C. Sabba, M. Serra, C. Shovlin, D. Sprecher, A.J. White, I. Winship, R. Zarrabeitia.

Final approval of the article: M.E. Faughnan, J.J. Mager, S.W. Hetts, V.A. Palda, K. Lang-Robertson, E. Buscarini, E. Deslan-

dres, R.S. Kasthuri, A. Lausman, D. Poetker, F. Ratjen, M.S. Chesnutt, M. Clancy, K.J. Whitehead, H. Al-Samkari, M. Chakinala, M. Conrad, D. Cortes, C. Crocione, J. Darling, E. de Gussem, C. Derksen, S. Dupuis-Girod, P. Foy, U. Geisthoff, J.R. Gossage, A. Hammill, K. Heimdal, K. Henderson, V.N. Iyer, A.D. Kjeldsen, M. Komiyama, K. Korenblatt, J. McDonald, J. McMahon, J. McWilliams, M.E. Meek, M. Mei-Zahav, S. Olitsky, S. Palmer, R. Pantalone, J.F. Piccirillo, B. Plahn, M.E.M. Porteous, M.C. Post, I. Radovanovic, P.J. Rochon, J. Rodriguez-Lopez, C. Sabba, M. Serra, C. Shovlin, D. Sprecher, A.J. White, I. Winship, R. Zarrabeitia.

Provision of study materials or patients: F. Ratjen, E. de Gussem, J. McDonald.

Obtaining of funding: M.E. Faughnan, M. Clancy, J. McMahon, I. Winship.

Administrative, technical, or logistic support: M.E. Faughnan, S.W. Hetts, V.A. Palda, C. Derksen, S. Olitsky, B. Plahn.

Collection and assembly of data: M.E. Faughnan, J.J. Mager, S.W. Hetts, K. Lang-Robertson, E. Buscarini, E. Deslandres, R.S. Kasthuri, A. Lausman, D. Poetker, F. Ratjen, H. Al-Samkari, M. Chakinala, D. Cortes, E. de Gussem, J.R. Gossage, A. Hammill, K. Henderson, V.N. Iyer, A.D. Kjeldsen, M. Komiyama, J. McDonald, J. McWilliams, M. Mei-Zahav, M.C. Post, I. Radovanovic, P.J. Rochon, J. Rodriguez-Lopez, M. Serra, C. Shovlin, A.J. White, I. Winship, R. Zarrabeitia.