

СОЮЗДЕ НАН
ОБЪЕДИНЕННОГО
ИНСТИТУТА
ЯДЕРНЫХ
ИССЛЕДОВАНИИ
ДУБНА



E2 - 7890

N.B.Skachkov

**A NEW INVARIANT DESCRIPTION
OF THE NUCLEON MEAN-SQUARE RADIUS**

1974

**ЛАБОРАТОРИЯ
ТЕОРЕТИЧЕСКОЙ ФИЗИКИ**

We regret that some of the pages in the microfiche copy of this report may not be up to the proper legibility standards, even though the best possible copy was used for preparing the master fiche.

RANGE OF JINR PUBLICATIONS

The preprints and communications of the Joint Institute for Nuclear Research (JINR) are considered to be original publications. They are published in accordance with Article 4 of the JINR Statut. Difference between the preprints and communications consists in that text of the preprint will be published in future in one of scientific journals or in aperiodic collections.

Indexing

The preprints, communications and deposited publications of the JINR have a single numbering (four last figures of the index).

The first sign of the index - a letter, denotes the language the paper is published in:

"P" - published in Russian;

"E" - published in English;

"D" - published both in Russian and English.

The preprints and communications, which are distributed only to the Member States of the JINR, have no letter-index.

The figure following the letter denotes the subject category of the given publication. List of subject categories of the JINR publications is sent periodically to the recipients.

Above-described index is placed in the right upper corner of the cover and on the title-page of each publication.

References

In bibliographical references to the JINR preprints and communications we recommend to indicate: author's initials and name, an abbreviation of the name of the institute, index, place and year of publication.

An example of bibliographical reference:

1. I. Ivanov. JINR, P2-4985, Dubna, 1971.

E2 - 7890

N.B.Skachkov

**A NEW INVARIANT DESCRIPTION
OF THE NUCLEON MEAN-SQUARE RADIUS**

The nucleon mean-square radius is defined from the expansion of form factor $F(t)$ for small momentum transfer $t = q^2 = (p-k)^2$

$$F(t) = F(0) + \frac{1}{6} \langle r_0^2 \rangle t + \dots \quad (1)$$

or, identically

$$\langle r_0^2 \rangle = \frac{6 \left. \frac{\partial F(t)}{\partial t} \right|_{t=0}}{F(0)} \quad (2)$$

To make correspond the mean-square radius thus defined to the spatial distribution of a nucleon it is usual to choose a special reference frame, the Breit frame. In this system $\vec{p} = -\vec{k}$ because of that the time component of momentum is zero $q_0 = p_0 - k_0 = 0$ and the four-dimensional Fourier transformation reduces to the three-dimensional one

$$f(r) = \frac{1}{(2\pi)^3} \int F(-\vec{q}^2) e^{i\vec{q}\vec{r}} d\vec{q} \quad (3)$$

By means of thus defined spatial distribution for a nucleon $\langle r_0^2 \rangle$ in the Breit system is determined by the formula

$$\langle r_0^2 \rangle = \frac{\left\{ \left(i \frac{\partial}{\partial \vec{q}} \right)^2 F(-\vec{q}^2) \right\} \Big|_{\vec{q}^2=0}}{F(0)} = \frac{-6 \left. \frac{\partial F(-\vec{q}^2)}{\partial \vec{q}^2} \right|_{\vec{q}^2=0}}{F(0)} = \frac{\int r^2 f(r) d\vec{r}}{\int f(r) d\vec{r}} \quad (4)$$

However, such a description of the spatial distribution of a nucleon is not satisfactory since in the Breit system the nucleon itself is moving.

In the present note we will show that this difficulty can be overcome if in describing the nucleon spatial distribution one makes use of the introduced in ^{/1/} relativistic configurational representation which, as will be seen below, provides the invariant description of the nucleon spatial distribution.

To this end we note that on the mass shell $p_0^2 - \vec{p}^2 = M^2$ the momentum transfer squared $t = q^2 = (p-k)^2$ can be expressed through the 3-vector $\vec{\Lambda} = \vec{p}(-) \vec{k}$ which is the difference of two vectors in the Lobachevsky space ^{/1,2/}:

$$\vec{\Lambda} = \vec{p}(-) \vec{k} = (\gamma_p^{-1} \vec{k}) = \vec{p} - \frac{\vec{k}}{M} (p_0 - \frac{\vec{p} \cdot \vec{k}}{k_0 + M}), \quad (5)$$

$$\Lambda_0 = (p(-)k) = \frac{p_0 k_0 - \vec{p} \cdot \vec{k}}{M} = \sqrt{M^2 + \vec{\Lambda}^2} \quad (6)$$

in the following way

$$t = (p-k)^2 = 2M^2 - 2pk = 2M^2 - 2M \sqrt{M^2 + \vec{\Lambda}^2}. \quad (7)$$

Consequently, in any reference frame form factor $F(t)$ is parametrized by the square of 3-vector of momentum transfer that belongs to the Lobachevsky space ^{**}

$$F(t) = F(\vec{\Lambda}^2). \quad (8)$$

^{*} The equation $p_0^2 - \vec{p}^2 = M^2$ defines in the Minkowski space the surface of hyperboloid on the upper sheet of which the Lobachevsky space is realized ^{/4,5/}.

^{**} The Lobachevsky space first was applied for description of form factors in ^{/3/}.

The group of motion of the Lobachevsky space is the Lorentz group. Therefore in ^{/1/} it has been suggested to introduce the coordinate representation in the relativistic theory by means of expansions over the unitary irreducible representations of the Lorentz group

$$F(\mathbf{r}) = \frac{1}{(2\pi)^3} \int \xi(\vec{\Delta}, \vec{r}) F(\vec{\Delta}^2) \frac{d\vec{\Delta}}{\Delta_0} \quad (9)$$

which are realized by the functions:

$$\xi(\vec{\Delta}, \vec{r}) = \left(\frac{\Delta_0 - \vec{\Delta} \cdot \vec{n}}{M} \right)^{-1 - i\mathbf{r} \cdot \mathbf{M}} \quad (10)$$

$$\vec{r} = \mathbf{r} \cdot \vec{n}; \quad \vec{n}^2 = 1; \quad 0 \leq r < \infty$$

explicitly found in ^{/6/}. In doing so, as is shown in ^{/1/}, the parameter r plays the role of modulus of the radius-vector of the relativistic relative distance, since in the nonrelativistic limit when $\vec{\Delta} \approx \vec{p} (-) \vec{k} \rightarrow \vec{q} = \vec{p} (-) \vec{k}$ the functions (10) turn into the usual plane waves:

$$\xi(\vec{\Delta}, \vec{r}) \xrightarrow{c \rightarrow \infty} e^{i\vec{q} \cdot \vec{r}}$$

The modulus of the relativistic relative coordinate r is the relativistic invariant because it determines the eigenvalues of the invariant operator, the Casimir operator of the Lorentz group $\hat{C} = \vec{N}^2 - \vec{M}^2$, on functions (10):

$$\hat{C} \xi(\vec{\Delta}, \vec{r}) = (\vec{N}^2 - \vec{M}^2) \xi(\vec{\Delta}, \vec{r}) = \left(\frac{1}{M^2} + r^2 \right) \xi(\vec{\Delta}, \vec{r}). \quad (11)$$

Thus, the function $F(r)$ defined by (9) provides in the relativistic configurational representation the invariant description of the nucleon spatial distribution in any reference frame including the rest frame.

In the nonrelativistic limit the Casimir operator of the Lorentz group becomes the Casimir operator of the group of motion of three-dimensional Euclidean momen-

tum space, i.e., the operator $(i \frac{\partial}{\partial \vec{q}})^2$ in (14). The eigenvalue of the operator $(i \frac{\partial}{\partial \vec{q}})^2$ on the functions $e^{i\vec{q}\vec{r}}$, as is seen from (3), is the square of nonrelativistic relative coordinate r^2 .

It appears that the invariant description of the nucleon mean-square radius (1), (2) has a group-theoretical interpretation. Employing the fact that in the spherical coordinates

$$\chi_0 = M \operatorname{ch} \chi; \quad \vec{\chi} = M \operatorname{sh} \chi (\sin \theta \sin \phi, \sin \theta \cos \phi, \cos \theta)$$

the Casimir operator of the Lorentz group has the form of the Laplace operator on hyperboloid

$$\hat{C} = \vec{N}^2 - \vec{M}^2 = \frac{1}{M^2} \frac{\partial^2}{\partial \chi^2} - 2 \frac{\operatorname{cth} \chi}{M^2} \frac{\partial}{\partial \chi} - \frac{\chi^2}{M^2 \operatorname{sh}^2 \chi} \quad (12)$$

(the hyperbolic angle χ in the Lobachevsky geometry is called "rapidity" $\frac{v}{c}$) it is easy to verify that in virtue of the spherical symmetry one has $F(t) = -F | 2M^2 (1 - \operatorname{ch} \chi) |$

$$\frac{\{ \hat{C} F(t) \} |_{\chi=0}}{F(0)} = \frac{-\frac{6}{M^2} \frac{\partial F(t)}{\partial \chi^2} |_{\chi=0}}{F(0)} = \frac{6 \frac{\partial F(t)}{\partial t} |_{t=0}}{F(0)} \quad (13)$$

Expression (13) coincides with the invariant definition of the mean-square radius (2) which can now be connected with the invariant function $F(r)$ describing, in accordance with (9), the nucleon spatial distribution in an arbitrary reference frame \vec{r}^1

$$\langle r_0^2 \rangle = \frac{6 \frac{\partial F(t)}{\partial t} |_{t=0}}{F(0)} = \frac{\{ (\vec{N}^2 - \vec{M}^2) F(t) \} |_{t=0}}{F(0)} =$$

$$= \frac{\int \left(\frac{\hbar^2}{M^2 c^2} + r^2 \right) F(r) d\vec{r}}{\int F(r) d\vec{r}} = \frac{\hbar^2}{M^2 c^2} + \langle r^2 \rangle. \quad (14)$$

Hence it is seen that the relativistic relative coordinate describes the nucleon structure beyond its Compton wave length.

Due to the spherical symmetry of the form factor $F(r)$ transformation (9) and its inverse are as follows:

$$F(r) = \frac{1}{2\pi^2} \int \frac{\sin rM\lambda}{rM \operatorname{sh} \lambda} F(\lambda^2) \operatorname{sh}^2 \lambda d\lambda. \quad (15)$$

$$F(\lambda^2) = 4\pi \int \frac{\sin rM\lambda}{rM \operatorname{sh} \lambda} F(r) r^2 dr. \quad (16)$$

For the form factor $\Phi(\lambda) = \frac{\operatorname{sh} \lambda}{\lambda} F(r)$ with the same

normalization condition as $F(r)$, i.e., $\Phi(0) = F(0) = 1$ transformations (15) and (16) take the form of the conventional transformation with the zero-order Bessel functions, where, however, the relativistic coordinate is conjugated not to the momentum transfer but to the

"rapidity" $\lambda = \operatorname{Ar ch} \left(\frac{2M^2 - t}{2M^2} \right)$, corresponding to this momentum transfer.

Choosing different forms of the distribution $F(r)$, we obtain various expressions for the form factors $\Phi(\lambda)$ and $F(t)$. In particular, to the localized in

r -space particle $F(r) = \frac{\delta(r)}{r^2}$ (i.e., localized in the size of Compton wave length) there corresponds, according to (17), the form factor

$$F(t) = \frac{\chi}{\text{sh}\chi} = 2M^2 \frac{\text{Ar ch} \left(\frac{2M^2 - t}{2M^2} \right)}{\sqrt{t(t - 4M^2)}}, \quad (17)$$

decreasing at large $|t|$ by the law

$$F(t) \sim 2M^2 \frac{\ln \frac{|t|}{2M^2}}{|t|} \quad (18)$$

To the Gaussian distribution $F(r) = e^{-ar^2}$ there corresponds the form factor

$$F(t) = \frac{\pi}{a} \sqrt{\frac{\pi}{2}} \cdot \frac{\chi}{\text{sh}\chi} \cdot e^{-\frac{\chi^2 M^2}{4a}} \quad (19)$$

which has been obtained earlier by analogy with the nonrelativistic form factor in^{/7/} and whose asymptotic expression

$$F(t) \sim \frac{\ln \frac{|t|}{2M^2}}{|t|} e^{-M^2 \ln^2 \frac{|t|}{2M^2}} \quad (20)$$

coincides with that of the form factors introduced in^{/8/}. In^{/7,9/} it has been shown that the form factors of the type of Gaussian distribution over rapidity describe, for certain values of parameters, the experimental data better than the dipole formula.

Thus, we see that the relativistic configurational representation introduced in^{/1/} allows one to describe in the invariant way, the nucleon spatial distribution. In doing so, the nucleon mean-square radius in an arbitrary reference frame can be connected with the spatial distribution in the new coordinate space.

In conclusion the author thanks sincerely V.M.Dubovik, A.V.Efremov, V.G.Kadyshevsky, V.A.Matveev and V.A.Meshcheryakov for useful discussions.

Reference

1. V.G.Kadyshevsky, R.M.Mir-Kasimov, N.B.Skachkov. *Nuovo Cim.*, 55A, 233 (1968).
2. V.G.Kadyshevsky, R.M.Mir-Kasimov, N.B.Skachkov. "Particles and Nucleus", vol. 2, part 3, p. 635. Atomizdat (1972).
3. A.V.Efremov. *Z. Eksper. Teoret. Fiz.*, 53, 732 (1967).
4. V.A.Fock. "The Theory of Space, Time and Gravitation", Pergamon Press, Oxford (1967).
5. N.A.Chernikov. *Dokl. Akad. Nauk SSSR*, 114, 530 (1957); *Lectures in International Winter School on Theoretical Physics, Dubna (1964)*. "Particles and Nucleus". vol. 4, part. 3, 773 (1973).
6. I.S.Shapiro. *Dokl. Akad. Nauk SSSR*, 106, 647 (1956); *Soviet Phys. Doklady*, 1, 91 (1956).
7. M.I.Pavkovic. *Phys.Rev.*, 4D, 1724 (1971).
8. G.Mack. *Phys.Rev.*, 154, 1617 (1967).
9. Minh Duong-van. Preprint SLAC-1383 (1974).

Received by Publishing Department
on April 23, 1974.

Subject Categories of the JINR Publications

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Gryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids.
15.	Experimental physics of nuclear reactions at low energies.
16.	Health physics. Shieldings
17.	Solid state physics theory.

Will you fill blank spaces in your library?

You can receive by post the books listed below. Prices - in US \$,
including the packing and registered postage

I6-4888	Dosimetry of Radiation and Charged Particle Protection Physics. Dubna, 1969.	250 pg	4.85
D-6004	Binary Reactions of Hadrons at High Energies. Dubna, 1971.	768 pg	12.59
D10-6142	Proceedings of the International Symposium on Data Handling of Bubble and Spark Chambers. Dubna, 1971.	564 pg	9.96
D13-6210	Proceedings of the Vth International Symposium on Nuclear Electronics. Warsaw, 1971.	372 pg	6.36
D-6465	International School on Nuclear Structure, Alushta, 1972.	525 pg	9.54
D- 6840	Proceedings of the Second International Symposium on High Energy and Elementary Particle Physics (Štrbské Pleso, ČSSR, 1972).	398 pg	5.78
D2-7161	Proceedings of the III International Seminar on Non-Local Quantum Field Theory. Alushta, 1973.	280 pg	5.00
	Proceedings of the VI European Conference on Controlled Fusion and Plasma Physics. Moscow, 1973. v. 1.	666 pg	21.00
D1.2-7411	Deep Inelastic and Many-Body Processes. Dubna, 1973.	507 pg	8.76
P1.2-7642	Proceedings of the International School of Young Physicists on High Energy Physics. Gornji, 1973.	623 pg	10.53

- D13-7616 Proceedings of the VII International Symposium on Nuclear Electronics. Budapest, 1973. 372 pg
- D10-7707 Meeting on Programming and Mathematical Methods for Solving the Physical Problems. Dubna, 1973. 564 pg 8.23

Orders for the above-mentioned books can be sent at the address:
Publishing Department, JINR
Head Post Office, P.O. Box 79
101000 Moscow, USSR.



Conditions of Exchange

The preprints and communications of the Joint Institute for Nuclear Research are distributed free of charge on the mutual exchange basis to the universities, institutes, libraries, scientific groups and individual scientists of more than 50 countries.

We expect that the recipients of the JINR publications will have a possibility to display initiative in sending free of charge publications to Dubna. We receive, on the exchange basis, scientific books, journals, preprints and other types of publications on the subject categories of our Institute.

The only kind of publications that we do not require are the reprints of the articles already published in scientific journals.

In a number of cases we address to scientific institutions - the most prominent recipients of our publications - with the request to send us free of charge some books or subscribe for our library to the scientific journals published in their countries.

Requests

The Publishing Department fulfils annually about 3000 individual requests for our preprints and communications. The index of our publication must be obligatory indicated in such requests.

Addresses

Letters on all the questions concerning the exchange of publications as well as requests for individual publications are to be sent at the address:

**Publishing Department
Joint Institute for
Nuclear Research
Head Post Office,
P.O. Box 79
101000 Moscow,
U.S.S.R.**

We kindly ask to send all the publications on the exchange basis and also free of charge subscriptions to scientific journals at the address:

**Scientific-Technical Library
Joint Institute for
Nuclear Research
Head Post Office
P.O. Box 79
101000 Moscow,
U.S.S.R.**

Издательский отдел Объединенного института ядерных исследований.
Заказ 17844. Тираж 690. Уч.-изд. листов 0,5.

Редактор Э.В. Ивашевич. . Подписано к печати 25/1У-74 г.



74.10.8