

ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ

ДУБНА



E6 - 7762

R.Arlt, A.Jasiński, W.Neubert, H.-G.Ortlepp

ON THE DECAY SCHEMES

OF ^{123}Ba AND ^{125}Ba

~~B13, A34, D13~~

1974

ЛАБОРАТОРИЯ ЯДЕРНЫХ ПРОБЛЕМ

We regret that some of the pages in the microfiche copy of this report may not be up to the proper legibility standards, even though the best possible copy was used for preparing the master fiche.

RANGE OF JINR PUBLICATIONS

The preprints and communications of the Joint Institute for Nuclear Research (JINR) are considered to be original publications. They are published in accordance with Article 4 of the JINR Statut. Difference between the preprints and communications consists in that text of the preprint will be published in future in one of scientific journals or in aperiodic collections.

Indexing

The preprints, communications and deposited publications of the JINR have a single numbering (four last figures of the index).

The first sign of the index - a letter, denotes the language the paper is published in:

"P" - published in Russian;

"E" - published in English;

"D" - published both in Russian and English.

The preprints and communications, which are distributed only to the Member States of the JINR, have no letter-index.

The figure following the letter denotes the subject category of the given publication. List of subject categories of the JINR publications is sent periodically to the recipients.

Above-described index is placed in the right upper corner of the cover and on the title-page of each publication.

References

In bibliographical references to the JINR preprints and communications we recommend to indicate: author's initials and name, an abbreviation of the name of the institute, index, place and year of publication.

An example of bibliographical reference:

1. I. Ivanov. JINR, P2-4985, Dubna, 1971.

R.Arlt,¹ A.Jasiński,² W.Neubert,³ H.-G.Ortlepp¹

ON THE DECAY SCHEMES

OF ^{123}Ba AND ^{125}Ba

Submitted to *Acta Physica Polonica*

¹ Technische Universität, Dresden, GDR.

² Institut Badań Jądrowych, Świerk
k/Otwocka, Poland.

³ Zentralinstitut für Kernforschung,
Rossendorf, GDR.

Арлыт Р., Ясински А., Нойберт В., Ортелеп Х.-Г.

E6 - 7762

О схемах распада ^{123}Ba и ^{125}Ba

Нейтрондефицитные изотопы ^{123}Ba и ^{125}Ba , полученные в реакциях $^{114}\text{Sn}(^{12}\text{C}, 3n)^{123}\text{Ba}$ и $^{117}\text{Sn}(^{12}\text{C}, 4n)^{125}\text{Ba}$, исследовались с помощью Ge(Li) -детекторов. Задержанные бета-гамма совпадения были измерены NE102 и Ge(Li) -детекторами. На базе относительных интенсивностей и энергии гамма-переходов предложены схемы распада ^{123}Ba и ^{125}Ba .

Рез.

Препринт Объединенного института ядерных исследований.

Дубна, 1974

Arlt R., Jasiński A., Neubert W., Ortlepp H.-G.

E6 - 7762

On the Decay Schemes of ^{123}Ba and ^{125}Ba

The neutron deficient ^{123}Ba and ^{125}Ba isotopes obtained in the $^{114}\text{Sn}(^{12}\text{C}, 3n)^{123}\text{Ba}$ and $^{117}\text{Sn}(^{12}\text{C}, 4n)^{125}\text{Ba}$ reactions have been studied with Ge(Li) detectors. Beta-gamma delayed coincidences have been measured using NE 102 Ge(Li) detectors. On the basis of gamma intensities and energy sums the decay schemes are proposed for ^{123}Ba and ^{125}Ba .

Preprint. Joint Institute for Nuclear Research.

Dubna, 1974

1. Introduction

During the last few years a considerable interest is observed in the predicted ^{11/} new nuclear region of permanent deformation containing neutron-deficient nuclides with $50 \leq Z \leq 82$. This new region of deformation was confirmed both by experimental data obtained from in-beam spectroscopy ^{12-14/} (even-even nuclei) and by theoretical calculations ^{15, 16/}. The latter suggest that oblate shapes would be rather more stable than the prolate ones. Data of the excited states of odd-A nuclei may serve to test not only the existence of deformation but also to establish the sign of it.

The object of the present work was to obtain further information on the low-lying levels of ¹²³Cs and ¹²⁵Cs excited in the beta decay of ¹²³Ba and ¹²⁵Ba. These isotopes, about whose excited states not very much is known, belong to this new region of interest.

2. Experiment and Results

2.1. Barium-123

The sources used in this experiment were obtained in the ¹¹⁴Sn(¹²C, 3n)¹²³Ba reaction from a target, prepared of commercial separated ¹¹⁴Sn isotope, irradiated for about 1 min. with carbon ions of optimal energy ^{17/} (for this reaction) on the external beam of the heavy ion U-300 cyclotron. The carbon ion energy was reduced to the proper one by means of an aluminium foil set.

The gamma-spectra were studied with a 2.5 cc Ge(Li)-detector with a 0.7 keV resolution at 100 keV. One of these spectra is shown in Fig. 1. The observed gamma-

Fig. 1. Single gamma-ray spectrum of ^{123}Ba observed from a ^{117}Sn target bombarded with ^{12}C ions. Two long-lived gamma-rays marked by asterisk have not been assigned.




Fig. 2. Decay curves of the gamma-rays assigned to ^{123}Ba

rays were identified by comparing their energies and lifetimes with those of known isotopes which could appear in the spectra as well as a decay product of ^{123}Ba and as reaction products with an admixture of other tin isotopes in the target material. Seven of these gamma-rays (Fig. 2) decayed with the same half-life of 2.7 min which one could not assign to any of known isotopes. This value has been adopted as the half-life of ^{123}Ba which is in agreement with the reported ^{8/} one of (2 ± 0.5) min.

The energies of gamma-rays assigned to ^{123}Ba and their relative intensities are given in Table 1.

2.2. Barium-125

The ^{125}Ba sources were obtained with the $^{117}\text{Sn}(C, 4n)^{125}\text{Ba}$ reaction. The experimental procedure used here was the same as in the case of ^{123}Ba .

An example of gamma-ray spectra obtained with the 2.5 cc Ge(Li) detector is shown in Fig. 3. In contrast to ^{123}Ba these spectra are relatively simple. A careful analysis revealed a group of gamma-rays with a half-life of 3.5 min (Fig. 4) which could not be identified with the other known isotopes. To check that these gamma-rays belong to ^{125}Ba , supplementary measurements of gamma-ray spectra were performed. In this case ^{125}Ba sources were obtained by the spallation reaction from CeO_2 and Ta targets irradiated with 660 MeV protons on the external beam of the JINR synchrocyclotron. After the chemical separation the ^{125}Ba source was prepared by electromagnetic isotope separation of the barium fraction using a surface ionization ion source ^{9/}. All the prominent gamma-rays of the group mentioned above were confirmed in the spectra obtained both with the mass-separated ^{125}Ba and with the mass-separated isobars of $A = 125$ obtained from a tantalum target ^{29/} (intense γ -transitions, in the high energy region, were not observed). This fact allows the authors to conclude that the found group of the gamma-transitions, listed in Table 2, belongs to ^{125}Ba .

Table 1
 Gamma-rays observed in the decay of ^{123}Ba . Energy
 uncertainties do not exceed 0.6 keV

Energy							
$E_{\gamma}(\text{keV})$	63.9	92.7	94.5	116.1	120.0	123.5	137.0
Rel. int.	14 \pm 4	51 \pm 5	100	54 \pm 8	27 \pm 4	69 \pm 6	23 \pm 7
I_{γ}							

◀ Fig. 3. Singles gamma-ray spectrum of ^{125}Ba observed from a ^{114}Sn target bombarded with ^{12}C ions.




Fig. 4. Decay curves of the gamma-rays assigned to ^{125}Ba .

Table 2
Gamma-rays observed in the decay of ^{125}Ba . Energy
uncertainties in the present work do not exceed 0.6 keV

Present work		Reference /10/			
$T_{\frac{1}{2}} = (3.5 \pm 0.4) \text{ min}$		$T_{\frac{1}{2}} = (3 \pm 0.5) \text{ min}$		$T_{\frac{1}{2}} = (8 \pm 1) \text{ min}$	
Energy $E_{\gamma}(\text{keV})$	Rel.intensity I_{γ}	Energy $E_{\gamma}(\text{keV})$	Rel.intens. I_{γ}	Energy $E_{\gamma}(\text{keV})$	Rel.intens. I_{γ}
45.0	~ 3			20	
55.6	48 ± 4	56 ± 3	5		
63.1	8 ± 1				
77.6	100	76 ± 2	100		
85.4	82 ± 8	84 ± 2	86		
100.1	6 ± 3				
108.0	8 ± 2				
140.9	86 ± 8	141 ± 2	42		

The value of 3.5 min adopted here as the half-life of ^{125}Ba should be compared with the values of (3 ± 0.5) and (8 ± 1) min published in ref. /10/ as the half-lives of two isomeric states of ^{125}Ba . The first one is in agreement with the value obtained in the present work. The 8 min activity (in the 5-200 keV energy region) in this experiment, was not observed.

3. Decay Schemes and Discussion

3.1. Barium -123

In experiments with heavy ion reactions an isomeric states with half-life of 1.6 sec was revealed in ^{123}Cs . This state is depopulated by two gamma-transitions of energies (63 ± 0.5) and (95.5 ± 0.5) keV of the same type of M1 + E2 multipolarity /11/. A careful analysis of the energy and intensity of the observed gamma- and KX -rays in the decay of ^{123}Ba (Table 1) makes it possible to conclude that very likely the 63.9 keV gamma-transition can be identified with the gamma-ray of energy 63 keV observed in ref. /11/. Basing on this information we assumed that the first excited state of ^{123}Cs is depopulated by this transition. The two additional levels shown in Fig. 5 were included in the decay scheme on the basis of the intensity and energy sum rules only.

The spin and parity of the ground state of the neighbouring odd- A cesium isotopes is $1/2^+$ /12-14/. By analogy to that one can expect that ^{123}Cs nucleus will continue this trend. Such a conclusion seems to be confirmed experimentally by works devoted to the investigations of the decay of ^{123}Cs /15/ and the excited states of ^{123}Xe /16/. However, in our gamma-spectra of ^{123}Ba obtained both in the (H1, xn) reaction and in the spectra obtained with mass-separated isobars of $A=123$ there appeared, beside the strong 97.3 keV gamma-ray, a 83.3 keV gamma-transition which depopulates the second excited state of ^{123}Xe of the $5/2^+$ spin value /16/. Because the assignment of this transition has not yet been established (work continued), the $3/2^+$ value for the ground state of ^{123}Cs cannot be excluded.




Fig. 5. Decay scheme proposed for ^{123}Ba .

Including the present data one can conclude that the isomeric state $^{11/2-}$ is associated with the cascade given on the left in Fig. 5. The spin assignments for the excited states follow from the multipolarity and a weak cross-over transition (unobserved in ref. $^{11/}$).


In the last years an $^{11/2-}$ isomeric state was found in ^{127}Cs $^{18/}$. Such a state could explain the existence of 1.6 sec activity in ^{123}Cs if one assumed the spin sequence given on the left in Fig. 5.

3.2. Barium-125

In the case of ^{125}Ba β^+ -gamma coincidence experiments were carried out using a NE 102 (2.5 x 1 cm) plastic detector as the gating crystal and a 27 cc Ge(Li) one to obtain the coincidence gamma-spectra ^{/19/}. Fig. 6 shows coincidence spectra with an integral gate set above 500 keV. The upper (A) spectrum represents a prompt- and the lower one (B) a 10 ns delayed coincidence spectrum, respectively. From these spectra it is seen that only one of the observed gamma-rays of energy 85.4 keV is delayed relative to the β^+ -decay. This implies the existence of an excited state of the same energy. Furthermore, from the relative intensities of the observed two gamma-rays (77.6 and 85.4 keV) in these spectra the half-life of this level was estimated roughly to be about 20 ns what is in agreement with transitions of the E2 type in the neighbouring odd-mass Cs isotopes ^{/20,23/}. Basing on these facts and taking into consideration the energy and intensity balance, a decay scheme for ^{125}Ba was constructed (Fig. 7).

The spin and parity of the ground state of ^{125}Cs has been found to be $1/2^+$ ^{/13,14/}. According to systematics of the excited states of odd- A cesium nuclei ^{/21-24/} and the present data the spin for the 85.4 keV is very likely to be $5/2^+$. A rough estimation of the log ft values (Q β^+ value was taken from ref. ^{/17/}) for the beta-decay to the proposed levels indicates that these transitions are consistent with an allowed character in this mass region. The transition intensities calculated on the basis of theoretical conversion coefficients ^{/25/} and gamma-ray intensities (assuming M1 character as the lowest of the possible multipolarities for the 55.6 and E2 one for 85.4 keV transitions) indicate that the second excited state, within the statistical error limits, is not populated in the β^+ -decay. From all these facts it follows that the spins of the 140.9 keV levels and of the ground state of ^{125}Ba are likely to be $3/2^+$ and $1/2^+$, respectively.

Taking into account a general trend of the levels given by pairing + quadrupole force model for odd-mass



CHANNEL NUMBER

Fig. 6. Gamma-ray spectra in coincidence with a continuous β^+ -spectrum at the energy > 500 keV. A) the prompt coincidence spectrum, B) 10 ns delayed coincidence spectrum.





Fig. 7. Decay scheme proposed for ^{125}Ba .

cesium isotopes 136 , one can see that this model is able to explain the $1/2^-$ spin of the ground state of 135 Cs. The experimental data of the ground states for odd Cs isotopes are also in good agreement with the theoretical ones 14,17 predicted by the Nilsson model 1,28 .

However, a more detailed calculation 6,9 gives a level sequence, for the few excited states of 135 Cs, far from being in satisfactory agreement with the observed one.

The authors would like to thank Academician G.N.Fleurov for his interest in the works in the mass region and Professor K.Ya.Gromov for valuable comments concerning the manuscript. They wish to thank P.M.Gopytch for his help and G.Beyer, M.Jachym and A.Latuszyński for preparing the mass-separated sources.

References

1. K.Sheline, T.Sikkeland, R.N.Chanda. Phys.Rev.Lett., 7, 446 (1961).
2. J.E.Clarkson, R.M.Diamond, F.S.Stephens, J.Perlman. Nucl.Phys., A93, 272 (1967).
3. D.Ward, R.M.Diamond, F.S.Stephens. Nucl.Phys., A117, 309 (1968).
4. J.Bergstrom, C.J.Herrlander, A.Kerek, A.Luukko. Nucl.Phys., A123, 99 (1969).
5. K.Kumar, M.Baranger. Phys.Rev.Lett., 12, 73 (1964).
6. D.A.Arseniev, A.Sobiczewski, V.G.Soloviev. Nucl.Phys., A126, 15 (1969).
7. W.Neubert. Nucl.Instr. Meth., 93, 473 (1971).
8. Nucl. Data Sheet, B7, No. 2 (1972).
9. G.I.Beyer, E.Herrmann, A.Piotrowski, V.J.Raiko, H.Tyroff. Nucl.Instr.Meth., 96, 437 (1971).
10. J.M.D'Auria, H.Bakru, I.L.Preiss. Phys.Rev., 172, 1176 (1969).
11. Ch.Droste, W.Neubert, S.Chojnacki, T.Morek, K.F.Alexander, Z.Wilhelmi. Nucl.Phys., A192, 595 (1972).
12. W.A.Nierenberg, H.A.Shugart, H.B.Silbsbee, R.J.Sunderland. Phys.Rev., 104, 1380 (1956) and Phys.Rev., 112, 186 (1958).
13. S.Jha, N.F.Peak, W.J.Knox, E.C.May. Phys.Rev., C6, 2196 (1972).
14. O.B.Dabousi, M.H.Prior, H.A.Shugart. Phys.Rev., C3, 1326 (1971).

15. J.M.D'Auria, T.L.Preiss. Nucl.Phys., 84, 37 (1966).
16. A.Kerek, C.J.Herrlander, A.Luukko, M.Grecescu. Phys.Lett., 31B, 10 (1970).
17. N.Zeldes, A.Grill, A.Simievic. Mat.Fys.Skr.Dan. Vid.Selsk., 3, No. 5 (1967).
18. T.W.Conlon. Nucl.Phys., A161, 289 (1971).
19. Yu.K.Akimov, K.Andert, A.I.Kalinin, H.-G.Ortlepp. Nucl.Instr. Meth., 104, 581 (1972).
20. H.Haupt, H.-G.Ortlepp, A.Jasinski. M.Jahim. Prog. Thes. XXIV Conf. on Nucl.Spectr. Nuclear Structure, Kharkov, 1964, 86.
21. K.Ishi, A.Aoki, S.Kageyama. J.Phys.Soc.Japan, 34, 285 (1973).
22. H.W.Taylor, B.Sing, F.S.Prato, J.D.King. Nucl. Phys., A179, 417 (1972).
23. R.Arlt, G.Beyer, H.-G.Ortlepp, H.Tyroff, E.Herrmann, H.Haupt, A.Jasinski. Conf. on Nucl.Spectroscopy Nucl. Theory, p. 98, Dubna, 1973.
24. R.Arlt, G.Beyer, H.Haupt, E.Herrmann, A.Jasinski, G.Musiol, W.Neubert, H.-G.Ortlepp, H.Tyroff. Proc. Int. Conf. Nucl.Phys., Munich, August 27 - September 1, 1973, p. 694.
25. R.S.Hager, E.C.Seltzer. Nucl. Data, A4 (1968).
26. L.S.Kisslinger, R.A.Sorensen. Rev.Modd.Phys., 35, 853 (1963).
27. I.V.Goldstein, A.G. de Pinho. Phys.Rev., C4, 653 (1971).
28. S.G.Nilsson, Kgl. Dansk. Vid. Selsk. Mat. Fys. Medd., 29, No. 16 (1955).
29. A.Latuszynski, K.Zuber, J.Zuber, A.Potempa, W.Żuk. JINR, 6-7469, Dubna, 1973.

Received by Publishing Department
on February 11, 1974.

Subject Categories of the JINR Publications

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids.
15.	Experimental physics of nuclear reactions at low energies.
16.	Health physics. Shieldings

Will you fill blank spaces in your library?

You can receive by post the books listed below. Prices in US \$, including the packing and registered postage

13-3700	Proceedings of the Symposium on Nanosecond Nuclear Electronics. Dubna, 1967.	726 pages	16.70
D-3893	Contributions. International Symposium on Nuclear Structure. Dubna, 1968.	192 pg	6.49
P1-3971	Nucleons and Pions. Proceedings of the First International Colloquium on Nucleon-Nucleon and Pion-Nucleon Interactions. Dubna, 1968.	294 pg	5.63
2-4816	Vector Mesons and Electromagnetic Interactions. Dubna, 1969.	588 pg	9.76
16-4888	Dosimetry of Radiation and Charged Particle Protection Physics. Dubna, 1969.	250 pg	4.85
3-4891	Some Lectures on Neutron Physics. Summer School, Alushta, May, 1969.	428 pg	9.0
D-5805	International Conference on Instrumentation for High Energy Physics. Dubna, 1971, 2 volumes.	882 pg	23.52
D1-5969	Proceedings of the International Symposium on High Energy Physics. Dresden, 1971.	772 pg	13.17
D-6004	Binary Reactions of Hadrons at High Energies. Dubna, 1971.	768 pg	12.59
D10-6142	Proceedings of the International Symposium on Data Handling of Bubble and Spark Chambers. Dubna, 1971.	564 pg	9.96
D13-6210	Proceedings of the 11th International Symposium on Nuclear Electronics. Warsaw, 1971.	372 pg	6.36
D1-6349	Proceedings of the 10th International Conference on High Energy Physics and Nuclear Structure. Dubna, 1971.	670 pg	11.65
D-6465	International School on Nuclear Structure. Alushta, 1972.	525 pg	9.54

D- 6840	Proceedings of the Second International Symposium on High Energy and Elementary Particle Physics (Strbské Pleso, CSSR, 1972).	398 pg.	6.78
P2 - 6867	Lectures on High Energy Physics. (Dubna School, Sukhumi, 1972).	506 pg	8.80
D6-7094	XIII Symposium on Nuclear Spectroscopy and Nuclear Theory. Dubna, 1973.	193 pg	3.71
D2-7161	Proceedings of the III International Seminar on Non-Local Quantum Field Theory. Alushta, 1973	280 pg	5.00
	Proceedings of the VI European Conference on Controlled Fusion and Plasma Physics. Moscow, 1973. v. I.	666 pg	21.00

Orders for the above-mentioned books can be sent at the address:
Publishing Department, JINR
Head Post Office, P.O. Box 79,
101000 Moscow, USSR.

Conditions of Exchange

The preprints and communications of the Joint Institute for Nuclear Research are distributed free of charge on the mutual exchange basis to the universities, institutes, libraries, scientific groups and individual scientists of more than 50 countries.

We expect that the recipients of the JINR publications will have a possibility to display initiative in sending free of charge publications to Dubna. We receive, on the exchange basis, scientific books, journals, preprints and other types of publications on the subject categories of our Institute.

The only kind of publications that we do not require are the reprints of the articles already published in scientific journals.

In a number of cases we address to scientific institutions - the most prominent recipients of our publications - with the request to send us free of charge some books or subscribe for our library to the scientific journals published in their countries.

Requests

The Publishing Department fulfils annually about 3000 individual requests for our preprints and communications. The index of our publication must be obligatory indicated in such requests.

Addresses

Letters on all the questions concerning the exchange of publications as well as requests for individual publications are to be sent at the address:

**Publishing Department
Joint Institute for
Nuclear Research
Head Post Office,
P.O. Box 79
101000 Moscow,
U.S.S.R.**

We kindly ask to send all the publications on the exchange basis and also free of charge subscriptions to scientific journals at the address:

**Scientific-Technical Library
Joint Institute for
Nuclear Research
Head Post Office
P.O. Box 79
101000 Moscow,
U.S.S.R.**

Издательский отдел Объединенного института ядерных исследований.
Заказ 17661. Тираж 454. Уч.-изд. листов 0,95.

Редактор Э.В.Ивашкевич. Подписано к печати 22/III-74 г.



75.04.25