

СООБЩЕНИЯ
ОБЪЕДИНЕННОГО
ИНСТИТУТА
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ

ДУБНА



547703452

E1 - 9368

A21

I.V.Boguslavsky, I.M.Gramenitsky, V.Karimaki,
I.A.Korzhevina, R.Orava, L.A.Tikhonova

**A METHOD FOR ESTIMATING
THE ANNIHILATION CROSS SECTION BASED
ON MEASUREMENT
OF SECONDARY INTERACTIONS**

1975

RANGE OF JINR PUBLICATIONS

The preprints and communications of the Joint Institute for Nuclear Research (JINR) are considered to be original publications. They are published in accordance with Article 4 of the JINR Statut. Difference between the preprints and communications consists in that text of the preprint will be published in future in one of scientific journals or in aperiodic collections.

Indexing

The preprints, communications and deposited publications of the JINR have a single numbering (four last figures of the index).

The first sign of the index - a letter, denotes the language the paper is published in:

"P" - published in Russian;

"E" - published in English;

"D" - published both in Russian and in English.

The figure following the letter denotes the subject category of the given publication. List of subject categories of the JINR publications is sent periodically to the recipients.

Above-described index is placed in the right upper corner of the cover and on the title-page of each publication.

References

In bibliographical references to the JINR preprints and communications we recommend to indicate: author's initials and name, an abbreviation of the name of the institute, index, place and year of publication.

An example of bibliographical reference:

I.Ivanov. JINR, P2-4985, Dubna, 1971.

E1 - 9368

**I.V.Boguslavsky, I.M.Gramenitsky, V.Karimaki,¹
I.A.Korzhevina,² R.Orava,¹ L.A.Tikhonova**

**A METHOD FOR ESTIMATING
THE ANNIHILATION CROSS SECTION BASED
ON MEASUREMENT
OF SECONDARY INTERACTIONS**

¹University of Helsinki, Helsinki, Finland.

²Institute for Nuclear Physics Research of
the Moscow State University, Moscow, USSR.

1. INTRODUCTION

The determination of the annihilation cross section is an important problem in anti-proton-proton experiments (see, e.g.,^{/1/}). At low energies the annihilation and non-annihilation channels can be separated rather well by means of kinematic fitting, whereas at high energies ($P \geq 10$ GeV/c) this method is practically impossible due to an increasing number of neutral and charged secondary particles.

A few attempts to separate annihilation events have been made at higher energies. Everett et al. have described technique^{/2/} which makes use of the charge conjugation symmetry of antiproton-proton interactions.

The method was applied to antiproton-proton experiments at 4.6 and 9.2 GeV/c.

In this article we discuss a completely different approach to the problem. The described method is based on the observation of secondary collisions in primary $\bar{p}p$ -interactions in the bubble chamber. The method is based on the fact that the rate of secondary interactions per event is a function of the average number of nucleons and anti-nucleons per event, and at rather high energies the

N_p and \bar{N}_p cross sections are considerably higher than the πp and Kp ones (see fig. 1).




Fig. 1. Total cross sections of different particles as a function of primary momentum.

Having determined the average multiplicity of nucleons and antinucleons from the number of secondary interactions, one can then calculate the annihilation cross section.

2. OUTLINES OF THE METHOD

For simplicity, consider first a monochromatic beam of antiprotons and π^- -mesons colliding with a hydrogen target. The fraction of antiprotons in the beam is assumed to be a constant value denoted by α . Then the effective interaction cross section of such a mixed beam on hydrogen can be written as

$$\sigma_{\text{eff}} = \alpha \sigma_{\bar{p}p} + (1-\alpha) \sigma_{\pi^-p} = \sigma_{\pi^-p} + \alpha \Delta\sigma, \quad (1)$$

where $\sigma_{\bar{p}p}$ and σ_{π^-p} are the total antiproton and π^- -meson cross sections on hydrogen, respectively, and $\Delta\sigma$ is their difference. Measuring σ_{eff} and knowing the cross sections $\sigma_{\bar{p}p}$ and σ_{π^-p} , one can determine the fraction

$$\alpha = \frac{1}{\Delta\sigma} (\sigma_{\text{eff}} - \sigma_{\pi^-p}). \quad (2)$$

It is obvious that the accuracy of determination of α depends on the difference $\Delta\sigma$. In the extreme case, when $\Delta\sigma = 0$, one cannot at all obtain information on the beam structure in this way.

Equations (1) and (2) can be used to determine the fraction of antiprotons in negative secondary tracks created in $\bar{p}p$ -interactions. Of course, the idea can be also applied to protons, as according to charge conjugation symmetry of $\bar{p}p$ -interactions, the average antiproton multiplicity equals the average proton one. But we restrict ourselves to consideration of only negative tracks as motivated by the fact that the difference $\Delta\sigma$ for proton and π^+ -meson is

smaller than the difference for their anti-particles (see fig. 1).

The value σ_{eff} for secondary negative tracks can be obtained from the observed number of interactions among negative secondaries according to the formula

$$N_s^- = \rho L \sigma_{\text{eff}} N^- = \rho L \sigma_{\text{eff}} \langle N^- \rangle N_{\text{ev}}, \quad (3)$$

where ρ is the number of target protons per cm^3 , L is the potential length of the secondary track (see fig. 2), N^- is the total number of negative secondary particles, N is the total number of primary $\bar{p}p$ -interactions, and $N^- = N^- / N_{\text{ev}}$ is the average number of negative secondaries per event.

From (1) and (3) we readily derive

$$\alpha N^- = \frac{1}{\rho L \Delta \sigma} N_s^- - \frac{\sigma_{\bar{n}p}}{\Delta \sigma} N^- . \quad (4)$$

In the real situation we have not the monochromatic beam of secondary particles, but the momentum spectrum. This is the reason why we have to take into account the fact that the quantities σ_{eff} , L , α in formula (4) depend on the track momentum \vec{p} as well as on the position \vec{r} of the primary interaction vertex. Then we have

$$\alpha(\vec{p}) dN^-(\vec{p}, \vec{r}) = \frac{dN_s^-(\vec{p}, \vec{r})}{\rho L(\vec{p}, \vec{r}) \Delta \sigma(\vec{p})} - \frac{\sigma_{\bar{n}p}(\vec{p})}{\Delta \sigma(\vec{p})} dN^-(\vec{p}, \vec{r}) . \quad (5)$$




Fig. 2. Secondary interactions in the bubble chamber pictures.

Now considering $dN^-(\vec{p}, \vec{r})$ and $dN_s^-(\vec{p}, \vec{r})$ as six-dimensional probability densities normalized to N^- and N_s^- , respectively, we obtain by integrating eq. (5)

$$\langle a \rangle = \frac{N_s^-}{N_{eV} \langle N^- \rangle_\rho} \left\langle \frac{1}{L \Delta \sigma} \right\rangle_s - \left\langle \frac{\sigma \vec{n}^- \cdot \vec{p}}{\Delta \sigma} \right\rangle, \quad (6)$$

where $\langle \rangle_s$ and $\langle \rangle$ mean the averaging over the probability densities $dN_s^-(\vec{p}, \vec{r})$ and

$dN_s^-(\vec{p}, \vec{r})$, respectively. In connection with a finite number of observed events in an experiment, the equation (6) leads to the following simple expression for the mean antiproton multiplicity:

$$\begin{aligned} \langle N_p^- \rangle &= \langle \alpha \rangle \langle N^- \rangle = \\ &= \frac{1}{N_{eV}} \left\{ \frac{1}{\rho} \sum_s [L(\vec{p}, \vec{r}) \Delta\sigma(\vec{p})]^{-1} - \sum_i [\sigma_{\pi p}^-(\vec{p}) / \Delta\sigma(\vec{p})] \right\}, \end{aligned} \quad (7)$$

where \sum_s means a sum taken over the negative tracks with observed secondary interactions and \sum_i a sum over all the negative secondary tracks.

Let us now consider all inelastic channels which make a main contribution to the $\bar{p}p$ - cross section

$$\bar{p}p \longrightarrow \bar{p}p + \dots \quad (8a)$$

$$\longrightarrow \bar{p}n + \dots \quad (8b)$$

$$\longrightarrow \bar{n}n + \dots \quad (8c)$$

$$\longrightarrow \bar{n}p + \dots \quad (8d)$$

$$\longrightarrow m\pi + ck \quad (\text{annihilation})(8e)$$

Having determined $\langle N_p^- \rangle$ for inelastic reaction by the formula (7), one can estimate the sum of the cross sections (8a) and (8b). Thus the upper limit for the annihilation cross section is derived as follows

$$\sigma_{\text{ann}} < \sigma_{\text{tot}}^{\text{in}} - \sigma(8a+8b) = \sigma_{\text{tot}}^{\text{in}} (1 - \langle N_p^- \rangle), \quad (9)$$

The accuracy of the derived cross section σ (8a + 8b) depends on the number of secondary interactions observable in the bubble chamber. This will be discussed in more detail in chapter 3.

In principle, use could be made of the same technique for determining the average multiplicity of neutrons and antineutrons by observing neutral secondary interactions or neutral stars. In the same way, as we derived eq. (7), we obtain

$$\langle N_n^- \rangle = \langle N_n^+ \rangle = \frac{1}{\rho N_{eV}} \frac{\sum_s L(\vec{p}, \vec{r})^{-1}}{\langle \sigma_n^- \rangle + \langle \sigma_n^+ \rangle} \quad (10)$$

where the quantities ρ , N_{eV} and L are defined as in eq. (3) and $\langle \sigma_n^- \rangle$, $\langle \sigma_n^+ \rangle$ are the $\bar{n}p$ and np cross sections averaged over the laboratory momentum spectrum of antineutrons and neutrons, respectively. The annihilation cross section can finally be calculated from the formula

$$\sigma_{ann} = \sigma_{tot}^{in} [1 - (\langle N_n^- \rangle + \langle N_n^+ \rangle)] \quad (11)$$

However, many practical difficulties and complications arise, especially in the application of formula (10). This matter is also considered in the following chapter.

3. DISCUSSION OF THE METHOD

In order to have an idea of the statistical accuracy of formula (7), we consider its application to the $\bar{p}p$ -experiment at 22.4 GeV/c in the "Ludmila" bubble chamber^{/3/}.

In this experiment, typical values for L and $\langle N^- \rangle$ are $L = 50$ cm and $\langle N^- \rangle = 2.5$. For simplicity, we assume $\Delta\sigma = \sigma_{\bar{p}p} - \sigma_{\pi^-p} \approx 20$ mb. Furthermore, assuming a constant relative error for the fraction a , $\delta a = 10\%$, we compute the numbers listed in the table for different a -values.

Table

	$a = 0,3$	$a = 0,2$	$a = 0,1$	$a = 0,05$
1) $\sigma_{\text{eff}}(\text{mb})$	36	34	32	31
2) $\delta\sigma_{\text{eff}}(\%)$	1.7	1.2	0.6	0.3
3) $\delta\sigma$ (8a + 8b) (%)	30	10	3.3	1.4
4) $N_s \cdot 10^3$	3.6	7.0	25.0	31.0
5) $N_e \nu 10^3$	22.0	46.0	172.0	220.0

The statistical accuracy of formula (10) appears to be always better than that of formula (7). This is connected with that neutral pions do not contribute to neutral secondary interactions due to their very short life time. On the other hand, there exist a few practical complications when we want to apply the formula (10). First of all, the momentum spectra of neutrons and antineutrons are unknown, as a rule, and it is necessary to make some assumptions about their forms. Secondly, there may be some admixture of neutrons connected with the primary antiproton beam. Both these facts lead to sizable uncertainty in $\langle N_n^- \rangle$.

Another complication of the method consists in the fact that not all the secondary particles are nucleons and pions. A part of antibaryon - baryon pairs in the final state involves hyperons or antihyperons and a fraction of mesons are kaons. Fortunately, this is not a serious problem, since hyperon production in the cross section is rather small (roughly 10%) and may be neglected in the first approximation. On the other hand, hyperons can be identified by their visible decays and can therefore be taken into account in more accurate calculations.

As for kaons their influence on formula (7) is also negligible because the Kp -cross section is only slightly different from the πp -cross section.

ACKNOWLEDGEMENTS

All the authors thank R.Lednický for helpful advices and two of us (V.Karimaki and R.Orava) would like to thank the colleagues from the Helsinki group for inspiring discussions.

REFERENCES

1. H.I.Miettinen. Proceedings of the Symposium on Antinucleon-Nucleon Interactions, Liblice-Prague, 1974, p. 405.
H.Murhead. Proceedings of the Symposium on Antinucleon-Nucleon Interactions, Liblice-Prague, 1974, p. 488.

2. P.Everett, P.Gregory, P.Grossman. Proceedings of the Symposium on Antinucleon-Nucleon Interactions, Liblice-Prague, 1974, p. 510.
3. L.N.Abesalashvili et al. Phys.Lett., v. 52B, No. 2, 236 (1974).

Received by Publishing Department
on December 9, 1975.



Conditions of Exchange

The preprints and communications of the Joint Institute for Nuclear Research are distributed free of charge on the mutual exchange basis to the universities, institutes, libraries, scientific groups of more than 50 countries.

Besides the regular distribution on the exchange basis, the Publishing Department fulfils annually about 4000 individual requests for our preprints and communications. The index of our publication must be obligatory indicated in such requests.

Addresses

Letters on all the questions concerning the exchange of publications as well as requests for individual publications are to be sent at the address:

*Publishing Department
Joint Institute for
Nuclear Research
Head Post Office,
P.O. Box 79
101000 Moscow,
U.S.S.R.*

We kindly ask to send all the publications on the exchange basis and also free of charge subscriptions to scientific journals at the address:

*Scientific-Technical Library
Joint Institute for
Nuclear Research
Head Post Office
P.O. Box 79
101000 Moscow,
U.S.S.R.*

Издательский отдел Объединенного института ядерных исследований.
Заказ 20747. Тираж 655. Уч.-изд. листов 0,63.

Редактор Э.В.Ивашкевич.

Подписано к печати 25.12.75 г.

Корректор Р.Д.Фомина