

5:17400 386

E7 - 9029

**W.Frank, P.Gippner, K.-H.Kaun, H.Sodan,
Yu.P.Tretyakov**

**QUASIMOLECULAR KX-RAY EXCITATION
BY BOMBARDING La TARGETS
WITH La AND Xe IONS**

E7 - 9029

**W.Frank, P.Gippner, K.-H.Kaun, H.Sodan,
Yu.P.Tretyakov**

**QUASIMOLECULAR KX-RAY EXCITATION
BY BOMBARDING La TARGETS
WITH La AND Xe IONS**

Submitted to *Physics Letters*

Франк В., Гиппнер П., Каун К.-Г., Зодан Х.,
Третьяков Ю.П.

E7 - 9029

Возбуждение квазимолекулярного КX-излучения при облучении
мишени La ионами La и Xe

Изучено рентгеновское излучение при облучении толстых мишеней La ионами Xe с энергией 85-150 МэВ и ионами La с энергией 115 МэВ. Измерены спектры X-лучей и определены их выходы. Обнаружены непрерывные распределения X-лучей с энергиями выше характеристического излучения мишеней и падающих частиц. Высокоэнергетические части сплошных спектров интерпретируются как КX-излучение квазимолекул с эффективными атомными номерами $Z=111$ и $Z=114$ соответственно.

Работа выполнена в Лаборатории ядерных реакций ОИЯИ.

Препринт Объединенного института ядерных исследований
Дубна 1975

Frank W., Gippner P., Kaun K.-H.,
Sodan H., Tret'yakov Yu.P.

E7 - 9029

Quasimolecular KX-Ray Excitation by Bombarding
La Targets with La and Xe Ions

req.
The X-ray emission in the 85-to-150 MeV Xe and 115 MeV La bombardments of thick natural La targets has been measured. The spectra and yields of X-ray emission are obtained. Continuous X-ray distributions have been found to lie beyond the target and projectile characteristic X-ray energies. The high energy parts of these continua are interpreted as K-radiation of quasimolecules with effective atomic numbers $Z = 111$ and $Z = 114$, respectively.

The investigation has been performed at the Laboratory of Nuclear Reactions, JINR.

Preprint of the Joint Institute for Nuclear Research
Dubna 1975

1. Introduction


The possibility of observing new fundamental processes in quantum electrodynamics proposed by Greiner ^{/1/} and Popov ^{/2/}, has motivated a recent search for K α -ray transitions between the transient molecular orbitals (MO) formed during collisions of heavy ions with heavy atoms.

Hitherto, quasimolecular KX-ray emission up to systems Nb + Nb (Z=82) have been investigated ^{/3-7/}. Experiments with high-energy La and Xe ions, described in this paper, first enable one to observe the MO KX-radiation of superheavy systems with Z = 114 and Z = 111, respectively.

2. Experimental Methods and Results

At the U-300 heavy ion cyclotron of the JINR Laboratory of Nuclear Reactions, ¹³⁹La⁸⁺, ¹³²Xe⁸⁺ and ¹³⁶Xe⁹⁺ ions were accelerated to an energy of 115, 120 and 150 MeV, respectively. In a subsequent measurement the energy of the Xe projectiles was decreased from 120 to 85 MeV by means of a 2.15 mg cm⁻² thick Au foil exposed near the reaction chamber.

Thick metallic natural La targets were used. They were irradiated at an angle of 45° with respect to the beam direction, whereas the X-ray emission was measured at 90°. The emitted X-rays were detected by an intrinsic Ge detector with a resolution of about 450 eV at 100 KeV, and two Ge(Li) detectors, whose resolution was better



X-ray spectra measured by bombarding thick natural La targets with (a) 115 MeV $^{139}\text{La}^{8+}$, (b) 120 MeV $^{132}\text{Xe}^{8+}$ and (c) 150 MeV $^{136}\text{Xe}^{9+}$ ions. The dashed lines denote the delayed background.

than 1.5 keV at 100 keV X-ray energy. In order to reduce the high rate of low-energy X-rays, Al foils were inserted between the target and detector. Thus, the coun-

ting rates (max, 60 s^{-1}) were low enough to avoid considerable pile-up contributions.

The beam pulsing of 2 ms beam-on and 2 ms beam-off time offered the possibility of reducing the background by measuring the spectra in a prompt-delayed regime.

The figure shows our experimental spectra without any corrections made. In the *table* the absolute and relative yields of the measured characteristic atomic X-rays and continuous components C1 and C2 are presented. The yields are determined after the subtraction of the background and after the correction made for the detector efficiency.

The figure shows the prompt X-ray spectra obtained by the La^{8+} , Xe^{8+} and Xe^{9+} ion bombardments of thick natural targets. The spectrum (a) was measured in a 8.5 h run using a $2.0 \text{ cm}^3 \text{ Ge(Li)}$ planar detector and a 0.5 mm Al absorber. The ion current amounted to about $0.1 \mu\text{A}$, corresponding to 7.8×10^{10} particles per second. The spectrum (b) was obtained using a 25 cm^3 coaxial Ge(Li) detector and a 0.5 mm Al absorber. The ion current was about $0.4 \mu\text{A}$ corresponding to 3×10^{11} particles per second. Finally, we obtained the spectrum (c) using the same detector and Al absorber, but with an ion current of about $0.1 \mu\text{A}$ which corresponds to 7×10^{10} particles per second.

As the figures show, the low-energy regions of all the spectra contain the characteristic KX-radiation of the projectiles and target materials, while in the region of higher energies two X-ray continua C1 and C2 can clearly be distinguished. The separation of these continua was carried out as shown schematically in the figure (dotted-dashed lines). The background shown in the figure consists of the delayed background measured.

The spectra (b) and (c) contain the fluorescence KX-radiation of Pb, which is produced in a cylindrical lead shielding surrounding the $25 \text{ cm}^3 \text{ Ge(Li)}$ detector. Further, a Coulomb excitation line (C.E.) of ^{139}La and those of the ^{19}F deexcitation are seen in the spectra. Before the yields of the continua C2 were calculated, the spectra were corrected for these effects. The dipole component of nuclear bremsstrahlung, which occurs as a

Table
Characteristic atomic KX-ray and continuous X-ray
yields obtained by bombardment of thick La targets with
La and Xe ions

IONS	E ₀ (MeV)	TAR- GET	ABSOLUTE X-RAY YIELDS PER 10 ⁹ PROJECTILES				RELATIVE YIELDS	
			Y(KX _α -La)	Y(KX _η -La)	Y(KX _α -Xe)	Y(C1)①	Y(C2)②	Y(C2)/Y(KX _α -PROJECTILE)
¹³² Xe 8+	85	La	—	(1,5±0,3) × 10 ¹	(1,1±0,2) × 10 ³	—	—	—
¹³⁹ La 8+	115	La	* (2,9±0,5) × 10 ³	* (3,6±0,6) × 10 ²	—	(5,7±1,7) × 10 ¹	0,4±0,1	(1,4±0,6) × 10 ⁻⁴
¹³² Xe 8+	120	La	—	(2,0±0,5) × 10 ²	(8,5±3,3) × 10 ³	(4,0±1,2) × 10 ¹	0,5±0,2	(5,9±4,1) × 10 ⁻⁵
¹³⁶ Xe 9+	150	La	—	(3,9±0,6) × 10 ²	(1,8±0,6) × 10 ⁴	(7,9±2,4) × 10 ¹	2,5±0,6	(1,4±1,0) × 10 ⁻⁴

* the La KX-yields are divided by 2

① E_x ≥ 40 keV

② E_x ≥ 58 keV

competing effect to the quasimolecular X-ray continuum searched, was calculated^{/8/}. The results of this calculation show that this component can be neglected under the present experimental conditions.

3. Discussion

At the impact energies used the orbital velocities of the K-shell electrons are large compared with the velocity of the incident particles. For example, at the 115 MeV impact energy the orbital velocity of the L_a-K electrons is about 10 times larger than that of the L_a ions. Hence, the spectra can be discussed in the framework of the molecular orbital model. In accordance with our results obtained previously^{/3-5/}, we conclude that the continua C2 for all the colliding systems investigated are mainly formed by KX-radiation of the quasimolecules with the atomic number $Z = Z_1 + Z_2$.

The results (see the table) evaluated from the measured spectra demonstrate that the yields of the characteristic X-rays ($Y(K_{\beta}X-L_a)$, ($Y(K_{\alpha}X$ -projectile)) as well as the continuous quasimolecular yields $Y(C2)$ increase with increasing ion energy, whereas the relative yields $Y(C2) / Y(K_{\alpha}X$ -projectile) remain nearly constant. This behaviour is in agreement with the results of the measurements of Arrabruster et al.^{/9/} and with the calculations of Burch^{/10/}, which show that the cross sections for both characteristic and quasimolecular X-ray excitation increase with increasing incident energies.

As in the X-ray spectra measured in the reactions Ge + Ge and Nb + Nb described elsewhere^{/3,4/}, we found again continuous parts of X-radiation, which we denote C1 (shown in the figure). These continua C1 lie beyond the energies of the target and projectile characteristic X-rays. Their origin is still an open question. Electron bremsstrahlung cannot account for the continua C1, because the calculated yield integrals are four orders of magnitude smaller than the values measured^{/8/}. Further, the models of Briggs and Macek^{/11/} and Müller^{/12/} pre-

dict for heavy ion collisions quasimolecular X-ray continua, described by a static approximation, with an endpoint energy near the united atomic limit and a smearing of the distribution beyond this "classical" limit, caused by dynamic effects. Generally, the continua C1 measured in our experiments look like the "static" distributions (in particular, they have shoulders as predicted in ref. ¹¹) but their end energies lie much lower than the united atomic limit.

In order to study the shape of the X-ray continua in dependence of the united atomic number $Z = Z_1 + Z_2$, we have recently carried out measurements of the colliding systems Ni + Ni ($Z = 56$)¹³. The X-ray continuum obtained extends beyond the united atomic limit but it does not permit the separation into two continuous parts either before or after making correction for the detector efficiency. On the other hand, in the Ge + Ge ($Z=64$) colliding system separation can be done after this correction³ and in the heavier systems such as Nb+Nb ($Z = 82$)⁴ and La + Xe ($Z = 111$) (this work) two continua can clearly be seen. So we conclude that the formation of two separated X-ray continua is mainly caused by the increasing atomic number Z of the colliding system. In the future, investigations of the angular asymmetry will be performed to obtain further information about the origin and character of these continua C1.

The authors wish to express their gratitude to Academician G.N.Flerov for his interest in and support of the work, and to the cyclotron staff for their fruitful cooperation.

References

1. B.Müller, J.Rafelski, W.Greiner. *Z.Phys.*, 257, 183 (1972).
B.Müller, J.Rafelski, W.Greiner. *Phys.Lett.*, 47B, 5 (1973).
2. V.S.Popov. *Sov. Phys. JETP*, 32, 526 (1971).
3. P.Gippner, K.-H.Kaun, F.Stary, W.Schulze, Yu.P.Tretyakov. *Nucl.Phys.*, A230, 509 (1974).

4. P.Gippner, K.-H.Kaun, H.Sodan, F.Stary, W.Schulze, Yu.P.Tretyakov. *Phys.Lett.*, 52B, 183 (1974).
5. W.Frank, P.Gippner, K.-H.Kaun, H.Sodan, W.Schulze, Yu.P.Tretyakov. *JINR, E7-8616, Dubna, 1975.*
6. W.E.Meyerhof, T.K.Saylor, S.M.Lazarus, W.A.Little, B.B.Triplett. *Phys.Rev.Lett.*, 30, 1279 (1973).
W.E.Meyerhof, T.K.Saylor, S.M.Lazarus, W.A.Little, B.B.Triplett. *Phys.Rev.Lett.*, 32, 1279 (1974).
7. J.S.Greenberg, C.K.Davis, P.Vincent. *Phys.Rev.Lett.*, 33, 473 (1974).
8. P.Gippner. *JINR E7-8843, Dubna, 1975.*
9. P.Armbruster, G.Kraft, P.Mokler, B.Fricke, H.J.Stein. *Proc. on the NSSE, June 11 (1974), Ronneby, Sweden.*
10. D.Burch. *VIII ICPEAC, Invited Lectures and Progress Reports, Belgrade (1973).*
11. J.S.Briggs. *J.Phys. B: Atom.Molec. Phys.*, vol. 7, 1, 47 (1974).
J.H.Macek, J.S.Briggs. *J.Phys. B: Atom.Molec. Phys.*, vol. 7, 11, 1312 (1974).
12. B.Müller, W.Greiner. *Phys.Rev.Lett.*, 33, 469 (1974).
13. W.Frank, P.Gippner, K.-H.Kaun, P.Manfrass (to be published).

*Received by Publishing Department
on June 27, 1975.*

Will you fill blank spaces in your library?

You can receive by post the books listed below. Prices - in US \$,
including the packing and registered postage

D1-5969	Proceedings of the International Symposium on High Energy Physics. Dresden, 1971.	773pg	10.00
D-6004	Binary Reactions of Hadrons at High Energies. Dubna, 1971.	768 pg	12.59
D10-6142	Proceedings of the International Symposium on Data Handling of Bubble and Spark Chambers. Dubna, 1971.	564 pg	9.96
D13-6210	Proceedings of the Vth International Symposium on Nuclear Electronics. Warsaw, 1971.	372 pg	6.36
D1-6349	Proceedings of the IVth International Conference on High Energy Physics and Nuclear Structure. Dubna, 1971.	670pg	9.92
D-6465	International School on Nuclear Structure. Alushta, 1972.	525 pg	9.54
D- 6840	Proceedings of the Second International Symposium on High Energy and Elementary Particle Physics (Strbske Pleso, CSSR, 1972).	398 pg	6.78
D2-7161	Proceedings of the III International Seminar on Non-Local Quantum Field Theory. Alushta, 1973.	280 pg	5.00
D1.2-7411	Deep Inelastic and Many-Body Processes. Dubna, 1973.	507 pg	8.76
D13-7616	Proceedings of the VII International Symposium on Nuclear Electronics. Budapest, 1973.	372 pg	5.62
P1.2-7642	Proceedings of the International School of Young Physicists on High Energy Physics. Gomel, 1973.	623 pg	10.53
D10-7707	Meeting on Programming and Mathematical Methods for Solving the Physical Problems. Dubna, 1973.	564 pg	8.23

D1,2-7781	Proceedings of the III International Symposium on High Energy and Elementary Particle Physics. Sinaia, Romania, 1973.	478pg 7.29
D3-7991	Proceedings of the II International School on Neutron Physics. Alushta, 1974.	554pg 4.35
D1,2-8405	Proceedings of the IV International Symposium on High Energy and Elementary Particle Physics. Varna, 1974.	376pg 3.89
D10,11-8450	International School on Use of Computers in Nuclear Research. Tashkent, 1974.	465 pg 4.52
P1,2-8529	The International School-Seminar of Young Scientists. Actual Problems of Elementary Particle Physics. Sochi, 1974.	582 pg 4.70
	Proceedings of the VI European Conference on Controlled Fusion and Plasma Physics. Moscow, 1973. v.I.	666 pg 21.00
	Proceedings of the VI European Conference on Controlled Fusion and Plasma Physics. Moscow, 1973. v.II.	466 pg 15.00

**Orders for the above-mentioned books can be sent at the address:
Publishing Department, JINR
Head Post Office, P.O. Box 79
101000 Moscow, USSR.**

Subject Categories of the JINR Publications

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Gryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids.
15.	Experimental physics of nuclear reactions at low energies.
16.	Health physics. Shieldings
17.	Theory of condensed matter.



Издательский отдел Объединенного института ядерных исследований.
Заказ 20037. Тираж 410. Уч.-изд. листов 0,5.
Редактор Э.В. Пвашкевич. Подписано к печати 30.07.75.
Корректор Р.Д. Фомина