


SM 8006805 ✓



ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА

E2 - 12885

A.V.Sidorov, N.B.Skachkov

DESCRIPTION OF WIDTH
AND SPECTRA
OF TWO RELATIVISTIC FERMIONS
BOUND STATES

1979

E2 - 12885

A.V.Sidorov, N.B.Skachkov

**DESCRIPTION OF WIDTH
AND SPECTRA
OF TWO RELATIVISTIC FERMIONS
BOUND STATES**

Submitted to TMΦ

Сидоров А.В., Скачков Н.Б.

E2 - 12885

Описание ширины и спектров связанных состояний
двух релятивистских фермионов

Целью работы является построение формализма для релятивистского описания системы двух частиц со спином $1/2$. Используется двухчастичное трехмерное ковариантное уравнение, полученное в квазипотенциальном подходе. Квазипотенциальное уравнение в релятивистском конфигурационном представлении сведено в приближении ОБЕР к системе разностных парциальных уравнений, которая является релятивистским аналогом нерелятивистской системы Хамеды-Джонстона. Методом ВКБ решена задача о спектре масс связанных состояний. С использованием точного релятивистского кулоновского решения двухчастичной задачи получены выражения для ширины лептонных распадов в кварковой модели.

Работа выполнена в Лаборатории теоретической физики, ОИЯИ.

Препринт Объединенного института ядерных исследований, Дубна 1979

Sidorov A.V., Skachkov N.B.

E2 - 12885

Descriptions of Width and Spectra of Two
Relativistic Fermions Bound States

The aim of this article is the construction of the formalism for the relativistic description of two particles with spin $1/2$. We use the two-particle three-dimensional equation, obtained by quasipotential approach. Quasipotential equation in the relativistic configurational space with OBEP potential is reduced to the system of partial equations which is the analog of nonrelativistic Hamada-Jonston system. WKB approach is used to calculate mass spectra and leptonic width of mesons in quark model.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna 1979

I. Introduction

The present paper may be treated as a sequel to papers^{1-3/} on the three-dimensional relativistic description of two-particle system with spin 1/2 in the quantum field theory. Principal formulae, allowing the practical application of our mathematical technique are obtained herein. It will be applicable to such attractive problems as the calculation of levels and width of systems like positronium, muon and baryon as well as bound states in a quark - antiquark system, that is old ρ, ω, φ and new $\psi/\psi, \gamma$ -mesons. Lately a nonrelativistic quark model based on the Schrödinger equation is widely used for the mass spectrum description of hard vector mesons of the ψ/ψ -particle type. For this purpose a confining potential is chosen and added by the Breit-Fermi potential accounting for spin effects.

$$V(r) = V_{\text{conf}}(r) + V_{\text{spin}}(r). \quad (1.1)$$

The Breit-Fermi potential is known to represent v^2/c^2 - expansion of Feynman matrix element of one-boson (OBE) or one-gluon exchange. However, such a quasirelativistic approach applied to ordinary light vector ρ, ω, φ mesons is not self-consistent. It was shown in paper^{4/} that in this case the contribution of relativistic corrections appears to be of the same order as the contribution of the nonrelativistic Hamiltonian,

which was initially taken as the principal one. Therefore an essentially relativistic approach is needed here.

In the present paper for the mass spectrum description of a system consisting of two fermions (for example, quark and antiquark), we use the equation of quasipotential approach ^{15/}. As in previous papers we use the relativistic three-dimensional two-particle quasipotential equation, obtained by Kadyshevsky, on the basis of the covariant Hamiltonian formulation of field theory and later in papers ^{16/} on the basis of the covariant technique of equating times in the two-particle wave function of Bethe-Salpiter equation. In the case of equal masses $m_1 = m_2 = m$ this equation, written in the centre-of-mass-system ($\vec{p}_1 = -\vec{p}_2 = \vec{p}$) has the form

$$(2E_q - 2E_p) \Psi_{g_1 g_2}(\vec{p}) = (2\pi)^{-3} \int \frac{d\vec{k}}{E_k} \sum_{g'_1 g'_2} V(\vec{p}, \vec{k}; E_q) \Psi_{g'_1 g'_2}(\vec{k}) \quad (1.2)$$

Now consider in more detail the part of quasipotential (1.1), containing spin effects. In the case of vector-boson exchange the quasipotential in r.h.s. of equation (1.1) represents the Feynman matrix element ^{11/}

$$\langle \vec{p}_1, \vec{g}_1; -\vec{p}_2, \vec{g}_2 | T_V^{(2)} | \vec{k}_1, \vec{g}_1; -\vec{k}_2, \vec{g}_2 \rangle = g_V^2 \frac{\bar{u}(\vec{p}) \gamma_\mu u(\vec{k}) \bar{u}(\vec{k}) \gamma_\mu u(-\vec{p})}{\mu^2 - (\vec{p} - \vec{k})^2} = \sum_{g_1 p, g_2 p} \xi_{g_1 p}^* \xi_{g_2 p}^* T_V^{(2)}(\vec{k}(-); \vec{p}; \vec{p}) \xi_{g_1 p} \xi_{g_2 p} \quad (1.3)$$

μ is the mass of a boson, ξ^G are two-component spinors.

Applying to (1.3) the suggested transformation ^{11/} from the four-dimensional representation to the three-dimension one ¹⁾ and separating the Wigner matrix of spin rotation, we obtain:

¹⁾ This transformation (1) plays the same role as the Foldy-Wenthuysen one ^{18/}, allowing one to pass to the three-dimensional description of the spin. However, in contrast to (8) the discussed in (1) transformation does not deal with the expansion of interaction terms in powers of v^2/c^2 .

$$\begin{aligned}
T_V^{(2)}(\vec{k}(-)\vec{p}; \vec{p}) = & -\frac{gV^2}{\mu^2 + 4\vec{\alpha}^2} \left[4m^2 + 4(\vec{G}_1 \vec{\alpha})(\vec{G}_2 \vec{\alpha}) - (\vec{G}_1 \vec{G}_2) \vec{\alpha}^2 + \right. \\
& + 8\rho_0 \alpha_0 i(\vec{G}_1 + \vec{G}_2)[\vec{p} \times \vec{\alpha}] + \frac{g}{m^2} (\rho_0^2 \alpha_0^2 + 2\rho_0 \alpha_0 (\vec{p} \vec{\alpha}) - m^4) + \\
& \left. + \frac{g}{m^2} (\vec{G}_1 \vec{p})(\vec{G}_2 \vec{\alpha})(\vec{G}_2 \vec{p})(\vec{G}_1 \vec{\alpha}) \right]. \quad (1.4)
\end{aligned}$$

Here the quantity $\vec{\alpha}$ means a half-transfer of momentum, that can be expressed through the momentum transfer in the Lobachevsky space introduced into \mathcal{M} :

$$\vec{\Delta} = \vec{k}(-)\vec{p} = \Lambda_p^{-1} \vec{k} = \vec{k} - \frac{\vec{p}}{m} \left(k_0 - \frac{\vec{k} \vec{p}}{\rho_0 + m} \right) \quad (1.5)$$

by the formula $\vec{\alpha} = \vec{\Delta} \sqrt{m/2(\Delta_0 + m)}$ \mathcal{M} .

Expression (1.4) has a form of the geometrical generalization of the Breit-Fermi potential taken in the momentum representation.

It is convenient to solve equation (1.3) by transformation to the relativistic configurational representation (RCR), introduced initially in \mathcal{M} . To obtain this, instead of the Fourier-transformation there is used an expansion over the complete and orthogonal system of functions in the Lobachevsky space^{9/}.

$$\begin{aligned}
\xi(\vec{p}; \vec{n}, r) = & \left(\frac{\rho_0 - \vec{p} \vec{n}}{m} \right)^{-1 - i r m} \\
\rho_0 = & \sqrt{m^2 + \vec{p}^2}, \quad \vec{n}^2 = 1. \quad (1.6)
\end{aligned}$$

For these functions in papers ^{16,12/} we found an expansion over spherical harmonics:

$$\begin{aligned}
\xi(\vec{p}; \vec{n}, r) = & \sqrt{\frac{\pi}{2 \operatorname{sh} x}} \sum_{\ell=0}^{\infty} (2\ell+1) \frac{\Gamma(i\frac{r}{x} + \ell + 1)}{\Gamma(i\frac{r}{x} + 1)} P_{\ell}^{-i\frac{r}{x} - \ell}(ch x) P_{\ell} \left(\frac{\vec{p} \vec{n}}{\rho} \right) = \\
= & \sum_{\ell=0}^{\infty} (2\ell+1) i^{\ell} P_{\ell}(ch x, r) P_{\ell} \left(\frac{\vec{p} \vec{n}}{\rho} \right) \quad (1.7)
\end{aligned}$$

as well as the condition of completeness and orthogonality for radial parts

$$\frac{2shxshx'}{\pi\lambda^3} \int_0^\infty r^2 dr \rho_e(chx, r) \rho_e^*(chx', r) = \delta(x-x') \quad (1.8)$$

$$\frac{2r'r'}{\pi\lambda^3} \int_0^\infty sh^2x dx \rho_e(chx, r) \rho_e^*(chx, r') = \delta(r-r').$$

In [2,3] there was discussed a problem on the form of the second and third terms (1.4) in the relativistic configurational representation. There are transforms of all the five spin structures present in (1.4) in this paper.

Equation (1.2) transformed into RCR is a finite-difference one and may be reduced to the system of partial equations, being a relativistic analog of the nonrelativistic Hamada-Jonston system [11].

To solve partial finite-difference equations we use the suggested analog [12] of WKB method for RCR.

Calculation of the electromagnetic decay width of mesons is another important problem. To determine lepton width of vector mesons we use the V.Royen-Weisscopff formula (with account of colour).

$$\Gamma_{V \rightarrow e^+e^-} = 16\pi\alpha^2 M_V^{-2} e_q^2 |\Psi_{ne}(0)|^2, \quad (1.9)$$

and for the two-photon decay of pseudoscalar mesons we use formula [14]

$$\Gamma_{P \rightarrow \gamma\gamma} = 12\pi\alpha^2 m^{-2} e_q^4 |\Psi_{ne}(0)|^2. \quad (1.10)$$

In these formulae M_V is the mass of a meson; m and e_q are the mass and charge of quarks, and $\Psi_{ne}(0)$ is WF of system of a quark and anti-quark. Usually instead of this WF the nonrelativistic WF is used in the Coulomb field neglecting the rest of the spin part of the OBEF. The width of states with $\ell \neq 0$ in terms of the relativistic theory is known to turn into zero due to the behaviour of the WF at the origin of coordinates as $\Psi_e^{nonrel} \xrightarrow{r \rightarrow 0} r^e$. This behaviour is of a kinematical character and is connected with the behaviour of free solutions $j_\ell(xr) \rightarrow (xr)^\ell$ for $x \rightarrow 0$. In a relativistic equation there is used another

complete system of functions (1.6) having an essentially different behaviour as $\Gamma \rightarrow 0$, so that $\psi_e^{r+0} \neq 0$ at $e \neq 0$.

In section 2 a quasipotential equation will be transformed to the system of partial finite-difference equations, in the relativistic configurational representation; section 3 is devoted to the solution of unbounded equations of the system through the WKB method; section 4 gives formulae for the width of vector meson decays and pseudoscalar mesons, using the exact Coulomb solution of the quasipotential equation at $\Gamma = 0$.

2. System of Partial Two-Particle Equations in the Relativistic Configurational Representation

For the quasipotential and WF the transformation into the relativistic configurational representation is performed by formulae:

$$V(\Gamma, \vec{n}; \vec{p}) = (2\pi)^{-3} \int d\Omega_\Delta \xi^*(\vec{\Delta}; \vec{\tau}) V(\vec{\Delta}; \vec{p}), \quad (2.1)$$

$$\Psi_{G_1 G_2}(r) = (2\pi)^{-3} \int d\Omega_p \xi(\vec{p}; \vec{n}, r) \Psi_{G_1 G_2}(\vec{p}). \quad (2.2)$$

In ^{15/} the physical meaning of the parameter r present in the function is discussed in detail.

Applying (2.1) to (1.4) we obtain:

$$V(\Gamma, \vec{n}; \vec{p}) = V_1(\Gamma, p_0) + V_2(\Gamma, \vec{n}; p_0, \vec{p}) \quad (2.3)$$

$$V_1(\Gamma, p_0) = -g_V^2 \left\{ (2p_0^2 - 1) \left[\frac{V(r)}{k_{VK}} + \frac{16\pi}{r} \frac{\delta'(r^2 - 1)}{r} \delta'(\vec{n}) \right] + \left[\frac{M^2}{3} V_{kVK}(r) - \frac{8\pi}{3} \frac{\delta'(r^2 - 1)}{r} \delta'(\vec{n}) \right] \cdot [(\vec{G}_1 \vec{G}_2) - (\vec{S} \vec{p})^2 + \vec{p}^2 \vec{S}^2] \right\}$$

$$V_2(\Gamma, \vec{n}; p_0, \vec{p}) = -g_V^2 \left\{ (\vec{S} \vec{L}) \frac{4}{r} [p_0 A(r) + (\vec{p} \vec{n}) \varphi(r)] - 4i p_0 (\vec{p} \vec{n}) A(r) + [\vec{S}_{1,2} - (\vec{S} \vec{L})^2 \frac{4}{r^2}] \varphi(r) \right\}. \quad (2.5)$$

The following notations are introduced here:

$$\vec{S} = \frac{\vec{G}_1 + \vec{G}_2}{2}; \vec{L} = [\vec{p} \times \vec{r}]; S'_{12} = 3(\vec{G}_1 \vec{n})(\vec{G}_2 \vec{n}) - (\vec{G}_1 \vec{G}_2). \quad (2.6)$$

$$A(r) = \frac{1}{r(r-i)} (r \operatorname{th} \alpha \sin \alpha + \cos \alpha) V_{\text{Yuk}}(r) \quad (2.7)$$

$$\varphi(r) = \frac{r^2}{3(r-i)(r+i)} \left[\mu^2 + 3 \frac{\mu}{r} \left(1 - \frac{\mu^2}{2}\right) \frac{\operatorname{th} \alpha}{\sqrt{1 - \frac{\mu^2}{4}}} + \right. \\ \left. + \frac{3 - 2\mu^2 \left(1 - \frac{\mu^2}{4}\right) - \frac{3}{2} \operatorname{ch} \alpha}{1 - \frac{\mu^2}{4}} \right] V_{\text{Yuk}}(r) \quad (2.8)$$

$V_{\text{Yuk}}(r)$ - the Yukawa potential transform: $4/(\mu^2 + 4\vec{x}^2)$

$$V_{\text{Yuk}}(r) = \frac{1}{4\pi r} \frac{\operatorname{ch} \alpha}{\operatorname{sh} \alpha} \frac{1}{r}, \alpha = \arccos \frac{\mu^2 - 2}{\mu^2} \quad (\mu^2 < 4). \quad (2.9)$$

In formulae (2.3)-(2.9) we assume the mass of interacting particles m to be equal to 1.

Formula (2.3) is splitted into two parts $V_1(r, p_0)$ independent of the single vector "relativistic coordinate" direction \vec{n} and $V_2(r, \vec{n}; p_0, \vec{p})$ containing the dependence on \vec{n} through expressions of the type

$$(\vec{p} \vec{n}), [\vec{p} \times \vec{n}], S'_{12}. \quad (2.10)$$

The necessity of such a splitting becomes obvious in the course of the transformation of the complete equation (1.2) into RCR using formulae (2.1), (2.2).

$$(2E_1 - 2\hat{H}) \Psi_{G_1 G_2}(\vec{r}) = \int d\Omega_p d\Omega_k d\vec{r}_1 \xi(\vec{p}; \vec{n}, \kappa) \times \\ \times \xi^*(\vec{k}; \vec{n}_k, \kappa) \sum_{G_1' G_2'} V_{(G_1, \vec{p}, E)}^{G_1' G_2'} \Psi_{G_2' G_1'}(\vec{r}_1) \quad (2.11)$$

$$\hat{H} = \operatorname{ch} \left(i \frac{\lambda}{r} \frac{d}{dr} \right) + i \frac{\lambda}{r} \operatorname{sh} \left(i \frac{\lambda}{r} \frac{d}{dr} \right) + \frac{\Delta \theta \theta}{(r/\lambda)^2} \exp \left(i \frac{\lambda}{r} \frac{d}{dr} \right) \quad (2.12)$$

$\Delta_{\theta\phi}$ - angular part of the Laplas operator, $\lambda = \frac{\hbar}{mc}$. Due to the character of the addition theorem of the relativistic plane waves [16]:

$$\xi(\vec{k}; \vec{n}, r) = \xi(\vec{k}(-)\vec{p}; \vec{n}_{\Lambda p}, z) \xi(\vec{p}; \vec{n}, r) \quad (2.13)$$

$$\vec{n}_{\Lambda p} = [m\vec{n} - \vec{p}(1 - \frac{\vec{p}\vec{n}}{\rho_0 + m})] / (\rho_0 - \vec{p}\vec{n}) \quad (2.14)$$

$V_2(r, \vec{n}; \rho_0, \vec{p})$ enters into equation (2.12) in a nonlocal way:

$$(2E_9 - 2A) \Psi_{\vec{G}_1 \vec{G}_2}(\vec{r}) = \sum_{\vec{G}'_1 \vec{G}'_2} V_2(\vec{r}, \rho) \Psi_{\vec{G}'_1 \vec{G}'_2}(\vec{r}) + \quad (2.15)$$

$$+ \int d\vec{r}_1 \sum_{\vec{G}'_1 \vec{G}'_2} \int d\Omega_p \xi(\vec{p}; \vec{n}, r) \xi^*(\vec{p}; \vec{n}_1, z_1) V_{\vec{G}'_1 \vec{G}'_2}(\vec{r}_1, \vec{n}_{\Lambda p}; \vec{p}) \Psi_{\vec{G}'_1 \vec{G}'_2}(\vec{r}_1)$$

since substituting it into the equation the dependence on vector \vec{n}_1 turns to the dependence on vector $\vec{n}_{\Lambda p}$. The vector rotation $\vec{n}_1 \rightarrow \vec{n}_{\Lambda p}$ taken into account gives the following dependence of expressions (2.10) on the vector \vec{p} and ρ_0 :

$$(\vec{p} \vec{n}_{\Lambda p}) = m^2 / (\rho_0 - \vec{p}\vec{n}_1) - \rho_0 \quad (2.16)$$

$$[\vec{p} \vec{n}_{\Lambda p}] = m [\vec{p} \times \vec{n}_1] / (\rho_0 - \vec{p}\vec{n}_1) \quad (2.17)$$

$$(\vec{G}_1 \vec{n}_{\Lambda p})(\vec{G}_2 \vec{n}_{\Lambda p}) = Z_1^T + Z_2^T \quad (2.18)$$

$$Z_1^T = m^2 (\vec{G}_1 \vec{n}_1)(\vec{G}_2 \vec{n}_1) / (\rho_0 - \vec{p}\vec{n}_1)^2$$

$$Z_2^T = \frac{m^2}{(\rho_0 - \vec{p}\vec{n}_1)^2} \left\{ -\frac{1}{m} [(\vec{G}_1 \vec{n}_1)(\vec{G}_2 \vec{p}) + (\vec{G}_2 \vec{p})(\vec{G}_1 \vec{n}_1)] (1 - \frac{\vec{p}\vec{n}_1}{\rho_0 + m}) + \right. \\ \left. + \frac{1}{m^2} (\vec{G}_2 \vec{p})(\vec{G}_1 \vec{p}) (1 - \frac{\vec{p}\vec{n}_1}{\rho_0 + m})^2 \right\}$$

However, with the help of the correlation:

$$\exp\left(i\frac{1}{m}\frac{d}{dr_2}\right)\xi^*(\vec{p}; \vec{n}_2, r_2) = \frac{m}{p_0 - \vec{p}\vec{n}_2}\xi^*(\vec{p}; \vec{n}_2, r_2) \quad (2.19)$$

expressions (2.16), (2.17) and the first term Z_1^T from (2.18) can be localized. Z_1^T is a relativistic generalization of the expression $(\vec{\sigma}_1 \vec{n})(\vec{\sigma}_2 \vec{n})$ present in the operator of tensor forces. Confining ourselves to Z_1^T in accounting tensor forces we obtain the r.h.s. of equation (2.15) to be of a local form:

$$(2E - 2\hat{H})\Psi_{\vec{\sigma}_1 \vec{\sigma}_2}(\vec{r}) = \sum_{\vec{\sigma}'_1 \vec{\sigma}'_2} \hat{V}(\vec{r}, \vec{p}) \Psi_{\vec{\sigma}'_1 \vec{\sigma}'_2}(\vec{r}) \quad (2.20)$$

The rotation $\vec{n} \rightarrow \vec{n}_{1p}$ in tensor forces taken into account gives an additional spin-spin interaction

$$\begin{aligned} & \varphi(r) [3(\vec{\sigma}_1 \vec{n}_{1p})(\vec{\sigma}_2 \vec{n}_{1p}) - (\vec{\sigma}_1 \vec{\sigma}_2)] \rightarrow \\ & \rightarrow \varphi(r-2i\lambda) S_{12} + [\varphi(r-2i\lambda) - \varphi(r)] (\vec{\sigma}_1 \vec{\sigma}_2), \end{aligned} \quad (2.21)$$

An angular and spin dependences of equation potential (2.20) are determined by five structures, having the same form as in the nonrelativistic theory:

$$\hat{V}(\vec{r}, \vec{p}) = V_S + V_T S_{12} + V_{LS} (\vec{L} \vec{S}) + V_G (\vec{\sigma}_1 \vec{\sigma}_2) + V_{(LS)} (\vec{S} \vec{L})^2 \quad (2.22)$$

Functions $V_i(\vec{r})$ ($i = S, S_{12}, LS, G, (LS)^2$) depend on vector modulus \vec{r} and difference operators \vec{p} and $\vec{p}_0 = \hat{A}$ in the following way ($\lambda=1$):

$$\begin{aligned} V_S = & (\lambda p_0^2 - 1) \left[V_{YUK} (r) + \frac{16\sqrt{\pi}}{r} \mathcal{S}(r^2) - \right. \\ & \left. - 4i f_0 \left[\left(\frac{r-i}{r}\right)^2 \exp(-i\frac{d}{dr}) - p_0 \right] A(r) \right] \end{aligned} \quad (2.23)$$

$$V_T = \left(\frac{r-2i}{r}\right)^2 \exp(-2i\frac{d}{dr}) \quad (2.24)$$

$$V_{LS} = 4P_0 \frac{r}{(r+i)^2} A(r) + \left[\frac{r(r-i)}{(r+i)^2} \exp(-i \frac{d}{dr}) - P_0 \right] \frac{r}{(r+i)^2} \varphi(r) \quad (2.25)$$

$$V_G = \left[\left(\frac{r-2i}{r} \right)^2 \exp(-2i \frac{d}{dr}) - 1 \right] \varphi(r) - \frac{\mu^2}{3} V_{YUK}(r) - \frac{g}{3} \frac{\hat{d}}{r} \delta(r+1) \quad (2.26)$$

$$V_{(LS)^2} = - \frac{4r}{(r+i)(r+2i)} \varphi(r) \quad (2.27)$$

We introduce a radial function $R_{e's'e_s}^j(r)$

$$\Psi_{g, G_1, G_2}^S(\vec{r}) = \sum_{j e' e' e_s} R_{e's'e_s}^j(r) \left\{ \Omega_{j e' e'}^{*S}(\vec{n}) \right\} \left\{ \Omega_{j e' e'}^S(\vec{n}) \right\}_{G_1 G_2} \quad (2.28)$$

The substitution of (2.28) into equation (2.27) gives us a system of four partial equations. Two of the system of equations, obeying the case of $S=0$, $e'_i = i$ and $S=1$, $e'_i = j$ are uncoupled:

$$\left[2\hat{H}_e - 2E_{g,e} + V_{e'_i s}^{-1}(r) \right] R_{e'_i s}^S(r) = 0, \quad e'_i = j, \quad S=0, 1. \quad (2.29)$$

$$V_{e'_i s}^{S=0}(r) = -g_V^2 (V_S - 3V_G) \quad (2.30)$$

$$V_{e'_i s}^{S=1}(r) = -g_V^2 (V_S + 2V_T + V_G - V_{LS} + V_{(LS)^2}) \quad (2.31)$$

Hamiltonian \hat{H}_e is obtained from expression (2.12) for the \hat{H} through the substitution of the operator $\Delta_{\Theta\Phi}$ for its eigenvalue $l(l+1)$. The other two WF describing states with a spin $S=1$ are defined by the system of two coupled equations

$$\begin{cases} (2E_q - 2H_{e'=j+1}) R_{e'=j+1}(r) = V(r) R(r) + \tilde{V}_T(r) R_{e'=j-1}(r) \\ (2E_q - 2H_{e'=j-1}) R_{e'=j-1}(r) = V(r) R(r) + \tilde{V}_T(r) R_{e'=j+1}(r) \end{cases} \quad (2.32)$$

$$V_{e'=j+1}(r) = g_V^2 [V_S - (j+2) V_{LS} + (j+2)^2 V_{LS}^2 - \frac{2(j+1)}{2j+1} V_T + V_G] \quad (2.33)$$

$$V_{e'=j-1}(r) = g_V^2 [V_S + (j-1) V_{LS} + (j-1)^2 V_{LS}^2 - \frac{2(j-1)}{2j+1} V_T + V_G] \quad (2.34)$$

$$\tilde{V}_T(r) = -g_V^2 \frac{6\sqrt{j(j+1)}}{2j+1} V_T(r) \quad (2.35)$$

Writing down system (2.29), (2.32) we follow the approximation on the phenomenological use of the OBEP, accepted in paper /11/ we neglect the \vec{P} quadratic terms in the spin-spin interaction $\vec{P}^2 \vec{S}^2 - (\vec{P} \cdot \vec{S})^2$ and confine ourselves to the account of the terms proportional to $m^2 (\vec{G}_1 \vec{G}_2)$. In expressions (2.24)-(2.27), as it has already been noticed, ρ_0 is understood as a difference operator \hat{H} . However, in obtaining numerical evaluations the reasonable approximation which is not reduced to the nonrelativistic one is the use, instead of the operator \hat{H} , of the eigenvalue of energy ρ_0 (as in the problem of a positronium /17/) of one bound particle in the part of potential (1.1), consisting of the sum $V_{conf}(r)$ and the scalar part of potential (1.4): $V = \frac{4m^2}{\mu^2 + 4\vec{x}^2}$.

3. Quasiclassical Solution of Partial Equations

We write down uncoupled equations (2.25) in such a form:

$$[\hat{H}_e - X(r)] R_e(r) = 0, \quad (3.1)$$

where $X(r) = (2m + E_{bound} - V(r)) / 2m$,

and the potential $V(r)$ is defined through expressions (2.30), (2.31).

Regular in the zero free solution ($V(r) = 0$) of equation (3.1), according to Ψ has the form:

$$\Psi_e^{\text{free}}(r, x) = \sqrt{\frac{\pi}{2}} \operatorname{sh} x (-1)^{l+1} \frac{(-r/x)^{(e+1)}}{r} P_{(-\frac{1}{2}-e, -\frac{1}{2}+i\sqrt{x}}(chx)) \quad (3.2)$$

$$chx = (2m + E_{\text{bound}}) / 2m \quad (3.3)$$

where $P_{(-\frac{1}{2}-e, -\frac{1}{2}+i\sqrt{x}}(chx))$ is the Legendre function and $(-r/x)^{(e+1)}$ is the generalized degree defined by the expression: $(\frac{r}{x})^{(e+1)} = i \frac{(e+1)\Gamma(i\frac{r}{x} + e+1)}{\Gamma(i\frac{r}{x})}$.

The Legendre function $P_{(-\frac{1}{2}-e, -\frac{1}{2}+i\sqrt{x}}(chx))$ is real. Thus in expression (3.2) only the factor $(\frac{r}{x})^{(e+1)}$ appears to be complex.

We separate this factor by the substitution:

$$\Psi_e(r, x) = \frac{(-r/x)^{(e+1)}}{r} K_e(r, x). \quad (3.4)$$

We obtain the following equation for the function $K_e(r, x)$:

$$\left[2ch\left(i\lambda \frac{d}{dr}\right) - \frac{2(e+1)}{r\lambda} \operatorname{sh}\left(i\lambda \frac{d}{dr}\right) - X(r) \right] K_e(r, x) = 0. \quad (3.5)$$

Note, that the Hamiltonian of the equation (3.5) in contrast to (3.5) appears to be a real operator. In the case of the potential is real, $K_e(r, x)$ is also a real function. When the interaction is absent, the Legendre function appears to be a solution of equation (3.5)

$$K_e(r, x) = P_{(-\frac{1}{2}-e, -\frac{1}{2}+i\sqrt{x}}(chx) = \left(\frac{\operatorname{sh}x}{2}\right)^{e+\frac{1}{2}} \frac{\exp[-\lambda(i\frac{r}{x} + e+1)]}{r^{(\frac{3}{2}+e)}} \times \quad (3.6)$$

$$\times F\left(i\frac{r}{x} + e+1, e+1; 2e+2; 1 - \exp(-2x)\right).$$

The conditions of the orthogonality for these functions will be written down analogously to (1.6b):

$$\frac{1}{\lambda} \int_0^{\infty} \text{sh}^2 x \, dx \, P_{(-\frac{1}{\lambda})}^{-\frac{1}{2}-e} P_{(-\frac{1}{\lambda})}^{-\frac{1}{2}-e} = \frac{S(r-r')}{\left(-\frac{r}{\lambda}\right)^{e+1} \left(-\frac{r'}{\lambda}\right)^{e+1}} \quad (3.7)$$

$$\frac{1}{\lambda} \int_0^{\infty} dr \, P_{(-\frac{1}{\lambda})}^{-\frac{1}{2}-e} P_{(-\frac{1}{\lambda})}^{-\frac{1}{2}-e} \left[\left(-\frac{r}{\lambda}\right)^{e+1} \right]^* \left(-\frac{r'}{\lambda}\right)^{e+1} = \frac{S(x-x')}{\sqrt{\text{sh} x \, \text{sh} x'}}.$$

We solve expressions (3.5) by the WKB method. Suppose:

$$K_e(r) = \exp \frac{i}{\hbar} g(r); \quad g(r) = g_0(r) + \frac{\hbar}{i} g_1(r) \dots \quad (3.8)$$

Inserting (3.8) into (3.5) and keeping terms of the zero order in \hbar , we derive a differential equation for $g_0(r)$:

$$\frac{\lambda}{\hbar} g_0'(r) = \text{arcch} X_{\Lambda}(r) - i \text{arctg} \Lambda \frac{\lambda}{r} \quad (3.9)$$

$$X_{\Lambda}(r) = X(r) \cdot \left[1 + \left(\Lambda \frac{\lambda}{r} \right)^2 \right]^{-1/2}, \quad \Lambda = l + 1.$$

The imaginary part in (3.9) contributes into the preexponential factor. With the account of the term of the order $\frac{\hbar}{i}$, we arrive at the equation for $g_1'(r)$:

$$g_1'(r) = - \frac{X_{\Lambda}'(r) + i \frac{\lambda \Lambda}{r^2 + \lambda^2 \Lambda^2} X_{\Lambda}(r)}{2(X_{\Lambda}^2(r) - 1)} X_{\Lambda}(r). \quad (3.10)$$

Thus we obtain for the function $K_e(r)$:

$$K_{ge}(r) = [X_{\Lambda}^2(r) - 1]^{-1/4} (r^2/\lambda^2 + \Lambda^2)^{-\frac{\Lambda}{2}} \times \quad (3.11)$$

$$\times \exp \frac{i}{\lambda} \int_{r_-}^r dr' \text{arcch} X_{\Lambda}(r') - \frac{X_{\Lambda}(r)}{2[X_{\Lambda}(r) - 1]} \cdot \frac{\lambda \Lambda}{r^2 + \lambda^2 \Lambda^2} \Big|.$$

Here r_- is a turning point, defined from the condition $X_{\Lambda}(r_-) =$

1 . In the case $r \gg r_- \gg \lambda$ that is characteristic of energies

$E \ll 1mc^2$ ¹⁾ formula (3.11) gets simplified. We write it down in the form of a standing wave

$$K_{qe}(r)_{r \gg \lambda} = [X_A^2(r) - 1]^{-1/4} \left(\frac{E}{\lambda}\right)^{e-1} \sin \frac{1}{\lambda} \int_{r-}^r dr' \operatorname{arccch} X_A(r'). \quad (3.12)$$

The factor r^{-e-1} compensates the separated expression $\left(\frac{E}{\lambda}\right)_{r \rightarrow \infty}^{(e+1)}$
 $\rightarrow (r/\lambda)^{e+1}$. With the help of (3.12) we obtain a quasi-classical condition of quantization

$$\int_{r-}^{r+} dr' \operatorname{arccch} X_A(r') = \lambda \pi \left(n + \frac{1}{2}\right). \quad (3.13)$$

For the majority of potentials a spectrum is formed at the distances $r \gg \lambda$. This allows us to neglect imaginary terms in $V_i(r)$ (2.23-2.27). In such an approach, potentials (2.30, 2.31) become real and we can find energy levels for a singlet ($S=0$) and triplet ($S=1, e=j$) states.

4. Calculation of Electromagnetic Decay Width of Meson

To illustrate the use of quasi-classical formulas, obtained in §3, we calculate the $|\psi(a)|^2$ -value of WF quadratic at $r=0$ ²⁾ that is present in formulas for electromagnetic width of mesons (1.9), (1.10). In the case of the massless-particle (gluon) exchange to the first term of the quasi-potential (1.3)

$V = -g_V^2 \frac{4m^2}{3r^2}$ there corresponds the potential $V(r) = \frac{c\hbar (rm\pi)}{r}$

in the r -space. There have been obtained in (19) exact solutions and a spectrum in the field of such a potential. The function $c\hbar (rm\pi)$ changes essentially only at a distance of the order of λ from the origin of coordinate and, as is seen in ¹⁷⁾, its presence does not influence the form of the spectrum.

¹⁾ For the free motion, for example: $\frac{r-}{\lambda} \approx \frac{2mc^2\lambda}{E} \gg 1$.

²⁾ Note $r=0$ means that particles are at the Compton wave length distance ¹⁵⁾. A possibility of substitution of nonrelativistic WF in (1.9) for $|\psi^{nonrel}|^2$ has been discussed in ¹⁵⁾.

Consider a potential, being a combination of the OREP transform in RCR, taking into account only the Coulomb part and a confining potential. The value of energy levels is in general influenced by the potential behaviour at large distances $r \gg \lambda$. In this region we may neglect a hypergeometrical factor in the Coulomb potential

$$V(r) = -\frac{\alpha}{r} + V_{conf}(r), \quad (4.1)$$

at small r a Coulomb term dominates (region I at the Figure). In this region a WF $\Psi(r)$ coincides with the exact regular solution at $r=0$ in the potential $-\alpha/r$:

$$\Psi_e(r) = C_e \frac{\Gamma(i\frac{\alpha}{\lambda} + e + 1)}{r \Gamma(i\frac{\alpha}{\lambda})} K_e(r) \quad (4.2)$$

$$K_e(r) = \exp(-i\frac{\alpha}{\lambda} - i\alpha x) F(\ell + 1 + i\frac{\alpha}{\lambda}; \ell + 1 + i\alpha, 2\ell + 2; 1 - e^{-2ix}) \quad (4.3)$$

$$x = \alpha/\lambda \operatorname{sh} x,$$

where C_e is an unknown normalizing factor. This factor enters into the definition $\Psi(0)$.

To obtain this we consider an asymptotic of the exact solution (4.2) at large Γ , written in the form of a standing wave

$$\Psi_e(r) \xrightarrow{r \rightarrow \infty} C_e \frac{2 \Gamma(2\ell + 2) \exp[-\frac{\pi}{2} x + (e + 1)x]}{(2 \operatorname{sh} x)^{e+1} \operatorname{Re}[\Gamma(\ell + 1 - i\alpha)]} \times \quad (4.4)$$

$$\times \frac{1}{r} \sin\left[xr + \alpha \ln(2r \operatorname{sh} x) + \eta_e - \frac{\pi}{2} e\right],$$

where η_e is the relativistic Coulomb phase:

$$\eta_e = \arg \Gamma(\ell + 1 - i\alpha). \quad (4.5)$$

The Coulomb phase can be also calculated through the WKB method:

$$\eta_e^{WKB} = \frac{1}{\lambda} \lim_{\Gamma \rightarrow \infty} \left[\int_{z_-}^z \text{arccosh } X_\lambda^{coul}(r') dr' - \int_{z_-}^z \text{arccosh } X_\lambda^{free}(r') dr' \right]. \quad (4.6)$$

Assume the Coulomb constant to be small

$d \ll 1$.

This is characteristic for the description of the J/ψ and δ -particles in the potential model^[20] and agrees with the hypothesis of asymptotic freedom. Further with the neglect of the terms of the first order in d and by integration in (4.6) we arrive at the following result:

$$\eta_e^{WKB} = x - x \ln \sqrt{x^2 + \Lambda^2} - \Lambda \arcsin \frac{x}{\sqrt{x^2 + \Lambda^2}}. \quad (4.7)$$

The forms of expressions (4.5) and (4.7) coincide with their nonrelativistic analogs. The difference is that x is determined by relation (4.3), that in the nonrelativistic limit turns into a dimensionless quantity, used in the Schrödinger equation $x \rightarrow \frac{e^2}{2V}$. Phases η_e and η_e^{WKB} for $x \gg 1$ differ only by terms of the order x^{-1} ^[21]. Hence in the Coulomb field the exact solution $\psi(r)$ and the quasiclassical wave function $\psi^{WKB}(r)$ coincide up to a factor in the region of large r (range II, figure 1) under the condition $d \ll 1$, $x \gg 1$. If we can neglect a confining part of the potential (4.1) at such a large r , then in region II it is possible to equate an exact Coulomb solution (it is true in regions II and III^[22]) and quasi-classical solution obtained for regions II and III^[22]. Thus, we obtain the following normalizing constants:

$$C_e \lambda \Gamma(2e+2) \frac{\exp[-\frac{1}{2}x + (e+1)x]}{(2 \operatorname{sh} x)^{e+1}} = \frac{C^{WKB}}{[X_\lambda^2(r) - 1]^{1/4}}. \quad (4.8)$$

The constant C^{WKB} is derived in the same way as in nonrelativistic case

$$\int |R_{ne}(r)|^2 r^2 dr = \frac{|C^{WKB}|^2}{\lambda} \int_{z_-}^z \frac{dr'}{\sqrt{X_\lambda^2(r') - 1}} = 1. \quad (4.9)$$

Integral in (4.10) is easily calculated by differentiation of quantization conditions (3.13) over n :

$$|C^{WIB}|^2 \approx \frac{2}{\pi \lambda} \frac{dX_{n,e}}{dn} ; X_{n,e} = (2m + E_{n,e})/2m \quad (4.10)$$


Calculating to exact Coulomb WF in the limit of $r \rightarrow 0$ and obtaining C_0^+ value from (4.8) and (4.10), we finally obtain:

$$|\Psi_{n,e}(0)|^2 = \lambda^{-3} \prod_{e \neq 0}^e (e^2 + x^2) \times \frac{\exp \pi x}{4x \operatorname{sh} \pi x} \left| P_{-\frac{1}{2}-e}^{(ix)} \right|_{-\frac{1}{2}+ix}^2 \left| \frac{dX_{n,e}}{dn} \right|^2 \quad (4.11)$$

This expression differs from zero even for states with $l \neq 0$. So, using (4.11) we can apply formulae (1.9), (1.10) to the description of electromagnetic width of states with any values of l .

In this paper a system has been obtained of partial equations describing two particles with spin 1/2, interacting through the one-vector-boson exchange. There have been solved uncoupled equations of the system by WKB method for a wide class of confining potentials and a condition of quantization and the expression for $|\Psi_{n,e}(0)|^2$ have been found. These results can be applied to the calculation of mass spectrum and width of electromagnetic decays of systems of e^+e^- , $\mu^+\mu^-$, $c\bar{c}$, $b\bar{b}$, $N\bar{N}$ type.

The authors are grateful to V.G.Kadyshevsky, S.P.Kuleshov for fruitful discussions, and are thankful to V.Kapshay, S.G.Kovalenko and I.L.Solovtsov for interest in the work.



References:

1. Skachkov N.B. *Theor. i Math.Fiz.*, 1975, 22, p. 213.
2. Skachkov N.B., Solovtsov I.L. *JINR Preprint*, E2-9763, E2-10026, Dubna, 1976.
3. Skachkov N.B., Solovtsov I.L. *JINR Communication*, P2-10320, Dubna, 1976.
4. Barbieri R. et al. *Nucl.Phys.*, 1976, B105, p. 122.
5. Logunov A.A., Tavkhelidze A.N. *Nuovo Cim.*, 1963, 29, p. 380.
6. Faustov R.N. *Ann.Phys.*, 1973, 78, p. 176.
7. Kadyshevsky V.G., Mir-Kasimov R.M., Skachkov N.B. *Nuovo Cim.*, 1968, 58, p. 233.
8. Foldy L.L., Wouthuysen S.A. *Phys.Rev.*, 1950, 78, p.29.
9. Shapiro I.S. *Dokl. Akad.Nauk SSSR*, 1956, 106, p. 647.
10. Kadyshevsky V.G., Mir-Kasimov R.M., Skachkov N.B. *Soviet Journal of Particles and Nuclei*, 1972, v.2, p. 635.
11. Hamada T., Johnston I.P. *Nucl.Phys.* 1962, 34, p. 382.
12. Donkov A.D. et al. *Int. Conf. on Hadron Interactions at High Energies*, Baku, 1971.
13. Van Royen R., Weisskopf V.F. *Nuovo Cim.*, 1957, 50, p. 617.
14. Pirenne J. *These Univ. of Paris*, 1944;
Wheller J.A. *Ann. N.Y. Acad. Sci.*, 1946, 48, p. 219.
Pomeranchuk J.Ya. *Doklady Akademii Nauk SSSR*, 1948, 60, p. 263.
15. Skachkov N.B. *JINR Preprint*, E2-7892, Dubna, 1974.
16. Mavrodiev S.Sch., Skachkov N.B. *Theor. i Math.Fiz.*, 1975, 23, p.32.
17. Berestecky V.B., Zifshic E.N., Pitaevsky A.P. *Relat' istic Quantum Theory*, v. 1, Fizmatgiz, 1968.
18. Poggio T.C., Schnitzer H.J. *Brandeis Univ.*, 1979.
19. Freeman M., Mateev M.P., Mir-Kasimov R.M. *Nucl.Phys.*, 1969, B12, p. 197.
20. Appelquist T., Barnett M.R., Lane K., *SLAC-PUB-2100*, 1978.
21. Flückge S., *Practical Quantum Mechanics*, I. Springer-Verlag, 1971.
22. Bell J.S., Pasupathy J. *Ref.TH 2649-CERN*, 1979.


Received by Publishing Department
on October 25 1979.

**SUBJECT CATEGORIES
OF THE JINR PUBLICATIONS**

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and techn que
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids
15.	Experimental physics of nuclear reactions at low energies
16.	Health physics. Shieldings
17.	Theory of condensed matter
18.	Applied researches

Издательский отдел Объединенного института ядерных исследований.
Заказ 27269. Тираж 680. Уч.-изд. листов 1,14.
Редактор Э.В.Ивашкевич.
Макет В.А.Халлякиной. Подписано к печати 13.11.79 г.

SM 8006805 ✓



ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА

E2 - 12885

A.V.Sidorov, N.B.Skachkov

DESCRIPTION OF WIDTH
AND SPECTRA
OF TWO RELATIVISTIC FERMIONS
BOUND STATES

1979

E2 - 12885

A.V.Sidorov, N.B.Skachkov

**DESCRIPTION OF WIDTH
AND SPECTRA
OF TWO RELATIVISTIC FERMIONS
BOUND STATES**

Submitted to TMΦ

Сидоров А.В., Скачков Н.Б.

E2 - 12885

Описание ширины и спектров связанных состояний
двух релятивистских фермионов

Целью работы является построение формализма для релятивистского описания системы двух частиц со спином $1/2$. Используется двухчастичное трехмерное ковариантное уравнение, полученное в квазипотенциальном подходе. Квазипотенциальное уравнение в релятивистском конфигурационном представлении сведено в приближении ОБЕР к системе разностных парциальных уравнений, которая является релятивистским аналогом нерелятивистской системы Хамеды-Джонстона. Методом ВКБ решена задача о спектре масс связанных состояний. С использованием точного релятивистского кулоновского решения двухчастичной задачи получены выражения для ширины лептонных распадов в кварковой модели.

Работа выполнена в Лаборатории теоретической физики, ОИЯИ.

Препринт Объединенного института ядерных исследований, Дубна 1979

Sidorov A.V., Skachkov N.B.

E2 - 12885

Descriptions of Width and Spectra of Two
Relativistic Fermions Bound States

The aim of this article is the construction of the formalism for the relativistic description of two particles with spin $1/2$. We use the two-particle three-dimensional equation, obtained by quasipotential approach. Quasipotential equation in the relativistic configurational space with OBEP potential is reduced to the system of partial equations which is the analog of nonrelativistic Hamada-Jonston system. WKB approach is used to calculate mass spectra and leptonic width of mesons in quark model.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna 1979

I. Introduction

The present paper may be treated as a sequel to papers^{1-3/} on the three-dimensional relativistic description of two-particle system with spin 1/2 in the quantum field theory. Principal formulae, allowing the practical application of our mathematical technique are obtained herein. It will be applicable to such attractive problems as the calculation of levels and width of systems like positronium, muon and baryon as well as bound states in a quark - antiquark system, that is old ρ, ω, φ and new $\psi/\psi, \gamma$ -mesons. Lately a nonrelativistic quark model based on the Schrödinger equation is widely used for the mass spectrum description of hard vector mesons of the ψ/ψ -particle type. For this purpose a confining potential is chosen and added by the Breit-Fermi potential accounting for spin effects.

$$V(r) = V_{\text{conf}}(r) + V_{\text{spin}}(r). \quad (1.1)$$

The Breit-Fermi potential is known to represent v^2/c^2 - expansion of Feynman matrix element of one-boson (OBE) or one-gluon exchange. However, such a quasirelativistic approach applied to ordinary light vector ρ, ω, φ mesons is not self-consistent. It was shown in paper^{4/} that in this case the contribution of relativistic corrections appears to be of the same order as the contribution of the nonrelativistic Hamiltonian,

which was initially taken as the principal one. Therefore an essentially relativistic approach is needed here.

In the present paper for the mass spectrum description of a system consisting of two fermions (for example, quark and antiquark), we use the equation of quasipotential approach ^{15/}. As in previous papers we use the relativistic three-dimensional two-particle quasipotential equation, obtained by Kadyshevsky, on the basis of the covariant Hamiltonian formulation of field theory and later in papers ^{16/} on the basis of the covariant technique of equating times in the two-particle wave function of Bethe-Salpiter equation. In the case of equal masses $m_1 = m_2 = m$ this equation, written in the centre-of-mass-system ($\vec{p}_1 = -\vec{p}_2 = \vec{p}$) has the form

$$(2E_q - 2E_p) \Psi_{g_1 g_2}(\vec{p}) = (2\pi)^{-3} \int \frac{d\vec{k}}{E_k} \sum_{g'_1 g'_2} V(\vec{p}, \vec{k}; E_q) \Psi_{g'_1 g'_2}(\vec{k}) \quad (1.2)$$

Now consider in more detail the part of quasipotential (1.1), containing spin effects. In the case of vector-boson exchange the quasipotential in r.h.s. of equation (1.1) represents the Feynman matrix element ^{11/}

$$\langle \vec{p}, \vec{g}_1; -\vec{p}, \vec{g}_2 | T_V^{(2)} | \vec{k}, \vec{g}'_1; -\vec{k}, \vec{g}'_2 \rangle = g_V^2 \frac{\bar{u}(\vec{p}) \gamma_\mu \bar{u}(\vec{k}) \gamma_\mu \bar{u}(-\vec{p}) \gamma_\mu \bar{u}(-\vec{k})}{\mu^2 - (\vec{p} - \vec{k})^2} = \sum_{g_{1p} g_{2p}} \xi_{g_{1p}}^* \xi_{g_{2p}}^* T_V^{(2)}(\vec{k} \rightarrow \vec{p}; \vec{p}) \xi_{g_{1p}} \xi_{g_{2p}} \quad (1.3)$$

μ is the mass of a boson, ξ^G are two-component spinors.

Applying to (1.3) the suggested transformation ^{11/} from the four-dimensional representation to the three-dimension one ¹⁾ and separating the Wigner matrix of spin rotation, we obtain:

¹⁾ This transformation (1) plays the same role as the Foldy-Wenthuysen one ^{18/}, allowing one to pass to the three-dimensional description of the spin. However, in contrast to (8) the discussed in (1) transformation does not deal with the expansion of interaction terms in powers of v^2/c^2 .

$$\begin{aligned}
T_V^{(2)}(\vec{k}(-)\vec{p}; \vec{p}) = & -\frac{gV^2}{\mu^2 + 4\vec{x}^2} \left[4m^2 + 4(\vec{G}_1 \vec{x})(\vec{G}_2 \vec{x}) - (\vec{G}_1 \vec{G}_2) \vec{x}^2 + \right. \\
& + 8\rho_0 \alpha_0 i(\vec{G}_1 + \vec{G}_2)[\vec{p} \times \vec{x}] + \frac{g}{m^2} (\rho_0^2 \alpha_0^2 + 2\rho_0 \alpha_0 (\vec{p} \vec{x}) - m^4) + \\
& \left. + \frac{g}{m^2} (\vec{G}_1 \vec{p})(\vec{G}_2 \vec{x})(\vec{G}_2 \vec{p})(\vec{G}_1 \vec{x}) \right]. \quad (1.4)
\end{aligned}$$

Here the quantity \vec{x} means a half-transfer of momentum, that can be expressed through the momentum transfer in the Lobachevsky space introduced into \mathcal{M} :

$$\vec{\Delta} = \vec{k}(-)\vec{p} = \Lambda_p^{-1} \vec{k} = \vec{k} - \frac{\vec{p}}{m} \left(k_0 - \frac{\vec{k} \vec{p}}{\rho_0 + m} \right) \quad (1.5)$$

by the formula $\vec{x} = \vec{\Delta} \sqrt{m/2(\Delta_0 + m)}$ \mathcal{M} .

Expression (1.4) has a form of the geometrical generalization of the Breit-Fermi potential taken in the momentum representation.

It is convenient to solve equation (1.3) by transformation to the relativistic configurational representation (RCR), introduced initially in \mathcal{M} . To obtain this, instead of the Fourier-transformation there is used an expansion over the complete and orthogonal system of functions in the Lobachevsky space^{9/}.

$$\xi(\vec{p}; \vec{n}, r) = \left(\frac{\rho_0 - \vec{p} \vec{n}}{m} \right)^{-1 - i r m} \quad (1.6)$$

$$\rho_0 = \sqrt{m^2 + \vec{p}^2}, \quad \vec{n}^2 = 1.$$

For these functions in papers ^{16,12/} we found an expansion over spherical harmonics:

$$\begin{aligned}
\xi(\vec{p}; \vec{n}, r) = & \sqrt{\frac{\pi}{2 \operatorname{sh} x}} \sum_{\ell=0}^{\infty} (2\ell+1) \frac{\Gamma(i\frac{r}{x} + \ell + 1)}{\Gamma(i\frac{r}{x} + 1)} P_{-\frac{1}{2} - \ell}^{-\ell}(chx) P_{\ell} \left(\frac{\vec{p} \vec{n}}{\rho} \right) = \\
= & \sum_{\ell=0}^{\infty} (2\ell+1) i^{\ell} P_{\ell}(chx, r) P_{\ell} \left(\frac{\vec{p} \vec{n}}{\rho} \right) \quad (1.7)
\end{aligned}$$

as well as the condition of completeness and orthogonality for radial parts

$$\frac{2shxshx'}{\pi\lambda^3} \int_0^\infty r^2 dr \rho_e(chx, r) \rho_e^*(chx', r) = \delta(x-x') \quad (1.8)$$

$$\frac{2r'r'}{\pi\lambda^3} \int_0^\infty sh^2x dx \rho_e(chx, r) \rho_e^*(chx, r') = \delta(r-r').$$

In [2,3] there was discussed a problem on the form of the second and third terms (1.4) in the relativistic configurational representation. There are transforms of all the five spin structures present in (1.4) in this paper.

Equation (1.2) transformed into RCR is a finite-difference one and may be reduced to the system of partial equations, being a relativistic analog of the nonrelativistic Hamada-Jonston system [11].

To solve partial finite-difference equations we use the suggested analog [12] of WKB method for RCR.

Calculation of the electromagnetic decay width of mesons is another important problem. To determine lepton width of vector mesons we use the V.Royen-Weisscopff formula (with account of colour).

$$\Gamma_{V \rightarrow e^+e^-} = 16\pi\alpha^2 M_V^{-2} e_q^2 |\Psi_{ne}(0)|^2, \quad (1.9)$$

and for the two-photon decay of pseudoscalar mesons we use formula [14]

$$\Gamma_{P \rightarrow \gamma\gamma} = 12\pi\alpha^2 m^{-2} e_q^4 |\Psi_{ne}(0)|^2. \quad (1.10)$$

In these formulae M_V is the mass of a meson; m and e_q are the mass and charge of quarks, and $\Psi_{ne}(0)$ is WF of system of a quark and anti-quark. Usually instead of this WF the nonrelativistic WF is used in the Coulomb field neglecting the rest of the spin part of the OREP. The width of states with $\ell \neq 0$ in terms of the relativistic theory is known to turn into zero due to the behaviour of the WF at the origin of coordinates as $\Psi_e^{nonrel} \xrightarrow{r \rightarrow 0} r^e$. This behaviour is of a kinematical character and is connected with the behaviour of free solutions $j_\ell(xr) \rightarrow (xr)^\ell$ for $x \rightarrow 0$. In a relativistic equation there is used another

complete system of functions (1.6) having an essentially different behaviour as $\Gamma \rightarrow 0$, so that $\psi_e^{r+0} \neq 0$ at $e \neq 0$.

In section 2 a quasipotential equation will be transformed to the system of partial finite-difference equations, in the relativistic configurational representation; section 3 is devoted to the solution of unbounded equations of the system through the WKB method; section 4 gives formulae for the width of vector meson decays and pseudoscalar mesons, using the exact Coulomb solution of the quasipotential equation at $\Gamma = 0$.

2. System of Partial Two-Particle Equations in the Relativistic Configurational Representation

For the quasipotential and WF the transformation into the relativistic configurational representation is performed by formulae:

$$V(\Gamma, \vec{n}; \vec{p}) = (2\pi)^{-3} \int d\Omega_\Delta \xi^*(\vec{\Delta}; \vec{\tau}) V(\vec{\Delta}; \vec{p}), \quad (2.1)$$

$$\Psi_{G_1 G_2}(\Gamma) = (2\pi)^{-3} \int d\Omega_p \xi(\vec{p}; \vec{n}, \Gamma) \Psi_{G_1 G_2}(\vec{p}). \quad (2.2)$$

In ^{15/} the physical meaning of the parameter Γ present in the function is discussed in detail.

Applying (2.1) to (1.4) we obtain:

$$V(\Gamma, \vec{n}; \vec{p}) = V_1(\Gamma, p_0) + V_2(\Gamma, \vec{n}; p_0, \vec{p}) \quad (2.3)$$

$$V_1(\Gamma, p_0) = -g_V^2 \left\{ (2p_0^2 - 1) \left[\frac{V(\Gamma)}{k_{VX}} + \frac{16\pi}{r} \frac{\delta(\Gamma^2 - 1)}{r} \delta(\vec{n}) \right] + \right. \quad (2.4)$$

$$\left. + \left[\frac{f^2}{3} V_{k_{VX}}(\Gamma) - \frac{8\pi}{3} \frac{\delta(\Gamma^2 - 1)}{r} \delta(\vec{n}) \right] \left[(\vec{G}_1 \vec{G}_2) - (\vec{S} \vec{p})^2 + \vec{p}^2 \vec{S}^2 \right] \right\}$$

$$V_2(\Gamma, \vec{n}; p_0, \vec{p}) = -g_V^2 \left\{ (\vec{S} \vec{L}) \frac{4}{r} [p_0 A(\Gamma) + (\vec{p} \vec{n}) \varphi(\Gamma)] - \right. \quad (2.5)$$

$$\left. - 4i p_0 (\vec{p} \vec{n}) A(\Gamma) + \left[\vec{S}_{1,2} - (\vec{S} \vec{L})^2 \frac{4}{r^2} \right] \varphi(\Gamma) \right\}.$$

The following notations are introduced here:

$$\vec{S} = \frac{\vec{G}_1 + \vec{G}_2}{2}; \vec{L} = [\vec{p} \times \vec{r}]; S'_{12} = 3(\vec{G}_1 \vec{n})(\vec{G}_2 \vec{n}) - (\vec{G}_1 \vec{G}_2). \quad (2.6)$$

$$A(r) = \frac{1}{r(r-i)} (r \operatorname{th} \alpha \sin \alpha + \cos \alpha) V_{\text{Yuk}}(r) \quad (2.7)$$

$$\varphi(r) = \frac{r^2}{3(r-i)(r+i)} \left[\mu^2 + 3 \frac{\mu}{r} \left(1 - \frac{\mu^2}{2}\right) \frac{\operatorname{th} \alpha}{\sqrt{1 - \frac{\mu^2}{4}}} + \right. \\ \left. + \frac{3 - 2\mu^2 \left(1 - \frac{\mu^2}{4}\right) - \frac{3}{2} \operatorname{ch} \alpha}{1 - \frac{\mu^2}{4}} \right] V_{\text{Yuk}}(r) \quad (2.8)$$

$V_{\text{Yuk}}(r)$ - the Yukawa potential transform: $4/(\mu^2 + 4\vec{x}^2)$

$$V_{\text{Yuk}}(r) = \frac{1}{4\pi r} \frac{\operatorname{ch} \alpha}{\operatorname{sh} r \pi}, \alpha = \arccos \frac{\mu^2 - 2}{\mu^2} \quad (\mu^2 < 4). \quad (2.9)$$

In formulae (2.3)-(2.9) we assume the mass of interacting particles m to be equal to 1.

Formula (2.3) is splitted into two parts $V_1(r, p_0)$ independent of the single vector "relativistic coordinate" direction \vec{n} and $V_2(r, \vec{n}; p_0, \vec{p})$ containing the dependence on \vec{n} through expressions of the type

$$(\vec{p} \vec{n}), [\vec{p} \times \vec{n}], S'_{12}. \quad (2.10)$$

The necessity of such a splitting becomes obvious in the course of the transformation of the complete equation (1.2) into RCR using formulae (2.1), (2.2).

$$(2E_1 - 2\hat{H}) \Psi_{G_1 G_2}(\vec{r}) = \int d\Omega_p d\Omega_k d\vec{r}_1 \xi(\vec{p}; \vec{n}, \kappa) \times \\ \times \xi^*(\vec{k}; \vec{n}_k, \kappa) \sum_{G_1' G_2'} V_{(G_1, \vec{p}, E)}^{G_1' G_2'} \Psi_{G_1' G_2'}(\vec{r}_1) \quad (2.11)$$

$$\hat{H} = \operatorname{ch}\left(i\lambda \frac{d}{dr}\right) + i \frac{\lambda}{r} \operatorname{sh}\left(i\lambda \frac{d}{dr}\right) + \frac{\Delta \theta \theta}{(r/\lambda)^2} \exp\left(i\lambda \frac{d}{dr}\right) \quad (2.12)$$

$\Delta_{\theta\phi}$ - angular part of the Laplas operator, $\lambda = \frac{h}{mc}$. Due to the character of the addition theorem of the relativistic plane waves /16/:

$$\xi(\vec{k}; \vec{n}, r) = \xi(\vec{k}(-)\vec{p}; \vec{n}_{\Lambda p}, z) \xi(\vec{p}; \vec{n}, r) \quad (2.13)$$

$$\vec{n}_{\Lambda p} = [m\vec{n} - \vec{p}(1 - \frac{\vec{p}\vec{n}}{\rho_0 + m})] / (\rho_0 - \vec{p}\vec{n}) \quad (2.14)$$

$V_2(r, \vec{n}; \rho_0, \vec{p})$ enters into equation (2.12) in a nonlocal way:

$$(2E_9 - 2A) \Psi_{\vec{G}_1 \vec{G}_2}(\vec{r}) = \sum_{\vec{G}'_1 \vec{G}'_2} V_2(\vec{G}'_1 \vec{G}'_2; \vec{r}, \rho) \Psi_{\vec{G}'_1 \vec{G}'_2}(\vec{r}) + \quad (2.15)$$

$$+ \int d\vec{r}_1 \sum_{\vec{G}'_1 \vec{G}'_2} \int d\Omega_p \xi(\vec{p}; \vec{n}, r) \xi^*(\vec{p}; \vec{n}_1, z_1) V_{\vec{G}'_1 \vec{G}'_2}(\vec{n}_1, \vec{n}_{\Lambda p}; \vec{p}) \Psi_{\vec{G}'_1 \vec{G}'_2}(\vec{r}_1)$$

since substituting it into the equation the dependence on vector \vec{n}_1 turns to the dependence on vector $\vec{n}_{\Lambda p}$. The vector rotation $\vec{n}_1 \rightarrow \vec{n}_{\Lambda p}$ taken into account gives the following dependence of expressions (2.10) on the vector \vec{p} and ρ_0 :

$$(\vec{p} \vec{n}_{\Lambda p}) = m^2 / (\rho_0 - \vec{p}\vec{n}_1) - \rho_0 \quad (2.16)$$

$$[\vec{p} \vec{n}_{\Lambda p}] = m [\vec{p} \times \vec{n}_1] / (\rho_0 - \vec{p}\vec{n}_1) \quad (2.17)$$

$$(\vec{G}_1 \vec{n}_{\Lambda p})(\vec{G}_2 \vec{n}_{\Lambda p}) = Z_1^T + Z_2^T \quad (2.18)$$

$$Z_1^T = m^2 (\vec{G}_1 \vec{n}_1)(\vec{G}_2 \vec{n}_1) / (\rho_0 - \vec{p}\vec{n}_1)^2$$

$$Z_2^T = \frac{m^2}{(\rho_0 - \vec{p}\vec{n}_1)^2} \left\{ -\frac{1}{m} [(\vec{G}_1 \vec{n}_1)(\vec{G}_2 \vec{p}) + (\vec{G}_2 \vec{p})(\vec{G}_1 \vec{n}_1)] (1 - \frac{\vec{p}\vec{n}_1}{\rho_0 + m}) + \right. \\ \left. + \frac{1}{m^2} (\vec{G}_2 \vec{p})(\vec{G}_1 \vec{p}) (1 - \frac{\vec{p}\vec{n}_1}{\rho_0 + m})^2 \right\}$$

However, with the help of the correlation:

$$\exp\left(i\frac{1}{m}\frac{d}{dr_2}\right)\xi^*(\vec{p}; \vec{n}_2, r_2) = \frac{m}{p_0 - \vec{p}\vec{n}_2}\xi^*(\vec{p}; \vec{n}_2, r_2) \quad (2.19)$$

expressions (2.16), (2.17) and the first term Z_1^T from (2.18) can be localized. Z_1^T is a relativistic generalization of the expression $(\vec{\sigma}_1 \vec{n})(\vec{\sigma}_2 \vec{n})$ present in the operator of tensor forces. Confining ourselves to Z_1^T in accounting tensor forces we obtain the r.h.s. of equation (2.15) to be of a local form:

$$(2E - 2\hat{H})\Psi_{\vec{\sigma}_1 \vec{\sigma}_2}(\vec{r}) = \sum_{\vec{\sigma}'_1 \vec{\sigma}'_2} \hat{V}(\vec{r}, \vec{p}) \Psi_{\vec{\sigma}'_1 \vec{\sigma}'_2}(\vec{r}) \quad (2.20)$$

The rotation $\vec{n} \rightarrow \vec{n}_{1p}$ in tensor forces taken into account gives an additional spin-spin interaction

$$\begin{aligned} & \varphi(r) [3(\vec{\sigma}_1 \vec{n}_{1p})(\vec{\sigma}_2 \vec{n}_{1p}) - (\vec{\sigma}_1 \vec{\sigma}_2)] \rightarrow \\ & \rightarrow \varphi(r-2i\lambda) S_{12} + [\varphi(r-2i\lambda) - \varphi(r)] (\vec{\sigma}_1 \vec{\sigma}_2), \end{aligned} \quad (2.21)$$

An angular and spin dependences of equation potential (2.20) are determined by five structures, having the same form as in the nonrelativistic theory:

$$\hat{V}(\vec{r}, \vec{p}) = V_S + V_T S_{12} + V_{LS} (\vec{L} \vec{S}) + V_G (\vec{\sigma}_1 \vec{\sigma}_2) + V_{(LS)} (\vec{S} \vec{Z})^2 \quad (2.22)$$

Functions $V_i(\vec{r})$ ($i = S, S_{12}, LS, G, (LS)^2$) depend on vector modulus \vec{r} and difference operators \vec{p} and $\vec{p}_0 = \hat{A}$ in the following way ($\lambda=1$):

$$\begin{aligned} V_S = & (\lambda p_0^2 - 1) \left[V_{YUK} (r) + \frac{16\sqrt{\pi}}{r} \mathcal{S}(r^2) - \right. \\ & \left. - 4i f_0 \left[\left(\frac{r-i}{r}\right)^2 \exp(-i\frac{d}{dr}) - p_0 \right] A(r) \right] \end{aligned} \quad (2.23)$$

$$V_T = \left(\frac{r-2i}{r}\right)^2 \exp(-2i\frac{d}{dr}) \quad (2.24)$$

$$V_{LS} = 4P_0 \frac{r}{(r+i)^2} A(r) + \left[\frac{r(r-i)}{(r+i)^2} \exp(-i \frac{d}{dr}) - P_0 \right] \frac{r}{(r+i)^2} \varphi(r) \quad (2.25)$$

$$V_G = \left[\left(\frac{r-2i}{r} \right)^2 \exp(-2i \frac{d}{dr}) - 1 \right] \varphi(r) - \frac{\mu^2}{3} V_{YUK}(r) - \frac{g}{3} \frac{\hat{r}}{r} \delta^j(r+1) \quad (2.26)$$

$$V_{(LS)^2} = - \frac{4r}{(r+i)(r+2i)} \varphi(r) \quad (2.27)$$

We introduce a radial function $R_{e's'e_s}^j(r)$

$$\Psi_{g, G_1, G_2}^S(\vec{r}) = \sum_{j e' e' e_s} R_{e's'e_s}^j(r) \left\{ \Omega_{j e' e'}^{*S}(\vec{n}) \right\} \left\{ \Omega_{j e' e'}^S(\vec{n}) \right\}_{G_1 G_2} \quad (2.28)$$

The substitution of (2.28) into equation (2.27) gives us a system of four partial equations. Two of the system of equations, obeying the case of $S=0$, $e'_i = i$ and $S=1$, $e'_i = j$ are uncoupled:

$$\left[2\hat{H}_e - 2E_{g,e} + V_{e'_i s}^{-1}(r) \right] R_{e'_i s}^S(r) = 0; \quad e'_i = j; \quad S=0, 1. \quad (2.29)$$

$$V_{e'_i s}^{S=0}(r) = -g_V^2 (V_S - 3V_G) \quad (2.30)$$

$$V_{e'_i s}^{S=1}(r) = -g_V^2 (V_S + 2V_T + V_G - V_{LS} + V_{(LS)^2}) \quad (2.31)$$

Hamiltonian \hat{H}_e is obtained from expression (2.12) for the \hat{H} through the substitution of the operator $\Delta_{\Theta\Phi}$ for its eigenvalue $\ell(\ell+1)$. The other two WF describing states with a spin $S=1$ are defined by the system of two coupled equations

$$\begin{cases} (2E_q - 2H_{e'=j+1}) R_{e'=j+1}(r) = V(r) R(r) + \tilde{V}_T(r) R_{e'=j-1}(r) \\ (2E_q - 2H_{e'=j-1}) R_{e'=j-1}(r) = V(r) R(r) + \tilde{V}_T(r) R_{e'=j+1}(r) \end{cases} \quad (2.32)$$

$$V_{e'=j+1}(r) = g_V^2 [V_S - (j+2) V_{LS} + (j+2)^2 V_{LS}^2 - \frac{2(j+1)}{2j+1} V_T + V_G] \quad (2.33)$$

$$V_{e'=j-1}(r) = g_V^2 [V_S + (j-1) V_{LS} + (j-1)^2 V_{LS}^2 - \frac{2(j-1)}{2j+1} V_T + V_G] \quad (2.34)$$

$$\tilde{V}_T(r) = -g_V^2 \frac{6\sqrt{j(j+1)}}{2j+1} V_T(r) \quad (2.35)$$

Writing down system (2.29), (2.32) we follow the approximation on the phenomenological use of the OBEP, accepted in paper /11/ we neglect the \vec{P} quadratic terms in the spin-spin interaction $\vec{P}^2 \vec{S}^2 - (\vec{P} \cdot \vec{S})^2$ and confine ourselves to the account of the terms proportional to $m^2 (\vec{G}_1 \vec{G}_2)$. In expressions (2.24)-(2.27), as it has already been noticed, ρ_0 is understood as a difference operator \hat{H} . However, in obtaining numerical evaluations the reasonable approximation which is not reduced to the nonrelativistic one is the use, instead of the operator \hat{H} , of the eigenvalue of energy ρ_0 (as in the problem of a positronium /11/) of one bound particle in the part of potential (1.1), consisting of the sum $V_{conf}(r)$ and the scalar part of potential (1.4): $V = \frac{4m^2}{\mu^2 + 4\vec{x}^2}$.

3. Quasiclassical Solution of Partial Equations

We write down uncoupled equations (2.25) in such a form:

$$[\hat{H}_e - X(r)] R_e(r) = 0, \quad (3.1)$$

where $X(r) = (2m + E_{bound} - V(r)) / 2m$,

and the potential $V(r)$ is defined through expressions (2.30), (2.31).

Regular in the zero free solution ($V(r) = 0$) of equation (3.1), according to Ψ has the form:

$$\Psi_e^{\text{free}}(r, x) = \sqrt{\frac{\pi}{2}} \operatorname{sh} x (-1)^{l+1} \frac{(-r/x)^{(e+1)}}{r} P_{(-\frac{1}{2}-e, -\frac{1}{2}+i\sqrt{x}}(chx) \quad (3.2)$$

$$chx = (2m + E_{\text{bound}}) / 2m \quad (3.3)$$

where $P_{(-\frac{1}{2}-e, -\frac{1}{2}+i\sqrt{x}}(chx)$ is the Legendre function and $(-r/x)^{(e+1)}$ is the generalized degree defined by the expression: $(\frac{r}{x})^{(e+1)} = i \frac{(e+1)\Gamma(i\frac{r}{x} + e+1)}{\Gamma(i\frac{r}{x})}$.

The Legendre function $P_{(-\frac{1}{2}-e, -\frac{1}{2}+i\sqrt{x}}(chx)$ is real. Thus in expression (3.2) only the factor $(\frac{r}{x})^{(e+1)}$ appears to be complex.

We separate this factor by the substitution:

$$\Psi_e(r, x) = \frac{(-r/x)^{(e+1)}}{r} K_e(r, x). \quad (3.4)$$

We obtain the following equation for the function $K_e(r, x)$:

$$\left[2ch\left(i\lambda \frac{d}{dr}\right) - \frac{2(e+1)}{r\lambda} \operatorname{sh}\left(i\lambda \frac{d}{dr}\right) - X(r) \right] K_e(r, x) = 0. \quad (3.5)$$

Note, that the Hamiltonian of the equation (3.5) in contrast to (3.5) appears to be a real operator. In the case of the potential is real, $K_e(r, x)$ is also a real function. When the interaction is absent, the Legendre function appears to be a solution of equation (3.5)

$$K_e(r, x) = P_{(-\frac{1}{2}-e, -\frac{1}{2}+i\sqrt{x}}(chx) = \left(\frac{\operatorname{sh}x}{2}\right)^{e+\frac{1}{2}} \frac{\exp[-\lambda(i\frac{r}{x} + e+1)]}{r^{(\frac{3}{2}+e)}} \times \\ \times F\left(i\frac{r}{x} + e+1, e+1; 2e+2; 1 - \exp(-2x)\right). \quad (3.6)$$

The conditions of the orthogonality for these functions will be written down analogously to (1.6b):

$$\frac{1}{\lambda} \int_0^{\infty} \text{sh}^2 x \, dx \, P_{(-\frac{1}{\lambda})}^{-\frac{1}{2}-e} P_{(-\frac{1}{\lambda})}^{-\frac{1}{2}-e} = \frac{S(r-r')}{\left(-\frac{r}{\lambda}\right)^{e+1} \left(-\frac{r'}{\lambda}\right)^{e+1}} \quad (3.7)$$

$$\frac{1}{\lambda} \int_0^{\infty} dr \, P_{(-\frac{1}{\lambda})}^{-\frac{1}{2}-e} P_{(-\frac{1}{\lambda})}^{-\frac{1}{2}-e} \left[\left(-\frac{r}{\lambda}\right)^{e+1} \right]^* \left(-\frac{r'}{\lambda}\right)^{e+1} = \frac{S(x-x')}{\sqrt{\text{sh} x \, \text{sh} x'}}.$$

We solve expressions (3.5) by the WKB method. Suppose:

$$K_e(r) = \exp \frac{i}{\hbar} g(r); \quad g(r) = g_0(r) + \frac{\hbar}{i} g_1(r) \dots \quad (3.8)$$

Inserting (3.8) into (3.5) and keeping terms of the zero order in \hbar , we derive a differential equation for $g_0(r)$:

$$\frac{\lambda}{\hbar} g_0'(r) = \text{arccch} X_{\Lambda}(r) - i \text{arctg} \Lambda \frac{\lambda}{r} \quad (3.9)$$

$$X_{\Lambda}(r) = X(r) \cdot \left[1 + \left(\Lambda \frac{\lambda}{r} \right)^2 \right]^{-1/2}, \quad \Lambda = l+1.$$

The imaginary part in (3.9) contributes into the preexponential factor. With the account of the term of the order $\frac{\hbar}{i}$, we arrive at the equation for $g_1'(r)$:

$$g_1'(r) = - \frac{X_{\Lambda}'(r) + i \frac{\lambda \Lambda}{r^2 + \lambda^2 \Lambda^2} X_{\Lambda}(r)}{2(X_{\Lambda}^2(r) - 1)} X_{\Lambda}(r). \quad (3.10)$$

Thus we obtain for the function $K_e(r)$:

$$K_{ge}(r) = [X_{\Lambda}^2(r) - 1]^{-1/4} (r^2/\lambda^2 + \Lambda^2)^{-\frac{\Lambda}{2}} \times \quad (3.11)$$

$$\times \exp \frac{i}{\lambda} \int_{r_-}^r dr' \left[\text{arccch} X_{\Lambda}(r') - \frac{X_{\Lambda}(r')}{2[X_{\Lambda}(r') - 1]} \cdot \frac{\lambda \Lambda}{r'^2 + \lambda^2 \Lambda^2} \right].$$

Here r_- is a turning point, defined from the condition $X_{\Lambda}(r_-) =$

1 . In the case $r \gg r_- \gg \lambda$ that is characteristic of energies

$E \ll 1mc^2$ ¹⁾ formula (3.11) gets simplified. We write it down in the form of a standing wave

$$K_{qe}(r)_{r \gg \lambda} = [X_A^2(r) - 1]^{-1/4} \left(\frac{E}{\lambda}\right)^{e-1} \sin \frac{1}{\lambda} \int_{r-}^r dr' \operatorname{arccch} X_A(r'). \quad (3.12)$$

The factor r^{-e-1} compensates the separated expression $\left(\frac{E}{\lambda}\right)_{r \rightarrow \infty}^{(e+1)}$
 $\rightarrow (r/\lambda)^{e+1}$. With the help of (3.12) we obtain a quasi-classical condition of quantization

$$\int_{r-}^{r+} dr' \operatorname{arccch} X_A(r') = \lambda \pi \left(n + \frac{1}{2}\right). \quad (3.13)$$

For the majority of potentials a spectrum is formed at the distances $r \gg \lambda$. This allows us to neglect imaginary terms in $V_i(r)$ (2.23-2.27). In such an approach, potentials (2.30, 2.31) become real and we can find energy levels for a singlet ($S=0$) and triplet ($S=1, e=j$) states.

4. Calculation of Electromagnetic Decay Width of Meson

To illustrate the use of quasi-classical formulas, obtained in §3, we calculate the $|\psi(a)|^2$ -value of WF quadratic at $r=0$ ²⁾ that is present in formulas for electromagnetic width of mesons (1.9), (1.10). In the case of the massless-particle (gluon) exchange to the first term of the quasi-potential (1.3)

$$V = -g_V^2 \frac{4m^2}{3\pi^2} \text{ there corresponds the potential } V(r) = \frac{c \hbar (r m \pi)}{r}$$

in the r -space. There have been obtained in (19) exact solutions and a spectrum in the field of such a potential. The function $c \hbar (r m \pi)$ changes essentially only at a distance of the order of λ from the origin of coordinate and, as is seen in ¹⁷⁾, its presence does not influence the form of the spectrum.

¹⁾ For the free motion, for example: $\frac{r-}{\lambda} \approx \frac{1mc^2 \lambda}{E} \gg 1$.

²⁾ Note $r=0$ means that particles are at the Compton wave length distance ¹⁵⁾. A possibility of substitution of nonrelativistic WF in (1.9) for $|\psi^{nonrel}|^2$ has been discussed in ¹⁵⁾.

Consider a potential, being a combination of the OREP transform in RCR, taking into account only the Coulomb part and a confining potential. The value of energy levels is in general influenced by the potential behaviour at large distances $r \gg \lambda$. In this region we may neglect a hypergeometrical factor in the Coulomb potential

$$V(r) = -\frac{\alpha}{r} + V_{conf}(r), \quad (4.1)$$

at small r a Coulomb term dominates (region I at the Figure). In this region a WF $\Psi(r)$ coincides with the exact regular solution at $r=0$ in the potential $-\alpha/r$:

$$\Psi_e(r) = C_e \frac{\Gamma(i\frac{\alpha}{\lambda} + e + 1)}{r \Gamma(i\frac{\alpha}{\lambda})} K_e(r) \quad (4.2)$$

$$K_e(r) = \exp(-i\frac{\alpha}{\lambda} - i\alpha x) F(\ell + 1 + i\frac{\alpha}{\lambda}; \ell + 1 + i\alpha, 2\ell + 2; 1 - e^{-2ix}) \quad (4.3)$$

$$x = \alpha/\lambda \operatorname{sh} x,$$

where C_e is an unknown normalizing factor. This factor enters into the definition $\Psi(0)$.

To obtain this we consider an asymptotic of the exact solution (4.2) at large Γ , written in the form of a standing wave

$$\Psi_e(r) \xrightarrow{r \rightarrow \infty} C_e \frac{2 \Gamma(2\ell + 2) \exp[-\frac{\pi}{2} x + (e + 1)x]}{(2 \operatorname{sh} x)^{e+1} \operatorname{Re}[\Gamma(\ell + 1 - i\alpha)]} \times \quad (4.4)$$

$$\times \frac{1}{r} \sin\left[xr + \alpha \ln(2r \operatorname{sh} x) + \eta_e - \frac{\pi}{2} e\right],$$

where η_e is the relativistic Coulomb phase:

$$\eta_e = \arg \Gamma(\ell + 1 - i\alpha). \quad (4.5)$$

The Coulomb phase can be also calculated through the WKB method:

$$\eta_e^{WKB} = \frac{1}{\lambda} \lim_{\Gamma \rightarrow \infty} \left[\int_{z_-}^z \text{arccosh } X_\lambda^{coul}(r') dr' - \int_{z_-}^z \text{arccosh } X_\lambda^{free}(r') dr' \right]. \quad (4.6)$$

Assume the Coulomb constant to be small

$d \ll 1$.

This is characteristic for the description of the J/ψ and δ -particles in the potential model^[20] and agrees with the hypothesis of asymptotic freedom. Further with the neglect of the terms of the first order in d and by integration in (4.6) we arrive at the following result:

$$\eta_e^{WKB} = x - x \ln \sqrt{x^2 + \Lambda^2} - \Lambda \arcsin \frac{x}{\sqrt{x^2 + \Lambda^2}}. \quad (4.7)$$

The forms of expressions (4.5) and (4.7) coincide with their nonrelativistic analogs. The difference is that x is determined by relation (4.3), that in the nonrelativistic limit turns into a dimensionless quantity, used in the Schrödinger equation $x \rightarrow \frac{e^2}{2V}$. Phases η_e and η_e^{WKB} for $x \gg 1$ differ only by terms of the order x^{-1} ^[21]. Hence in the Coulomb field the exact solution $\psi(r)$ and the quasiclassical wave function $\psi^{WKB}(r)$ coincide up to a factor in the region of large r (range II, figure 1) under the condition $d \ll 1$, $x \gg 1$. If we can neglect a confining part of the potential (4.1) at such a large r , then in region II it is possible to equate an exact Coulomb solution (it is true in regions II and III^[22]) and quasi-classical solution obtained for regions II and III^[22]. Thus, we obtain the following normalizing constants:

$$C_e \lambda \Gamma(2e+2) \frac{\exp[-\frac{1}{2}x + (e+1)x]}{(2 \operatorname{sh} x)^{e+1}} = \frac{C^{WKB}}{[X_\lambda^2(r) - 1]^{1/4}}. \quad (4.8)$$

The constant C^{WKB} is derived in the same way as in nonrelativistic case

$$\int |R_{ne}(r)|^2 r^2 dr = \frac{|C^{WKB}|^2}{\lambda} \int_{z_-}^z \frac{dr'}{\sqrt{X_\lambda^2(r') - 1}} = 1. \quad (4.9)$$

Integral in (4.10) is easily calculated by differentiation of quantization conditions (3.13) over n :

$$|C^{WIB}|^2 \approx \frac{2}{\pi \lambda} \frac{dX_{n,e}}{dn} ; X_{n,e} = (2m + E_{n,e})/2m \quad (4.10)$$


Calculating to exact Coulomb WF in the limit of $r \rightarrow 0$ and obtaining C_0^+ value from (4.8) and (4.10), we finally obtain:

$$|\Psi_{n,e}(0)|^2 = \lambda^{-3} \prod_{e \neq 0}^e (e^2 x^2) \times \frac{\exp \pi x}{4x \operatorname{sh} \pi x} \left| P_{-\frac{1}{2}-e}^{(ix)} \right|_{-\frac{1}{2}+ix}^2 \left| \frac{dX_{n,e}}{dn} \right|^2 \quad (4.11)$$

This expression differs from zero even for states with $l \neq 0$. So, using (4.11) we can apply formulae (1.9), (1.10) to the description of electromagnetic width of states with any values of l .

In this paper a system has been obtained of partial equations describing two particles with spin 1/2, interacting through the one-vector-boson exchange. There have been solved uncoupled equations of the system by WKB method for a wide class of confining potentials and a condition of quantization and the expression for $|\Psi_{n,e}(0)|^2$ have been found. These results can be applied to the calculation of mass spectrum and width of electromagnetic decays of systems of e^+e^- , $\mu^+\mu^-$, $c\bar{c}$, $b\bar{b}$, $N\bar{N}$ type.

The authors are grateful to V.G.Kadyshevsky, S.P.Kuleshov for fruitful discussions, and are thankful to V.Kapshay, S.G.Kovalenko and I.L.Solovtsov for interest in the work.



References:

1. Skachkov N.B. Theor. i Math.Fiz., 1975, 22, p. 213.
2. Skachkov N.B., Solovtsov I.L. JINR Preprint, E2-9763, E2-10026, Dubna, 1976.
3. Skachkov N.B., Solovtsov I.L. JINR Communication, P2-10320, Dubna, 1976.
4. Barbieri R. et al. Nucl.Phys., 1976, B105, p. 122.
5. Logunov A.A., Tavkhelidze A.N. Nuovo Cim., 1963, 29, p. 380.
6. Faustov R.N. Ann.Phys., 1973, 78, p. 176.
7. Kadyshevsky V.G., Mir-Kasimov R.M., Skachkov N.B. Nuovo Cim., 1968, 58, p. 233.
8. Foldy L.L., Wouthuysen S.A. Phys.Rev., 1950, 78, p.29.
9. Shapiro I.S. Dokl. Akad.Nauk SSSR, 1956, 106, p. 647.
10. Kadyshevsky V.G., Mir-Kasimov R.M., Skachkov N.B. Soviet Journal of Particles and Nuclei, 1972, v.2, p. 635.
11. Hamada T., Johnston I.P. Nucl.Phys. 1962, 34, p. 382.
12. Donkov A.D. et al. Int. Conf. on Hadron Interactions at High Energies, Baku, 1971.
13. Van Royen R., Weisskopf V.F. Nuovo Cim., 1957, 50, p. 617.
14. Pirenne J. These Univ. of Paris, 1944;
Wheller J.A. Ann. N.Y. Acad. Sci., 1946, 48, p. 219.
Pomeranchuk J.Ya. Doklady Akademii Nauk SSSR, 1948, 60, p. 263.
15. Skachkov N.B. JINR Preprint, E2-7892, Dubna, 1974.
16. Mavrodiev S.Sch., Skachkov N.B. Theor. i Math.Fiz., 1975, 23, p.32.
17. Berestecky V.B., Zifshic E.N., Pitaevsky A.P. Relat' istic Quantum Theory, v. 1, Fizmatgiz, 1968.
18. Poggio T.C., Schnitzer H.J. Brandeis Univ., 1979.
19. Freeman M., Mateev M.P., Mir-Kasimov R.M. Nucl.Phys., 1969, B12, p. 197.
20. Appelquist T., Barnett M.R., Lane K., SLAC-PUB-2100, 1978.
21. Flückge S., Practical Quantum Mechanics, I. Springer-Verlag, 1971.
22. Bell J.S., Pasupathy J. Ref.TH 2649-CERN, 1979.

Received by Publishing Department
on October 25 1979.

**SUBJECT CATEGORIES
OF THE JINR PUBLICATIONS**

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and techn que
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids
15.	Experimental physics of nuclear reactions at low energies
16.	Health physics. Shieldings
17.	Theory of condensed matter
18.	Applied researches

Издательский отдел Объединенного института ядерных исследований.
Заказ 27269. Тираж 680. Уч.-изд. листов 1,14.
Редактор Э.В.Ивашкевич.
Макет В.А.Халлякиной. Подписано к печати 13.11.79 г.