

ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА

SD 824946c

E1-80-316

SCALING IN THE MEAN
AND ASSOCIATIVE MULTIPLICITIES FOR
THE INCLUSIVE REACTIONS

$\bar{p} + p \rightarrow K_s^0 + x$ AND $\bar{p} + p \rightarrow \Lambda + X$
AT 22.4 GeV/c

Dubna - Alma-Ata - Helsinki - Moscow -
Prague - Tbilisi Collaboration

1980

B.V.Batyunya, I.V.Boguslavsky, I.M.Gramenitsky, R.Iednicky,
L.A.Tikhonova, A.Valkarova, V.Vrba.
Joint Institute for Nuclear Research, Dubna, USSR.

D.I.Jermilova, V.V.Philippova, V.V.Samojlov, T.Temiraliev.
Institute of High Energy Physics, Alma-Ata, USSR.

S.Dumbrajs, J.Ervanne, E.Hannula, P.Villanen.
Department of Nuclear Physics of the Helsinki University,
Helsinki, Finland.

R.K.Dementiev, I.A.Korzhevina, E.M.Leikin, A.G.Pavlova,
V.I.Rud.
Institute for Nuclear Physics Research of the Moscow State
University, Moscow, USSR.

I.Herynek, V.Kol, J.Řídký, V.Šimák.
Institute of Physics, ČSAV, Prague, ČSSR.

L.Rob, M.Suk.
Nuclear Center, Charles University, Prague, ČSSR.

J.Patočka.
SHAFARIC University, Kosice, ČSSR.

G.O.Kuratashvili, A.M.Hudzhadze, V.D.Tsintsadze, T.P.Topuriya.
Tbilisi State University, USSR.

S.V.Levonjan.
Lebedev Institute of Physics, Moscow, USSR.

E1-80-316

**SCALING IN THE MEAN
AND ASSOCIATIVE MULTIPLICITIES FOR
THE INCLUSIVE REACTIONS**

**$\bar{p} + p \rightarrow K_s^0 + x$ AND $\bar{p} + p \rightarrow \Lambda + X$
AT 22.4 GeV/c**

**Dubna - Alma-Ata - Helsinki - Moscow -
Prague - Tbilisi Collaboration**

Submitted to JΦ

Исследование скейлинга в среднем и ассоциативной множественности в инклюзивных реакциях образования K_s^0 - и Λ -частиц в $\bar{p}p$ -взаимодействиях при 22,4 ГэВ/с

В работе представлены результаты экспериментального исследования скейлинга в среднем и ассоциативной множественности в реакциях образования K_s^0 - и Λ -частиц в $\bar{p}p$ -взаимодействиях при 22,4 ГэВ/с. Показано, что скейлинг в среднем выполняется для K_s^0 -мезонов и нарушается для Λ -гиперонов. Проверяется теоретическое предсказание о связи скейлинга в среднем с другими скейлинговыми законами.

Для Λ -частиц получена линейная зависимость ассоциативной множественности от логарифма квадрата недостающей массы с параметром наклона, близким по величине аналогичным параметрам, найденным в других экспериментах. Проверено выполнение ассоциативного KNO-скейлинга для реакций образования Λ -гиперонов.

Работа выполнена в Лаборатории высоких энергий ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна 1980

Scaling in the Mean and Associative Multiplicities for the Inclusive Reactions $\bar{p} + p \rightarrow K_s^0 + X$ and $\bar{p} + p \rightarrow \Lambda + X$ at 22.4 GeV/c

Scaling in the mean and associative multiplicities are studied for K_s^0 and Λ production in $\bar{p}p$ -interactions at 22.4 GeV/c. It is found that the scaling in the mean is valid for K_s^0 -mesons and not for Λ -hyperons. The connection of the scaling in the mean with other scaling laws is discussed. For Λ particles the associative multiplicities depend linearly on the logarithmic missing mass squared with a slope parameter comparable to those found in other experiments. The validity of the associative KNO-scaling has been tested for events with Λ -production.

The investigation has been performed at the Laboratory of High Energies, JINR.

1. SCALING IN THE MEAN

The hypothesis of scaling in the mean was introduced in paper ¹. From it a scaling equation follows for semi-inclusive one-dimensional distributions of particle c

$$\frac{d^2\sigma_n^c}{d^2p_i} = \phi_i \left(\frac{p_i}{p_{i,n}} \right) \quad (1)$$

Here $i = T, L$ for transverse (p_T) and c.m. longitudinal (p_L^*) momenta of particle c, $p_{i,n}$ are the mean values of these momenta, and σ_n^c is the inclusive cross section of particle c produced in association with n charged particles. The functions ϕ_i depend only on the scaling variable $p_i/p_{i,n}$, but neither on primary energy nor on charged multiplicity. It has been also assumed that the functions ϕ_i are independent of the initial state.

Equation (1) has been tested for π^0 -meson production in different reactions ^{1,2a} and for neutral strange particle production in pp-interactions at 19 GeV/c ^{2b} and 300 GeV/c ^{2c}. It has been found that scaling in the mean is valid for mesons, but for Λ -particles only the variable $p_T/p_{T,n}$ scales.

In this paper the semi-inclusive processes

$$\bar{p} + p \rightarrow K_s^0 + n + \text{neutrals} \quad (2)$$

$$\bar{p} + p \rightarrow \Lambda + n + \text{neutrals} \quad (3)$$

(n is the number of charged particles) are studied using equation (1).

The experimental data have been obtained in exposures of the hydrogen bubble chamber Ludmila to a RF separated beam of 22.4 GeV/c antiprotons at the Serpukhov accelerator. The amount of interactions is 1123 with K_s^0 -mesons and 454 with Λ -hyperons. The scanning, measuring and fitting procedures of the interactions with neutral strange particles have been described elsewhere ³.

The transverse and longitudinal distributions (equation(1)) are presented for K_s^0 -mesons in Figs.1 and 2 and for Λ -hyperons in Figs.3 and 4*. The data with $n \geq 6$ have been combined in

*For clarity our data with $n \geq 6$ and for all events have been shifted along the abscissa axis from the real positions (data at $n = 2,4$). The data for all events include zero-prong interactions as well.




Fig.1. Plot of $(\langle p_T \rangle_n / \sigma_n) \cdot (d\sigma_n / dp_T)$ versus $p_T / \langle p_T \rangle_n$ for K^0 -mesons. The solid line is the result of fitting the π^- -production in pp-interactions in an energy interval of 13-300 GeV/c.




Fig.2. Plot of $(\langle p_L^* \rangle_n / \sigma_n) (d\sigma_n / dp_L^*)$ versus $p_L^* / \langle p_L^* \rangle_n$ for K^0 -mesons. The solid line is the result of fitting the π^- -production in pp-interactions in an energy interval of 13-300 GeV/c.




Fig.3. Plot of $(\langle p_T \rangle_n / \sigma_n) (d\sigma_n / dp_T^*)$ versus $p_T / \langle p_T \rangle_n$ for Λ -particles. The solid line is the same as that in fig.1.




Fig.4. Plot of $(\langle p_L^* \rangle_n / \sigma_n) (d\sigma_n / dp_L^*)$ versus $p_L^* / \langle p_L^* \rangle_n$ for Λ -particles.

Table

Average values of p_T and p_L^* (in the c.m.s.) for K_s^0 and Λ - particles.

Particle	Multiplicity	Number of events	$\langle p_T \rangle_n$ GeV/c	$\langle p_L^* \rangle_n$ GeV/c
K_s^0	2	265	0.42 ± 0.02	0.65 ± 0.04
	4	375	0.43 ± 0.01	0.57 ± 0.03
	≥ 6	435	0.43 ± 0.01	0.44 ± 0.02
	all	1123	0.423 ± 0.008	0.56 ± 0.02
Λ	2	166	0.47 ± 0.02	-1.58 ± 0.08
	4	185	0.49 ± 0.02	-1.09 ± 0.07
	≥ 6	81	0.47 ± 0.03	-0.69 ± 0.08
	all	454	0.48 ± 0.01	-1.26 ± 0.05

order to improve statistics. The points from the pp-experiment at 19 GeV/c are given for comparison.

For K_s^0 -mesons the hypothesis of scaling in the mean is valid within errors for both transverse and longitudinal variables. For Λ -hyperons the scaling in the mean holds for the transverse variable, but for the longitudinal variable the shape of the distribution depends on multiplicity. For low multiplicities it is steeper. A similar effect has been also observed in the pp-experiment at 19 GeV/c.

The curves in Figs. 1, 2 and 3 describe the scaling in the mean for π^- -production in pp-interactions in an energy interval of 13-300 GeV/c^{1/}. One can see that the data points follow these curves closely. A similar feature has been observed for π^0 -mesons in π^-p -interactions at 5 GeV/c^{2a/}. Thus, one can see that scaling in the mean distributions for both p_T and p_L^* -variables are very similar for different types of mesons. In addition, for the p_T -variable these distributions are very alike for mesons and Λ -hyperons.

The mean values of the transverse and longitudinal momentum for K_s^0 and Λ are presented in the Table. As is seen, the absolute values of $\langle p_L^* \rangle_n$ for Λ -hyperons decrease with increasing n , i.e., the leading behaviour* of Λ 's is more pronounced for small multiplicities. So the kinematical limits for the variables $p_L^*/\langle p_L^* \rangle_n$ vary strongly for different n ** and scaling in the mean is violated at large negative values of $p_L^*/\langle p_L^* \rangle_n$ (for $p_L^*/\langle p_L^* \rangle_n \lesssim -1.8$ in Fig. 4). Thus, the violation of the law of scaling in the mean for Λ 's in this kinematical region is connected with the leading effect.

It has been proposed^{4/} that the scaling in the mean is a consequence of: a) Feynman scaling^{5/}, b) KNO-scaling^{6/} and c) factorization of the cross section over longitudinal and transverse momenta

$$\frac{d^2\sigma_n}{dp_L^* dp_T} = f_1(p_L^*, \frac{n}{\langle n \rangle}) \cdot f_2(p_T), \quad (4)$$

* The production of Λ 's mainly in the proton fragmentation region.

** The $p_L^*/\langle p_L^* \rangle_n$ kinematical limits are -1.94 for $n = 2$, -2.74 for $n = 4$, -4.22 for $n \geq 6$ and -2.43 for all events.




Fig.5. Distribution of the invariant function $\frac{1}{\sigma_{\text{tot}}} \frac{2}{\sqrt{s}} \int E + \frac{d^2\sigma}{dx dp_T^2} dp_T^2$ versus x for K_s^0 -mesons.

where $n/\langle n \rangle$ is the KNO-variable ^{6/}. In other words, it has been supposed that the hypothesis of scaling in the mean contains no new information. In order to check this assumption, we have tested the conditions a) and c) for K_s^0 inclusive production. Figure 5 shows the invariant normalized inclusive x -distributions (x is the Feynman variable) for K_s^0 production in our experiment and in pp -interactions at $19 \text{ GeV}/c$ ^{7/}. One can see that there is a significant difference between the presented distributions in the intermediate region of x , i.e., Feynman scaling does not hold for different initial states.

In fig.6 (a,b,c,d) we show the p_T^* -distributions of K_s^0 in different intervals of p_T for all events and separately for multiplicities 2, 4 and 6. It is seen that in all considered cases the shapes of the distributions change with p_T , i.e., the factorization condition (4) is not fulfilled. The same result has been found for pp -interactions at $19 \text{ GeV}/c$ ^{2b/}.

Thus, the results presented in figs.5 and 6 do not support the assumption that the scaling in the mean is only the consequence of the above laws a), b), c). Another picture of connection between the scaling laws is considered in papers ^{8/},




Fig. 6. p_L^* -distribution of K_B^0 for different intervals of p_T .

where it has been obtained that at asymptotic energies the validity of KNO-scaling and scaling in the mean leads to the violation of Feynman scaling. This violation in the central region (at $x = 0$) may correspond (with certain additional assumptions) to the rise of the cross section of meson production, which is also experimentally observed^{/9/} in pp-interactions up to ISR energies.

2. MULTIPLICITIES ASSOCIATED WITH STRANGE PARTICLES IN $\bar{p}p \rightarrow K_s^0 + X$ AND $\bar{p}p \rightarrow \Lambda + X$ REACTIONS AT 22.4 GEV/C

Associative multiplicities in inclusive processes have been studied as a function of missing mass M_x for this and other experiments^{/10/}. The following logarithmic dependence has been predicted^{/11/}.

$$n(M_x^2) = A + B \ln(M_x^2) \quad (5)$$

in the framework of the multiperipheral model for particle production in the fragmentation region. In eq.(5) B is the universal slope parameter. The prediction has been made assuming that a main contribution to these processes comes from the diagrams where system X is produced in the virtual interaction of some effective trajectory with one of the primary particles. The M_x^2 value is analogous to the c.m.s. energy squared for real processes.

In Fig.7 the average charged multiplicities $n(M_x^2)$ are presented as a function of M_x (to logarithmic scale) for inclusive production of K_s^0 and Λ . The fit to expression (5) (solid line in Fig.7) yields for the slope parameter of Λ 's $B = 1.26 \pm 0.14$ ($\chi^2/N_F = 1.5/6$). For K_s^0 -mesons the fit is not satisfactory ($\chi^2/N_F = 27/7$). For comparison with other results we have also made the same fit for Λ 's with $\cos \theta < -0.5$ ($\sim 85\%$ of the sample) where θ is the polar angle in the c.m.s. We got $B = 1.31 \pm 0.11$, which is almost the same as without the cut. Both values of the parameter B for Λ -particles agree rather well with the values obtained in other experiments^{/10/}.

Analogously to real processes, one can assume that for system X produced with Λ -particles the Wroblewski^{/12/} and KNO^{/13/} relations have been applied in the following way:

$$D(M_x^2) = a + b \ln(M_x^2) \quad (6)$$

* In Fig.7 these new data are denoted by circles if they do not coincide with those observed for all Λ 's (crosses in Fig.7).

and

$$\frac{\langle n(M_x^2) \rangle}{\sigma(M_x^2)} \sigma_n(M_x^2) = \Psi \left(\frac{n(M_x^2)}{\langle n(M_x^2) \rangle} \right), \quad (7)$$

where $D(M_x^2)$ is the square root of the dispersion of the associative multiplicity distribution, $\sigma(M_x^2)$ and $\sigma_n(M_x^2)$ are the total and topological cross sections for definite values of M_x^2 .

The experimental dependence of $D(M_x^2)$ on $\langle n(M_x^2) \rangle$ is shown in Fig.8 for different M_x^2 intervals. After fitting the experimental data by the function (6), we have obtained the parameters $a = 0.28 \pm 0.3$ and $b = 0.40 \pm 0.09$ (straight line in fig.8). In Fig.9 is shown the KNO dependence (7) for three intervals of M_x^2 . One can see that the KNO distributions for all M_x^2 intervals are similar.

An analytical expression for the associative KNO scaling has been obtained in the framework of renormalizable quantum field theory using the renormalization group method^{15/}. For system X with mass M_x we get

$$\Psi(Z(M_x^2)) = C \frac{\alpha^\alpha}{\Gamma(\alpha)} Z^{\alpha-1} (M_x^2) e^{-\alpha Z(M_x^2)}, \quad (8)$$

where $Z(M_x^2) = n(M_x^2)/\langle n(M_x^2) \rangle$, C is a normalization para-




Fig.7. Dependence of $\langle n(M_x^2) \rangle$ on M_x^2 (to logarithmic scale) for K_s^0 and Λ -particles. The straight line is the result of fitting the data for Λ -hyperons.

meter and $a = 1/b^2$ (b is the same as in eq.(6)). The curve in Fig.9 is the result of fitting our data by expression (8) with the parameter $C = 1.86 + 0.11 (\chi^2/N_F = 4.2/10)$. One can see that the data are quite well described by expression (8).




Fig.8. Dependence of $D(M_x^2)$ on $\langle n(M_x^2) \rangle$ for Λ 's for the intervals $\Delta M_x^2 = (4-8, 8-12, 12-16, 16-24, 24-30, 30-32)$ (GeV/c)². The straight line is the fitting result.




Fig.9. Plot of $\frac{\langle \sigma_n(M_x^2) \rangle \times \langle n(M_x^2) \rangle}{\sigma(M_x^2)}$ versus $n(M_x^2) / \langle n(M_x^2) \rangle$ for Λ particles. The curve is the result of fitting by the theoretical expression (8).

3. CONCLUSIONS

1. The scaling in the mean is shown to be valid in the variables $p_T / \langle p_T \rangle_n$, $p_L^* / \langle p_L^* \rangle_n$ for inclusive K_S^0 -production. The scaled distributions coincide with the data obtained in pp - and π^-p -interactions.
2. For Λ particles the scaling in the mean in the variable $p_L^* / \langle p_L^* \rangle_n$ is violated which can be due to the leading particle behaviour.
3. The dependence of the mean associative multiplicities on $\ln(M_x^2)$ for Λ -production is linear, the slope agreeing rather well with the values obtained in other experiments.
4. The associative KNO-scaling for Λ 's is observed in our experiment. The KNO distribution has the shape predicted by renormalizable quantum field theory.

The authors want to express their gratitude to the staff responsible for operation of the Serpukhov accelerator and of the beam channel number 9 and to the technical staff of the Ludmila HBC. We also thank the technicians and assistants at all laboratories for their excellent work.

REFERENCES

1. Dao F.T. et al. Phys.Rev.Lett., 1974, 33, p.389.
2. a) Amoglobeli N.S. et al. Yadernaya Fizika, 1978, 27, p.995; Angelov N.S. et al. Nucl.Phys., 1975, B97, p.219; b) Hagman V.-M. et al. Physica Scripta, 1976, 14, p.24; c) Sheng A. et al. Phys.Rev., 1975, 110, p.1733.
3. Batyunya B.V. et al. JINR, 1-11194, Dubna, 1977 (in Russian); Boos E.G. et al. JINR, E1-11665, Dubna, 1978; Yadernaya Fizika, 1979, 29, p.374.
4. Svensseon B.E.Y., Sollin L. Phys.Rev.Lett., 1975, 34, p.1199.
5. Feynman R.P. Phys.Rev.Lett., 1969, 23, p.1415.
6. Koba Z. et al. Phys.Lett., 1972, 38B, p.25.
7. Boggild H. et al. Nucl.Phys., 1973, B57, p.77.
8. Ernst W., Schmitt I. Lett. Nuovo Cim., 1976, 16, p.39; Vaes R.J. Phys.Rev.Lett., 1976, 36, p.821; Phys.Rev., 1977, D15, p.2627.
9. Chlyapnikov P.V. Plenary report at the 18th Int. Conf. on HEP. Tbilisi, 1976. JINR, D1,2-10400, Dubna, 1977.

10. a) Barish S.J. et al. Phys.Rev.Lett., 1973, 31, p.1080;
Winkelmann F.C. et al. Phys.Rev.Lett., 1974, 32, p.121;
Albrow M.G. et al. Nucl.Phys., 1976, B102, p.275;
Fong O. et al. Phys.Rev.Lett., 1976, 37, p.736;
Varreiro F. et al. Phys.Rev., 1978, D17, p.669;
b) Chlyapnicov P.V. et al. Phys.Lett., 1974, 52B, p.375;
Azhinenko I.V. et al. Yadernaya Fizika, 1977, 25, p.585;
Amaglobeli N.S. et al. Yadernaya Fizika, 1978, 28, p.1511;
Batyunya B.V. et al. JINR, E1-12117, Dubna, 1979.
11. Chan C.F. Phys.Rev., 1973, D8, p.179.
12. Wroblewski A. Acta Phys. Polon, 1973, 4B, p.857.
13. Kota Z., Nielsen H.B., Olesen P. Nucl.Phys., 1972, B40,
p.317.
14. Darbaidze Ya.Z. et al. JINR, P2-10429, Dubna, 1977.

Received by Publishing Department
on April 24 1980.

WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?

You can receive by post the books listed below. Prices - in US \$,
including the packing and registered postage

P1,2-7642	Proceedings of the International School of Young Physicists on High Energy Physics. Gomel, 1973.	10.53
D1,2-8405	Proceedings of the IV International Symposium on High Energy and Elementary Particle Physics. Varna, 1974.	3.89
P1,2-8529	The International School-Seminar of Young Scientists. Actual Problems of Elementary Particle Physics. Sochi, 1974.	4.70
D6-8846	XIV Symposium on Nuclear Spectroscopy and Nuclear Theory. Dubna, 1975.	3.70
E2-9086	Proceedings of the 1975 JINR-CERN School of Physics. Alushta, 1975.	16.00
D13-9164	Proceedings of the International Meeting on Proportional and Drift Chambers. Dubna, 1975.	10.00
D1,2-9224	Proceedings of the IV International Seminar on High Energy Physics Problems. Dubna, 1975.	4.80
	Proceedings of the VI European Conference on Controlled Fusion and Plasma Physics. Moscow, 1973, v.II.	15.00
D13-9287	Proceedings of the VIII International Symposium on Nuclear Electronics. Dubna, 1975.	6.6
D7-9734	Proceedings of the International School-Seminar on Reactions of Heavy ions with Nuclei and Synthesis of New Elements. Dubna, 1975.	4.86
D-9920	Proceedings of the International Conference on Selected Topics in Nuclear Structure. Dubna, 1976.	15.00
D2-9788	Nonlocal, Nonlinear and Nonrenormalizable Field Theories. Proceedings of the IVth International Symposium. Alushta, 1976.	4.20
09-10500	Proceedings of the Second Symposium on Collective Methods of Acceleration. Dubna, 1976.	11.00

D2-10533	Proceedings of the X International School on High Energy Physics for Young Scientists. Baku, 1976.	11.00
D13-11182	Proceedings of the IX International Symposium on Nuclear Electronics. Varna, 1977.	10.00
D10,11-11264	Proceedings of the International Meeting on Programming and Mathematical Methods for Solving the Physical Problems. Dubna, 1977.	13.00
D17-11490	Proceedings of the International Symposium on Selected Problems of Statistical Mechanics. Dubna, 1977.	18.00
D6-11574	Proceedings of the XV Symposium on Nuclear Spectroscopy and Nuclear Theory. Dubna, 1978.	4.70
D3-11787	Proceedings of the III International School on Neutron Physics. Alushta, 1978.	12.00
D13-11807	Proceedings of the III International Meeting on Proportional and Drift Chambers. Dubna, 1978.	14.00
	Proceedings of the VI All-Union Conference on Charged Particle Accelerators. Dubna, 1978. 2 volumes.	25.00
D1,2-12036	Proceedings of the V International Seminar on High Energy Physics Problems. Dubna, 1978.	15.00
D1,2-12450	Proceedings of the XII International School on High Energy Physics for Young Scientists. Bulgaria, Primorsko, 1978.	18.00
R2-12462	Proceedings of the V International Symposium on Nonlocal Field Theories. Alushta, 1979.	9.50
D2-11707	Proceedings of the XI International School on High Energy Physics and Relativistic Nuclear Physics for Young Scientists. Gomei, 1977.	14.00

Orders for the above-mentioned books can be sent at the address:

Publishing Department, JINR

Head Post Office, P.O. Box 79 101000 Moscow, USSR



Издательский отдел Объединенного института ядерных исследований.
Заказ 28153. Тираж 620. Уч.-изд. листов 1,33.
Редактор Э.В.Ивашевич.
Набор Л.М.Крыловой, Н.И.Коротковой.
Макет В.А.Халякиной. Подписано к печати 2.06.80 г.