


METHANE FORMATION FROM THE REACTIONS OF HYDROXYL RADICALS  
AND HYDROGEN ATOMS WITH DIMETHYL SULFOXIDE (DMSO).


A.M. KOULKES-PUJO, M. MOREAU and J. SUTTON  
CNRS and DPC/SCM/CEN.SACLAY, 91191 GIF SUR YVETTE CEDEX (France).

Miller conference 1981  
Windermere, UK 12 - 16 April 1981  
CEA-CONF-- 6297


DMSO is known to possess important radioprotective and antiinflammatory properties which are generally associated with its rapid reaction with OH radicals. These may originate in chemical processes, for example the Fenton reaction ( $\text{Fe}^{2+} + \text{H}_2\text{O}_2 \rightarrow \text{Fe}^{3+} + \text{OH} + \text{OH}^-$ ), the Haber-Weiss reaction ( $\text{O}_2^- + \text{H}_2\text{O}_2 \rightarrow \text{OH} + \text{OH}^- + \text{O}_2$ ), in biological processes such as microsomal electron transfer reactions(1,2,3), and in the radiolysis of aqueous systems. The rate constant of the reaction is  $4.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  in neutral media (4). It is generally accepted that the first step in the reaction mechanism is the formation of a radical adduct (5)



followed by, for example



or



where RH may be another molecule of DMSO.

If the stoichiometry is simple the methane yield should be equal to that of the OH radicals consumed. This formation of methane is important in biological research as it serves to estimate the OH production by liver microsomal reduction of oxygen and to determine the effect of other chemicals on this enzymatic process.

The radiolysis of aqueous DMSO solutions would appear to be a method well suited for establishing this stoichiometry : the yield of OH radicals in water is known ( $G_{OH} = 2.95$ ) (6) and by adding a second OH' scavenger such as the bromide ion it is possible to study the competition kinetics. The yield of methane should then vary according to the relation (7)

$$G(CH_4)^{-1} = G_{OH}^{-1} \left( 1 + \frac{k_1[Br^-]}{k_2[DMSO]} \right)$$

where  $k_1$  and  $k_2$  represent, respectively, the rate constants for the reactions of  $Br^-$  and DMSO with OH radicals.

It has been previously shown (8) that methane is formed in pure DMSO by the reaction of hydrogen atoms and the question arises : does this reaction also occur in the radiolysis of aqueous DMSO mixtures giving a second route to methane production ?

This question can be answered by studying the effect of a second H-atom scavenger on the methane yield. Ethanol has been selected in the present work since, in acidic media, it reacts with H to give  $H_2$  but no  $CH_4$  (9). Conversely, the radiolysis of acidic aqueous solutions of DMSO leads to methane formation with only a constant low yield of (molecular) hydrogen ( $G_{H_2} = 0,42$ ). Thus the hydrogen yield  $G(H_2)$  formed by the competition between the two solutes for the radiolytic yield of H atoms will be given by

$$\Delta G(H_2)^{-1} = G_H^{-1} \left( 1 + \frac{k_3[DMSO]}{k_4[EtOH]} \right)$$

where  $\Delta G(H_2) = G(H_2) - G_{H_2}$ ,  $k_3$  and  $k_4$  are the rate constants for the reaction of hydrogen with DMSO and EtOH respectively.

## 2 - EXPERIMENTAL

Acidic, aqueous, deaerated solutions of pure DMSO, of DMSO-KBr and of DMSO-EtOH in completely filled glass ampoules were irradiated with a  $^{60}Co$   $\gamma$  source. The dose rate was  $2.50 \times 10^{21} \text{ ev } \ell^{-1} \text{ h}^{-1}$ . The gases formed were extracted under vacuum and analysed by gas chromatography on 5 m. columns packed with molecular seive 5A or 13X or Porapak Q. Argon and helium were used as carrier gases and the principal gaseous products were  $CH_4, H_2, C_2H_6$  : traces of  $CO_2$  and  $C_2H_4$  were also detected.

3 - RESULTS

Figure 1 shows the  $\text{CH}_4$  yield from water-DMSO mixtures plotted against the DMSO concentration. One observes that  $G(\text{CH}_4)$  does not reach a true plateau but continues to increase gently at higher  $[\text{DMSO}]$ . The curve giving  $G(\text{C}_2\text{H}_6)$  is similar in form but the values are much lower and less precise as the yields decrease with the total dose absorbed. The plateau values are slightly lower than the values obtained by Hart (10) for neutral media,  $G(\text{CH}_4) = 2.25$  ;  $G(\text{C}_2\text{H}_6) = 0.45$ .

Both hydrogen and methane yields were measured in the DMSO-EtOH mixtures. In figure 2,  $\Delta G(\text{H}_2)^{-1}$  is plotted against  $[\text{DMSO}]/[\text{EtOH}]$ . The points are fairly well aligned and the plot leads to values of  $G_{\text{H}} = 3.65$  and  $k_3/k_4 = 0.57$ . The former agrees with the literature value for  $G_{\text{H}}$  in acid solution and the latter leads to a value of  $k_3 = 2.6 \pm 0.3 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$  taking  $k_4 = 4.6 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$  (11). Thus DMSO is a relatively efficient H-atom scavenger. However, it is evident from the difference between the plateau value  $G(\text{CH}_4) = 1.8$  (figure 1) and  $G_{\text{H}} = 3.65$  that all the hydrogen atoms captured by DMSO do not give rise to methane formation. Moreover, the plot of  $G(\text{CH}_4)^{-1}$  against  $[\text{EtOH}]/[\text{DMSO}]$  is not linear, showing that the methane production mechanism is more complex than the simple reaction of H atoms with DMSO.

The other possible source of  $\text{CH}_4$  being the OH radical reaction with DMSO (1,10) , in order to determine the contribution of this process the methane yields were measured at different DMSO concentrations in the presence of  $7 \times 10^{-2} \text{ M. KBr}$ . The rate constants of the reactions  $\text{OH} + \text{Br}^- \rightarrow \text{Br} + \text{OH}^-$  and  $\text{OH} + \text{DMSO} \rightarrow \text{products}$ , being respectively  $2 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$  in acid solution<sup>(12)</sup> and  $7.4 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  in neutral solution<sup>(12)</sup> the former process is probably largely favoured for  $[\text{DMSO}] \leq 6 \times 10^{-2} \text{ M}$ .

The results are shown in figure 1. The methane yield has been reduced to a constant value of 0.85 which therefore represents the contribution of the  $\text{H} + \text{DMSO}$  reaction to  $\text{CH}_4$  production. The difference between the total  $\text{CH}_4$  yields and those measured in the presence of  $7 \times 10^{-2} \text{ M KBr}$  corresponds to the contribution furnished by the  $\text{OH} + \text{DMSO}$  reaction. Thus in neither

\* extrapolated from values given in NSRDS-NBS 59 (1977) n°3-12.

case, H or OH, does the total radical yield lead to methane production, which would otherwise have a maximum value equal to  $G_H + G_{OH} = 2.95 + 3.65 = 6.6$ , much greater than the value  $G(CH_4) = 1.8$  observed.

In order to determine  $k_2$  in acid media methane yields were determined in solutions containing  $1.4 \times 10^{-2} M$  DMSO,  $1.5 \times 10^{-3} M < KBr^- < 7 \times 10^{-2} M$ , and  $5 \times 10^{-1} M$  sulfuric acid. Figure 3 represents the function  $\Delta G(CH_4)^{-1}$  plotted against  $[Br^-]/[DMSO]$  where  $\Delta G(CH_4) = G(CH_4) - 0.85$ . From the linear plot one obtains  $G(OH) = 0.69$  and  $k_4 = 2.0 \pm 0.2 \times 10^{10} M^{-1} s^{-1}$  in acid solution. This value of the rate constant is 5 times greater than that found in neutral solution: a similar increase in the rate constant of the reaction  $OH + Br^-$  is observed in going from neutral to acid medium ( $k_1 = 2 \times 10^9 M^{-1} s^{-1}$  at pH = 7:  $2 \times 10^{10} M^{-1} s^{-1}$  at pH = 0).

Thus whereas both H and OH radicals react with DMSO to give methane, only a fraction of each species captured leads to this product ( $0.69/2.95 = 23\%$  of OH radicals:  $0.85/3.65 = 23\%$  of H atoms). These results suggest that the radical adducts  $(CH_3)_2S \begin{matrix} \swarrow O^* \\ \searrow OH \end{matrix}$  and  $(CH_3)_2S^*OH$  react in several different ways, methane production being a relatively unimportant path.

It is also evident that the maximum methane yield deriving from the H atom and OH radical reactions ( $0.85 + 0.69 = 1.54$ ) is significantly less than the experimental value ( $G(CH_4) = 1.8$ ) suggesting the existence of another source of methane as has been previously suggested (8).

The present results thus show that, taken alone, the methane production is not a reliable or sufficient test for the identification of OH radicals in biological systems.

#### CONCLUSION

It has been shown that both hydrogen atoms and hydroxyl radicals react with dimethylsulfoxide in aqueous acid solution to produce methane. In both cases however only a fraction of the radicals

captured ( $\sim 23$  % of each under the experimental conditions employed) gives rise to this product. Thus methane formed in the reaction of DMSO with biological systems is not unequivocal proof of the presence of OH radicals ; nor is the yield of methane a direct measure of an OH yield.

The rate constants  $k(\text{H} + \text{DMSO}) = 2.6 \times 10^7 \text{M}^{-1} \text{s}^{-1}$  and  $k(\text{OH} + \text{DMCO}) = 2.1 \times 10^{10} \text{M}^{-1} \text{s}^{-1}$  have been determined in the presence of  $5 \times 10^{-1} \text{M}$  sulfuric acid.

#### ACKNOWLEDGEMENTS.

We wish to thank Dr. E.J. HART for useful discussion and for furnishing some unpublished results on the radiolysis of aqueous DMSO solutions.

R E F E R E N C E S

---

- (1) COHEN G., CEDERBAUM A.I. (1979) Science 204, 66-67.
- (2) COHEN G., CEDERBAUM A.I. (1980) Arch. Biophys. 199, 2, 438-447.
- (3) KLEIN S.M., COHEN G., CEDERBAUM A.I. (1980) FEBS Letters 116, 2, 220-222.
- (4) MEISSNER G., HENGLEIN A., BECK G. (1967) Z. Naturforschg. 22, B, 13-19.
- (5) GILBERT B.C., NORMAN R.O.C., SEALY R.C., J.C.S. Perkin II (1975) 303-308.  
SYMONS Martyn C.R. (1976) J.C.S. Perkin II (1976) 908-915.
- (6) HART E.J., ANBAR M. (1970) THE HYDRATED ELECTRON, WILEY Inter science Ed. p. 18.
- (7) ALLEN A.O. (1961) THE RADIATION CHEMISTRY OF WATER AND AQUEOUS Solutions, VAN NOSTRAND COMPANY Ed. p. 29.
- (8) KOULKES-PUJO A.M., BERTHOU M. (1969) J. Chim. Phys. 66, 1178-1179.
- (9) SEDDON W.A., Allen H.O. J. Phys. Chem. (1967) 71, 1914-1918.
- (10) HART E. private communication from BROWN W.G., HART E.J. and CLARKE R.M.
- (11) NSRDS-NBS 51 (1975) ref. 68.0525 HENTZ R.R., FARHATAZIZ M.D.J. (1968) J. Chem. Phys. 49, 2153.
- (12) NSRDS-NBS 59 (1977) n° 3 - 12.
- (13) VETWISER D, JANATA E, ASMUS v. D. (1980) J.C.S. Perkin II, 146-153

L E G E N D S

Fig. 1 - ● CH<sub>4</sub> yield  
+ CH<sub>4</sub> yield (in presence of  $7 \times 10^{-2} \text{M Br}^-$ )  
● C<sub>2</sub>H<sub>6</sub> yield  
versus DMSO concentration.

Fig. 2 - Reciprocal of  $G(\text{H}_2) - 0,42$  versus  
[DMSO] / [EtOH]

Fig. 3 - Reciprocal of  $G(\text{CH}_4) - 0.85$  versus  
[Br<sup>-</sup>] / [DMSO].




Fig.1



