



(11) (A) No. 1 142 915

(45) ISSUED 830315

(52) CLASS 260-28.2

(51) INT. CL. C07J 1/00³

(19) (CA) **CANADIAN PATENT** (12)

(54) STEROID DERIVATIVES AND THEIR USE IN
RADIOIMMUNOASSAYS

(72) Varma, Ravi K.;
Chao, Sam T.,
U.S.A.

(73) Granted to Squibb (E. R.) & Sons, Inc.
U.S.A.

(21) APPLICATION No. 326,279

(22) FILED 790425


(30) PRIORITY DATE U.S.A. (908,294) 780522

No. OF CLAIMS 11 - NO DRAWING

Canada

ABSTRACT
STEROID DERIVATIVES AND THEIR USE
IN RADIOIMMUNOASSAYS

Radiolabeled steroid derivatives having the formula




wherein St is a 6-dehydro derivative of a 3-oxo-4,5-dehydro steroid intended for radioimmunoassay, said steroid being saturated in the 1,2-position; R is hydrogen or alkyl of 1 to 3 carbon atoms; n is 0, 1, 2, 3 or 4; and the asterisk (*) indicates tagging with a radioisotope, are useful as tracers in radioimmunoassays. The unlabeled analogs can be coupled with an immunogenic carrier and used to induce antibody formation in animals.

STEROID DERIVATIVES AND THEIR USE
IN RADIOIMMUNOASSAYS

Radioimmunoassay techniques have been used to determine the concentration in body fluids of various endogenous and exogenous steroids. In the development of radioimmunoassays for the various steroids, the preparation of a radiolabeled antigen is of primary concern. Possible radioisotope labels are tritium, carbon-14, iodine-125, iodine-131 and others. However, because tritium and carbon-14 must be counted by liquid scintillation (a time-consuming and expensive process), iodine-125 and iodine-131 are more desirable. For reasons well-recognized in the art (e.g., half-life, radiation hazard, counting efficiency and others) iodine-125 has become the radioisotope of choice for use in steroid radioimmunoassays.

The chemical structure of steroids is such that it is generally not possible to radioiodinate them directly. It is necessary, therefore, to utilize as a precursor of the radiolabeled antigen a derivative of the steroid to be assayed which can be readily iodinated. In choosing or developing such a derivative, the primary concern is the affinity of the derivative for the antibodies of the steroid to be assayed; the affinity of the derivative for the antibodies should, of course, be as close to the affinity of the steroid for the antibodies as possible.

Compounds having the formula




are readily tagged with a radioisotope and can be used (when radiolabeled) as a tracer in radioimmunoassay procedures for the determination of steroid levels in a body fluid. The unlabeled compounds can be coupled with
 5 an immunogenic carrier and suitable adjuvant and injected into animals to induce antibody formation. In formula I, and throughout the specification, R is hydrogen or an alkyl group of 1 to 3 carbon atoms; n is 0, 1, 2, 3 or 4; and St is a 6-dehydro derivative of a 3-oxo-4,5-dehydro
 10 steroid intended for radioimmunoassay, said steroid being saturated in the 1,2-position.

Exemplary of 3-oxo-4,5-dehydro steroids intended for radioimmunoassay which can be modified structurally as shown in formula I are cortisol, cortisone, corticosterone,
 15 testosterone, 19-nortestosterone and methyltestosterone.

The steroids of formula I can be prepared using the corresponding 6-hydrosteroid as the starting material. Many of these starting steroids (St-H) will be 17 α ,21-dihydroxy pregnenes. In these instances, it will be
 20 necessary to protect the substituents in the 17-, 20- and 21-positions prior to running the reactions needed to prepare the steroids of formula I. Various methods for protecting these substituents will be apparent to the practitioner of this invention. An exemplary method
 25 comprises the reaction of the unprotected steroid with paraformaldehyde in the presence of hydrochloric acid to yield the corresponding 17,20:20,21-bis-methylenedioxy derivative.


Reaction of the 6-hydrosteroid starting material (protected, if necessary) with formaldehyde in the
 30 presence of a secondary amine, e.g., pyrrolidine, affords a mixture of 6 α -and 6 β -hydroxymethyl steroids having the formula



these derivatives can be separated using conventional techniques. In formula II, and throughout the specification, St' can be St or the corresponding protected steroid moiety. The above reaction can be carried out in an alcohol solvent, preferably in an inert atmosphere, e.g., nitrogen or argon. The absence of oxygen during the reaction minimizes the formation of significant amounts of 6 α - and 6 β - hydroxy steroids as by-products.


Alternatively, the 6-hydrosteroid starting material (protected if necessary) can be reacted with pyrrolidine in an alcoholic solvent to yield III 3-[(1-pyrrolidinyl)] St'''-3,4,5,6-dehydro, wherein St''' is a steroid saturated in the A and B rings. Compounds of this type are known; see, for example, Helvetica Chimica Acta, 56(7):2396-2404 (1973). An intermediate of formula III can then be reacted with formaldehyde in an alcoholic solvent to yield the corresponding 6 α - and 6 β - hydroxymethyl steroids of formula II.

The 6 α - and 6 β - hydroxymethyl steroids of formula II can be reacted with a glutaric anhydride derivative having the formula



30 to yield the corresponding steroid having the formula

V




5

The reaction can be run in the presence of an organic base (e.g., a nitrogen containing heterocyclic such as pyridine or a tertiary amine such as triethylamine), preferably at an elevated temperature.

Removal of the phenolic hydroxyl protecting group in a compound of formula V (and, where necessary, removal of the protecting group from the 17 β , 20- and 21-positions) yields the corresponding steroid product of formula I.


The anhydrides of formula IV can be prepared by first reacting a 4-methoxyphenyl aldehyde having the formula

VI



with at least 2 molar equivalents of cyanoacetic acid in the presence of a base (e.g., sodium hydroxide) to yield, on acid hydrolysis, a compound having the formula


VII



30

An alternative preparation for the compound of formula VII wherein n is 0 and R is hydrogen, *i.e.*, 3-(4-methoxyphenyl)glutaric acid, is disclosed by Smith et al., *J.A.C.S.*, 72, 1877 (1950). In that procedure, anisaldehyde is condensed with ethyl acetoacetate in the presence of piperidine to give ethyl anisal-bisacetoacetate. Cleavage of this product to give the desired 3-(4-methoxyphenyl)glutaric acid can be accomplished with boiling alcoholic sodium hydroxide solution.

Demethylation of the glutaric acid derivatives of formula VII results in glutaric acid derivatives having the formula




and can be accomplished by following one of the several procedures known in the art for the demethylation of aryl methyl ethers. One such procedure, described by Feutrill et al., *Aust. J. Chem.*, 25, 1719 (1972), involves the treatment of the aryl methyl ether with thioethoxide ion (readily prepared *in situ* from ethanethiol and sodium hydride) in a polar aprotic solvent, preferably dimethylformamide.

The phenolic hydroxy group of a compound of formula VIII can be protected with an alkanoyl group using art-recognized procedures. One such procedure comprises reacting the glutaric acid derivative with the appropriate acid anhydride (acetic anhydride is preferred). The preferred method of preparing a glutaric anhydride derivative of formula IV from the glutaric acid derivative of formula VIII

is to combine the conversion of the acid to anhydride and the protection of the phenolic hydroxy group into a single step. When the R₁ protecting group is acetyl, this would involve heating a glutaric acid derivative of formula VIII in acetic anhydride.

The compounds of formula I can be coupled with an immunogenic carrier, such as a high molecular weight protein of which bovine serum albumin and thyroglobulin are exemplary, and if necessary an adjuvant in order to produce a substance capable of inducing antibody formation in animals. Procedures for such couplings are well known in the art; see, for example, Parker, "Radioimmunoassay of Biologically Active Compounds", Prentice-Hall, Inc., New Jersey (1976).

The compounds of formula I can be labeled ("tagged") with a radioisotope, preferably iodine-125 or iodine-131, and most preferably iodine-125, using procedures well known in the art, to yield a radio-labeled hapten having the formula



The asterisk (*) in formula IX indicates tagging with a radioisotope. Exemplary of the methods known in the art is the method of Hunter and Greenwood; see

Nature, 194:495 (1962). The radiolabeled compounds of formula IX form an integral part of this invention.

5 The radiolabeled compounds of formula IX can be used as tracers in radioimmunoassay procedures following the general principles set forth in the Background of the Invention, supra. Exemplary detailed procedures are described in Jaffe et al., "Methods of Hormone Radioimmunoassay", Academic Press, 10 New York (1974) and Berson et al., "Methods in Investigative and Diagnostic Endocrinology", Vol. 3 on "Steroid Hormones", North Holland, Amsterdam (1975). The radiolabeled compounds of this invention may also be used as reagents in the automated radio- 15 immunoassay system of Brooker et al. disclosed in United States patent 4,022,577, issued May 10, 1977.

The following examples are specific embodiments of this invention.

Example 1

(6 α , 11 β)-6-[[DL-4-Carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]methyl]-11,17,21-trihydroxypregn-4-ene-3,20-dione (cortisol derivative)

- 5 A) 11 β -Hydroxy-6-(hydroxymethyl)-17,20:20,21-bis-[methylenebis(oxy)]pregn-4-en-3-one (6 α and 6 β -isomers)

10 A suspension of 11 β -hydroxy-17,20:20,21-bis[methylenebis(oxy)]pregn-4-en-3-one (1.6g) in methanol (50 ml) containing pyrrolidine (1.1g) is refluxed until a homogeneous solution is obtained. The solution is then cooled to room temperature and a 37% formaldehyde solution (5.0 ml) is added. The solution is then refluxed for 15 minutes and the solvents are evaporated to afford 1.7g of a gummy residue.

15 The above reaction is repeated two more times on 3.6g each of the starting steroid. The materials from the three runs are then pooled and chromatographed on a silica gel column to isolate (i) 6.8g of the starting steroid, (ii) 1.05g of a mixture of the 6 α and 6 β -isomers of the title compound, and (iii) 0.8g of a mixture of the 6 α and 6 β -isomers of 6,11 β -dihydroxy-17,20:20,21-bis-[methylenebis(oxy)]pregn-4-en-3-one.

25 The 6 α and 6 β -isomers of the 6-hydroxymethyl steroid (1.05g) are separated by repeated thin-layer chromatography on silica gel plates to afford 0.25g of the 6 α -isomer, melting point 228-245°C, and 0.74g of the 6 β -isomer, melting point 175-185°C.

30

B) (6 α ,11 β)-6-[[DL-3-[4-(Acetyloxy)phenyl]-4-carboxy-1-oxobutoxy]methyl]-11-hydroxy-17,20:20,21-bis-[methylenebis(oxy)]pregn-4-en-3-one

5 A solution of 105 mg of 11 β -hydroxy-6 α -
(hydroxymethyl)-17,20:20,21-bis[methylenebis(oxy)]-
pregn-4-en-3-one in 0.8 ml of dry pyridine containing
180 mg of 3-[4-(acetyloxy)phenyl]glutaric anhydride
is heated in a bath at 70-80°C for 3.0 hours. The
pyridine is then evaporated in vacuo, the residue
10 is dissolved in ethyl acetate, washed with 5%
hydrochloric acid and water, and evaporated to a
residue. This is subjected to thin-layer chroma-
tography on silica gel plates to isolate 113 mg of
the title compound, melting point 117-124°C.

15 C) (6 α ,11 β)-6-[[DL-4-Carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]methyl]-11-hydroxy-17,20:20,21-bis-[methylenebis(oxy)]pregn-4-en-3-one

A solution of 110 mg of (6 α ,11 β)-6-[[DL-3-
[4-(acetyloxy)phenyl]-4-carboxy-1-oxobutoxy]methyl]-
20 11-hydroxy-17,20:20,21-bis[methylenebis(oxy)]pregn-
4-en-3-one in 8 ml of methanol is adjusted to pH 9.0
with triethylamine and refluxed in an oil bath for
22 hours. The mixture is then cooled, acidified
with 5% hydrochloric acid, concentrated in vacuo
25 to remove most of the methanol, diluted with water,
and extracted with ethyl acetate. The ethyl acetate
solution is washed with a saturated sodium chloride
solution, dried and evaporated to afford 95 mg of
the title compound, melting point 105-118°C. A
30 thin-layer chromatographic examination shows that
this material contains only traces of impurities.

D) (6 α ,11 β)-6-[[DL-4-Carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]methyl]-11,17,21-trihydroxypregn-4-ene-3,20-dione

5 A solution of 60 mg of (6 α ,11 β)-6-[[DL-4-carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]methyl]-11-hydroxy-17,20:20,21-bis[methylenebis(oxy)]pregn-4-en-3-one in 4 ml of 50% acetic acid is heated at 100°C for 4.0 hours under nitrogen. The solvents are then evaporated in vacuo and the residue is
10 subjected to preparative thin-layer chromatography on silica gel plates to afford 30 mg of the title compound, melting point 103-128°C.

Example 2

15 (6 β ,11 β)-6-[[DL-4-Carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]methyl]-11,17,21-trihydroxypregn-4-ene-3,20-dione

A) (6 β ,11 β)-6-[[DL-3-[4-(Acetyloxy)phenyl]-4-carboxy-1-oxobutoxy]methyl]-11-hydroxy-17,20:20,21-bis-[methylenebis(oxy)]pregn-4-en-3-one

20 A solution of 11 β -hydroxy-6 β -(hydroxymethyl)-17,20:20,21-bis[methylenebis(oxy)]pregn-4-en-3-one (300 mg) is reacted with 3-[4-(acetyloxy)phenyl]glutaric anhydride (515 mg) following the procedure described in Example 1B to afford 295 mg of the title compound,
25 melting point 99-119°C.

B) (6 β ,11 β)-6-[[DL-4-Carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]methyl]-11-hydroxy-17,20:20,21-bis-[methylenebis(oxy)]pregn-4-en-3-one

30 (6 β ,11 β)-6-[[DL-3-[4-(Acetyloxy)phenyl]-4-carboxy-1-oxobutoxy]methyl]-11-hydroxy-17,20:20,21-bis-[methylenebis(oxy)]pregn-4-en-3-one (325 mg) in

methanol (18ml is reacted following the procedure described in Example 1C to afford 305 mg of the title compound, melting point 103-128°C.

5 C) (6 β ,11 β)-6-[[DL-4-Carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]methyl]-11,17,21-trihydroxypregn-4-ene-3,20-dione

10 A solution of 100 mg of (6 β ,11 β)-6-[[DL-4-carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]methyl]-11-hydroxy-17,20:20,21-bis[methylenebis(oxy)]pregn-4-en-3-one is reacted in 4.0 ml of 50% acetic acid following the procedure described in Example 1D to afford 26 mg of the title compound, melting point 177-205°. Spectral examination shows that this material is contaminated with a small amount of the 15 6 α -isomer.

Examples 3-9

Following the procedures of Examples 1 and 2, but substituting the starting steroid listed in column I for the cortisol derivative of Example 1A and the 20 anhydride reagent listed in column II for 3-[4-(acetyloxy)phenyl] glutaric anhydride, yields the α - and β -isomers compound listed in column III. In those instances wherein the starting steroid is not a bis-methylenedioxy derivative, the procedures of 25 Examples 1D and 2C will, of course, not be necessary.

	Column I	Column II	Column III
3	17,20:20,21-bis-[methylene-bis(oxy)]pregn-4-ene-3,11-dione	3-[4-(acetyloxy)phenyl]-glutaric anhydride	6-[[DL-4-carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]-methyl]-17,21-dihydroxypregn-4-ene-3,11,20-trione
4	11 β ,21,dihydroxypregn-4-ene-3,20-dione	3-[4-(acetyloxy)phenyl]-glutaric anhydride	6-[[DL-4-carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]-methyl]-11 β ,21-dihydroxypregn-4-ene-3,20-dione
5	pregn-4-ene-3,20-dione	3-[4-(acetyloxy)-3-methylphenyl]glutaric anhydride	6-[[DL-4-carboxy-3-(4-hydroxy-3-methylphenyl)-1-oxobutoxy]methyl]pregn-4-ene-3,20-dione
6	17 β -hydroxyandrost-4-en-3-one	3-[[4-(acetyloxy)phenyl]-methyl]glutaric anhydride	6-[[DL-4-carboxy-3-[(4-hydroxyphenyl)methyl]-1-oxobutoxy]-methyl]-17 β -hydroxyandrost-4-en-3-one

1142915


-12-

RB53

7	19-nor-17 β -hydroxyandrost-4-en-3-one	3-[2-[4-(acetyloxy)phenyl]-ethyl]glutaric anhydride	19-nor-6-[[DL-4-carboxy-3-[2-(4-hydroxyphenyl)-ethyl]-1-oxobutoxy]-methyl]-17 β -hydroxyandrost-4-en-3-one
8	17 β -hydroxy-17 α -methyl-androst-4-en-3-one	3-[3-[4-(acetyloxy)phenyl]-propyl]glutaric anhydride	6[[DL-4-carboxy-3-[3-(4-hydroxyphenyl)propyl]-1-oxobutoxy]methyl]-17 β -hydroxy-17 α -methyl-androst-4-en-3-one
9	11 β -hydroxy-17,20:20,21-bis[methylenebis(oxy)]-pregn-4-en-3-one	3-[4-[4-(acetyloxy)phenyl]-butyl]glutaric anhydride	6-[[DL-4-carboxy-3-[4-(4-hydroxyphenyl)butyl]-1-oxobutoxy]methyl]-11 β ,17,21-trihydroxypregn-4-ene-3,20-dione

What is claimed is:


1. A steroid having the formula




wherein St is a 6-dehydro derivative of a 3-oxo-4,5-dehydro steroid intended for radioimmunoassay, said steroid being saturated in the 1,2-position; R is hydrogen or alkyl of 1 to 3 carbon atoms; and n is 0, 1, 2, 3 or 4; and radiolabeled derivatives thereof.

2. A steroid in accordance with claim 1 wherein R is hydrogen.
3. A steroid in accordance with claim 1 wherein R is alkyl of 1 to 3 carbon atoms.
4. A steroid in accordance with claim 1 wherein n is 0.
5. The steroid in accordance with claim 1 having the name (6 α ,11 β)-6-[[DL-4-carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]methyl]-11,17,21-trihydroxypregn-4-ene-3,20-dione.
6. The steroid in accordance with claim 1 having the name (6 β ,11 β)-6-[[DL-4-carboxy-3-(4-hydroxyphenyl)-1-oxobutoxy]methyl]-11,17,21-trihydroxypregn-4-ene-3,20-dione.
7. A radiolabeled derivative in accordance with claim 1.
8. A radiolabeled derivative in accordance with claim 1 wherein the radiosotope is iodine-125.


9. A process for preparing a steroid having the formula



wherein St is a 6-dehydro derivative of a 3-oxo-4,5-dehydro steroid intended for radioimmunoassay, said steroid being saturated in the 1,2-position; R is hydrogen or alkyl of 1 to 3 carbon atoms; and n is 0, 1, 2, 3 or 4, characterized by reacting a steroid having the formula



wherein St is a steroid as hereinabove defined, or in a protected form, with a compound of the formula



wherein the phenolic hydroxy group is protected, and removing said protecting groups by conventional methods.

10. A process according to claim 9 wherein the product is radiolabeled.

11. A process according to claim 9 wherein the product is radiolabeled with iodine-125.



SUBSTITUTE

REPLACEMENT

SECTION is not Present

Cette Section est Absente