



INTERNATIONAL ATOMIC ENERGY AGENCY

**FOURTEENTH INTERNATIONAL CONFERENCE ON PLASMA  
PHYSICS AND CONTROLLED NUCLEAR FUSION RESEARCH**

Würzburg, Germany, 30 September - 7 October 1992

IAEA-CN-56/E-2-3

**The New Features of Ion Bernstein Wave Heating  
in JIPP T-IIU Tokamak**

T. Watari, R. Kumazawa, T. Seki, Y. Yasaka, A. Ando, Y. Oka, O. Kaneko, K. Adati,  
R. Akiyama, Y. Hamada, S. Hidekuma, S. Hirokura, K. Ida, K. Kawahata,  
T. Kawamoto, Y. Kawasumi, S. Kitagawa, M. Kojima, T. Kuroda, K. Masai,  
S. Morita, K. Narihara, Y. Ogawa, K. Ohkubo, S. Okajima, T. Ozaki, M. Sakamoto,  
M. Sasao, K. Sato, K.N. Sato, F. Shimpo, H. Takahashi, S. Tanahasi, Y. Taniguchi,  
K. Toi, T. Tsuzuki and M. Ono

(Received - Sep. 7, 1992)

NIFS-181

Sep. 1992

This is a preprint of a paper intended for presentation at a scientific meeting. Because of the provisional nature of its content and since changes of substance or detail may have to be made before publication, the preprint is made available on the understanding that it will not be cited in the literature or in any way be reproduced in its present form. The views expressed and the statements made remain the responsibility of the named author(s); the views do not necessarily reflect those of the government of the designating Member State(s) or of the designating organization(s). In particular, neither the IAEA nor any other organization or body sponsoring this meeting can be held responsible for any material reproduced in this preprint.

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Nagoya 464-01, Japan.



INTERNATIONAL ATOMIC ENERGY AGENCY

**FOURTEENTH INTERNATIONAL CONFERENCE ON PLASMA  
PHYSICS AND CONTROLLED NUCLEAR FUSION RESEARCH**

Würzburg, Germany, 30 September – 7 October 1992

---

IAEA-CN-56/E-2-3

**THE NEW FEATURES OF ION BERNSTEIN WAVE HEATING  
IN JIPP T-IIU TOKAMAK**

T. WATARI, R. KUMAZAWA, T. SEKI, Y. YASAKA\*, A. ANDO, Y. OKA,  
O. KANEKO, K. ADATI, R. AKIYAMA, Y. HAMADA, S. HIDEKUMA,  
S. HIROKURA, K. IDA, K. KAWAHATA, T. KAWAMOTO, Y. KAWASUMI,  
S. KITAGAWA, M. KOJIMA, T. KURODA, K. MASAI, S. MORITA,  
K. NARIHARA, Y. OGAWA\*\*\*, K. OHKUBO, S. OKAJIMA\*\*, T. OZAKI,  
M. SAKAMOTO, M. SASAO, K. SATO, K. N. SATO, F. SHIMPO,  
H. TAKAHASHI, S. TANAHASI, Y. TANIGUCHI, K. TOI, T. TSUZUKI

National Institute for Fusion Science, Nagoya 464-01, Japan

\* Department of Engineering, Kyoto University, , Kyoto, 606, Japan

\*\* Department of Applied Physics, Chubu University, Kasugai, 487, Japan

M. ONO,

Princeton Plasma Physics Laboratory, Princeton, New Jersey, 08543, U.S.A.

THE NEW FEATURES OF ION BERNSTEIN WAVE HEATING  
IN JIPP T-IIU TOKAMAK

T. WATARI, R. KUMAZAWA, T. SEKI, Y. YASAKA\*, A. ANDO, Y. OKA,  
O. KANEKO, K. ADATI, R. AKIYAMA, Y. HAMADA, S. HIDEKUMA,  
S. HIROKURA, K. IDA, K. KAWAHATA, T. KAWAMOTO, Y. KAWASUMI,  
S. KITAGAWA, M. KOJIMA, T. KURODA, K. MASAI, S. MORITA,  
K. NARIHARA, Y. OGAWA\*\*\*, K. OHKUBO, S. OKAJIMA\*\*, T. OZAKI,  
M. SAKAMOTO, M. SASAO, K. SATO, K. N. SATO, F. SHIMPO,  
H. TAKAHASHI, S. TANAHASI, Y. TANIGUCHI, K. TOI, T. TSUZUKI

National Institute for Fusion Science, Nagoya 464-01, Japan

\* Department of Engineering, Kyoto University, , Kyoto, 606, Japan

\*\* Department of Applied Physics, Chubu University, Kasugai, 487, Japan

M. ONO,

Princeton Plasma Physics Laboratory, Princeton, New Jersey, 08543, U.S.A.

Key words: Ion Bernstein Wave Heating, high frequency, impurity problem, third cyclotron harmonic heating, profile peaking, bulk heating

THE NEW FEATURES OF ION BERNSTEIN WAVE HEATING  
IN JIPP T-IIU TOKAMAK

ABSTRACT

Ion Bernstein Wave Heating experiment was conducted in JIPP T-II U tokamak. A relatively high frequency, 130MHz, was used to reduce impurity influx and IBW power up to 400kW was injected without plasma disruption. It was found that the radial profiles of electron density, electron temperature, and ion temperature are all peaked during the IBWH. It was also found that ion distribution function does not have high energy tail above certain critical energy. These are favorable and useful features in optimizing fusion reactivity in reactor applications.

I. INTRODUCTION

Several experiments were conducted for demonstration of IBW heating on tokamaks in the past several years[1-7]. Some of these experiments provide proofs that IBWH works and an unique impact on the plasma confinement has been reported[3-4]. However, other recent experiments did not always reproduce those results[6-7]. Further experimental studies are therefore required in order to improve understanding of physics of IBW Heating. The difficulty in IBWH may partly be attributed to the delicate physics of wave launching[8-10]. We further assume, in this paper, that the impurity problem, which were to some degree problematical in these experiments, may be the other cause of inconsistency between the previous experiments. This thought, combined with the following model of impurity generation[11], made us choose IBWH experiment at a high frequency of 130MHz: A particle in an electric field with frequency  $\omega$  has velocity  $v$ , equivalent energy  $\epsilon$ , and excursion length  $\xi$  in the following form:

$$v = \frac{qE}{m\omega}, \quad \epsilon = \frac{1}{2} m \left( \frac{qE}{m\omega} \right)^2, \quad \xi = \frac{qE}{m\omega^2} \quad (1)$$

These expressions, having inverse power of  $\omega$ , suggests a prescription for reducing impurity problem in favor of higher frequencies. High frequency also reduces non-linear effects such as ponderomotive potential and ionization.

## II. EXPERIMENTAL RESULT

The toroidal magnetic field,  $B_t$ , at the plasma center is 3T and hydrogen gas is used as working gas. With this experimental setup, the third cyclotron frequency locates at the plasma axis and fourth harmonic layer behind the antenna. We are thus exploring a new regime of the IBW Heating (3rd hydrogen cyclotron IBW branch) which is another interest we have in the present experiment. The antennas are three T-shaped Nagoya Type-III coils in a toroidal array. The experiment was conducted at density range of  $n_e = 2 \sim 6 \times 10^{13} \text{ cm}^{-3}$ .

The experiment demonstrated that IBW heating also works in the new regime of IBWH. To make this point firm, efforts are made for elaboration of diagnostics. Detailed ion temperature profile was measured by the use of charge-exchange recombination spectroscopy[12]. A significant rise of the central ion temperature occurs during IBWH even with the simultaneous doubling of the central electron density (from  $3.7 \times 10^{13} \text{ cm}^{-3}$  to  $7.5 \times 10^{13} \text{ cm}^{-3}$ ). As shown in Fig.1, the central ion temperature rises from that of NBI heated target plasma by significant amount. As clearly seen in Fig.1, the ion temperature profile of the IBW heated plasma is peaked. The peaking of the profile is not limited to that of ion temperature. Detailed profile measurement revealed similar peaking features of both plasma density and electron temperature profiles. Particularly, density peaking is noted in the comparison of two FIR interferometer signals viewing center and periphery of the plasma(see Fig.2). Improvement of particle confinement on the application of IBW has been reported in the PLT[3] and ALCATOR[4] experiments. The present experiment in JIPP T-II U confirmed that  $H_\alpha$  decreases at least around the antenna. The peaking of the electron density profiles described above indicates an improvement of particle confinement at interior of the plasma.

In the past experiments[1-2:40MHz IBWH], the impurity accumulation occurred causing the plasma disruption and prevented them from successful run at higher power level than 100kW. In the present experiment, injection power of 400 kW was readily achieved. Their difference by a factor of four indicates a better performance of high frequency IBWH with regard to impurity problem. The index of the density peakedness,  $k$ , defined as some power of the parabolic profile is plotted in Fig.3. The sequential change of  $k$  shows that density profile broadens on application of NBI to the Ohmic plasma, peaks when IBW is superposed, and peaks further in the IBW only phase after NBI

is terminated.

As another important observation in the experiment, we note that the high energy ion tail is not produced during IBWH. The energy distribution measured with FNA is shown in Fig.4. Though the tail grows with increasing injection power the distribution function drops above a certain energy level. This feature is quite different from those of other ICRF heating regimes and may be explained in terms of the quasi-linear Fokker-Planck RF diffusion model which accommodates third cyclotron harmonic damping, i.e.,

$\partial f / \partial t = C(f) + Q(f)$  , with

$$Q(f) = \frac{\rho Z^2 e^2}{2m^2 |k_{\parallel}|} |E_x|^2 \sum_n (\partial / v_{\perp} \partial v_{\perp}) \left( \frac{v_{\perp}^2 n^2}{Z^2} |J_n(Z)|^2 \delta(v_{\parallel}(\text{res})) (\partial f / v_{\perp} \partial v_{\perp}) \right) \quad (2)$$

Here, C(f) is the collision operator and  $Z = (k_{\perp} v_{\perp} / \omega_i)^2$ . For higher harmonic ion Bernstein wave ( $n=3$ ),  $k_{\perp} \geq \rho_i$  holds and the Bessel function involved in eq.(2) is better approximated by an asymptotic expansion rather than by a power expansion, i.e., the strength of the wave particle interaction reduces with increasing ion energy. Therefore, high energy tail drops off above a certain critical energy.

## REFERENCES

- [1] ONO, M., WATARI, T., ANDO, R., et al. Phys. Rev. Lett. 54(1985) 2239
- [2] OGAWA, Y., KAWAHATA, K., ANDO, R., et al., Nucl. Fusion 27 (1987) 1379
- [3] ONO, M., BEIERSDORFER, P., BELL, R., et al., in Plasma Physics and Controlled Nuclear Fusion Research (Proc. 11th Int. Conf. Kyoto, 1986), Vol. 1, IAEA, Vienna (1987) 477.
- [4] PORKOLAB, M., BONOLI, P., CHEN, KUOIN, et al., in Plasma Physics and Controlled Nuclear Fusion Research (Proc. 11th Int. Conf. Kyoto, 1986), Vol. 1, IAEA, Vienna (1987) 509.
- [5] SHINOHARA, S., NAITO, O., TOYAMA, H., et al., in Application of Radio-Frequency Power to Plasmas (Proc. 7th Top. Conf. Kissimmee, FL, 1987) 330.
- [6] TAMAI, H., OGAWA, T., MATSUMOTO, H., et al., in Radio-Frequency Power in Plasmas

- (Proc. 8th Top. Conf. Irvine, CA, 1989), American Physical Society, New York (1989) 350.
- [7] PINSKER, R.I., MAYBERRY, M.J., PORKOLAB, M., et al., in Radio-Frequency Power in Plasmas (Proc. 8th Top. Conf. Irvine, CA, 1989), American Physical Society, New York (1989) 314.
- [8] ONO, M., review paper ; to be published in Physics of Fluids
- [9] BRAMBILLA, M., Nuclear Fusion, 28(1988)549,
- [10] CHIU, S.C., MAYBERRY, M.J., PINSKER, R.I., et al., Rep. GA-A20608, General Atomics, San Diego, CA (1991).
- [11] ITOH, K., FUKUYAMA, A., ITOH, S., et al., Comments on Plasma Phys. and Controlled Fusion , 10(1986)91.
- [12] IDA, K., HIDEKUMA, S., Rev. Sci. Instrum. 60 (1989) 876.

#### FIGURE CAPTIONS

- Fig. 1. Ion temperature profiles of NBI heated target plasma and IBW heated plasma.
- Fig. 2. Line-integrated density of various lines of sight measured with FIR interferometers; central to peripheral chords from top to bottom.
- Fig. 3. The peakedness index,  $k$ , of electron density: the labels p-1, p-2, p-3, p-4 refers to Ohmic, NBI, NBI+IBW, and IBW phases, respectively.
- Fig. 4. The ion energy distribution function of IBW heated plasma.






Fig.1




Fig.2




Fig.3




Fig.4

## Recent Issues of NIFS Series

- NIFS-132 K. Sato and F. Miyawaki, *Formation of Presheath and Current-Free Double Layer in a Two-Electron-Temperature Plasma* ; Feb. 1992
- NIFS-133 T. Maruyama and S. Kawata, *Superposed-Laser Electron Acceleration* Feb. 1992
- NIFS-134 Y. Miura, F. Okano, N. Suzuki, M. Mori, K. Hoshino, H. Maeda, T. Takizuka, JFT-2M Group, S.-I. Itoh and K. Itoh, *Rapid Change of Hydrogen Neutral Energy Distribution at LH-Transition in JFT-2M H-mode* ; Feb. 1992
- NIFS-135 H. Ji, H. Toyama, A. Fujisawa, S. Shinohara and K. Miyamoto *Fluctuation and Edge Current Sustainment in a Reversed-Field-Pinch*; Feb. 1992
- NIFS-136 K. Sato and F. Miyawaki, *Heat Flow of a Two-Electron-Temperature Plasma through the Sheath in the Presence of Electron Emission*; Mar. 1992
- NIFS-137 T. Hayashi, U. Schwenn and E. Strumberger, *Field Line Diversion Properties of Finite  $\beta$  Helias Equilibria*; Mar. 1992
- NIFS-138 T. Yamagishi, *Kinetic Approach to Long Wave Length Modes in Rotating Plasmas*; Mar. 1992
- NIFS-139 K. Watanabe, N. Nakajima, M. Okamoto, Y. Nakamura and M. Wakatani, *Three-dimensional MHD Equilibrium in the Presence of Bootstrap Current for Large Helical Device (LHD)*; Mar. 1992
- NIFS-140 K. Itoh, S. -I. Itoh and A. Fukuyama, *Theory of Anomalous Transport in Toroidal Helical Plasmas*; Mar. 1992
- NIFS-141 Y. Kondoh, *Internal Structures of Self-Organized Relaxed States and Self-Similar Decay Phase*; Mar. 1992
- NIFS-142 U. Furukane, K. Sato, K. Takiyama and T. Oda, *Recombining Processes in a Cooling Plasma by Mixing of Initially Heated Gas*; Mar. 1992
- NIFS-143 Y. Hamada, K. Masai, Y. Kawasumi, H. Iguchi, A. Fijisawa and JIPP T-IIU Group, *New Method of Error Elimination in Potential Profile Measurement of Tokamak Plasmas by High Voltage Heavy Ion Beam Probes*. Apr. 1992
- NIFS-144 N. Ohyabu, N. Noda, Hantao Ji, H. Akao, K. Akaishi, T. Ono, H. Kaneko,

- T. Kawamura, Y. Kubota, S. Morimoto, A. Sagara, T. Watanabe, K. Yamazaki and O. Motojima, *Helical Divertor in the Large Helical Device*; May 1992
- NIFS-145 K. Ohkubo and K. Matsumoto, *Coupling to the Lower Hybrid Waves with the Multijunction Grill*; May 1992
- NIFS-146 K. Itoh, S.-I. Itoh, A. Fukuyama, S. Tsuji and Allan J. Lichtenberg, *A Model of Major Disruption in Tokamaks*; May 1992
- NIFS-147 S. Sasaki, S. Takamura, M. Ueda, H. Iguchi, J. Fujita and K. Kadota, *Edge Plasma Density Reconstruction for Fast Monoenergetic Lithium Beam Probing*; May 1992
- NIFS-148 N. Nakajima, C. Z. Cheng and M. Okamoto, *High- $n$  Helicity-induced Shear Alfvén Eigenmodes*; May 1992
- NIFS-149 A. Ando, Y. Takeiri, O. Kaneko, Y. Oka, M. Wada, and T. Kuroda, *Production of Negative Hydrogen Ions in a Large Multicusp Ion Source with Double-Magnetic Filter Configuration*; May 1992
- NIFS-150 N. Nakajima and M. Okamoto, *Effects of Fast Ions and an External Inductive Electric Field on the Neoclassical Parallel Flow, Current, and Rotation in General Toroidal Systems*; May 1992
- NIFS-151 Y. Takeiri, A. Ando, O. Kaneko, Y. Oka and T. Kuroda, *Negative Ion Extraction Characteristics of a Large Negative Ion Source with Double-Magnetic Filter Configuration*; May 1992
- NIFS-152 T. Tanabe, N. Noda and H. Nakamura, *Review of High Z Materials for PSI Applications*; Jun. 1992
- NIFS-153 Sergey V. Bazdenkov and T. Sato, *On a Ballistic Method for Double Layer Regeneration in a Vlasov-Poisson Plasma*; Jun. 1992
- NIFS-154 J. Todoroki, *On the Lagrangian of the Linearized MHD Equations*; Jun. 1992
- NIFS-155 K. Sato, H. Katayama and F. Miyawaki, *Electrostatic Potential in a Collisionless Plasma Flow Along Open Magnetic Field Lines*; Jun. 1992
- NIFS-156 O.J.W.F.Kardaun, J.W.P.F.Kardaun, S.-I. Itoh and K. Itoh, *Discriminant Analysis of Plasma Fusion Data*; Jun. 1992
- NIFS-157 K. Itoh, S.-I. Itoh, A. Fukuyama and S. Tsuji, *Critical Issues and Experimental Examination on Sawtooth and Disruption Physics*; Jun. 1992

- NIFS-158 K. Itoh and S.-I. Itoh, *Transition to H-Mode by Energetic Electrons*; July 1992
- NIFS-159 K. Itoh, S.-I. Itoh and A. Fukuyama, *Steady State Tokamak Sustained by Bootstrap Current Without Seed Current*; July 1992
- NIFS-160 H. Sanuki, K. Itoh and S.-I. Itoh, *Effects of Nonclassical Ion Losses on Radial Electric Field in CHS Torsatron/Heliotron*; July 1992
- NIFS-161 O. Motojima, K. Akaishi, K. Fujii, S. Fujiwaka, S. Imagawa, H. Ji, H. Kaneko, S. Kitagawa, Y. Kubota, K. Matsuoka, T. Mito, S. Morimoto, A. Nishimura, K. Nishimura, N. Noda, I. Ohtake, N. Ohyaibu, S. Okamura, A. Sagara, M. Sakamoto, S. Satoh, T. Satow, K. Takahata, H. Tamura, S. Tanahashi, T. Tsuzuki, S. Yamada, H. Yamada, K. Yamazaki, N. Yanagi, H. Yonezu, J. Yamamoto, M. Fujiwara and A. Iiyoshi, *Physics and Engineering Design Studies on Large Helical Device*; Aug. 1992
- NIFS 162 V. D. Pustovitov, *Refined Theory of Diamagnetic Effect in Stellarators*; Aug. 1992
- NIFS-163 K. Itoh, *A Review on Application of MHD Theory to Plasma Boundary Problems in Tokamaks*; Aug. 1992
- NIFS-164 Y. Kondoh and T. Sato, *Thought Analysis on Self-Organization Theories of MHD Plasma*; Aug. 1992
- NIFS-165 T. Seki, R. Kumazawa, T. Watari, M. Ono, Y. Yasaka, F. Shimpō, A. Ando, O. Kaneko, Y. Oka, K. Adati, R. Akiyama, Y. Hamada, S. Hidekuma, S. Hirokura, K. Ida, A. Karita, K. Kawahata, Y. Kawasumi, Y. Kitoh, T. Kohmoto, M. Kojima, K. Masai, S. Morita, K. Narihara, Y. Ogawa, K. Ohkubo, S. Okajima, T. Ozaki, M. Sakamoto, M. Sasao, K. Sato, K. N. Sato, H. Takahashi, Y. Taniguchi, K. Toi and T. Tsuzuki, *High Frequency Ion Bernstein Wave Heating Experiment on JIPP T-IIU Tokamak*; Aug. 1992
- NIFS-166 Vo Hong Anh and Nguyen Tien Dung, *A Synergetic Treatment of the Vortices Behaviour of a Plasma with Viscosity*; Sep. 1992
- NIFS-167 K. Watanabe and T. Sato, *A Triggering Mechanism of Fast Crash in Sawtooth Oscillation*; Sep. 1992
- NIFS-168 T. Hayashi, T. Sato, W. Lotz, P. Merkel, J. Nührenberg, U. Schwenn and E. Strumberger, *3D MHD Study of Helias and Heliotron*; Sep. 1992
- NIFS-169 N. Nakajima, K. Ichiguchi, K. Watanabe, H. Sugama, M. Okamoto, M. Wakatani, Y. Nakamura and C. Z. Cheng, *Neoclassical Current and*

*Related MHD Stability, Gap Modes, and Radial Electric Field Effects in Heliotron and Torsatron Plasmas; Sep. 1992*

- NIFS-170 H. Sugama, M. Okamoto and M. Wakatani, *K- $\epsilon$  Model of Anomalous Transport in Resistive Interchange Turbulence* ; Sep, 1992
- NIFS-171 H. Sugama, M. Okamoto and M. Wakatani, *Vlasov Equation in the Stochastic Magnetic Field* ; Sep. 1992
- NIFS-172 N. Nakajima, M. Okamoto and M. Fujiwara, *Physical Mechanism of  $E_{\phi}$ -Driven Current in Asymmetric Toroidal Systems* ; Sep.1992
- NIFS-173 N. Nakajima, J. Todoroki and M. Okamoto, *On Relation between Hamada and Boozer Magnetic Coordinate System* ; Sep. 1992
- NIFS-174 K. Ichiguchi, N. Nakajima, M. Okamoto, Y. Nakamura and M. Wakatani, *Effects of Net Toroidal Current on Mercier Criterion in the Large Helical Device* ; Sep. 1992
- NIFS-175 S. -I. Itoh, K. Itoh and A. Fukuyama, *Modelling of ELMs and Dynamic Responses of the H-Mode* ; Sep. 1992
- NIFS-176 K. Itoh, S.-I. Itoh, A. Fukuyama, H. Sanuki, K. Ichiguchi and J. Todoroki, *Improved Models of  $\beta$ -Limit, Anomalous Transport and Radial Electric Field with Loss Cone Loss in Heliotron / Torsatron* ; Sep. 1992
- NIFS-177 N. Ohyabu, K. Yamazaki, I. Katanuma, H. Ji, T. Watanabe, K. Watanabe, H. Akao, K. Akaishi, T. Ono, H. Kaneko, T. Kawamura, Y. Kubota, N. Noda, A. Sagara, O. Motojima, M. Fujiwara and A. Iiyoshi, *Design Study of LHD Helical Divertor and High Temperature Divertor Plasma Operation* ; Sep. 1992
- NIFS-178 H. Sanuki, K. Itoh and S.-I. Itoh, *Selfconsistent Analysis of Radial Electric Field and Fast Ion Losses in CHS Torsatron / Heliotron* ; Sep. 1992
- NIFS-179 K. Toi, S. Morita, K. Kawahata, K. Ida, T. Watari, R. Kumazawa, A. Ando, Y. Oka, K. Ohkubo, Y. Hamada, K. Adati, R. Akiyama, S. Hidekuma, S. Hirokura, O. Kaneko, T. Kawamoto, Y. Kawasumi, M. Kojima, T. Kuroda, K. Masai, K. Narihara, Y. Ogawa, S. Okajima, M. Sakamoto, M. Sasao, K. Sato, K. N. Sato, T. Seki, F. Shimpo, S. Tanahashi, Y. Taniguchi, T. Tsuzuki, *New Features of L-H Transition in Limiter H-Modes of JIPP T-HU* ; Sep. 1992
- NIFS-180 H. Momota, Y. Tomita, A. Ishida, Y. Kohzaki, M. Ohnishi, S. Ohi, Y. Nakao and M. Nishikawa, *D-<sup>3</sup>He Fueled FRC Reactor "Artemis-L"* ; Sep. 1992