

HETEROGENEOUS SURFACE FLUXES AND THEIR EFFECTS  
ON THE SGP CART SITE

J. C. Doran  
Q. Hu  
J. M. Hubbe  
J. C. Liljegren

W. J. Shaw  
S. Zhong  
G. J. Collatz<sup>(a)</sup>

March 1995

Presented at the  
ARM Science Team Meeting  
March 20-23, 1995  
San Diego, California

Prepared for  
the U.S. Department of Energy  
under Contract DE-AC06-76RLO 1830

Pacific Northwest Laboratory  
Richland, Washington 99352

(a) Goddard Space Flight Center/NASA, Greenbelt, Maryland

**DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

GH

**MASTER**

## **DISCLAIMER**

**Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.**

J.C. Doran, Q. Hu, J.M. Hubbe, J.C. Liljegren, W.J. Shaw, and S. Zhong  
Pacific Northwest Laboratory  
Richland, WA 99352

G. J. Collatz  
Goddard Space Flight Center/NASA  
Greenbelt, MD 20771

### Introduction

The treatment of subgrid-scale variations of surface properties and the resultant spatial variations of sensible and latent heat fluxes has received increasing attention in recent years. Mesoscale numerical simulations of highly idealized conditions, in which strong flux contrasts exist between adjacent surfaces, have shown that under some circumstances the secondary circulations induced by land-use differences can significantly affect the properties of the planetary boundary layer (PBL) and the region of the atmosphere above the PBL. At the Southern Great Plains (SGP) Cloud and Radiation Testbed (CART) site, the fluxes from different land-surface types are not expected to differ as dramatically as those found in idealized simulations. Although the corresponding effects on the atmosphere should thus be less dramatic, they are still potentially important. From an ARM perspective, in tests of single column models (SCMs) it would be useful to understand the effects of the lower boundary conditions on model performance. We describe here our initial efforts to characterize the variable surface fluxes over the CART site and to assess their effects on the PBL that are important for the performance of SCMs.

### Fluxes Over the CART site

The SiB2 model (Sellers et al. 1986) was chosen to help produce a flux map of the CART site. The SiB2 model makes use of soil, plant, and meteorological data to calculate the sensible and latent heat fluxes over vegetated or bare surfaces. Estimates of leaf area index for vegetation are obtained from satellite-derived Normalized Difference Vegetation Index values, while meteorological fields are computed using data from the Oklahoma Mesonet, National Weather Service stations, the Kansas State University Mesonet, and the CART central and extended sites. The multiquadric basis function interpolation scheme of Nuss and Titley (1994) is used to obtain values for precipitation, temperature, humidity, incoming solar radiation, and wind speeds over the CART domain, and these variables are then mapped onto an array of 50 x 58 grid cells, each 6.25 km on a side. A new set of fields is computed every 30 minutes for the time periods of interest.

Initial soil moisture values are computed using the approach of Liston et al. (1993). The SiB2 model is then run for each grid cell in the array to obtain the sensible and latent heat fluxes at each grid point. If there are several principal vegetation types

found in a given grid cell, SiB2 can be run for each vegetation type, and the net fluxes are then computed from a weighted average of the contributions from each type. An example of the fluxes computed by this methodology is shown in Figure 1. The gridded flux values are then used to obtain a time series of spatially averaged fluxes over the domain, at resolutions ranging from  $(6.25 \text{ km})^2$  on up to domain-wide averages.

Such flux calculations are subject to several sources of uncertainty. The largest appears to be the difficulty in specifying the soil moisture profiles accurately for the whole site. At this time soil moisture measurements are quite limited, and although this situation will improve over the next two years with the addition of soil moisture probes at the extended sites, soil moisture descriptions are expected to remain problematic. A second difficulty is determining the accuracy of the interpolated fields, particularly for variables such as solar radiation, which can change rapidly and irregularly under partly cloudy conditions. A third problem is the specification of the plant characteristics on a fine enough scale, and with sufficient differentiation among the various plant types, to allow SiB2 to compute fluxes accurately.

Some degree of tuning is possible by comparing modeled flux values with fluxes measured at the extended sites and making some adjustments in the plant parameters or soil moisture values to obtain improved agreement. This procedure must be used with care, however, to avoid forcing agreement between modeled and measured flux values for the wrong reason. To date, flux measurements have not been available from wheat fields, which constitute a major crop in the CART site, but this situation is also expected to improve substantially by late in 1995.

### Mesoscale Modeling

Using data from the 1992 Boardman experiment (Doran et al., 1995) and applying the RAMS (Regional Atmospheric Modeling System [Pielke et al. 1992]) mesoscale model as an analysis tool, it has been shown that spatially varying surface heat fluxes can result in varying mixed-layer depths over the domain (Doran and Zhong 1995). A similar result is anticipated for the CART region. In addition, differences in column-integrated moisture, temperature, and the probability of cloud formation are also expected to be found. To examine these features, the RAMS model is being used to simulate conditions over the SGP CART site for a number of selected cases.

Initial tests of the RAMS model for this domain do show effects in the PBL arising from spatially-varying heat fluxes caused by differences in soil moisture. However, the parameterizations used in version 3a of RAMS show a sensitivity of sensible and latent heat fluxes to soil moisture that appears to be greater than that actually found from CART data. As a result, the soil moisture in RAMS must currently be treated as a tuning parameter that can be adjusted to produce flux fields similar to those obtained from the SiB2 modeling exercise described above. Figure 2 shows the variations in mixed-layer depth computed for a flux distribution similar but not identical to that given in Figure 1. The variations across the domain are significant and are not currently

accounted for in SCMs.

### Acknowledgment

This research was supported by the U.S. Department of Energy under Contract DE-AC06-76RLO 1830 at Pacific Northwest Laboratory under the auspices of the Atmospheric Radiation Measurement Program. Pacific Northwest Laboratory is operated for the U. S. Department of Energy by Battelle Memorial Institute.

### References

Doran, J.C., and S. Zhong, 1995: Variations in mixed layer depths arising from inhomogeneous surface conditions. *J. Climate*, in press.

Doran, J.C., W.J. Shaw, and J.M. Hubbe, 1995: Boundary layer characteristics over areas of inhomogeneous surface fluxes. *J. Appl. Meteor.*, **34**, 559-571.

Liston, G.E., Y.C. Sud, and G.K. Walker, 1993: Design of a global soil moisture initialization procedure for the Simple Biosphere Model. NASA Technical Memorandum 104590, 24pp.

Nuss, W.A., and D.W. Titley, 1994: Use of multiquadric interpolation for meteorological objective analysis. *Mon. Wea. Rev.*, **122**, 1611-1631.

Pielke, R. A., W. R. Cotton, R. L. Walko, C. J. Tremback, W. A. Lyons, L. D. Grasso, M. E. Nicholls, M. D. Moran, D. A. Wesley, and J. H. Copeland, 1992: A comprehensive meteorological modeling system - RAMS. *Meteorol. Atmos. Phys.*, **49**, 69-91.

Sellers, P.J., Y. Mintz, Y.C. Sud, and A. Dalcher, 1986: A simple biosphere model (SiB) for use within general circulation models. *J. Atmos. Sci.*, **43**, 505-531.




Figure 1. Heat fluxes derived from the SiB2 model for July 28, 1994.




Figure 2. Mixing heights derived from the RAMS model for fluxes similar to those in Figure 1.