

PPPL-3135 - Preprint: September 1995, UC-420, 427

Alpha Power Channeling with Two Waves

N.J. Fisch and M.C. Herrmann
Princeton Plasma Physics Laboratory
P.O. Box 451
Princeton, NJ 08543

MAR 11 1995

OSTI

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

AT

Alpha Power Channeling with Two Waves

**N. J. Fisch and M. C. Herrmann
Princeton Plasma Physics Laboratory
Princeton, NJ 08543, USA**

ABSTRACT. The complete channeling of energy from alpha particles is likely to be realized only through the excitation of a variety of waves, rather than by one wave alone. While one wave constrains more firmly the direction of the energy transfer, the necessary wave characteristics are far more easily achieved through a combination of waves, even at the expense of less restrictive motion of the α -particles.

1. INTRODUCTION

Fusion reactors burning deuterium and tritium (DT) fuel are sustained by α -particle heating. Generally, the α -particles heat mainly electrons through Coulomb collisions. However, in the presence of appropriate waves, the α -particles might amplify these waves on a collisionless time scale. If the waves then are absorbed either by electrons traveling in one direction to accomplish current drive [1] or are absorbed by fuel ions for increased reactivity [2], the α -particle power has been effectively "channeled" to accomplish more useful functions than mere electron heating. The amplification of the waves can occur if there is a population inversion in the α -particles along the wave diffusion path, where the diffusion path points from high energy α -particles in the plasma center to low energy α -particles at the plasma periphery. Complete channeling of the α -particle power to ions would result in about a factor of two in the fusion power density, where most of this increase is due to the operation in the so-called "hot-ion" mode [3,4].

Although, in principle, substantial channeling of α -particle energy can occur, exactly how to accomplish this effect is a challenging problem. Wave interactions that tend to drive particles in velocity space only tend not to extract most of the recoverable energy [5-7]. A poloidally and toroidally propagating lower hybrid wave was proposed [1] to make use of the free energy available in diffusing α -particles also in space, but this wave alone appears to be difficult to excite with the required wave characteristics [8, 9]. One of the more promising wave candidates is the short wavelength mode-converted ion Bernstein wave [10, 11]. However, this wave is hard to excite with sufficient toroidal wavenumber to deliver α -particles to the plasma periphery.

This work shows that qualitatively different behavior occurs when α -particles are resonant with more than one kind of wave. In particular, we show that waves in the ion cyclotron range of frequencies, such

as the ion Bernstein wave (IBW), may be combined to good effect with low frequency waves, such as the toroidal Alfvén eigenmode (TAE). For certain α -particles, it can be seen that the presence of only either of these waves leads to almost no energy extracted, while almost complete energy extraction can occur when both waves are present. The extent to which substantially all the α -particles can be controlled in a similar fashion will determine the utility of using such waves.

The paper is organized as follows: In Sec. 2, we review the issues in energy extraction by one wave. In Sec. 3, we show how qualitatively different behavior can occur with more than one wave. In Sec. 4, we show as an example how an ion Bernstein wave and a TAE-like wave can possess complementary functions. In Sec. 5, we discuss the limitations and implications of this work.

2. CONSTRAINING DIFFUSION TO ONE DIMENSION

The energy of the α -particles is best tapped by diffusing them in energy and space. The channeling occurs when the diffusion path connects high energy α -particles at the plasma center to low energy α -particles at the plasma periphery. When one wave is utilized, there can be only one diffusion path, so that there are stringent constraints on the α -particle motion. If the path is chosen appropriately, then, if an α -particle gains energy, it must diffuse to the tokamak periphery; conversely, if it were to lose energy in interacting with the wave, it must diffuse to the tokamak center. The α -particle motion is constrained to lie on a one dimensional curve, a line. Thus, particles diffused by the wave to the tokamak periphery give up a precise amount of energy to the wave which is proportional to their distance traveled in reaching the periphery.

For simplicity in explaining the effect, consider a slab geometry, where one can write the ratio of dis-

placement of the guiding center in the x -direction, Δx_{gc} , to energy absorbed as

$$\Delta x_{gc} = -k_y \Delta \epsilon / m \Omega \omega. \quad (1)$$

This quantity is determined by wave and particle parameters only; ω is the wave frequency, k_y is the wavenumber in the y -direction, m is the α -particle mass and $\Omega \equiv qB/m$ is the α -particle gyrofrequency. In the slab case, upon repeated interactions with the wave, a particle will trace a line in $\epsilon - x_{gc}$ space.

Suppose that the plasma boundary is at $x = a$, i.e., α -particles can only leave at $x = a$. The plasma center is at $x = 0$, by which is meant that no α -particles can leave at $x = 0$. For efficient channeling, one would then require $\Delta x_{gc}/\Delta \epsilon \sim a/\epsilon_\alpha$, where a is the extent of the plasma and ϵ_α is the α -particle birth energy. If instead $\Delta x_{gc}/\Delta \epsilon \gg a/\epsilon_\alpha$, then the α -particle would be extracted from the center with almost all its energy intact, whereas if $\Delta x_{gc}/\Delta \epsilon \ll a/\epsilon_\alpha$, then the α -particles are not extractable from the plasma center. In this case, a population inversion is not likely to occur, and the wave will not be amplified.

In toroidal geometry, particles interacting with one wave trace a line in $\epsilon - \mu - P_\phi$ space, where $\mu = mv_\perp^2/2B$ is the magnetic moment, $\epsilon = \mu B + mv_\parallel^2/2$ is the kinetic energy, and $P_\phi = R(mB_\phi v_\parallel/B - qA_\phi)$, is the canonical angular momentum, and where A_ϕ is the vector potential. Each point in $\epsilon - \mu - P_\phi$ space represents a single guiding center orbit for trapped particles, and, for each sign of v_\parallel , a passing particle orbit. Given ϵ , μ , and P_ϕ , and the sign of v_\parallel for passing orbits, it may be determined if the orbit intersects the plasma periphery, thus losing the particle [12, 13]. Particles tend to follow closely the magnetic surfaces, i.e., surfaces of constant RA_ϕ , so that the particle position is largely determined by P_ϕ , especially for low energy particles.

Upon interaction with a wave with toroidal mode number n_ϕ , and absorbing energy $\Delta \epsilon$, P_ϕ changes by

$$\Delta P_\phi = (n_\phi/\omega) \Delta \epsilon. \quad (2)$$

Assume that the exchange of energy occurs only for particles satisfying the resonance condition

$$\omega - k_\parallel v_\parallel = n\Omega, \quad (3)$$

where n is an integer, then, upon absorbing energy $\Delta \epsilon$, μ changes by

$$\Delta \mu = (nZe/m\omega) \Delta \epsilon, \quad (4)$$

where e is the charge on an electron and, for α -particles, $Z = 2$. Thus, upon repeated interaction with one wave, the constants of motion, ϵ , μ , and P_ϕ , trace a line.

Waves with the appropriate characteristics must be excited both to extract energy and to deliver the α -particle to the periphery. A particularly promising wave in a tokamak is the mode converted IBW wave, excited with high k_θ , where θ is the poloidal mode number [10]. This wave breaks the μ invariance of the α -particle motion, but its excursion is limited, by Eq.(2), if n_ϕ is not sufficiently large. In Fig. 1, we show the orbit of an α -particle interacting with an IBW wave with $n_\phi = 50$. The outer orbit is the largest orbit the particle can sample and still be resonant with the wave. In this orbit, the α -particle has lost about 2/3 of its initial energy and essentially all of its perpendicular energy. At this point, constrained through Eq.(4), the α -particle can lose no more energy and so can not quite make it to the periphery. The magnetic geometry for this case is TFTR-like, with concentric circular flux surfaces, 2 MA toroidal current, and 5.3 T toroidal field. The major radius at the magnetic axis is $R_0 = 2.6$ m, and the minor radius is $a = 1$ m. The mode-converted wave, at 32 MHz, exists near the mode conversion layer in a vertical slab between 1.05 and 1.15 R/R_0 . The vertical extent of the slab is the full tokamak minor cross-section, while the horizontal extent is 26 cm. The α -particle could be extracted either in a lower current equilibrium or in the presence of a higher mode number IBW wave. However, in practice, higher mode numbers in a TFTR-size tokamak would be difficult to excite efficiently.




FIG. 1. Poloidal cross section of orbit of a cogoing α particle interacting solely with a Mode Converted Ion Bernstein Wave (MCIBW). Magnetic axis is at $(1,0)$. If the particle were to lose energy under interaction with this wave it would move from the central orbit to the outermost orbit, eventually losing 2.5 MeV.

3. DIFFUSION ALONG A TWO DIMENSIONAL SURFACE

There are a number of obstacles in channeling α -particle power with just one wave. First, the necessary wave characteristics are difficult to obtain. In the case of Fig. 1, these high- n_ϕ modes are not easily excited in a tokamak. Second, even if such a wave could be excited, it tends mainly to extract the perpendicular energy only of the α -particle. Third, there is the possibility of "hard landings" [11]; if the α -particle were born away from the plasma center, it would have a shorter distance to travel to the periphery, and, hence, it would leave with much of its energy intact.

A qualitatively different picture emerges when more than one wave interacts with the α -particle. Consider, again for simplicity, a slab geometry. If the waves induce different ratios of $\Delta x_{gc}/\Delta\epsilon$, the α -particles are no longer constrained to a line in $x-\epsilon$

space. If an α -particle encounters the periphery, its energy is no longer immediately determined from its initial coordinates; its energy at the wall would depend on the history of how it received kicks from each of the waves with which it was resonant.

However, by interacting with more than one wave, the constraints on each of the waves for expelling the α -particle are relaxed. In addition, depending on the waves involved, it may be that, even though the α -particle is not constrained to lose energy upon expulsion from the plasma, the α -particle may still be vastly predisposed to lose energy to the waves rather than to gain energy. To illustrate this, consider the interaction of α -particles with two waves in a slab, as sketched in Fig. 2. Here, $x = 0$ depicts the plasma center, while the plasma periphery is at $x = a$. Alpha particles can leave the plasma at $x = a$ only. In Fig. 2a, we depict by horizontal lines diffusion paths in energy only, such as might result from a Landau-type resonance. These diffusion paths can have finite extent in the horizontal direction, and these line segments, of course, need not be the same length, since the resonance condition may be energy and space dependent. By diffusing along these paths only, α -particles born near the plasma center can not reach the plasma boundary, as all α -particles are effectively confined to horizontal line segments in $x-\epsilon$ space. Similarly, we depict, by vertical line segments, diffusion paths in space only, such as might result from interactions with a magnetic ripple. By diffusing along these paths only, α -particles born near the plasma center reach the plasma boundary, but without any loss in energy.

Clearly, in the presence of either resonance only, α -particles sample only line segments, whereas in the presence of both resonances at once, α -particles can sample a two-dimensional volume in $x-\epsilon$ space. In the presence of just one wave, the α -particle may not be able to access the plasma boundary, but if it does come into contact with the boundary, it is constrained to do so at one precise energy. On the other hand, when diffusion along multiple diffusion path occurs, it is appar-




FIG. 2-a. Schematic diffusion paths in energy and space. Diffusion in energy ϵ is exhibited by horizontal paths; diffusion in space x by vertical paths. Plasma periphery is at $x=a$; plasma center is at $x=0$.

ent that α -particles leaving the plasma are not constrained to do so at a single energy.

There are, however, a number of ways in which control can be exercised in the presence of multiple waves. Let us imagine particles diffusing along multiple diffusion paths as receiving random kicks in a 2-D web. Strands within the web may be removed or strengthened by means of resonance conditions. Thus, although the kicks are random, structure in the web or surface itself can lead to diffusion in a definite direction. As an example, consider Fig. 2b, where we imagine that diffusion in energy is of finite extent, whereas the diffusion in space is energy-dependent, say, with significant diffusion only at low energy. Such an energy-dependent diffusion in space only has, in fact, been investigated [14, 15]. Clearly, all α -particles must leave the plasma at low energy, so that there is net extraction of the α -particle energy.

In toroidal geometry, the details are somewhat more complicated, but the idea is the same – namely, one wave diffuses along a line, while several waves diffuse along a web in ϵ - μ - P_ϕ space. Again, structure in the web can lead to usefully constrained motion of the particles.




FIG. 2-b. Schematic diffusion paths in energy and space. Diffusion in space has strong energy dependence.

4. DIFFUSION BY TWO WAVES IN A TORUS

In optimizing energy extraction through diffusion by two waves, one strategy would be to have the two waves play complementary roles. For example, choose one wave in the ion cyclotron range of frequencies in order to break the invariance of μ and thereby access the particle's perpendicular energy. The mode converted ion Bernstein wave [16] could play this role admirably. What the IBW does not do easily is to move the α -particles large distances. The second wave then need not break the μ -invariance, but ought to move the α -particles large distances with little energy exchange. Low frequency modes, such as the toroidal Alfvén eigenmode (TAE), that do not break the adiabatic motion of the α -particles, should serve well the complementary role.

The IBW wave may be excited by launching a fast wave that mode converts at the ion-ion hybrid resonance layer [17], or through other mode conversion techniques [18]. This wave has been excited with toroidal phasing in TFTR in order to drive plasma current [19]. In addition, it appears to interact with fusion byproducts [20]. The TAE mode [21, 22] has been extensively analyzed theoretically, including models for its excitation and saturation [23–25]. It has

been predicted to interact effectively with α -particles, expelling them with little energy loss [26]. This mode has been driven directly in JET using a saddle coil antenna [27].

In this section, we demonstrate the qualitatively different motion that results when such waves are used in combination. In order to demonstrate this difference we employ highly simplified models that capture the essential features of both the IBW wave and a TAE-like wave in a TFTR-size tokamak. The model that we use is that if a particle is resonant with a wave at any point in its orbit, i.e., if it satisfies locally Eq.(3), then it experiences a jump to a nearby orbit, or point in ϵ - μ - P_ϕ space. The nearby point meets the criteria, $\epsilon \rightarrow \epsilon + \Delta\epsilon$, where $\Delta\epsilon = mv\Delta v + m\Delta v^2/2$, where Δv is a random variable with equal probability of being positive or negative. The magnitude of Δv is a parameter that depends on the wave amplitude. The excursions in μ and P_ϕ follow from Eqs. (2) and (4).

The IBW wave is modeled as described in the text accompanying Fig. 1. The mode that we call "TAE-like," has certain characteristics in common with a TAE mode, but we do not attempt to be precise either with respect to the mode or the wave-particle interaction. Rather, key features of the interaction with the TAE mode, or other similar modes, are retained. Specifically, we imagine this mode to be sufficiently below the ion cyclotron frequency that the invariance of μ is not broken, i.e, the only resonances encountered are Landau resonances, $n = 0$, in Eq.(3). Hence, in interacting with the wave, the α -particle experiences kicks in ϵ - μ - P_ϕ space with μ conserved.

While the diffusion path for the TAE-like mode is exact for any wave with frequency ω and toroidal mode number n_ϕ , the probability of a kick and the possible correlation between kicks depend on more precise wave models than are employed here. Our assumption, which may not model precisely any low frequency mode, is that the mode is global, but a significant wave-particle interaction occurs only where there is a local resonance. This model TAE-like mode

interacts with α -particles in such a way as to retain certain key features of the TAE mode interaction with α -particles. First, α -particles are expelled by the model TAE-like mode with the correct energy, since the orbits are precisely constrained by Eq.(2) and not affected by the details of the model. Second, although there is no threshold for stochastic interaction in our model mode, there is a threshold for expelling α -particles, since there must be overlap of the TAE modes to deliver an α -particle to the boundary. This can be modeled through a resonance broadening – at very small wave amplitudes, the wave-particle resonances given by Eq.(3) must be satisfied exactly, while for larger amplitudes, this condition acquires a finite width, the extent of which depends on the wave amplitude. In practice, only a finite width resonance is effective at expelling the α -particles. Thus, the model TAE-like mode at a given n_ϕ and at a given ω expels α -particles with the same exit energy as a TAE mode with the same wave characteristics, and, like the TAE mode, expels these α -particles only if a threshold condition is met.

Recall Fig. 1, where the interaction of an α -particle with only an IBW wave is shown. The succession of orbits depicted assumes that between orbits the α -particle has lost energy to the wave according to Eqs.(2) and (4). The result is that the α -particle can not reach the plasma periphery by interacting only with this wave. Now consider instead the interaction of this α -particle with only a TAE-like mode; in Fig. 3a such an interaction is depicted, where, upon repeated interactions, the α -particle orbit can explore the set of orbits shown. The α -particle begins near the plasma center at 3.5 MeV; it can then explore the outer orbits, with the outermost orbit at 2.7 MeV. None of these orbits, however, intersects the plasma boundary, so the α -particle remains bracketed in both position and energy.

The simultaneous presence of both of these waves, however, leads to different behavior. In Fig. 3b, we depict the α -particle succeeding in reaching the




FIG. 3-a. Poloidal cross section of co-going α particle interacting solely with a TAE like mode. Parameters as in Figure 1. If the particle were to lose 0.8 MeV under interaction with this wave it would move from the central orbit to the outermost trapped orbit. The particle cannot proceed further because it would no longer be resonant with the wave.

plasma periphery, if both waves are present, and if the α -particle always loses energy upon resonating with either wave. In this case, where all kicks are to lower energy, the α -particle loses 2.3 MeV to the waves before exiting at the plasma periphery.

In the case that only one wave is present, the complete set of orbits accessible to the α -particle is a line in ϵ - μ - P_ϕ space and can be generated simply: Generate one set of orbits by assuming that between orbits the α -particle only loses energy and generate a second set of orbits by assuming that between orbits the α -particle only gains energy; the accessible space is the union of these two sets. Thus, in Fig. 2 and in Fig. 3a, it is possible to conclude that if the α -particle loses energy, it must be found somewhere in the set of orbits depicted. In the case of two waves, however, the accessible region is a surface in ϵ - μ - P_ϕ space, and Fig. 3b represents only one possible realization. Although in the realization depicted the α -particle did




FIG. 3-b. Same as Figure 3a except the α -particle interacts with both the MCIBW and a TAE like mode, exiting at the periphery with 1.2 MeV.

reach the periphery while losing energy, there is no guarantee that this need occur most of the time for random kicks. What has been shown, however, is that there is at least a realizable pathway to the periphery and to lower energy.

It is clear, in addition, that if there is one sequence of kicks that results in the α -particle exiting the plasma, then, if the α -particle receives random kicks from the wave, it must eventually exit the plasma.

In Fig. 4, we show the exit energy as a function of exit time for the same α -particle subjected to random sequences of kicks. Interestingly, the sequence that leads to the most direct exit, namely the sequence depicted in Fig. 3b in which energy is always extracted, does not lead to the largest extracted energy at the particle exit. Note too that, although the α -particle can wander around for long times before exiting, nonetheless, substantial energy is still extracted. In Fig. 5, we integrate over the exit times to show the distribution of exit energies for random sequences of kicks.




FIG. 4. Energy extracted vs. time to extract for particle in Figure 3b




FIG. 5. Distribution of energy extracted in Figure 4.

The wave-particle interaction depicted in Figs. 3–5 is not necessarily optimized for energy extraction. In Fig. 6, by modifying somewhat the wave characteristics, we demonstrate how substantially all of the energy might be extracted from the same α -particle. Here, there is substantially less overlap in space between the IBW waves and the TAE-like mode, so that the α -particle first interacts with the TAE-like mode, essentially diffusing in one dimension, and then subsequently interacts with the IBW wave, also essentially diffusing on a line. In this case, as in the previous one, the presence of both waves is required to expel the α -particle. In Fig. 7, we show the distribution of exit energies for random sequences of kicks and for waves as given in Fig. 6. Note that the α -particle tends to exit in a band of energies even narrower than in




FIG. 6. Poloidal cross section for cogoing α particle as in Figure 3b, interacting with different MCIBW and TAE waves, and exiting at the periphery with 0.3 MeV.




FIG. 7. Distribution of energy extracted, under interaction with waves as given in Figure 6.

the first case. As demonstrated by these examples, the presence of more than one wave creates possibilities in expelling α -particles that would not be present with only either wave. At the same time, the relaxation of the constraint that the α -particle diffusion occur only on a line is, at least in these examples, not significant enough to affect the tendency of the α -particles to be

cooled by the wave.

5. CONCLUSIONS

In developing a scenario for channeling α -particle power, the first step must be to imagine how α -particles might diffuse to the periphery while releasing nearly all of their energy. This paper offers how this might come about. The interaction of several waves with an energetic ion population can lead to behavior radically different from what can be accomplished with one wave only.

Of course, several significant caveats must be offered. First, the waves here are only coarsely modeled. The models employed here of the IBW wave and the TAE mode retain only the main features of these waves, not the details of their propagation and damping. Hence, the waves as described here are not expected to be realizable in detail. In addition, the wave-particle interaction is also only coarsely modeled. In particular, the use of a local resonance condition is likely a good model for the IBW wave, but a poor model for at least the TAE mode interaction with trapped α -particles. Further, while we have reason to believe that energy extracted by either the TAE mode or the IBW wave could be damped on ions, this has not been shown for the model waves employed here.

Even if the IBW wave and the TAE mode could be excited in the plasma to accomplish exactly what has been illustrated in this paper, it must be realized that what may be the perfect combination of waves for some α -particles, may not be effective at all in channeling energy from other α -particles. Thus, the next step must be to show that many α -particles can be controlled effectively by the same set of waves. Also, the time scale for producing the channeling effect must be shorter than the α -particle slowing down time. In the simulations described here, this is assumed to be the case. The wave amplitudes required to produce this effect are probably realizable in smaller tokamak

reactors if substantially all the α -particle power is channeled into the waves, but, until a better model of the waves is employed, this is by no means certain. Finally, the concentric circular flux surfaces used here are usefully simple, but neither an optimal nor even a realizable magnetic geometry for a tokamak reactor.

Nonetheless, as a first step to the development of the channeling scenario, this paper does demonstrate possibilities inherent in injecting several different kinds of waves. The combination of waves put forth here, one in the ion cyclotron range of frequencies or of higher frequency to break the μ -invariant, and one lower frequency wave to move α -particles large distances with relatively little energy exchange, appears to work well. The exploration of TAE-like modes is already quite developed, because of the potentially deleterious effects associated with the ejection of energetic α -particles by these waves. Now there may be a further impetus, associated with possibly advantageous effects, for studying and controlling these modes.

6. ACKNOWLEDGEMENTS

This work was supported by the United States Department of Energy under contract number DE-AC02-76-CHO3073. One of the authors (MCH) acknowledges the support of the Fannie and John Hertz Foundation.

REFERENCES

- [1] N. J. FISCH, J. M. RAX, *Phys. Rev. Lett.* **69**, 612 (1992).
- [2] N. J. FISCH, J. M. RAX, *Plasma Physics and Controlled Nuclear Fusion Research Vol. 1*, 769 (1993).
- [3] J. F. CLARKE, *Nucl. Fusion* **20**, 563 (1980).
- [4] N. J. FISCH, M. C. HERRMANN, *Nucl. Fusion* **34**, 1541 (December, 1994).
- [5] D. J. SIGMAR, in *Proc. Physics of Plasmas Close to Thermo-nuclear Conditions*, Varenna, Italy (Commission of the European Communities, Brussels, Belgium, 1979) p. 271.

- [6] W. R. SUTTON III, D. J. SIGMAR, G. H. MILEY, *Fusion Tech.* **7**, 374 (1985).
- [7] K. R. CHEN, *Phys. Rev. Lett.* **72**, 3534 (1994).
- [8] L. KRLÍN, P. PAVLO, *Nucl. Fusion* **34**, 1517 (1994).
- [9] K. KUPFER, S. C. CHIU, V. S. CHAN, *Nucl. Fusion* **35**, 163 (1995).
- [10] E. J. VALEO, N. J. FISCH, *Phys. Rev. Lett.* **73**, 3536 (1994).
- [11] N. J. FISCH, *Phys. Plasmas* **2**, 2375 (1995).
- [12] J. A. ROME, Y-K. M. PENG, *Nucl. Fusion* **19**, 1193 (1979).
- [13] C. T.HSU, D.T. SIGMAR, *Phys. Fluids B* **4**, 1492 (1992).
- [14] H. E. MYNICK, *Phys. Fluids* **5**, 1471 (1993).
- [15] H. E. MYNICK, N. POMPHREY, *Nucl. Fusion* **34**, 1277 (1994).
- [16] I. B. BERNSTEIN, *Phys. Rev.* **109**, 10 (1958).
- [17] F. W. PERKINS, *Nucl. Fusion* **17**, 1197 (1977).
- [18] M. ONO, *Phys. Fluids B* **5**, 241 (1993).
- [19] R. MAJESKI, C. K. PHILLIPS, J. R. WILSON
Phys. Rev. Lett. **73**, 2204 (1994).
- [20] D. DARROW et al., to be submitted to *Phys. Rev. Lett.* (1995).
- [21] C. Z. CHENG, L. CHEN, M. S. CHANCE, *Ann. Physics* **161**, 21 (1985).
- [22] C. Z. CHENG, M. S. CHANCE, *Phys. Fluids* **29**, 3695 (1986).
- [23] H. L. BERK, B. N. BREIZMAN, H. YE, *Phys. Rev. Lett.* **68**, 3563 (1992).
- [24] G. Y. FU, J. W. VAN DAM, *Phys. Fluids B* **1**, 1949 (1989).
- [25] H. L. BERK, B. N. BREIZMAN, *Phys. Fluids B* **2**, 2246 (1990).
- [26] D.T. SIGMAR, C. T.HSU, R. B. WHITE, C. Z. CHENG,
Phys. Fluids B **4**, 1506 (1992).
- [27] G. T. A. HUYSMANS, W. KERNER, D. BORBA, H. A. HOL-
TIES J. P. GOEDBLOED, *Phys. Plasmas* **2**, 1605 (1995).