

E4-96-19

E.A.Kolganova*, A.K.Motovilov**

USE OF THE FADDEEV DIFFERENTIAL
EQUATIONS FOR COMPUTATIONS
OF THREE-BODY RESONANCES

Submitted to «Ядерная физика»

*E-mail: KEA@THSUNI.JINR.DUBNA.SU

**E-mail: MOTOVILV@THSUNI.JINR.DUBNA.SU

VOL 27 № 18

I. INTRODUCTION

Resonances are one of most interesting phenomena in scattering of quantum particles. The problem of definition and studying resonances is paid a lot of attention both in physical and mathematical literature (see, e. g., the books [1] – [8]). Main difficulties connected with a rigorous definition of resonance are explicitly emphasized by B. Simon in his survey [9]. A presence of such difficulties is obliged, first of all, to the fact that in a contrast to the usual spectrum, resonances are not an unitary invariant of an operator (Hamiltonian of a quantum system). The generally accepted interpretation of resonance as a complex pole of the scattering matrix continued analytically on unphysical sheet(s) of the energy plane, goes back to the known paper by G. Gamow [10]. For radially symmetric potentials, such an interpretation of the two-body resonances has been rigorously approved by R. Jost [11]. Beginning from E.C. Titchmarsh [12] resonances are considered as well as poles of analytic continuation of the Green function (or its matrix elements between suitable states [6], [7]). A survey of different physical approaches to studying three-body resonances may be found e. g., in [5] and [13].

At the moment, one of the most effective approaches to practical calculation of resonances is the complex scaling method [14] (see also [7], [9]). This method is applicable to the few-body problem in the case where interaction potentials between particles are analytic functions of coordinates. The complex scaling gives a possibility to rotate the continuous spectrum of the N -body Hamiltonian in such a way that certain sectors become accessible for observation on unphysical sheets neighboring with the physical one. At the same time, the real discrete spectrum of the Hamiltonian stays fixed during all the scaling transformation. Resonances in the sectors above turn out to be extra discrete eigenvalues of the scaled Hamiltonian [7]. Thereby, when searching for resonances one may use standard methods to find discrete spectrum. Practical applications of the complex scaling method to concrete problems may be found, in particular, in the recent papers [15] – [17]. Alongside with the complex scaling, another methods are used for calculations of three-body resonances which are based in particular on solving the momentum space Faddeev integral equations [18], [19] continued through the cut (see the survey [13] by K. Möller and Yu.V. Orlov and the literature cited therein). In this approach, resonances are searched for as poles of the T-matrix.

The present paper is devoted to developing a method to calculate three-body resonances using the recently found explicit representations disclosing a structure of the T-matrix on unphysical sheets as well as analogous representations for the scattering matrix and resolvent [20], [21]. These representations were obtained in supposition that the interaction potentials were pairwise and falling-off in the coordinate representation not slower than exponentially. According to the representations [20], [21], the matrix $M(z) = \{M_{\alpha\beta}(z)\}$,

$\alpha, \beta = 1, 2, 3$, constructed of the operator $T(z)$ Faddeev components [18], [19], is explicitly expressed on unphysical sheet Π_l of the energy z plane in terms of this matrix itself taken on the physical sheet and a certain truncation $S_l(z)$ of the total three-body scattering matrix $S(z)$. Character of the truncation is determined by the index (number) l of the unphysical sheet concerned. Respective representations for analytic continuation of the matrix $S(z)$ and resolvent $R(z)$ follow immediately from the representations for $M(z)|_{\Pi_l}$. A main consequence of the representations admitting direct practical applications, is the fact that the T-matrix and resolvent as well as the scattering matrix have nontrivial singularities on unphysical sheet Π_l exactly at those values of the energy z where the corresponding matrix $S_l(z)$ has zero as eigenvalue. It is important that $S_l(z)$ is considered on the physical sheet only. Therefore, one can provide a search for resonances (poles of $M(z)$, $S(z)$ and $R(z)$) on a certain unphysical sheet Π_l keeping z always on the physical one and calculating only a position of zeros of the operator-valued function $S_l(z)$. For all this, one can use any method allowing to calculate (on the physical sheet) amplitudes of the processes necessary to construct the truncation $S_l(z)$.

In the present paper, the matrices $S_l(z)$ are computed on the base of the numerical algorithm [22] elaborated to solve the Faddeev differential equations in configuration space (see the book [19], survey [23] and references therein). Certainly, when computing the amplitudes on the physical sheet one has to extend the Faddeev differential formulation of the scattering problem as well on the complex values of z . It should be noted that, in the holomorphy domain (see [21]) of the amplitudes, the differential formulation stays to be correct.

Unfortunately, the algorithm [19], [22], [23] (see also [24] – [27]) has been worked out in details only for the processes ($2 \rightarrow 2, 3$). Thus, there may be computed in practice only the amplitudes of elastic scattering and rearrangement for the processes ($2 \rightarrow 2$) and the breakup amplitude into three particles. A knowledge of these amplitudes is sufficient to compute those truncations $S_l(z)$ of the three-body scattering matrix $S(z)$, zeros of which are “responsible” for resonances situated on the so-called two-body unphysical sheets, i. e. those sheets of the energy Riemann surface where the parameter z may be guided going around the pair thresholds only. As a concrete application of the method concerned we make a search for resonances in the nnp system and in a model system of three bosons with the nucleon masses.

Let us describe shortly structure of the paper.

In Sec. II, we introduce main notations and formulate the explicit representations [21] for the unphysical-sheet three-body T-matrix, scattering matrix and resolvent which are used then to approve the numerical method of the work.

In Sec. III the system nnp and a three-boson system are considered. Formulations of the boundary-value problems [19], [22], [23] are given for the Faddeev partial differential equations, corresponding to the processes ($2 \rightarrow 2, 3$)

in these systems and going out to a domain of complex energy values in the physical sheet. Numerical method to solve these problems is described. Truncated (partial) scattering matrices are constructed in terms of the amplitudes for elastic scattering ($2 \rightarrow 2$). Zeros of these matrices represent resonances (including virtual poles) on the unphysical sheet connected with the physical one by crossing the continuous spectrum interval between the deuteron energy and three-body threshold. Results of numerical computations are exposed.

We conclude this introduction with some notation. Throughout the paper, we understand by $\sqrt{z - \lambda}$, $z \in \mathbb{C}$, $\lambda \in \mathbb{R}$, the main branch of the function $(z - \lambda)^{1/2}$. By \hat{p} we denote the unit vector in the direction $p \in \mathbb{R}^n$. $\hat{p} = \frac{p}{|p|}$ and by S^{n-1} , the unit sphere in \mathbb{R}^n , $\hat{p} \in S^{n-1}$.

II. EXPLICIT REPRESENTATIONS FOR T-MATRIX, SCATTERING MATRIX AND RESOLVENT ON UNPHYSICAL SHEETS

The scattering matrix, T-matrix and Hamiltonian resolvent (Green function) for a quantum-mechanical system are stringently connected with each other. Therefore, all these three objects, considered as functions of energy, have usually the same Riemann surface. Such a fact takes place at least in the multichannel scattering problem with binary channels and in the three-body problem with quickly decreasing interactions [28], [21]. In a "first approximation", the structure of the Riemann surfaces in these problems coincides. The thing is that the branching points (in real axis) are stipulated to these surfaces in the both problems by the same reason, namely by a presence of the Cauchy-type integrals in the Lippmann-Schwinger or Faddeev equations. In the equations considered in the momentum representation, the Cauchy-type integrals are engendered by the kernels $\frac{\delta(p - p')}{\lambda + p^2 - z}$ with λ , the channel thresholds and $p, p' \in \mathbb{R}^n$, the respective channel momentum variables. In the case of the channels ($2 \rightarrow 2, 3$) in three-body problem and the odd-dimensional channels (i. e., with n odd) in the matrix multichannel problem, the thresholds λ turn out to be the branching points of the second order. Even-dimensional channels in the multichannel problem as well as the channel ($3 \rightarrow 2, 3$) in the three-body problem give logarithmic branching points (see [21], [28], [13]).

The method used for calculation of resonances in the present work, is based on the explicit representations [21] for analytic continuation of the T-matrix, scattering matrices and Green function on unphysical sheets keeping true at least for a part of the three-body Riemann surface. To describe this part we introduce the auxiliary vector-function $f(z) = (f_0(z)), f_{1,1}(z), \dots, f_{1,n_1}(z), f_{2,1}(z), \dots, f_{2,n_2}(z), f_{3,1}(z), \dots, f_{3,n_3}(z))$ with $f_0(z) = \ln z$ and $f_{\alpha,j}(z) = (z - \lambda_{\alpha,j})^{1/2}$. Here, by $\lambda_{\alpha,j}$ we understand respective bound-state energies

of pair subsystems α , $\alpha = 1, 2, 3$, $j = 1, 2, \dots, n_\alpha$, numerated taking into account their multiplicities. It is supposed that $n_\alpha < \infty$. The sheets Π_l of the vector-function $f(z)$ Riemann surface \mathfrak{R} are numerated via multi-index $l = (l_0, l_{1,1}, \dots, l_{1,n_1}, l_{2,1}, \dots, l_{2,n_2}, l_{3,1}, \dots, l_{3,n_3})$, where $l_{\alpha,j} = 0$ if the sheet Π_l corresponds to the main (arithmetic) branch of the square root $(z - \lambda_\alpha)^{1/2}$. Otherwise, $l_{\alpha,j} = 1$ is assumed. Value of l_0 coincides with the number of branch of the function $\ln z$, $\ln z = \ln |z| + i2\pi l_0 + i\phi$ where $\phi = \arg z$. For the physical sheet identified by $l_0 = l_{\alpha,j} = 0$, $\alpha = 1, 2, 3$, $j = 1, 2, \dots, n_\alpha$, we use the notation Π_0 . A "sticking" of the sheets Π_l of the surface \mathfrak{R} is realized in intervals between neighboring thresholds along rims of the cut along the continuous spectrum. A detailed description of the surface \mathfrak{R} may be found in [28].

The type \mathfrak{R} surfaces without extra branching points arise in the multi-channel problems with binary channels only. Structure of the total three-body Riemann surface is essentially more complicated. For instance, the sheets Π_l with $l_0 = \pm 1$ have additional branching points corresponding to resonances of the two-body subsystems. In the sheets Π_l with $l_0 = 0$, one can discover (in the left half-plane) logarithmic branching points of a kinematical origin. The part $\mathfrak{R}^{(3)}$ of the total three-body Riemann surface where the representations [21] are valid, consists of the surface \mathfrak{R} sheets Π_l identified by $l_0 = 0$ (such unphysical sheets are called *two-body* sheets) and two *three-body* sheets identified by $l_0 = \pm 1$ и $l_{\alpha,j} = 1$, $\alpha = 1, 2, 3$, $j = 1, 2, \dots, n_\alpha$. Note that $\mathfrak{R}^{(3)}$ includes all the unphysical sheets neighboring with the physical sheet Π_0 .

In the case of the nnp -system and a model three-boson system considered below, the surface $\mathfrak{R}^{(3)}$ is shown schematically in Fig. 1. In this case a single pair threshold, the deuteron energy $\lambda = E_d$ is present only. Therefore, the index l of the sheets Π_l consists of two components only: $l = (l_0, l_1)$. In the terminology accepted, the sheet $\Pi_{(0,1)}$ (see Fig. 1) is a two-body sheet, but the sheets $\Pi_{(-1,1)}$ и $\Pi_{(1,1)}$ are three-body ones.

Construction of the representations for the T-matrix consists of the following stages. At the first step, one carries out analytic continuation on unphysical sheets, of the absolute terms and kernels of the Faddeev integral equations for the components $M_{\alpha\beta}(z)$ (the continuation is understood in the sense of distributions). As the absolute terms as the kernels after continuation are expressed in terms of the pair T-matrices and scattering matrices taken on the physical sheet. Transforming the Faddeev equations continued, one finds the kernels $M_{\alpha\beta}(P, P', z)|_{z \in \Pi_l}$ can be explicitly expressed in terms of these kernels themselves taken on the physical sheet Π_0 in their off-shell and/or half-on-shell variants. It is supposed in the last case that the first argument P of the kernels $M_{\alpha\beta}(P, P', z)|_{z \in \Pi_l}$ is taken on the energy ("mass") shells $|P|^2 = z$ or $|p_\alpha|^2 = z - \lambda_{\alpha,j}$, $j = 1, 2, \dots, n_\alpha$. Here, we use the notations $P = \{k_\alpha, p_\alpha\}$, где k_α, p_α , $\alpha = 1, 2, 3$, of the standard relative momenta [19]. Transferring in the expressions obtained all the off-shell terms to the l.h. part and inverting




FIG. 1. Physical, Π_0 , and neighboring unphysical, $\Pi_{(0,1)}$, $\Pi_{(1,1)}$ and $\Pi_{(-1,1)}$, sheets of the Riemann surface in the three-nucleon (nnp) and model three-boson problems. The near (in figure) rim of the cut in the sheet Π_0 is identified in the interval between the deuteron energy E_d and three-body threshold $z = 0$ with the remote rim of the cut in the sheet $\Pi_{(0,1)}$. On the contrary, the remote rim in the sheet Π_0 is identified in this interval with the near rim in the sheet $\Pi_{(0,1)}$. In the interval $(0, +\infty)$, one identifies the near rim of the cut in the sheet Π_0 with the remote rim in the sheet $\Pi_{(1,1)}$. Respectively, the remote rim in the sheet Π_0 is identified in this interval with the near rim in the sheet $\Pi_{(-1,1)}$. On the sheet $\Pi_{(0,1)}$, the virtual level $z_{res}^{(3H)}$ in the s -state of the nnp -system is marked, and the resonance $z_{res}^{(3B)}$ in the model three-boson system. Also, the complex conjugate pole $\bar{z}_{res}^{(3B)}$ is marked.

an operator arising there, one comes to a closed system of equations for the half-on-shell components $M_{\alpha\beta}(P, P', z)|_{z \in \Pi_l}$. This system admits an explicit solution using the terms of the physical sheet only. As a result one gets the following representations* for the matrix $\mathbf{M}(z) = \{M_{\alpha\beta}(z)\}$, $\alpha, \beta = 1, 2, 3$, continued on the sheet Π_l :

$$\mathbf{M}(z)|_{\Pi_l} = \mathbf{M}(z) - \mathbf{B}^\dagger(z)A(z)LS_l^{-1}(z)\tilde{L}\mathbf{B}(z). \quad (1)$$

Here, the factor $A(z)$ is the diagonal matrix,

$$A(z) = \text{diag}\{A_0(z), A_{1,1}(z), \dots, A_{1,n_3}(z)\},$$

combined of the functions $A_0(z) = -\pi iz^2$ and $A_{\alpha,j} = -\pi i\sqrt{z - \lambda_{\alpha,j}}$. Notations L and \tilde{L} are used for diagonal number matrices whose nontrivial elements are the sheet Π_l indices:

$$L = \text{diag}\{l_0, l_{1,1}, \dots, l_{3,n_3}\} \text{ и } \tilde{L} = \text{diag}\{\tilde{l}_0, \tilde{l}_{1,1}, \dots, \tilde{l}_{3,n_3}\}.$$

By $S_l(z)$ we understand a truncation of the three-body scattering matrix: $S(z), S_l(z): \hat{\mathcal{G}} \rightarrow \hat{\mathcal{G}}, \hat{\mathcal{G}} = L_2(S^5) \bigoplus_{\alpha=1}^3 \bigoplus_{j=1}^{n_\alpha} L_2(S^2)$, defined by the equation

$$S_l(z) = \hat{I} + \tilde{L}[S(z) - \hat{I}]L$$

where \hat{I} is the identity operator in $\hat{\mathcal{G}}$. Also, we use the notations

$$\mathbf{B}(z) = \begin{pmatrix} J_0\Omega\mathbf{M} \\ J_1\Psi^*[\Upsilon\mathbf{M} + \mathbf{v}] \end{pmatrix} \text{ and } \mathbf{B}^\dagger(z) = \begin{pmatrix} \mathbf{M}(z)\Omega^\dagger J_0^\dagger, [\mathbf{v} + \mathbf{M}\Upsilon]\Psi J_1^\dagger \end{pmatrix}.$$

Here, $\mathbf{v} = \text{diag}\{v_1, v_2, v_3\}$ with v_α , the pair potentials, $\alpha = 1, 2, 3$. At the same time,

$$\Omega = (1, 1, 1), \quad \Upsilon = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} \text{ and } \Psi = \text{diag}\{\Psi_1, \Psi_2, \Psi_3\}$$

where Ψ_α , $\alpha = 1, 2, 3$, are operators acting on $f = (f_1, f_2, \dots, f_{n_\alpha}) \in \bigoplus_{j=1}^{n_\alpha} L_2(\mathbb{R}^3)$ as

*For the sake of simplicity, we write here these representations as well as representations for the scattering matrix and resolvent (see below Eqs. (2) and (3), respectively) for the case of spinless particles only. A direct generalization of the representations on the case of spin particles causes no difficulties.

$$(\Psi_\alpha f)(P) = \sum_{j=1}^{n_\alpha} \psi_{\alpha,j}(k_\alpha) f_j(p_\alpha),$$

where, in turn, $\psi_{\alpha,j}$ is the bound state wave function of the pair subsystem α corresponding to the level $\lambda_{\alpha,j}$. By Ψ^* we denote operator adjoint to Ψ . Notation $J_0(z)$ is used for operator restricting a function on the energy-shell $|P|^2 = z$. The diagonal matrix-valued function

$$J_1(z) = \text{diag}\{J_{1,1}(z), \dots, J_{3,n_3}(z)\},$$

consists of the operators $J_{\alpha,j}(z)$ of restriction on the energy surfaces $|p_\alpha|^2 = z - \lambda_{\alpha,j}$. The operators Ω^\dagger , $J_0^\dagger(z)$ и $J_1^\dagger(z)$ represent the "transposed" matrices Ω , $J_0(z)$ and $J_1(z)$, respectively. Operators $J_0^\dagger(z)$ и $J_1^\dagger(z)$ act in the expression for \mathbf{B}^\dagger (as if) to the left.

Representations for the scattering matrix and resolvent on unphysical sheets are an immediate consequence of the representations (1) for the matrix $\mathbf{M}(z)|_{\Pi_l}$.

With some stipulations (see [21]) the representations for the scattering matrix $S(z)$ look as

$$S(z)|_{\Pi_l} = \mathcal{E}(l) \left\{ \hat{I} + S_l^{-1}(z)[S(z) - \hat{I}]e(l) \right\} \mathcal{E}(l). \quad (2)$$

Here, $\mathcal{E} = \text{diag}\{\mathcal{E}_0, \mathcal{E}_{1,1}, \dots, \mathcal{E}_{3,n_3}\}$ where \mathcal{E}_0 is the identity operator in $L_2(S^5)$ if $l_0 = 0$ and \mathcal{E}_0 , the inversion, $(\mathcal{E}_0 f)(\hat{P}) = f(-\hat{P})$, if $l_0 = \pm 1$. Analogously, $\mathcal{E}_{\alpha,j}$ is the identity operator in $L_2(S^2)$ for $l_{\alpha,j} = 0$ and inversion for $l_{\alpha,j} = 1$. Notation $e(l)$ is used for diagonal number matrix $e(l) = \text{diag}\{e_0, e_{1,1}, \dots, e_{3,n_3}\}$ with nontrivial elements $e_{\alpha,j} = 1$ if $l_{\alpha,j} = 0$ and $e_{\alpha,j} = -1$ if $l_{\alpha,j} = 1$; for all the cases $e_0 = 1$.

Analytic continuation $R(z)|_{\Pi_l}$ of the resolvent $R(z) = (H - z)^{-1}$ of the three-body Hamiltonian H admits the representation

$$R(z)|_{\Pi_l} = R + \mathbf{Q}^\dagger A L S_l^{-1} \tilde{L} \mathbf{Q}. \quad (3)$$

Here,

$$\mathbf{Q} = \begin{pmatrix} J_0[I - V R] \\ J_1 \Psi^* [I - \Upsilon \mathbf{M} R_0] \Omega^\dagger \end{pmatrix}, \quad \mathbf{Q}^\dagger = ([I - R V] J_0^\dagger, \Omega [I - R_0 \mathbf{M} \Upsilon] \Psi J_1^\dagger)$$

with $V = v_1 + v_2 + v_3$ and $R_0(z) = (H_0 - z)^{-1}$, the resolvent of the kinetic energy operator H_0 for the system under consideration.

There were holomorphy domains $\Pi_l^{(\text{hol})}$ found in [21] for the truncated scattering matrices $S_l(z)$ in the physical sheet. Representations (1) and (3) are valid in the same domains.

It follows from the representations (1)—(3) that the resonances (the non-trivial poles of $M(z)|_{\Pi_l}$, $S(z)|_{\Pi_l}$ and $R(z)|_{\Pi_l}$) situated on the unphysical sheet Π_l are in fact those points $z = z_{\text{res}}$ on the physical sheet where the matrix $S_l(z)$ has zero as eigenvalue. Therefore, a calculation of resonances on the unphysical sheet Π_l is reduced to a search for zeros of the truncation $S_l(z)$ of the total three-body scattering matrix $S(z)$ in the physical sheet. At the numerical search for the resonances, one can use any method allowing to find analytic continuation on the physical sheet, of the elastic scattering, rearrangement or breakup amplitudes necessary to construct the respective truncation $S_l(z)$.

III. RESULTS OF SEARCH FOR RESONANCES IN THREE-NUCLEON (nnp) AND THREE-BOSON SYSTEMS

One of the most effective methods for a study of concrete three-particle systems is the numerical algorithm [22] (see also [23], [19] and references therein) based on the Faddeev differential equations for the wave function components in the configuration space. This method gives opportunity of comparatively easy calculations of the scattering wave functions and respective amplitudes for $(2 \rightarrow 2, 3)$ processes. An extension of the differential formulation to a domain of complex energies enables us to calculate analytic continuation of the $(2 \rightarrow 2, 3)$ amplitudes on the physical sheet. This is quite enough for us to construct the truncated s -state scattering matrices $S_l(z)$ whose zeros are resonances on the respective *two-body* unphysical sheets Π_l with $l_0 = 0$.

As to a search for resonances on the *three-body* unphysical sheets Π_l with $l_0 \neq 0$ (in particular on the sheets $\Pi_{(-1,1)}$ and $\Pi_{(1,1)}$ in the three-nucleon problem), the situation is much more complicated. To construct $S_l(z)$ in this case, one has, alongside with $(2 \rightarrow 2)$, to calculate amplitudes for the processes $(3 \rightarrow 2, 3)$ with three asymptotically free particles in an initial state. Unfortunately, the reliable practical methods for calculation of the processes $(3 \rightarrow 2, 3)$ are not developed so far even for the real energies. In particular, when using the differential formulation of the scattering problem, one has from the very beginning to separate explicitly contributions to the Faddeev components not only from single-rescattering but also from double-rescattering processes [19]. In the last case, one has to take into account explicitly (see [19]) a presence of the "light" and "shadow" zones for the correspondent waves, and to use the Fresnel integral for description of intermediate regimes. Moreover, computations of the $(3 \rightarrow 2, 3)$ amplitudes have to be carried out for many different directions of the incident momentum P . This circumstance is itself a factor enlarging numerical complexity as compared with the case of the processes $(2 \rightarrow 2, 3)$.

So, in the present work we restrict ourselves with a search for the nnp system resonances situated on the unphysical sheet $\Pi_{(0,1)}$ only, connected with the physical one by crossing the continuum spectrum interval $(E_d, 0)$ between

the deuteron energy $z = E_d$ and breakup threshold $z = 0$ (see Fig. 1). To construct the truncated scattering matrix $S_{(0,1)}(z)$ (in accordance with (1)–(3) just its zeros represent resonances on the sheet $\Pi_{(0,1)}$), it suffices to calculate the elastic nd scattering amplitudes only.

To find these amplitudes we use the two-dimensional Faddeev integro-differential equations [19] being a result of the partial and angular analysis of the Faddeev differential equations. In addition we make an assumption, rather usual in nuclear physics, that the nucleons involved interact in the s -state only. With this assumption the partial equations become exact. The Faddeev component U_L^q corresponding to the total spin $\mathbf{S} = 3/2$ and total orbital momentum L , satisfies the equation

$$(H_L - z)U_L^q(x, y) = V_t(x)\Psi_L^q(x, y) \quad (4)$$

with $H_L = -\frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} + \frac{L(L+1)}{y^2}$, the partial Laplacian and x, y , the absolute values of the Jacobi vectors [19]. Respective partial component $\Psi_L^q(x, y)$ of the total wave function for the npn -system is expressed by the function $U_L^q(x, y)$ as

$$\Psi_L^q(x, y) = U_L^q(x, y) - \frac{1}{2} \int_{-1}^1 du h^L(x, y, u) U_L^q(x', y') \quad (5)$$

where $x' = \sqrt{\frac{1}{4}x^2 + \frac{3}{4}y^2 - \frac{\sqrt{3}}{2}xyu}$ and $y' = \sqrt{\frac{3}{4}x^2 + \frac{1}{4}y^2 + \frac{\sqrt{3}}{2}xyu}$. The geometric function h^L is given by

$$h^L = \frac{4}{\sqrt{3}} \frac{xy}{x'y'} \left(\frac{-1}{2 \sin \theta'} \right)^L \sum_{k=0}^L \frac{L!}{k!(L-k)!} P_k(u) \left(\sqrt{3} \cos \theta \right)^k (\sin \theta)^{L-k}$$

with $P_k(u)$, the Legendre polynomial of the order k and $\theta = \arctg \frac{y}{x}$, $\theta' = \arctg \frac{y'}{x'}$. The factor $V_t(x)$ represents a triplet part of the nucleon-nucleon interaction potential.

Function $U_L^q(x, y)$ satisfies the boundary conditions

$$U_L^q(x, y)|_{x=0} = 0 \quad \text{и} \quad U_L^q(x, y)|_{y=0} = 0. \quad (6)$$

It satisfies also the asymptotical conditions

$$U_L^q(x, y) \underset{\rho \rightarrow \infty}{\sim} \psi_d(x) \left[j_L(\sqrt{z - E_d} y) + a_L^q(z) \exp \left\{ i\sqrt{z - E_d} y + i\frac{\pi L}{2} \right\} \right] + A_L^q(z, \theta) \frac{\exp \left\{ i\sqrt{z} \rho + i\frac{\pi L}{2} \right\}}{\sqrt{\rho}} \quad (7)$$

where j_L is the Bessel spherical function of the order L and $\rho = \sqrt{x^2 + y^2}$. By $\psi_d(x)$ we denote the deuteron wave function and by $a_L^d(z)$, the quartet amplitude for elastic nd -scattering in a state with angular momentum L . The function $A_L^q(z, \theta)$ represents the partial Faddeev component of the breakup amplitude of this system into three particles.

The Faddeev integro-differential equations for the doublet ($S = 1/2$) nd -scattering look as

$$(H_L - z)U_{i(2),L}^d(x, y) = V_{i(s)}(x)\Psi_{i(2),L}^d(x, y) \quad (8)$$

where $\Psi_L^d, \Psi_L^q = \begin{pmatrix} \Psi_{1,L}^d \\ \Psi_{2,L}^d \end{pmatrix}$, is expressed by the vector $U_L^d = \begin{pmatrix} U_{1,L}^d \\ U_{2,L}^d \end{pmatrix}$ as

$$\Psi_L^q = U_L^q + \frac{1}{2} \int_{-1}^1 du h^L(u) B U_L^d, \quad B = \begin{pmatrix} \frac{1}{4} & -\frac{3}{4} \\ -\frac{3}{4} & \frac{1}{4} \end{pmatrix}. \quad (9)$$

By V_s we understand a singlet part of the nucleon-nucleon interaction potential.

The Faddeev partial components $U_{i,L}^d$ satisfy as U_L^q , the boundary conditions

$$U_{i,L}^d(x, y)|_{x=0} = 0, \quad U_{i,L}^d(x, y)|_{y=0} = 0, \quad (10)$$

and have the asymptotics

$$U_{i,L}^d(x, y) \underset{\rho \rightarrow \infty}{\sim} \delta_{i1} \psi_d(x) \left[j_L(\sqrt{z - E_d} y) + a_L^d(z) \exp \left\{ i\sqrt{z - E_d} y + i\frac{\pi L}{2} \right\} \right] + A_{i,L}(z, \theta) \frac{\exp \left\{ i\sqrt{z} \rho + i\frac{\pi L}{2} \right\}}{\sqrt{\rho}}, \quad i = 1, 2. \quad (11)$$

Here, $a_L^d(z)$ is the doublet elastic nd -scattering amplitude in the state of the nnp -system with angular momentum L . At the same time, the functions $A_{1,L}^d$ and $A_{2,L}^d$ represent the partial Faddeev components of the breakup amplitude for this state. Remind that the physical breakup amplitudes in quartet and doublet states are expressed by the amplitudes respectively, A_L^q and $A_{1,L}^d, A_{2,L}^d$ via relations analogous to (5) and (9) (see e. g., [19]).

A component of the truncated scattering matrix $S_{(0,1)}$ in the state with a fixed angular momentum L is diagonal. Its nontrivial quartet, $s_L^q(z)$, and doublet, $s_L^d(z)$, elements are given by

$$s_L^q(z) = 1 + 2ia_L^q(z),$$

TABLE I. Convergence of the nnp -system virtual level $z_{res}(^3H)$

N	$N_\theta^{(1)}$	$N_\theta^{(2)}$	$N_\theta^{(3)}$	ρ_1 , fm	ρ_{max} , fm	$z_{res}(^3H)$, MeV
40	10	20	30	0.40	19.0	2.9296
60	15	30	45	0.30	23.9	2.8229
80	20	40	60	0.25	29.9	2.7565
120	30	60	90	0.15	34.0	2.7434
160	40	80	120	0.10	39.1	2.7282
180	45	90	135	0.08	39.5	2.7275

$$s_L^d(z) = 1 + 2ia_L^d(z).$$

Therefore, the sheet $\Pi_{(01)}$ resonances in the state L of the nnp -system are in fact, zeros of the *scalar* functions $s_L^q(z)$ and $s_L^d(z)$.

When solving the boundary-value problems (4), (6), (7) and (8), (10), (11) numerically at complex energies z , we use the same algorithm [19], [22], [23] as at real z . First, we make a finite-difference approximation of the problems above in polar coordinates ρ , θ . As mesh points we take the intersection points of the arcs $\rho = \rho_i$, $i = 1, 2, \dots, N_\rho$, and rays $\theta = \theta_j$, $j = 1, 2, \dots, N_\theta$, $\theta_j < \theta_{j+1}$. At $\rho_0 = 0$ and given ρ_1 (see Table I), the mesh points in ρ for $i \geq 2$ are chosen in such a way that $\rho_{i+1} = \rho_i + a(\rho_i - \rho_{i-1})$ with a parameter a (acceleration) not depending on the number of a mesh point. All the results expressed below, are related to $a = 1,01$. The interval $[0, \pi/2]$ where the variable θ changes, is divided by special points with numbers $N_\theta^{(k)}$, $k = 1, 2, 3$, into three subintervals, inside of which the mesh points in θ are distributed uniformly. When going from one interval to another (in direction of the parameter θ rise), the mesh step is divided by two. The choice of the mesh described, is explained by a necessity to take into account essentially more quick change of the Faddeev component values in the domain where ρ and/or x [19] are small. Usually, we chose the numbers of points in θ and ρ the same, $N_\theta = N_\rho$. A maximal value of the parameters N_θ , N_ρ has equaled to 180. With these values, the cut-off radius $\rho_{max} = \rho_{N_\rho}$ has reached 39 fm. Typical values of the mesh parameters are given in Table I where $N = N_\theta = N_\rho$.

As a nucleon-nucleon interaction, the Malfliet-Tjon potential MT I III is used in its initial version [29].

Having solved the problems (4), (6), (7) and (8), (10), (11) we calculate the functions $s_L^q(z)$ and $s_L^d(z)$. Resonances, considered as roots of these functions in the complex plane, are found using the Newton method with a three-point approximation of derivative.

As a test of the computer code we have calculated the bound-state energy E_t of the 3H nucleus as a pole of the function $s_L^d(z)$. More precisely, this pole

was found as a root of the inverse amplitude $1/a_L^q(z)$. Beginning with mesh dimension 80×80 , we obtained for E_i the value -8.55 MeV. Hereafter all the energies are given with respect to the breakup threshold. Note that the value stated for E_i is in a good agreement with known results on E_i in the MT I-III model (see [26]).

When searching for the np resonances on the sheet $\Pi_{(0,1)}$ at $L = 0, 1$, we have inspected a domain of a range about 10 MeV in vicinity of the segment $[E_d, 0]$ in the complex z plane. Especially carefully we studied a vicinity of the points $z = -1.5 \pm 0.3 + i(0.6 \pm 0.3)$ MeV interpreted in the recent works [30], [31] as a location of an excited state energy of the 3H nucleus. Unfortunately we have succeeded to find only one root $z_{\text{res}}({}^3H)$ of the function $s_0^d(z)$, corresponding to the known virtual state of the np system at the total spin $S=1/2$. Position of this root for different meshes is shown in Table I illustrating a degree of convergence in the method used. As one can see from the table, for the maximal of the meshes examined, the mesh 180×180 , we have found $z_{\text{res}}({}^3H) = -2.728$ MeV. This means that the calculated virtual level is situated 0.504 MeV to the left from the nd threshold $E_d = -2.224$ MeV (in the MT I-III model [29]). Note that the shift $E_d - z_{\text{res}}({}^3H)$ found from experimental data on nd scattering, is 0.515 MeV (see [13]). Its value computed in a separabilized MT I-III model on the base of the momentum space Faddeev equations, equals to 0.502 MeV [13]. As to the resonance $z = -1.5 \pm 0.3 + i(0.6 \pm 0.3)$ MeV at $L = 0$ discussed in [30], [31], it is quite possible to be situated not on the *two-body* sheet $\Pi_{(0,1)}$ but on the *three-body* sheets $\Pi_{(-1,1)}$ or $\Pi_{(1,1)}$ (see Fig. 1). Due to the reasons mentioned at the beginning of the section, we may unfortunately neither approve nor disprove this hypothesis. As it should be expected (see the data on three-nucleon resonances in [13]), we have failed to find any resonances in the quartet state at $L = 0$ as well as at $L = 1$.

Also, we have studied a model three-body system including identical spin-zero bosons with masses of the nucleon. Remind that the Faddeev integro-differential equations for such a three-boson system look exactly as the equations (4) except a necessity to replace the factor $-1/2$ in the expression (5) with unity. The boundary conditions for respective partial Faddeev components $U_L^{3B}(x, y)$ have the form (6) and (7) where instead of $a_L^q(z)$ and $A_L^q(\theta, z)$, one has to substitute $a_L^{3B}(z)$ and $A_L^{3B}(\theta, z)$.

Component of the truncated scattering matrix $S_{(0,1)}$ for the three-boson system is given in the state with the angular momentum L by

$$s_L^{3B}(z) = 1 + 2ia_L^{3B}(z).$$

Resonances on the sheet $\Pi_{(0,1)}$ in this state are roots of the equation $s_L^{3B}(z) = 0$ considered on the physical sheet.

In the three-boson problem we restrict ourselves with a treating the s -state only and thereby, with a searching for zeros of the function $s_L^{3B}(z)$ at $L = 0$.




FIG. 2. Surface of the function $|s_0^{3B}(z)|$ in the model system of three bosons with the nucleon masses. The potential $V^G(r)$ is used with the barrier $V_b = 1.5$ MeV. Position of the resonance $z_{\text{res}}(3B)$ corresponds to the minimal (zero) value of $|s_0^{3B}(z)|$.




FIG. 3. Trajectory of the resonance $z_{\text{res}}(3B)$ on the sheet $\Pi_{(0,1)}$ in the model system of three bosons with the nucleon masses. The potential $V^G(r)$ is used. Values of the barrier V_b in MeV are given near the points marked on the curve. These points are distributed between $V_b = 0.85$ MeV and $V_b = 1.6$ MeV with the same step 0.05 MeV in V_b .




FIG. 4. Dependence of the "deuteron" energy E_d (curve 1) and real part of the resonance $z_{\text{res}}(3B)$ (curve 2) on the barrier value V_b .

As a pairwise interaction between the bosons we have used the Gauss-type potential supplied with an additional Gauss repulsive barrier term,

$$V^G(r) = V_0 \exp[-\mu_0 r^2] + V_b \exp[-\mu_b (r - r_b)^2]$$

where the values $V_0 = -55$ MeV, $\mu_0 = 0.2$ fm $^{-2}$, $\mu_b = 0.01$ fm $^{-2}$, $r_b = 5$ fm have been fixed and the barrier amplitude V_b varied. A resonance (with non-zero imaginary part) on the sheet $\Pi_{(0,1)}$ arises in the system concerned just due to the presence of the barrier term. Example of a surface of the $s_0^{3B}(z)$ absolute value for the barrier amplitude $V_b = 1.5$ MeV is shown in Fig. 2 (for a 80×80 mesh). A trajectory of the resonance $z_{\text{res}}(3B)$ (a zero of the function $s_0^{3B}(z)$) is shown for the changing barrier V_b in Fig. 3. This trajectory was watched for the barrier V_b decreasing in the interval between 1.6 MeV and 0.84 MeV. When drawing the trajectory, we have used a 160×160 mesh. It can be seen from Fig. 3 that a behavior of the resonance concerned turns out to be rather unexpected: with monotonously decreasing real part, the imaginary part of the resonance changes not monotonously and at $V_b \cong 1.15$ MeV it has an obvious minimum. There is this value of V_b when real part of the resonance $z_{\text{res}}(3B)$ coincides with the "deuteron" energy E_d (see Fig. 4). When $\text{Re } z_{\text{res}}(3B)$, decreasing with V_b , leaves E_d behind, the value of $\text{Im } z_{\text{res}}(3B)$ increases at the beginning but then, after $V_b \cong 1$ MeV falls off again until the resonance crossing at $V_b \cong 0.84$ MeV the real axis. Trajectory of the resonance

concerned in the lower complex half-plane is symmetric to the curve shown in Fig. 3 with respect to the real axis. Respective points, symmetric to those marked in Fig. 3, correspond to the same values of V_6 .

ACKNOWLEDGMENTS

The authors are grateful to Professor V.B. Belyaev for valuable discussions and to Dr. F.M. Penkov for useful remarks. One of the authors (A.K.M.) is indebted to the International Science Foundation and Russian government for financial support of the work (Grant RFB300).

-
- [1] *Baz A., Zeldovich Ya., Perelomov A.* Scattering, reactions and decays in non-relativistic quantum mechanics. Jerusalem: Israel Program for Scientific Translations, 1969.
 - [2] *Newton R.G.*, Scattering theory of waves and particles. McGraw Hill, New York, 2nd ed., 1982.
 - [3] *Alfaro V. de, Regge T.* Potential scattering [Russian translation]. Mir, Moscow, 1986.
 - [4] *Böhm A.* Quantum Mechanics: Foundations and Applications. Springer-Verlag, 1986.
 - [5] *Kukulin V.I., Krasnopol'sky V.M., Horáček J.* Theory of resonances: Principles and applications. — Praha: Academia, 1989.
 - [6] *Reed M., Simon B.* Methods of modern mathematical physics. III: Scattering theory. — N.Y.: Academic Press, 1979.
 - [7] *Reed M., Simon B.* Methods of modern mathematical physics. IV: Analysis of operators. — N.Y.: Academic Press, 1978.
 - [8] *Albeverio S., Gesztesy F., Høegh-Krohn R., Holden H.* Solvable models in quantum mechanics. Springer-Verlag, 1988.
 - [9] *Simon B.*// Intern. J. Quant. Chem. 1978. V. 14. P. 529-542.
 - [10] *Gamow G.*// Z.Phys. 1928. V. 51. P. 204-212.
 - [11] *Jost R.*// Helv. Phys. Acta. 1947. V. 20. P. 250-266.
 - [12] *Titchmarsh E.C.* Eigenfunction expansions associated with second order differential equations, Vol. II. — London: Oxford U.P., 1946.
 - [13] *Möller K., Orlov Yu. V.* //Fiz. Elem. Chast. At. Yadra. 1989. V. 20. P. 1341-1395.
 - [14] *Balslev E., Combes J.M.*// Commun.Math.Phys. 1971. V. 22. P. 280-294.
 - [15] *Hu C.-Y., Bhatia A.K.*// Muon Catalyzed Fusion. 1990/91. V. 5/6. P. 439-444.
 - [16] *Korobov V.I.*// Intern. Sympos. on Muon Catalyzed Fusion, Dubna June 19-24, 1995. Book of Abstracts. Dubna: JINR, 1995. P. 77.

- [17] *Szótó A.*// Contributed papers to 14th Intern. Conf. on Few Body Problems in Physics. — Williamsburg: CEBAF, 1994. P. 777-780; *Szótó A., Oberhummer H., Pichler R.* Searching for three-nucleon resonances.// LANL E-print nucl-th/9510017, 1995.
- [18] *Faddeev L.D.*// Trudy Mat. In-ta AN SSSR. 1963. V. 69. P. 1-125 [English translation: Mathematical aspects of the three-body problem in quantum mechanics. Israel Program for Scientific Translations, Jerusalem, 1965].
- [19] *Faddeev L.D., Merkuriev S.P.* Quantum scattering theory for several particle systems. — Dodrecht: Kluwer Academic Publishers, 1993.
- [20] *Motovilov A.K.*// Contributed papers to 14th Intern. Conf. on Few Body Problems in Physics. — Williamsburg: CEBAF, 1994. P. 816-819.
- [21] *Motovilov A.K.* Representations for three-body T-matrix on unphysical sheets, I, II.// Preprints JINR E5-95-45 (LANL E-print nucl-th/9505028) and JINR E5-95-46 (LANL E-print nucl-th/9505029). JINR, Dubna, 1995.
- [22] *Merkuriev S.P., Gignoux C., Laverne A.*// Ann. Phys. (N.Y.). 1976. V. 99. P. 30-71.
- [23] *Kvitsinsky A.A., Kuperin Yu.A., S.P.Merkuriev S.P., Motovilov A.K., Yakovlev S.L.*// Fiz. Elem. Chast. At. Yadra. 1986. V. 17. P. 267-317.
- [24] *Benayoun J.J., Gignoux C., Chauvin J.*// Phys. Rev. C. 1981. V. 23. P. 1854-1857.
- [25] *Chen C.R., Payne G.L., Friar J.L., Gibson B.F.*// Phys. Rev. C. 1989. V. 39. P. 1264-1268; Phys. Rev. C. 1991. V. 44. P. 50-59.
- [26] *Carbonell J., Gignoux C., Merkuriev S.P.*// Few-Body Systems. 1993. V. 15. P. 15-23.
- [27] *Yakovlev S.L., Filikhin I.N.*// Yad. Fiz. 1993. V. 56. P. 98-106.
- [28] *Motovilov A.K.*// Teor. Mat. Fiz. 1993. V. 95. P. 427-438.
- [29] *Malfliet R.A., Tjon J.A.*// Nucl. Phys. A. 1969. V. 127. P. 161-168.
- [30] *Alexandrov D.V., Nikolsky E.Yu., Novatsky B.G., Stepanov D.N.*// Pis'ma ZhETF. 1994. V. 59. P. 301-304.
- [31] *Barabanov A.L.* Does the excited state of the ^3H nucleus exist? // Preprint IAE-5811/2. Moscow: RRC "Kurchatov Institute", 1994.

Received by Publishing Department
on January 23, 1996

Использование дифференциальных уравнений Фаддеева для расчетов трехчастичных резонансов

На основе явных представлений для аналитического продолжения компонент Фаддеева трехчастичной T -матрицы на нефизические листы римановой поверхности энергии разработан алгоритм расчета резонансов в задаче трех квантовых частиц. Согласно представлениям полюса T -матрицы матрицы рассеяния и функции Грина на нефизических листах, интерпретируемые как резонансы, совпадают с теми значениями энергии, для которых подходящие усечения матрицы рассеяния на физическом листе имеют собственное число нуль. Амплитуды процессов на физическом листе, необходимые для построения матрицы рассеяния, вычисляются с помощью дифференциальных уравнений Фаддеева. Разработанный алгоритм применяется для поиска резонансов в системе $m\bar{p}$ и модельной системе трех бозонов.

Работа выполнена в Лаборатории вычислительной техники и автоматизации и Лаборатории теоретической физики им. Н.Н.Боголюбова ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 1996

Use of the Faddeev Differential Equations for Computations of Three-Body Resonances

Algorithm, based on explicit representations for analytic continuation of the T -matrix Faddeev components on unphysical sheets, is worked out for calculations of resonances in the three-body quantum problem. According to the representations, poles of T -matrix, scattering matrix and Green function on unphysical sheets, interpreted as resonances, coincide with those complex energy values where appropriate truncations of the scattering matrix have zero as eigenvalue. Scattering amplitudes on the physical sheet, necessary to construct scattering matrix, are calculated on the basis of the Faddeev differential equations. The algorithm developed is applied to search for the resonances in the $m\bar{p}$ system and in a model three-boson system.

The investigation has been performed at the Laboratory of Computing Techniques and Automation and Bogoliubov Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research, Dubna, 1996

Макет Т.Е.Попеко

Подписано в печать 12.02.96
Формат 60 × 90/16. Офсетная печать. Уч.-изд.листов 1,71
Тираж 370. Заказ 48874. Цена 2052 р.

Издательский отдел Объединенного института ядерных исследований
Дубна Московской области