

AE-414

Radiolysis of Aqueous Toluene Solutions

H.C. Christensen and
R. Gustafson

This report is intended for publication in a periodical. References may not be published prior to such publication without the consent of the author.



AKTIEBOLAGET ATOMENERGI

STUDSVIK, NYKÖPING, SWEDEN 1971

RADIOLYSIS OF AQUEOUS TOLUENE SOLUTIONS

H. C. Christensen and R. Gustafson

ABSTRACT

Aqueous toluene solutions have been irradiated with ^{60}Co γ -rays. In unbuffered solutions the various cresol isomers are formed in a total yield of 0.45, 0.87 and 0.94 molecules/100 eV absorbed energy in argon-, N_2O - and air-saturated solutions, respectively. The yields are reduced in acid (pH 3) solutions ($G = 0.14, 0.14$ and 0.52 , respectively) but the reduction is compensated by the formation of 1,2-diphenylethane in yields of 0.49 and 1.60 in argon- and N_2O -saturated solutions, respectively. Benzyl radicals are formed through an acid catalysed water elimination reaction from the initially formed hydroxymethylcyclohexadienyl radical. Phenyltolylmethanes, dimethylbiphenyls and partly reduced dimers are also formed during the radiolysis. Hydrogen is formed in the same yield as the molecular yield, $g(\text{H}_2)$. Xylene isomers and benzene are formed in trace quantities. The most remarkable effects of the addition of Fe(III) ions to deaerated acid toluene solutions are the formation of benzyl alcohol and benzaldehyde and an increase in the yield of 1,2-diphenylethane.

LIST OF CONTENTS

	<u>Page</u>
1. INTRODUCTION	3
2. EXPERIMENTAL	4
2.1 Material	4
2.2 Preparation of samples and irradiation	4
2.3 Analysis	4
3. RESULTS AND DISCUSSION	5
3.1 Deaerated solutions	5
3.2 The effect of iron(III) ions	10
3.3 Aerated solutions	12
4. CONCLUSIONS	13
5. ACKNOWLEDGEMENT	14
6. REFERENCES	15

1. INTRODUCTION

The oxidation of toluene in aqueous solutions by OH radicals produced by ionizing radiation or by Fenton's reagent has been studied previously (1 - 6). Khenokh and Lapinskaia (1) reported the formation of formaldehyde ($G = 0.17$) and phenols ($G = 0.56$) after γ -irradiation of aerated aqueous toluene solutions. M. A. B. Manzitti et al. (2) found 1,2-diphenylethane as the main product but phenyltolylmethanes, 4,4'-dimethylbiphenyl and o- and m-cresols were also found after irradiation with γ -rays in oxygen-free solutions. The investigations of Hotta et al. (3 - 5) were mainly concerned with the autoxidation of toluene in aerated aqueous solutions catalysed by iron(II) ions at elevated temperatures ($\geq 70^{\circ}\text{C}$). The pH of the solution was found by Jefcoate et al. (6) to have a marked effect on the product distribution when toluene was oxidized by Fenton's reagent. The effect was explained as an acid catalysed water elimination reaction from the initially formed OH adduct, resulting in the formation of benzyl radicals.

In pulse radiolysis studies the initial radicals formed by reaction of OH radicals (7) and H atoms (8) with toluene in aqueous solutions have been studied and the radicals were interpreted as the OH and H adducts, respectively.

In the present investigation we have attempted a more complete product analysis both of the oxygen containing products and of the complex dimer group. We have also made an effort to find some of the reduced products formed, such as partly hydrated toluene and toluene dimers with the purpose of improving the balance between oxidized and reduced products.

2. EXPERIMENTAL

2.1 Material

Toluene (Merck p. a.) was further purified by recrystallisation twice. The water used for irradiations was quadruply distilled. For identification purposes the three isomers of phenyltolylmethane, which were not commercially available were synthesized by a Friedel-Crafts reaction described by Errede and Cassidy (9).

2.2 Preparation of samples and irradiation

Quadruply distilled water was deaerated, saturated with purified argon or nitrous oxide, transferred into carefully rinsed 50 ml syringes and subsequently saturated with toluene by procedures previously described (10). The samples were irradiated in the syringes at ambient temperature with ^{60}Co γ -rays at a dose rate of 11,800 rad/min and with a total dose normally within the range 20,000 - 220,000 rad.

2.3 Analysis

Hydrogen peroxide was determined by the method of Ghormley (11) and hydrogen by use of a van Slyke apparatus for the measurement of the gas-phase and subsequent analysis on a gas chromatograph using a column filled with molecular sieve. The yield of iron(II) ions was determined by the method of Hummel and Spinks (12). The organic products were extracted from a 50 ml aqueous solution with 3 x 8 ml n-hexane and then with 3 x 3 ml ether and analysed on a Perkin Elmer 880 gas chromatograph equipped with 5 m long 1/8" OD columns filled with 5 % of either Emulphor O or Silicone DC 710 on 80 - 100 mesh

Chromosorb W. These columns are the same as those used by Hoigné and Gäumann in their study of the radiolysis of toluene in liquid phase (13). The ether phase, which contained cresols and benzyl alcohol, was analysed on a 2 m long column filled with 5 % tri-(2,4-xylenyl) phosphate on 60 - 100 mesh Embacel. Analysis was also carried out on a combined gas chromatograph-mass spectrometer (LKB 9000) equipped with a 2.7 m long column filled with 1 % Silicone SE 30, in order to identify some of the partly hydrated products. For the quantitative analysis phenol and 4,4'-dimethylbiphenyl were used as internal standards.

Ultraviolet spectra of irradiated solutions were also recorded.

3. RESULTS AND DISCUSSION

3.1 Deaerated solutions

Deaerated, unbuffered aqueous toluene solutions saturated with argon or nitrous oxide were irradiated and the resulting yields are shown as a function of dose in Figs. 1 and 2. In Figs. 3 and 4 are shown the yields obtained in solutions acidified with perchloric acid to pH 3. The G values of the products of radiolysis under various conditions are collected in Table 1, which gives the values for the various isomers, whereas the figures only give the total amount of isomers within a group.

From the table and the figures it may be observed that in unbuffered solutions the N_2O saturated solutions give a two times higher yield of hydroxytoluenes than the argon-saturated solutions, whereas in acid solutions the yield is unaffected by the presence of N_2O and also considerably lower than in unbuffered solutions. The lower yield

Table 1. G values from radiolysis
of aqueous toluene solutions



	Unbuff. solutions saturated with		Acid solutions saturated with		1mM Fe(III), acid solutions sat. with		Aerated solutions	
	Ar	N ₂ O	Ar	N ₂ O	Ar	N ₂ O	Unbuff.	Acid
H ₂ O ₂	0	0.76	0.55	0.33	0	0	2.3/neg	3.0/0.16
H ₂	0.48	0.38	0.49	0.36	0.41	0.36	0.44	0.48
Acid	0	0	0	0			1.1/0.03	1.7/0
Benzaldehyde	0	0	0	0	0.14	0.16	0	0.33/0.03
o-Cresol	0.22	0.48	0.08	0.08	0.09	0.22	0.35/	0.22/
p-Cresol	0.14	0.26	0.03	0.04	0.05	0.11	0.34/	0.17/
m-Cresol	0.09	0.13	0.03	0.02	0.01	0.03	0.25/	0.13/
Σ Cresols	0.45	0.87	0.14	0.14	0.15	0.36	0.94/0.20	0.52/0.03
Benzyl alcohol	0.02	0.06	0.03	0.05	0.56	0.51	0.03	0.03
1,2-diphenyl-ethane	0.02	0.02	0.49	1.60	1.03	2.07	0/0.17	0/0.54
Phenyltolyl-methanes	0.08	0.46	0.18	0.36	0.06	0.23	0/0.41	0/0.15
Dimethyl-biphenyls	0.29	1.10	0.14	0.06	0	0	0/0.26	0/0.10
Reduced Dimers	0.05	0.01	0.57	0.22	0	0	0/0.11	0/0.40
Σ Dimers	0.44	1.59	1.38	2.24	1.09	2.30	0/0.95	0/1.19
2Σ Dimer+ other organic products	1.35	4.11	2.93	4.67	3.03	5.62	2.07/2.16	2.58/2.47
Decomp. of toluene	3.0	5.8	3.9	5.9	3.30	5.7	3.0	3.1








The results after the slant are the yield after the break point of the curves (i. e. the yields after consumption of the dissolved oxygen).

of cresols in acid solutions is compensated by a considerably higher yield of dimers, especially 1,2-diphenylethane. The yield of benzyl alcohol is low in all cases (~ 0.05). The yield of the various isomers of cresol was in the order ortho > para > meta. It may also be observed that the reduced dimers are formed preferentially in acid solutions, probably due to the reaction of H atoms.

The order of the yield of isomers of phenyltolylmethane was ortho > meta > para with a para isomer yield of 15 - 20 % of the total yield. The separation of the ortho and meta isomers on the gas chromatograph column was poor, so only the total yield was determined together with the yield of the para isomer. Of the possible dimethylbiphenyl isomers the 4,4' isomer was not produced and the 2,2' only in trace quantities (~ 5 % of the total yield of the isomers), while the 2,3' and 3,3' isomers were formed in equal amounts and with a yield of about 80 % of the total yield of isomers in unbuffered solutions. In acid solutions the 2,2' and 2,3' isomers predominated.

The following reaction scheme may account for the products





In pulse radiolysis investigations of aqueous toluene solutions Dorfman et al. (7) and Sauer and Ward (8) have shown that OH radicals and H atoms react with toluene to form the cyclohexadienyl radicals and they excluded any major contribution of the benzyl radical to their transient spectra. However, the presence of 1,2-diphenylethane among the radiolysis products, especially in acid solution, is a strong indication that the benzyl radical is formed. From the table it can be seen that the yield of 1,2-diphenylethane in nitrous oxide-saturated solutions is increased much more than the yield of H atoms when the solution is acidified. (At pH 3 about 80 % of the electrons react with N₂O in reaction 3 and only 20 % in reaction 2 to form H atoms (10).) So the increase in the yield of 1,2-diphenylethane is probably caused by an acid-catalysed process such as reaction 11, as proposed

by Jefcoate et al. in a study of the oxidation of toluene by Fenton's reagent (6). The reason why such a first order decay of the hydroxymethylcyclohexadienyl radical was not observed by Sauer and Ward could be the much higher dose rate used by them, favouring the bimolecular decay.

The benzyl radical is probably not formed through the reaction of H atoms, as seen by comparison of the yield of 1,2-diphenylethane, $G(B_2)$, in nitrous oxide-saturated ($G(B_2) = 1.60$) and in argon-saturated solutions ($G(B_2) = 0.49$), the yield of H atoms in the latter case being 3 times higher than in the first case.

The observed yield of reduced dimers (with mole weight = 184) is too low to balance the sum of oxidized products, indicating the presence of non-observed reduced products e.g. reduced monomers analogous to the formation of cyclohexadiene, cyclohexene and cyclohexane by irradiation of aqueous benzene solutions (14). An analysis using a 50 m capillary column in a gas chromatograph did not reveal any hydrogenated toluenes, but showed the formation of xylene isomers in trace quantities.


3.2 The effect of iron(III) ions

Deaerated aqueous toluene solutions saturated with argon or nitrous oxide and containing 10^{-3} M iron(III) ions at pH 3 were irradiated and the results are shown in Table 1 and in Figs. 5 and 6. The most remarkable effects of iron(III) are the tenfold increase in the yield of benzyl alcohol, the formation of benzaldehyde, the increase of the yield of 1,2-diphenylethane and the decrease to zero of the yield of reduced dimers.

The reduction of the yield of reduced dimers cannot be caused by the removal of H atoms by Fe(III) ions according to



because toluene [$c = 6 \cdot 10^{-3}$ M, $k_4 = 3.9 \cdot 10^9$ (8)] competes favourably with iron(III) ions [$c = 10^{-3}$ M, $k_{16} = 10^8$ (15)] for the reaction of H atoms. The formation of iron(II) ions in a yield of 1.3 and 2.3 in nitrous oxide- and argon-saturated solutions, respectively, is therefore probably caused by reaction of Fe(III) ions with toluene radicals, resulting in the formation of benzyl alcohol and benzaldehyde according to



if the $\text{OHCH}_3\text{C}_6\text{H}_5$ radical is formed. However, as the iron(III) ions increased the yield of 1,2-diphenylmethane and also eliminated the reduced dimers, the oxidation of $\text{CH}_3\text{C}_6\text{H}_6$ radicals by iron(III) ions may be possible:



As, according to the reaction mechanism, the hydroxymethyl-cyclohexadienyl radical causes the formation of cresols (Cr), benzyl alcohol (BA), benzaldehyde (BAld), 1,2-diphenylethane (B₂) and phenyltolymethanes (BT), the total yield, Σ, of these products was calculated, counting the yield of 1,2-diphenylmethane twice. The total yield together with the yields of H and OH in argon- and nitrous oxide-saturated solutions are shown in Table 2.

Table 2. Yields in the radiolysis of aqueous toluene solutions
(Σ = G (Cr) + G (BA) + G (BAld) + G (BT) + 2G (B₂))




	10 ⁻³ M HC10 ₄ + 10 ⁻³ M Fe(III) ions saturated with		10 ⁻³ M HC10 ₄ saturated with		Unbuffered saturated with	
	Ar	N ₂ O	Ar	N ₂ O	Ar	N ₂ O
H	3.15	1.07	3.15	1.07	0.55	0.55
OH	2.65	4.73	2.65	4.73	2.65	5.25
Σ	2.97	5.40	1.33	3.75	0.59	1.43
Σ/OH	1.12	1.14	0.50	0.79	0.22	0.27

The ratio Σ/OH is the same in argon- and nitrous oxide-saturated solutions containing iron(III) ions, indicating that H atoms do not contribute to the Σ yield. An effect of iron(III) ions is, hence, to remove the CH₃C₆H₆ radicals, a part of which in the absence of iron(III) ions could react with the CH₃C₆H₅OH radicals according to reaction 9.

3.3 Aerated solutions

The effect of oxygen on the radiolysis of aqueous toluene solutions has been studied by irradiation of aerated solutions [$c(O_2) \simeq \simeq 0.25 \text{ mM}$] and the results are shown in Figs. 7 and 8 and in Table 1. A high yield of acid may be noted both in acid and in unbuffered solutions. The increase in the cresol yield, caused by oxygen, is not so large as the corresponding increase in the phenol yield during the radiolysis of aqueous benzene solutions (16). On the other hand the yield of hydrogen peroxide is increased to the same value as in the benzene system and the yield of dimers, including reduced dimers, is zero until the oxygen is consumed, in agreement with the absence of biphenyl in the benzene system as long as oxygen is present. Consequently, the reaction of oxygen with the hydroxymethylcyclohexadienyl radical (reaction 22) competes favourably with the elimination of water from the same radical (reaction 11).

The following reactions, based on analogy with the benzene system, together with reactions 1, 2 and 5 can explain the mechanism



Thus the total yield of hydroxytoluenes, benzoic acid and benzaldehyde is 2.1 and 2.6 in unbuffered and acid solutions, respectively, and almost the same as the phenol yield in the benzene system (2.1 and 2.3, respectively).

The formation of benzoic acid and benzaldehyde may occur through intermediates like peroxides and hydroperoxides which may be formed in the toluene system as in the benzene system (16).

CONCLUSIONS

The reaction mechanism in the aqueous toluene system is analogous with the mechanism in the aqueous benzene system (16), except: In acid deaerated solutions the methylhydroxycyclohexadienyl radical splits off a water molecule to form the benzyl radical. A similar reaction has not been observed in the benzene system. In aerated solutions acid is formed in the toluene but not in the benzene system and consequently oxygen does not increase the cresol yield as much as the phenol yield in the benzene system.

Addition of iron(III) ions changes the distribution of hydroxytoluenes favouring the formation of benzyl alcohol.

In acid solutions containing Fe(III) ions or oxygen the agreement between the toluene decomposed $G(-To)$ and the total yield of organic products $G(Org)$ was good. In unbuffered argon-saturated solutions $G(-To)$ exceeded $2 \cdot G(Org)$ indicating the formation of non-detected organic products.

ACKNOWLEDGEMENT

The authors are greatly indebted to Elfin Larsen at the Chemistry Dept. of the Danish AEC Research Station at Risö for assistance with mass spectrometric analysis. Financial support from the Swedish Atomic Research Council and The Swedish Board for Technical Development has been obtained and is gratefully acknowledged.

REFERENCES

1. KHENOKH, M. A. and LAPINSKAIA, E. M.
Action of γ -radiation of radioactive cobalt (Co^{60}) on aqueous solutions of aromatic hydrocarbons.
J. gen. Chem. USSR. 28 (1958) p. 679.
2. MANCITTI, M. A. B., BERTOLACCINI, P. L. and PUCINI, L.
Chemical effects induced by Co^{60} γ -irradiation in aqueous solutions of toluene and phenol.
Energia Nucl. 8 (1961) p. 519.
3. SUZUKI, N. and HOTTA, H.
The autoxidation of toluene catalyzed by metal ions in aqueous solutions. I
Bull. Chem. Soc. Japan. 40 (1967) p. 1361.
4. HOTTA, H. and SUZUKI, N.
The termination processes of the autoxidations of benzene toluene and ethylbenzene in the aqueous phase as studied by oxygen-18.
Bull. Chem. Soc. Japan. 41 (1968) p. 1537.
5. HOTTA, H., SUZUKI, N. and ISODA, K.
The effect of organic additives on the autoxidation of benzene and toluene in the aqueous phase and the kinetic analysis.
Bull. Chem. Soc. Japan. 42 (1969) p. 3056.
6. JEFCOATE, C. R. E., SMITH, J. R. L. and NORMAN, R. O. C.
Hydroxylation. Part IV. Oxidation of some benzenoid compounds by Fenton's reagent and the ultraviolet irradiation of hydrogen peroxide.
J. Chem. Soc. (B). 1969 p. 1013.
7. DORFMAN, L. M., TAUB, I. A. and BÜHLER, R. E.
Pulse radiolysis studies. I Transient spectra and reaction-rate constants in irradiated aqueous solutions of benzene.
J. Chem. Phys. 36 (1962) p. 3051.
8. SAUER, M. C. Jr. and WARD, B.
The reactions of hydrogen atoms with benzene and toluene studied by pulsed radiolysis: Reaction-rate constants and transient spectra in the gas phase and aqueous solution.
J. Phys. Chem. 71 (1967) p. 3971.
9. ERREDE, L. A. and CASSIDY, J. P.
The chemistry of xylylenes. V The formation of anthracenes via fast flow pyrolysis of toluenes and related compounds.
J. Am. Chem. Soc. 82 (1960) p. 3653.
10. CHRISTENSEN, H. C. and GUSTAFSON, R.
Radiolysis of aqueous benzene solutions. Effect of nitrous oxide and iron ions.
Nukleonik 12 (1969) p. 49.

11. ALLEN, A.O. et al.
Decomposition of water and aqueous solutions under pile radiation.
1949 (ORNL-130) p. 9.
12. HUMMEL, R.W. and SPINKS, J.W.T.
Betatron irradiation of aqueous ferrous sulphate solutions.
Can. J. Chem. 31 (1953) p. 250.
13. HOIGNE, J. and GÄUMANN, T.
Strahlungschemie der Kohlenwasserstoffe. Toluol.
Helv. Chim. Acta 44 (1961) p. 2141.
14. STUDIER, M.H. and HART, E.J.
The reduction of benzene by hydrated electrons in γ -ray irradiated alkaline solutions.
J. Am. Chem. Soc. 91 (1969) p. 4068.
15. ANBAR, M. and NETA, P.
A compilation of specific biomolecular rate constants for the reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals with inorganic and organic compounds in aqueous solution.
1966. (IA-1079).
16. CHRISTENSEN, H.C.
Radiolysis of aqueous benzene solutions.
Nukleonik 7 (1965) p. 1.




Fig. 1 Yields in unbuffered argon-saturated solutions. +, Cresols; x, benzyl alcohol; O, 1,2-diphenylethane; ∇, phenyltolylmethanes; Δ, dimethylbiphenyls; □, reduced dimers; e, H₂.




Fig. 2 Yields in unbuffered N₂O-saturated solutions. +, Cresols; x, benzyl alcohol; O, 1,2-diphenylethane; ∇, phenyltolylmethanes; Δ, dimethylbiphenyls; □, reduced dimers; e, H₂; •, H₂O₂.




Fig. 3 Yields in acid argon-saturated solutions (pH 3).

+, Cresols; x, benzyl alcohol; ○, 1,2-diphenylethane;
 ▽, phenyltolylmethanes; △, dimethylbiphenyls; □, reduced
 dimers; ⊙, H₂; ●, H₂O₂.




Fig. 4 Yields in acid N₂O-saturated solutions (pH 3).

+, Cresols; x, benzyl alcohol; ○, 1,2-diphenylethane;
 ▽, phenyltolylmethanes; △, dimethylbiphenyls; □, reduced
 dimers; ⊙, H₂; ●, H₂O₂.




Fig. 5 Yields in acid argon-saturated solutions in the presence of 10^{-3} M Fe(III) ions (pH 3). +, Cresols; x, benzyl alcohol; ○, 1,2-diphenylethane; ⊙, H_2 ; ▼, benzaldehyde; ▲, Fe(II) ions.




Fig. 6 Yields in acid N_2O -saturated solutions in the presence of 10^{-3} M Fe(III) ions (pH 3). +, Cresols; x, benzyl alcohol; ○, 1,2-diphenylethane; ⊙, H_2 ; ∇, phenyltolylmethanes; ▼, benzaldehyde; ▲, Fe(II) ions.




Fig. 7 Yields in unbuffered aerated solutions. +, Cresols; x, benzyl alcohol; O, 1,2-diphenylethane; ▽, phenyltolylmethanes; Δ, dimethylbiphenyls; □, reduced dimers; ⊙, H₂; ●, H₂O₂; ▾, benzaldehyde; ■, acids.




Fig. 8 Yields in acid aerated solutions (pH 3). +, Cresols; x, benzyl alcohol; O, 1,2-diphenylethane; ▽, phenyltolylmethanes; Δ, dimethylbiphenyls; □, reduced dimers; ⊙, H₂; ●, H₂O₂; ▾, benzaldehyde; ■, acid.

LIST OF PUBLISHED AE-REPORTS

1-340 (See back cover earlier reports.)

341. Nonlinear dynamic model of power plants with single-phase coolant reactors. By H. Vollmer. 1968. 26 p. Sw. cr. 10:--.

342. Report on the personnel dosimetry at AB Atomenergi during 1967. By J. Carlsson and T. Wahlberg. 1968. 10 p. Sw. cr. 10:--.

343. Friction factors in rough rod bundles estimated from experiments in partially rough annuli - effects of dissimilarities in the shear stress and turbulence distributions. By B. Kjellström. 1968. 22 p. Sw. cr. 10:--.

344. A study of the resonance interaction effect between ²³⁵U and ²³⁹Pu in the lower energy region. By H. Häggblom. 1968. 48 p. Sw. cr. 10:--.

345. Application of the microwave discharge modification of the Wilzbach technique for the tritium labelling of some organics of biological interest. By T. Gosztonyi. 1968. 12 p. Sw. cr. 10:--.

346. A comparison between effective cross section calculations using the intermediate resonance approximation and more exact methods. By H. Häggblom. 1969. 64 p. Sw. cr. 10:--.

347. A parameter study of large fast reactor nuclear explosion accidents. By J. R. Wiesel. 1969. 34 p. Sw. cr. 10:--.

348. Computer program for inelastic neutron scattering by an anharmonic crystal. By L. Bohlin, I. Ebbsjö and T. Höjberg. 1969. 52 p. Sw. cr. 10:--.

349. On low energy levels in ²³⁵W. By S. G. Malmkog, M. Höjberg and V. Berg. 1969. 18 p. Sw. cr. 10:--.

350. Formation of negative metal ions in a field-free plasma. By E. Larsson. 1969. 32 p. Sw. cr. 10:--.

351. A determination of the 2 200 m/s absorption cross section and resonance integral of arsenic by pile oscillator technique. By E. K. Sokolowski and R. Bladh. 1969. 14 p. Sw. cr. 10:--.

352. The decay of ¹⁹¹Os. By S. G. Malmkog and A. Bäcklin. 1969. 24 p. Sw. cr. 10:--.

353. Diffusion from a ground level point source experiment with thermoluminescence dosimeters and Kr 85 as tracer substance. By Ch. Gyllander, S. Hollman and U. Widemo. 1969. 23 p. Sw. cr. 10:--.

354. Progress report, FFN, October 1, - September 30, 1968. By T. Wiedling. 1969. 35 p. Sw. cr. 10:--.

355. Thermodynamic analysis of a supercritical mercury power cycle. By A. S. Roberts, Jr. 1969. 25 p. Sw. cr. 10:--.

356. On the theory of compensation in lithium drifted semiconductor detectors. By A. Lauber. 1969. 45 p. Sw. cr. 10:--.

357. Half-life measurements of levels in ²¹⁰Pb. By M. Höjberg and S. G. Malmkog. 1969. 14 p. Sw. cr. 10:--.

358. A non-linear digital computer model requiring short computation time for studies concerning the hydrodynamics of the BWR. By F. Reisch and G. Vayssier. 1969. 38 p. Sw. cr. 10:--.

359. Vanadium beta emission detectors for reactor in-core neutron monitoring. By I. Ö. Andersson and B. Söderlund. 1969. 26 p. Sw. cr. 10:--.

360. Progress report 1968. Nuclear chemistry. 1969. 38 p. Sw. cr. 10:--.

361. A half-life measurement of the 343.4 keV level in ¹⁷⁵Lu. By M. Höjberg and S. G. Malmkog. 1969. 10 p. Sw. cr. 10:--.

362. The application of thermoluminescence dosimeters to studies of released activity distributions. By B.-I. Rudén. 1969. 36 p. Sw. cr. 10:--.

363. Transition rates in ¹⁶¹Dy. By V. Berg and S. G. Malmkog. 1969. 32 p. Sw. cr. 10:--.

364. Control rod reactivity measurements in the Ågesta reactor with the pulsed neutron method. By K. Björus. 1969. 44 p. Sw. cr. 10:--.

365. On phonons in simple metals II. Calculated dispersion curves in aluminium. By R. Johnson and A. Westin. 1969. 124 p. Sw. cr. 10:--.

366. Neutron elastic scattering cross sections. Experimental data and optical model cross section calculations. A compilation of neutron data from the Studsvik neutron physics laboratory. By B. Holmqvist and T. Wiedling. 1969. 212 p. Sw. cr. 10:--.

367. Gamma radiation from fission fragments. Experimental apparatus - mass spectrum resolution. By J. Higbie. 1969. 50 p. Sw. cr. 10:--.

368. Scandinavian radiation chemistry meeting, Studsvik and Stockholm, September 17-19, 1969. By H. Christensen. 1969. 34 p. Sw. cr. 10:--.

369. Report on the personnel dosimetry at AB Atomenergi during 1968. By J. Carlsson and T. Wahlberg. 1969. 10 p. Sw. cr. 10:--.

370. Absolute transition rates in ¹⁹¹Ir. By S. G. Malmkog and V. Berg. 1969. 16 p. Sw. cr. 10:--.

371. Transition probabilities in the 1/2+(631) Band in ²³⁵U. By M. Höjberg and S. G. Malmkog. 1969. 18 p. Sw. cr. 10:--.

372. E2 and M1 transition probabilities in odd mass Hg nuclei. By V. Berg, A. Bäcklin, B. Fogelberg and S. G. Malmkog. 1969. 19 p. Sw. cr. 10:--.

373. An experimental study of the accuracy of compensation in lithium drifted germanium detectors. By A. Lauber and B. Malmsten. 1969. 25 p. Sw. cr. 10:--.

374. Gamma radiation from fission fragments. By J. Higbie. 1969. 22 p. Sw. cr. 10:--.

375. Fast neutron elastic and inelastic scattering of vanadium. By B. Holmqvist, S. G. Johansson, G. Lodin and T. Wiedling. 1969. 48 p. Sw. cr. 10:--.

376. Experimental and theoretical dynamic study of the Ågesta nuclear power station. By P. Å. Bliselius, H. Vollmer and F. Åkerhielm. 1969. 39 p. Sw. cr. 10:--.

377. Studies of Redox equilibria at elevated temperatures 1. The estimation of equilibrium constants and standard potentials for aqueous systems up to 374°C. By D. Lewis. 1969. 47 p. Sw. cr. 10:--.

378. The whole body monitor HUGO II at Studsvik. Design and operation. By L. Devell, I. Nilsson and L. Venner. 1970. 26 p. Sw. cr. 10:--.

279. ATMOSPHERIC DIFFUSION. Investigations at Studsvik and Ågesta 1960-1963. By L.-E. Häggblom, Ch. Gyllander and U. Widemo. 1969. 91 p. Sw. cr. 10:--.

380. An expansion method to unfold proton recoil spectra. By J. Kockum. 1970. 20 p. Sw. cr. 10:--.

381. The 93.54 keV level ⁸⁷Sr, and evidence for 3-neutron states above N=50. By S. G. Malmkog and J. McDonald. 1970. 24 p. Sw. cr. 10:--.

382. The low energy level structure of ²³⁹U. By S. G. Malmkog, V. Berg, A. Bäcklin and G. Hedin. 1970. 24 p. Sw. cr. 10:--.

383. The drinking rate of fish in the Skagerack and the Baltic. By J. E. Larsson. 1970. 16 p. Sw. cr. 10:--.

384. Lattice dynamics of NaCl, KCl, RbCl and RbF. By G. Raunio and S. Rolandson. 1970. 26 p. Sw. cr. 10:--.

385. A neutron elastic scattering study of chromium, iron and nickel in the energy region 1.77 to 2.76 MeV. By B. Holmqvist, S. G. Johansson, G. Lodin, M. Salama and T. Wiedling. 1970. 26 p. Sw. cr. 10:--.

386. The decay of bound isobaric analogue states in ²⁸Si and ²⁸Si using (d, n-) reactions. By L. Nilsson, A. Nilsson and I. Bergqvist. 1970. 34 p. Sw. cr. 10:--.

387. Transition probabilities in ¹⁹¹Os. By S. G. Malmkog, V. Berg and A. Bäcklin. 1970. 40 p. Sw. cr. 10:--.

388. Cross sections for high-energy gamma transition from MeV neutron capture in ²⁰⁸Pb. By I. Bergqvist, B. Lundberg and L. Nilsson. 1970. 16 p. Sw. cr. 10:--.

389. High-speed, automatic radiochemical separations for activation analysis in the biological and medical research laboratory. By K. Samsahl. 1970. 18 p. Sw. cr. 10:--.

390. Use of fission product Ru-106 gamma activity as a method for estimating the relative number of fission events in U-235 and Pu-239 in low-enriched fuel elements. By R. S. Forsyth and W. H. Blackadder. 1970. 25 p. Sw. cr. 10:--.

391. Half-life measurements in ¹³⁴I. By V. Berg and A. Höglund. 1970. 16 p. Sw. cr. 10:--.

392. Measurement of the neutron spectra in FRO cores 5, 9 and PuB-5 using resonance sandwich detectors. By T. L. Andersson and M. N. Qazi. 1970. 30 p. Sw. cr. 10:--.

393. A gamma scanner using a Ge(Li) semi-conductor detector with the possibility of operation in anti-coincidence mode. By R. S. Forsyth and W. H. Blackadder. 1970. 22 p. Sw. cr. 10:--.

394. A study of the 190 keV transition in ¹⁴¹La. By V. Berg, A. Höglund and B. Fogelberg. 1970. 22 p. Sw. cr. 10:--.

395. Magnetoacoustic waves and instabilities in a Hall-effect-dominated plasma. By S. Palmgren. 1970. 20 p. Sw. cr. 10:--.

396. A new boron analysis method. By J. Weitman, N. Däverhög and S. Farvolden. 1970. 26 p. Sw. cr. 10:--.

397. Progress report 1969. Nuclear chemistry. 1970. 39 p. Sw. cr. 10:--.

398. Prompt gamma radiation from fragments in the thermal fission of ²³⁵U. By H. Albinsson and L. Lindow. 1970. 48 p. Sw. cr. 10:--.

399. Analysis of pulsed source experiments performed in copper-reflected fast assemblies. By J. Kockum. 1970. 32 p. Sw. cr. 10:--.

400. Table of half-lives for excited nuclear levels. By S. G. Malmkog. 1970. 33 p. Sw. cr. 10:--.

401. Needle type solid state detectors for in vivo measurement of tracer activity. By A. Lauber, M. Wolgast. 1970. 43 p. Sw. cr. 10:--.

402. Application of pseudo-random signals to the Ågesta nuclear power station. By P.-Å. Bliselius. 1970. 30 p. Sw. cr. 10:--.

403. Studies of redox equilibria at elevated temperatures 2. An automatic divided-function autoclave and cell with flowing liquid junction for electrochemical measurements on aqueous systems. By K. Johnson, D. Lewis and M. de Pourbaix. 1970. 38 p. Sw. cr. 10:--.

404. Reduction of noise in closed loop servo systems. By K. Nygaard. 1970. 23 p. Sw. cr. 10:--.

405. Spectral parameters in water-moderated lattices. A survey of experimental data with the aid of two-group formulae. By E. K. Sokolowski. 1970. 22 p. Sw. cr. 10:--.

406. The decay of optically thick helium plasmas, taking into account ionizing collisions between metastable atoms or molecules. By J. Stevetleit. 1970. 18 p. Sw. cr. 10:--.

407. Zooplankton from Lake Magelungen, Central Sweden 1960-63. By E. Almquist. 1970. 62 p. Sw. cr. 10:--.

408. A method for calculating the washout of elemental iodine by water sprays. By E. Bachofner and R. Hesböl. 1970. 24 p. Sw. cr. 10:--.

409. X-ray powder diffraction with Guinier-Hägg focusing cameras. By A. Brown. 1970. 102 p. Sw. cr. 10:--.

410. General physic section Progress report. Fiscal year 1969/70. By J. Braun. 1970. 92 p. Sw. cr. 10:--.

411. In-pile determination of the thermal conductivity of UO₂ in the range 500-2500 degrees centigrade. By J.-Å. Gyllander. 1971. 70 p. Sw. cr. 10:--.

412. A study of the ring test for determination of transverse ductility of fuel element canning. By G. Anevi and G. Östberg. 1971. 17 p. Sw. cr. 15:--.

413. Pulse radiolysis of Aqueous Solutions of aniline and substituted anilines. By H. C. Christensen. 1971. 40 p. Sw. cr. 15:--.

414. Radiolysis of aqueous toluene solutions. By H. C. Christensen and R. Gustafson. 1971. 20 p. Sw. cr. 15:--.

List of published AES-reports (In Swedish)

1. Analysis by means of gamma spectrometry. By D. Brune. 1961. 10 p. Sw. cr. 6:--.

2. Irradiation changes and neutron atmosphere in reactor pressure vessels - some points of view. By M. Grounes. 1962. 33 p. Sw. cr. 6:--.

3. Study of the elongation limit in mild steel. By G. Östberg and R. Attermo. 1963. 17 p. Sw. cr. 6:--.

4. Technical purchasing in the reactor field. By Erik Jonson. 1963. 64 p. Sw. cr. 8:--.

5. Ågesta nuclear power station. Summary of technical data, descriptions, etc for the reactor. By B. Lilliehöök. 1964. 336 p. Sw. cr. 15:--.

6. Atom Day 1965. Summary of lectures and discussions. By S. Sandström. 1966. 321 p. Sw. cr. 15:--.

7. Building materials containing radium considered from the radiation protection point of view. By Stig O. W. Bergström and Tor Wahlberg. 1967. 26 p. Sw. cr. 10:--.

Additional copies available from the Library of AB Atomenergi, Fack, S-611 01 Nyköping 1, Sweden.