

Growth Performances of BRR1 dhan55 under Integrated Nutrient Management

M. Das¹, M Hussain¹, P. Chakrabarty¹, A. Rahman¹, M. S. A. Talucder^{1,2*}, M. O. Sharif¹ and A.U. Khan¹

¹Climate-Smart Agriculture Lab, Department of Agroforestry and Environmental Science, Sylhet Agricultural University, Sylhet, Bangladesh

²Interdisciplinary Research for Future Agriculture, Sylhet Agricultural University, Sylhet-3100, Bangladesh.

*Corresponding author: talucdermsa.aes@sau.ac.bd, <https://orcid.org/0000-0002-6863-9586>, Tel: +8801711974124

Received: 19/12/2022

Accepted: 31/12/2022

Available online: 09/01/2023



Copyright: ©2022 by the author(s).

This work is licensed under a Creative Commons Attribution 4.0 License.

<https://creativecommons.org/licenses/by/4.0/>

Abstract: Integrated nutrient management is vital for the development and growth of rice (*Oryza sativa* L.), but they are rarely used nowadays due to the widespread use of inorganic fertilizers. So, the current research was executed to determine the effect of organic amendments on different growth parameters of BRR1 dhan55. A field investigation was managed at the Plant Pathology field lab, Sylhet Agricultural University, from November 2018 to April 2019. Different combinations of organic amendments (T₁= Recommended Fertilizer Dose, RFD which includes TSP 100g 10 m⁻², MoP 100g 10 m⁻², Gypsum 50g 10 m⁻², Zinc 15g 10 m⁻², T₂= RFD with vermicompost 2kg 10 m⁻², T₃= RFD with vermicompost 4 kg 10 m⁻², T₄= RFD with cowdung 2 kg 10 m⁻², T₅= RFD with cowdung 4 kg 10 m⁻², and T₆= RFD with Biochar 4 kg 10 m⁻²) were used in a RCBD (Randomized Complete Block Design) with three replications. The existing results showed that RFD with biochar treatments showed better performance in terms of the initial weight of grain (50.70 g), clean grain weight (48.55 g), and number of non-effective grains per panicle (27.67) and tiller per hill (12.07). T₂ (RFD with vermicompost 2 kg 10 m⁻²) treatments demonstrated better results in parameters such as weight of grain from a single panicle (4.04 g), single panicle grain weight (3.90 g), panicle length (24.77 cm), and grain per panicle (153.00). Biomass (34.56 g) and above-ground length parameters (90.00 cm) performed better with the T₄ treatment (RFD with 2 kg 10 m⁻² of cowdung). T₆ treatments were showed the highest grain yield (10.71 t ha⁻¹) followed by T₃ treatments. Therefore, the present results recommended that organic amendments could be a better supplement than inorganic amendments for the growth performance of BRR1 dhan55.

Keywords: Rice; *Oryza sativa*; Vermicompost; Biochar; Cowdung.

INTRODUCTION

In Asia, rice (*Oryza sativa* L.) is the primary staple meal (Rao et al., 2018). The yield of rice must increase above existing levels in order to feed the expanding population (Pan et al., 2013). Because of the recent rapid economic expansion, there are fewer workers available for agricultural output (Pan et al., 2017). With around 370 kcal of energy per 100 g of grains, rice is the main food grain crop of Bangladesh. With a total yearly production of 33.89 million metric tons, rice is grown on approximately 84% of the country's cropped land (BBS, 2015). Having an eight-month dry season and a four-month wet season, Bangladesh

features a monsoon climate in addition to a hot, humid subtropical climate (Ahmad et al., 2014). There are three main production seasons of rice in Bangladesh: Aus, which is the spring season, and Boro, which is the dry season and is fully irrigated. Aman is the wet season. Irrigated dry season Boro rice, the most fruitful of the three rice seasons, is the main crop (Ahmed et al., 2013). When Boro rice is cultivated with high-yielding conventional varieties, the optimum yield is 3.86 tons ha⁻¹, and when it is grown with hybrid kinds, the average yield is 4.75 tons ha⁻¹. Bangladesh produces a relatively small amount of rice on average, 3.0 t ha⁻¹ which is extremely little (BBS, 2015).

The most significant and widely grown food crop is rice which provides one in three individuals with half of their daily nutrition. The high yielding varieties, intensive farming, and inorganic fertilizers take the lead in providing food for these large populations. Our soil is degrading to an alarming level as a result of the excessive and improper use of complex organic substances in stabilized, humus-like chemical fertilizers in the soil, which is also contributing to ecosystem imbalance and environmental pollution (Satyanarayana et al., 2002; Mahajan et al., 2008; Hasanuzzaman et al., 2010). As a result of the stimulation of a downturn in soil properties (physical and chemical), as well as in soil microbial growth (Mahajan et al., 2008). Organic materials are better sources of plant nutrition because they don't harm crops or the soil (Pieters, 2004). It is known that using earthworms in vermicomposting produces products that resemble hummus and stabilizes complicated organic materials (Benitez et al., 2000). Vermicompost application to soil is recognized as a desirable strategy in any production system due to the enhancement of microbial activities of soil, plant nutrients mineralization, and rose soil fertility (Arancon et al., 2006, Ferraras et al., 2006). Vermicompost is a material created by earthworms using organic waste (Blouin et al., 2019). Vermicompost has good physical, chemical, and biological qualities that can increase the fertility of the soil and reduce crop diseases, according to a prior study (Patnaik et al., 2020). According to Fernandez et al. (2011), vermicompost has a high microbial functional diversity and the potential to be used for the treatment of chemical pollution in agricultural production. Jahanbakhshi et al. (2019) demonstrated the value of vermicompost as an organic fertilizer with a suitable CN (carbon to nitrogen) ratio, pH, and salinity. Jaouni et al. (2019) also suggested that vermicompost could enhance the jujube coconut fruit's chemical makeup and boost its nutritional and therapeutic potential. The productiveness of crop fields has been steadily deteriorating due to constant cropping, nutrient mining, and indiscriminate chemical fertilizer use in Bangladesh. In this light, biochar-primarily carbon-enriched materials with a negligible quantity of plant nutrients might be the finest organic manures for restoring depleted soils. Biochar is produced from organic waste under circumstances of high pyrolysis temperature and oxygen limitation. A product of anaerobic pyrolysis made from organic materials-biochar can be used to restore soil carbon for a prolonged period of time by lowering greenhouse gases emission from the soil to the troposphere. It is resistant to easy degradation. Additionally, using biochar will lessen the need for chemical fertilizers when growing rice, and changing the paddy ecosystem may reduce GHG emissions. In the highly sensitive dry tropical agroecosystem of India, the of FYM and rice husk biochar in combination with less inorganic fertilizer and fewer water inputs was found to help sustain wheat crop output (Singh et al., 2019). Additionally, Singh et al. (2021) found such appropriate farming techniques depending on certain agroecosystems may be successful in helping humans adapt to climate change.

A carbon-rich organic substance called biochar is produced by pyrolyzing biomass (Borchard et al., 2012). Another definition for it is "the carbonaceous residue from pyrolysis, especially naturally occurring burns under low oxygen conditions" (Xu et al., 2012). Biochar stores nutrients in the soil and absorbs moisture, which lowers the need for mineral fertilizer and shields crops from the effects of drought. Utilizing biochar in farming soils, such as paddy fields, reduces CH₄ emissions while boosting soil microbial activity. Additionally, because biochar-amended soils have higher germination rates, fewer seeds are needed. Notably, biochar is advantageous for all kinds of agricultural systems (Brunn et al., 2011). There was a lot of work done in rice fields to find out the effects of various organic and inorganic fertilizer combinations, but very few experiments were found in rice cultivation with only organic fertilizer. The goal of the current study was to determine how different organic fertilizer doses (including vermicompost, cowdung, and biochar in addition to the prescribed fertilizer dose) affected several yield-contributing factors in a rice field.

MATERIALS AND METHODS

Study area: The investigation was carried out between November 2018 and April 2019 at the Sylhet Agricultural University's Plant Pathology Field Lab in Sylhet, Bangladesh, which is located at 24°53'56.54"N latitude and 91°52'19.13"E longitude (Figure 1).

At higher elevations, Sylhet's characteristic tropical monsoon climate transitions into a humid subtropical climate. The rainy season starts in April and is extremely hot and humid with frequent, severe rains and thunderstorms, in contrast to the short dry season that starts in November and lasts until February. Between May and September, precipitation totals an average of 4,200 mm, or about 80% of the entire year (Talucder 2015).

Materials: A test crop of the rice cultivar BRRI dhan55 was utilized. The Bangladesh Rice Research Institute (BRRI) created this variety in 2011. The boro season is the best time to use it. The wet bed method was used to raise the rice seedlings. In November 2018, a hand spade was used to dig up the chosen seed bed. 1 kg Seeds bigha⁻¹ (95% germinated) were soaked and nurtured for 48 hours; 95% of them germinated. On January 10, 2019, seedlings that were thirty days old were painstakingly removed from the seedling nursery and transplanted. From the surrounding Sylhet area sawmills, sawdust was gathered. Biochar was prepared from the collected sawdust. Sawdust was measured out and let to dry for a few days in the sun to create biochar. After drying, 40 kg of dry material was recovered from about 50 kg of sawdust. These dry materials were pyrolyzed in a kiln for an extended period of time to create biochar. Twenty kg of biochar were produced after pyrolysis (Mustakim et al., 2022). Vermicompost (from *E. fetida*-cowdung) was collected from the soil science laboratory of Sylhet Agricultural University. Well decomposed cowdung was collected from Jonaki village near Sylhet Agricultural University.

Land preparation: Maintaining line to line space of 20 cm and hill to hill space of 15 cm, two seedlings per hill were used. After seven days of transplantation, all blocks were examined for misplaced hills, which were wherever necessary occupied in with more plants. A power tiller was used to plow the ground initially. The soil was eventually

prepared by serial plowing, cross plowing, and laddering after being sufficiently soaked with irrigation water. After 40 days, the unwanted residues were sorted out from the test plot. The terrain was eventually leveled. There were 18 plots (5 m x 2 m) on the experimental land. The spacing between blocks was 1 m, and 0.5 m separated the two plots.




Figure 1. Study area map.

Experimental design and data analysis: The experiment was done with RCBD (Randomized Complete Block Design) with three replications. There were six treatments in this research experiment namely, T_1 = Recommended Fertilizer Dose, RFD which includes TSP 100g 10 m⁻², MoP 100g 10 m⁻², Gypsum 50g 10 m⁻², Zinc 15g 10 m⁻², T_2 = RFD with vermicompost 2kg 10 m⁻², T_3 = RFD with vermicompost 4 kg 10 m⁻², T_4 = RFD with cowdung 2 kg 10 m⁻², T_5 = RFD with cowdung 4 kg 10 m⁻², and T_6 = RFD with Biochar 4 kg 10 m⁻². 100g Urea 10 m⁻² was applied three times at 10, 30, and 50 days after transplanting. The field data were statistically analyzed by using the statistical R-software program.

RESULTS AND DISCUSSION

This study's results showed that the initial weight of grain (50.70 g plant⁻¹) was observed to be highest in treatments RFD with biochar 4 kg 10 m⁻² (Table 1). RFD with vermicompost 2kg 10 m⁻² produced the highest weight of grain from a single panicle (4.04 g plant⁻¹). Clean grain weight (48.55 g plant⁻¹) showed better results in treatment RFD with biochar 4 kg 10 m⁻² (Table 1). RFD with vermicompost 2kg 10 m⁻² exhibited the highest result in single panicle grain weight (3.90 g plant⁻¹) (Table 1).

Sharma and Banik (2012) stated that organic manures from sources like FYM or vermicompost can replace 30% of the recommended doses of nitrogen without harming the system's productivity or soil fertility. According to Rahman and Barmon (2019), vermicompost application greatly boosted rice productivity and technical efficacy.

The no. of non-effective grains per panicle (27.67) was found to be highest in treatment RFD with biochar 4 kg 10 m⁻² (Table 1). RFD with cowdung 2 kg 10 m⁻² resulted in the greatest above-ground length (90.00 cm) (Table 1). According to Muhammad Aslam Ali et al., 2021 rice yield rose while employing the AWDI method with biochar amendments at 15-20 t ha⁻¹ with half the required inorganic fertilizers, especially when cultivating dry Boro rice.

Rice below ground length (18.73 cm) was the longest in RFD with vermicompost 4 kg 10 m⁻² (Table 2). RFD with vermicompost 2 kg 10 m⁻² showed the best performances in terms of panicle length (24.77 cm) (Table 2). At 2 kg 10 m⁻², grain per panicle (153.00) was discovered to be the most common in RFD with vermicompost (Table 2). Any fuel obtained from plants is referred to as biomass. This comprises plant and animal waste as well as crop residues and wood. RFD with cowdung 2 kg 10 m⁻² demonstrated the highest results in biomass (34.56 g) (Table 2).

Tillering is a crucial characteristic of grain production and, as a result, a crucial component in enhancing rice growth. Effective tillers and non-bearing tillers together make up the total number of tillers on hill⁻¹. The number of tillers per hill (12.07) exhibited the best results with RFD with Biochar 4 kg 10 m⁻² (Table 2). According to Shaoyi et al. (2021), the implementation of vermicompost accelerated

the length of the root, surface area, mean diameter, the volume of the root, and no. of root tip's fragrant rice seedlings, as well as increased root activity. The study's findings suggested that growing plants in nurseries with vermicompost would be an effective agronomic strategy for the cultivation of fragrant rice.

Table 1. Effect of various organic amendments along with RFD on yield contributing characters of Boro rice (BRRI [dhan55])

Treatments	Initial weight of grain (gm)	Weight of grain from single panicle (gm)	Clean grain weight (gm)	Single panicle grain weight (gm)	No. of non-effective grain per panicle	Above ground length (cm)
T ₁	34.68±18.93	3.23±1.17	33.16±18.26	3.08±1.13	23.00±11.14	84.90±0.96
T ₂	37.04±3.04	4.04±0.73	35.71±3.37	3.90±0.76	22.33±6.11	82.70±6.82
T ₃	42.04±16.52	2.88±0.49	39.67±15.74	2.68±0.49	22.00±4.58	86.10±2.01
T ₄	32.11±14.16	3.15±0.89	30.55±13.90	3.00±0.90	24.67±7.77	90.00±3.61
T ₅	31.97±8.84	3.94±0.62	28.19±5.64	3.80±0.66	24.00±7.21	86.67±8.74
T ₆	50.70±5.34	3.56±0.27	48.55±4.95	3.39±0.26	27.67±10.26	67.80±29.60
CV (%)	34.86	23.20	34.20	24.20	22.82	16.17

Note: T₁= Recommended Fertilizer Dose, RFD which includes TSP 100g 10 m⁻², MoP 100g 10 m⁻², Gypsum 50g 10 m⁻², Zinc 15g 10 m⁻², T₂= RFD with vermicompost 2kg 10 m⁻², T₃= RFD with vermicompost 4 kg 10 m⁻², T₄= RFD with cowdung 2 kg 10 m⁻², T₅= RFD with cowdung 4 kg 10 m⁻², T₆= RFD with Biochar 4 kg 10 m⁻²

Table 2. Effect of various organic amendments along with RFD on yield contributing characters of Boro rice (BRRI [dhan55])

Treatments	Below ground length (cm)	Panicle length (cm)	Grain per panicle	Biomass (gm)	No. of tillers per hill
T ₁	12.93±1.80	22.27±0.74	125.67±44.29	27.50±6.93	9.07±0.31
T ₂	13.60±4.33	24.77±1.72	153.00±18.25	28.73±5.65	8.57±0.72
T ₃	18.73±9.98	23.50±1.25	110.33±11.59	27.00±2.08	11.13±0.64
T ₄	14.50±3.91	22.33±2.66	124.67±28.59	34.56±15.36	10.13±1.79
T ₅	13.30±2.25	24.00±1.18	150.67±15.04	33.00±20.75	7.68±1.74
T ₆	13.13±4.90	23.90±1.73	141.33±12.42	28.14±6.79	12.07±3.18
CV (%)	39.68	7.16	19.38	35.43	17.64

Note: T₁= Recommended Fertilizer Dose, RFD which includes TSP 100g 10 m⁻², MoP 100g 10 m⁻², Gypsum 50g 10 m⁻², Zinc 15g 10 m⁻², T₂= RFD with vermicompost 2kg 10 m⁻², T₃= RFD with vermicompost 4 kg 10 m⁻², T₄= RFD with cowdung 2 kg 10 m⁻², T₅= RFD with cowdung 4 kg 10 m⁻², T₆= RFD with Biochar 4 kg 10 m⁻²




Figure 2. Effect of various organic amendments along with RFD on the grain yield of Boro rice (BRRI dhan55)

According to the results (Figure 2), the highest rice yield (10.71 t/ha) was obtained from T₆ treatments and the lowest rice yield (6.22 t/ha) was obtained from T₅ treatments. Recommended fertilizer dose (T₁) treatments gave 7.31 t/ha rice yield, T₂ treatments gave 7.88 t/ha rice yield, T₃ treatments gave 8.75 t/ha rice yield, and T₄ treatments gave 6.74 t/ha rice yield. According to the BRRI report, the yield of BRRI dhan55 is 7 t/ha. The study result was similar to different previous research. By increasing the number of panicles and grains, the use of biochar can increase rice yield (Zhang et al. 2014; Zhang et al. 2013). With the addition of more biochar, the number of rice panicles and grains rose, and the rice yield increased by 15.26-44.89% (Zhang et al. 2014). According to Morteza et al. (2011), plants treated with 2 t/ha of organic fertilizer had the highest grain yield in 2008 (4335.88 kg/ha), whereas plants treated with chemical fertilizer plus 1.5 tons/ha of organic fertilizer had the highest yield in 2009 (4662.71 kg/ha).

CONCLUSION

According to the findings, all treatments had a significant impact on the growth and development of BRRI dhan55 (Boro rice). The study also confirmed that different doses of vermicompost, cowdung, and biochar treatment along with RFD gave the best results compared to RFD alone. Biochar treatments showed the best results regarding the initial weight of grain, clean grain weight, no. of non-effective grains per panicle, and no. of tillers per hill. Treatments with different doses of vermicompost gave the best results in terms of weight of grain from a single panicle, below-ground length, panicle length, and grain per panicle. To obtain higher growth and yield of rice, organic amendments can be an excellent complement to inorganic fertilizers, as suggested by our results.

Acknowledgment

We thank the Plant Pathology and Seed Science Department, SAU, Sylhet-3100, for providing the field for this experiment. We also thank Md. Kamrujjaman, Md. Moudud Ahmod, and Shamim Reza for helping with biochar and site preparation.

Conflict of Interest

The authors have not declared any conflicts of interest.

REFERENCES

- Ahmad, M.D., Kirby, M., Islam, M.S., Hossain, M.J. and Islam, M.M. 2014. Groundwater Use for Irrigation and Its Productivity: Status and Opportunities for Crop Intensification for Food Security in Bangladesh. *Water Resource Management*, 28: 1415-1429.
- Ahmed, A.U., Ahmad, K., Chou, V., Hernandez, R., Menon, F.N., Naher, F. and Yu, B. 2013. The Status of Food Security in the Feed the Future Zone and Other Regions of Bangladesh: Results from the 2011-2012 Bangladesh Integrated Household Survey. International Food Policy Research Institute Report.
- Arancon, N.Q., Edwards, C.A. and Bierman, P. 2006. Influences of vermicomposts on field strawberries: Part 2. Effects on soil microbiological and chemical properties. *Bioresour. Technol.*, 97: 831-840.
- Bangladesh Bureau of Statistics (BBS). 2015. Statistical Year Book of Bangladesh 2015: Bangladesh Bureau of Statistics.
- Benitez, E., Nogales, R., Masciandaro, G. and Ceccanti, B. 2000. Isolation by isoelectric focusing of humic-urease complexes from earthworms (*Eisenia fetida*) processed sewage sludges. *Biol. Fertil. Soils.*, 31: 489-493.
- Blouin, M., Barrere, J., Meyer, N., Lartigue, S., Barot, S. and Mathieu, J. 2019. Vermicompost significantly affects plant growth. A meta-analysis. *Agronomy for Sustainable Development*, 39: 34. [CrossRef]
- Borchard, N., Prost, K., Kautz, T., Moeller, A. and Siemens, J. 2012. Sorption of copper (II) and sulphate to different biochars before and after composting with farmyard manure. *European Journal of Soil Science*, 63:399-409.
- Bruun, E.W., Hauggaard-Nielsen, H., Ibrahim, N., Egsgaard, H., Ambus, P., Jensen, P.A. and Dam-Johansen K. 2011. Influence of fast pyrolysis temperature on biochar labile fraction and short-term carbon loss in a loamy soil. *Biomass Bioenergy*, 35:1182-1189.
- Fernández-Gómez, M.J., Nogales, R., Insam, H., Romero, E. and Goberna, M. 2011. Role of vermicompost chemical composition, microbial functional diversity, and fungal community structure in their microbial respiratory response to three pesticides. *Bioresource Technology*, 102: 9638-9645. [CrossRef]
- Ferreras, L., Gomez, E., Toresani, S., Firpo, I. and Rotondo, R. 2006. Effect of organic amendments on some physical, chemical and biological properties in a horticultural soil. *Bioresource Technology*, 97: 635-640.
- Hasanuzzaman, M., Ahamed, K.U., Rahmatullah, N.M., Akhter N. and Rahman, M.L. 2010. Plant growth characters and productivity of wetland rice (*Oryza sativa* L.) as affected by application of different manures. *Emirates Journal of Food and Agriculture*, 22(1): 46-58.
- Jahanbakhshi, A. and Kheiralipour, K. 2019. Influence of vermicompost and sheep manure on mechanical properties of tomato fruit. *Food Sciences and Nutrition*, 7: 1172-1178. [CrossRef]
- Jaouni, A.S., Selim, S., Hassan, S.H., Mohamad, H.S.H., Wadaan, M.A.M., Hozzein, W.N., Asard, H. and AbdelGawad, H. 2019. Vermicompost Supply Modifies Chemical Composition and Improves Nutritive and Medicinal Properties of Date Palm Fruits from Saudi Arabia. *Front. Plant Sci.*, 10:424. [CrossRef]
- Lehmann, C.J. and Rondon, M. 2006. Bio-Char Soil Management on Highly-Weathered Soils in the Tropics. In N. T. Uphoff (Ed.), *Biological Approaches to Sustainable Soil Systems*. pp. 517-530. CRC Press.
- Mahajan, A., Bhagat R.M. and Gupta, R.D. 2008. Integrated nutrient management in sustainable rice-wheat cropping system for food security in India. *SAARC Journal of Agriculture*, 6: 29-32.
- Morteza, S., Alireza N., Shankar. L. L. 2011. Effect of Organic Fertilizer on Growth and Yield Components in Rice (*Oryza sativa* L.) *Journal of Agricultural Science*, 3(3). doi:10.5539/jas.v3n3p217
- Muhammad, A.A., Rajib, H., Zubair, A.I., Sanjit, C.B., Badiuzzaman, K., Rehana, K., Hafsa, J.H. and Touhidul, I. 2021. Development of Environment Friendly Paddy Ecosystem for Sustainable Rice Farming through Soil Amendments with Biochar and Alternate Wetting-Drying Irrigations. *American Journal of Climate Change*, 10: 581-596. <https://www.scirp.org/journal/ajcc>
- Mustakim, M.T., Talucder, M.S.A., Ruba, U.B., Islam, F., Rahman, A., Uddin, M.S. and Khan, A.U. 2022. Growth performance of black pepper (*P. nigrum*) cuttings in different

- rooting media and growth regulator. *Journal of Agroforestry and Environment*, 15(2): 63-68. doi: <https://doi.org/10.55706/jae1518> [CrossRef]
- Pan, S., Rasul, F., Li, W., Tian, H., Mo, Z., Duan, M. and Tang, X. 2013. Roles of plant growth regulators on yield, grain qualities and antioxidant enzyme activities in super hybrid rice (*Oryza sativa* L.). *Rice*, 6:9. [CrossRef]
- Pan, S., Wen, X., Wang, Z., Ashraf, U., Tian, H., Duan, M., Mo, Z., Fan, P. and Tang, X. 2017. Benefits of mechanized deep placement of nitrogen fertilizer in direct seeded rice in South China. *Field Crops Research*, 203: 139-149. [CrossRef]
- Patnaik, P., Abbasi, T. and Abbasi, S. 2020. Vermicompost of the widespread and toxic xerophyte prosopis (*Prosopis juliflora*) is a benign organic fertilizer. *Journal of Hazardous Materials*, 399:22864. [CrossRef]
- Pieters, A.J. 2004. *Green Manuring: Principles and Practice*. Agrobios, Jodhpur. pp: 356.
- Rahman, S. and Barmon, B. K. 2019. Greening Modern Rice Farming Using Vermicompost and Its Impact on Productivity and Efficiency: An Empirical Analysis from Bangladesh. *Agriculture*, 9:239; doi:10.3390/agriculture9110239
- Rao, G., Ashraf, U., Huang, S., Cheng, S., Abrar, M., Mo, Z., Pan, S. and Tang, X. 2018. Ultrasonic seed treatment improved physiological and yield traits of rice under lead toxicity. *Environmental Science and Pollution Research*, 25: 33637-33644. [CrossRef] [PubMed]
- Satyanarayana, V., Prasad, R.D., Murthy V.R.K. and Boote, K. J. 2002. Influence of integrated use of farmyard manure and inorganic fertilizers on yield and yield components of irrigated lowland rice. *Journal of Plant Nutrition*, 25(10):2081-2090.
- Shaoyi, R., Feida, W., Rifang, L., Xiangru, T., Haowen, L. and Longxin, H. 2021. Preliminary Application of Vermicompost in Rice Production: Effects of Nursery Raising with Vermicompost on Fragrant Rice Performances. *Agronomy*, 11:1253. <https://doi.org/10.3390/agronomy11061253>
- Sharma, R. C. and Banik, P. 2012. Effect of integrated nutrient management on baby corn–rice cropping system: economic yield, system productivity, nutrient-use efficiency and soil nutrient balance. *Indian Journal of Agricultural Sciences*, 82(3):220.
- Singh, R., Kumari, T., Verma, P., Singh, B.P. and Raghubanshi, A.K. 2021. Compatible Package-Based Agriculture Systems: An Urgent Need for Agro-Ecological Balance and Climate Change Adaptation. *Soil Ecology Letters*. <https://doi.org/10.1007/s42832-021-0087-1>
- Singh, R., Singh, P. and Singh, H. 2019. Impact of Sole and Combined Application of Biochar, Organic and Chemical Fertilizers on Wheat Crop Yield and Water Productivity in a Dry Tropical Agro-Ecosystem. *Biochar*, 1: 229-235. <https://doi.org/10.1007/s42773-019-00013-6>
- Talucder, M.S.A. 2015. Has climate change already affected the temporal trends of seasonal and annual precipitation in Sylhet district of Bangladesh? *Journal of Soil and Nature*. 8(3):21-27.
- Xu, G., L, Y., Sun, J., Shao, H. and Wei, L. 2012. Recent advances in biochar applications in agricultural soils: Benefits and environmental implications. *CLEAN Soil Air Water*, 40:1093-1098.
- Zhang, A.P., Liu, R.L., Gao, J., Zhang, Q.W., Xiao, J.N., Chen, Z., Yang, S.Q., Hui, J.Z. and Yang, Z.L. 2014. Effects of biochar on nitrogen losses and rice yield in anthropogenic-alluvial soil irrigated with yellow river water. *Journal of Agro-Environment Science*, 33, 2395-2403.
- Zhang, W.M., Meng, J., Wang, J.Y., Fan, S.X. and Chen, W.F. 2013. Effect of biochar on root morphological and physiological characteristics and yield in rice. *Acta Agronomica Sinica*. 39, 1445. [CrossRef]

