



***Corrigendum to***  
**“Geochemical signatures of pingo ice and its origin in Grøndalen,  
west Spitsbergen” published in The Cryosphere, 13, 3155–3169, 2019**

**Nikita Demidov<sup>1</sup>, Sebastian Wetterich<sup>2</sup>, Sergey Verkulich<sup>1</sup>, Aleksey Ekaykin<sup>1,3</sup>, Hanno Meyer<sup>2</sup>, Mikhail Anisimov<sup>1,3</sup>, Lutz Schirrmeister<sup>2</sup>, Vasily Demidov<sup>1</sup>, and Andrew J. Hodson<sup>4,5</sup>**

<sup>1</sup>Arctic and Antarctic Research Institute, Bering St. 38, 199397 St. Petersburg, Russia

<sup>2</sup>Alfred Wegener Institute Helmholtz Center for Polar and Marine Research, Telegrafenberg A45, 14473 Potsdam, Germany

<sup>3</sup>St. Petersburg State University, 10th Line 33–35, 199178 St. Petersburg, Russia


<sup>4</sup>University Centre in Svalbard, 9171 Longyearbyen, Norway

<sup>5</sup>Western Norway University of Applied Sciences, Røyrgata 6, 6856 Sogndal, Norway

**Correspondence:** Nikita Demidov (nikdemidov@mail.ru)

Published: 13 December 2019

Figure 4 is incorrectly displayed on page 3163. The correct version of Fig. 4 is given here.



**Figure 4.** Hydrochemical composition of water extracts from the sedimentary cores 9, 10 and 11. Data are given in Table 3. Note different axis scale for ion content in (a) reaching up to  $1335 \text{ mg L}^{-1}$ .