

# 1 **Presence of rapidly degrading permafrost plateaus in** 2 **southcentral Alaska**

3  
4 Benjamin M. Jones<sup>1</sup>, Carson A. Baughman<sup>1</sup>, Vladimir E. Romanovsky<sup>2,3</sup>, Andrew D. Parsekian<sup>4</sup>,  
5 Esther L. Babcock<sup>1</sup>, Eva Stephani<sup>1</sup>, Miriam C. Jones<sup>5</sup>, Guido Grosse<sup>6</sup>, and Edward E. Berg<sup>7</sup>  
6  
7

8 <sup>1</sup> Alaska Science Center, U.S. Geological Survey, Anchorage, AK 99508, USA

9 <sup>2</sup> Geophysical Institute, University of Alaska Fairbanks, Fairbanks, AK 99775, USA

10 <sup>3</sup> Earth Cryosphere Institute, 86 Malygina Street, Tyumen 625000, Russia

11 <sup>4</sup> Department of Geology and Geophysics, University of Wyoming, Laramie, WY 82070, USA

12 <sup>5</sup> Eastern Geology and Paleoclimate Science Center, U.S. Geological Survey, Reston, VA 20192, USA

13 <sup>6</sup> Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Potsdam, 14473, Germany

14 <sup>7</sup> Kenai National Wildlife Refuge, U.S. Fish and Wildlife Service, Soldotna, AK 99669, USA (retired)  
15

16 *Correspondence to:* Benjamin M. Jones (bjones@usgs.gov)

17 **Abstract.** Permafrost presence is determined by a complex interaction of climatic, topographic,  
18 and ecological conditions operating over long time scales. In particular, vegetation and organic  
19 layer characteristics may act to protect permafrost in regions with a mean annual air temperature  
20 (MAAT) above 0 °C. In this study, we document the presence of residual permafrost plateaus on  
21 the western Kenai Peninsula lowlands of southcentral Alaska, a region with a MAAT of 1.5±1 °C  
22 (1981–2010). Continuous ground temperature measurements between 16 September 2012 and 15  
23 September 2015, using calibrated thermistor strings, documented the presence of warm permafrost  
24 (-0.04 °C to -0.08 °C). Field measurements (probing) on several plateau features during the fall  
25 of 2015 showed that the depth to the permafrost table averaged 1.48 m but at some locations was  
26 as shallow as 0.53 m. Late winter surveys (augering, coring, and GPR) in 2016 showed that the  
27 average seasonally frozen ground thickness was 0.45 m, overlying a talik above the permafrost  
28 table. Measured permafrost thickness ranged from 0.33 m to >6.90 m. Manual interpretation of  
29 historic aerial photography acquired in 1950 indicates that residual permafrost plateaus covered

30 920 ha as mapped across portions of four wetland complexes encompassing 4,810 ha. However,  
31 between 1950 and ca. 2010, permafrost plateau extent decreased by 60.0 %, with lateral feature  
32 degradation accounting for 85.0 % of the reduction in area. Permafrost loss on the Kenai Peninsula  
33 is likely associated with a warming climate, wildfires that remove the protective forest and organic  
34 layer cover, groundwater flow at depth, and lateral heat transfer from wetland surface waters in  
35 the summer. Better understanding the resilience and vulnerability of ecosystem-protected  
36 permafrost is critical for mapping and predicting future permafrost extent and degradation across  
37 all permafrost regions that are currently warming. Further work should focus on reconstructing  
38 permafrost history in southcentral Alaska as well as additional contemporary observations of these  
39 ecosystem-protected permafrost sites lying south of the regions with relatively stable permafrost.

## 40 **1 Introduction**

41 Permafrost is a major component of the cryosphere in the northern hemisphere, covering ~24 %  
42 of the terrestrial landscape (Brown et al., 1998). Permafrost is defined as ground that remains at  
43 or below 0 °C for at least two consecutive years (Van Everdingen, 1998). Four zones describe the  
44 lateral extent of permafrost regions: continuous (90–100 %), discontinuous (50–90 %), sporadic  
45 discontinuous (10–50 %), and isolated discontinuous (<10 %). This zonation typically represents  
46 the north to south changes in spatial distribution for terrestrial permafrost in high latitudes. Mean  
47 annual ground temperatures (MAGT) in the continuous permafrost zone can be as cold at -15 °C,  
48 fall within a narrow range around -2 °C in the discontinuous permafrost zone, and can be warmer  
49 than -1 °C in sporadic and isolated permafrost zones (Smith and Riseborough, 2002; Romanovsky  
50 et al., 2010; Smith et al., 2010). In the absence of extensive ground temperature data, researchers  
51 have estimated the southern limit of permafrost in northern high latitudes with continental-scale  
52 patterns of air temperature isotherms (Brown, 1960, 1970; Ferrians, 1965; Brown et al., 1998).  
53 However, in reality complex interactions between climatic, topographic, hydrologic, and ecologic  
54 conditions operating over long time scales regulate permafrost presence and stability (Shur and

55 Jorgenson, 2007). Due to these interactions, permafrost may persist in regions with a mean annual  
56 air temperature (MAAT) above 0 °C, and it may degrade in regions with a MAAT below -10 °C  
57 (Jorgenson et al., 2010). Thus, the extent and dynamics of permafrost and permafrost-related  
58 landscape features remain poorly mapped and modelled at sufficiently fine resolution needed for  
59 predicting the impact of climate change on specific local landscapes, which is necessary for many  
60 decision makers.

61 Permafrost warming, degradation, and thaw subsidence can have significant implications  
62 for ecosystems, infrastructure, and climate at local, regional, and global scales (Jorgenson et al.,  
63 2001; Nelson et al., 2001; Schuur et al., 2008). In general, permafrost in Alaska has warmed  
64 between 0.3–6.0 °C since ground temperature measurements began between the 1950–1980s  
65 (Lachenbruch and Marshall, 1986; Romanovsky and Osterkamp, 1995; Romanovsky et al., 2002;  
66 Osterkamp, 2007; Romanovsky et al., 2010). Warming and thawing of near-surface permafrost  
67 may lead to widespread terrain instability in ice-rich permafrost in the Arctic (Jorgenson et al.,  
68 2006; Lantz and Kokelj, 2008; Gooseff et al., 2009; Jones et al., 2015; Liljedahl et al., 2016) and  
69 the sub-Arctic (Osterkamp et al., 2000; Jorgenson and Osterkamp, 2005; Lara et al., 2016). Such  
70 land surface changes can impact vegetation, hydrology, aquatic ecosystems, and soil-carbon  
71 dynamics (Grosse et al., 2011; Jorgenson et al., 2013; Kokelj et al., 2015; O’Donnell et al., 2011;  
72 Schuur et al., 2008; Vonk et al., 2015). For example, in boreal peatlands, thaw of ice-rich  
73 permafrost often converts forested permafrost plateaus into lake and wetland bog and fen  
74 complexes (Camill, 1999; Jorgenson et al., 2001; Payette et al., 2004; Sannel and Kuhry 2008;  
75 Sannel and Kuhry, 2011; Quinton et al., 2011; Jorgenson et al., 2012; Kanevskiy et al., 2014;  
76 Swindles et al., 2015; Lara et al., 2016). Furthermore, the transition from permafrost peatlands to  
77 thawed or only seasonally frozen peatlands can have a positive or a negative feedback on regional

78 and global carbon cycles depending on permafrost conditions and differential effects of thaw on  
79 net primary productivity and heterotrophic respiration (Turetsky et al., 2007; Swindles et al.,  
80 2015), as well as on the degree of loss of the former deep permafrost carbon pool (O'Donnell et  
81 al., 2012).

82 In Alaska, a variety of permafrost conditions shape roughly 80 % of the landscape (Jorgenson  
83 et al., 2008). Shur and Jorgenson (2007) proposed five classes of permafrost that describe the  
84 interaction of climatological and ecological processes. Arranged from coldest to warmest, these  
85 permafrost classes are as follows: climate-driven; climate-driven but ecosystem-modified;  
86 climate-driven but ecosystem-protected; ecosystem-driven; and ecosystem-protected. Ecosystem-  
87 protected permafrost is the warmest and most vulnerable of the five classes of permafrost and  
88 characterizes the sporadic and isolated permafrost zones. It comprises residual permafrost that  
89 persists due to favourable ecosystem factors under a climate that is not conducive to its formation.  
90 Press disturbances, associated with warming air temperatures and increases in precipitation  
91 (especially snow), and pulse disturbances, such as fire or human activities, can trigger immediate  
92 ecosystem modification and permafrost thaw in these regions (Shur and Jorgenson, 2007).

93 Since permafrost acts as a sentinel, integrator, and regulator of climate change, improved  
94 understanding of its distribution and dynamics is essential, particularly along the southern  
95 permafrost boundary (Lunardini, 1996). Southcentral Alaska, a region with a MAAT  $\sim 2$  °C, is  
96 typically mapped as being within the permafrost-free zone (Ferrians, 1965; Brown et al., 1998;  
97 Pastick et al., 2015). However, ecosystem-protected permafrost persists in southcentral Alaska in  
98 regions with present-day climatic conditions that are no longer conducive to its formation (Shur  
99 and Jorgenson, 2007). Isolated permafrost patches in southcentral Alaska exist on the western  
100 Kenai Peninsula Lowlands (Berg et al., 2009; Hopkins et al., 1955; Jorgenson et al., 2008) and in

101 the vicinity of Anchorage (Jorgenson et al., 2003; Riddle and Rooney, 2012; Kanevskiy et al.,  
102 2013). Enhanced insight into the resilience and vulnerability of ecosystem-protected permafrost  
103 is important due to its utility as a climate indicator and a forecaster of the environmental  
104 consequences expected to arise from permafrost thaw elsewhere in the boreal forest where MAAT  
105 is expected to warm beyond 0 °C in the coming decades (Beilman et al., 2001). Nevertheless, to  
106 date, detailed studies of these southcentral Alaska ecosystem-protected permafrost deposits have  
107 remained limited (Kanevskiy et al., 2013).

108         This study documents the presence of rapidly degrading permafrost plateaus on the western  
109 Kenai Peninsula lowlands of southcentral Alaska (Fig. 1), a region with a MAAT of  $1.5\pm 1$  °C (Fig.  
110 2). In mid-September 2012, we conducted field studies at several black spruce plateaus located  
111 within herbaceous wetland complexes. Continuous ground temperature measurements between  
112 16 September 2012 and 15 September 2015 confirmed the presence and degradation of permafrost.  
113 Probing, drilling, coring, and ground-penetrating radar surveys conducted in the summer, fall, and  
114 winter seasons provided additional information on the geometry of the frozen ground below the  
115 forested plateaus. Historic aerial photography and high-resolution satellite imagery from 1950,  
116 1984, 1996, and ca. 2010 were also used to map decadal-scale changes in the aerial extent of the  
117 residual permafrost plateaus in portions of four wetland complexes on the western Kenai  
118 Peninsula. This study aims to document and incorporate the loss of ecosystem-protected  
119 permafrost into the overall understanding of landscape dynamics on the western Kenai Peninsula  
120 lowlands. More importantly, insights into its stability will enhance mapping and predicting current  
121 and future permafrost extent along the southern fringe of the circumpolar permafrost region.

## 122 **2 Study Area**

123 The western Kenai Peninsula lowlands are located in southcentral Alaska, between 59.6–61.0 °N,  
124 and are generally less than 100 m above sea level (asl) (Fig. 1). The lowlands experience a semi-  
125 continental climate due to a rain shadow produced by the Kenai Mountains to the east and the  
126 presence of Cook Inlet to the west and north, and Kachemak Bay to the south (Jones et al., 2009).  
127 Regional MAAT for 1981–2010 was 1.5 °C, with a mean annual precipitation of 441 mm  
128 (<http://www.ncdc.noaa.gov/crn/observations.htm>) (Fig. 2). The lowlands represent a unique  
129 landscape where two major glacial ice fields converged during the Late Wisconsin, 25,000–21,000  
130 kya (Reger et al., 2007). The modern topography, composed of moraines, outwash fans, kettle  
131 lakes, kames, and eskers, is indicative of this glacial history (Hopkins et al., 1955). During the  
132 Holocene, the Kenai Peninsula lowlands have succeeded to boreal forest, muskeg, and wetlands  
133 laced with rivers and creeks and dotted with lakes (Anderson et al., 2006; Reger et al., 2007).  
134 Pastick et al. (2015) recently mapped this region as being permafrost-free in the upper one meter  
135 of the ground surface.

136 The Kenai lowlands are situated in an ecotone between the coastal temperate rainforest and  
137 interior boreal forest. Species assemblages depend on topography and disturbance history, as well  
138 as their location relative to the rain shadow. Black spruce (*Picea mariana*), white spruce (*Picea*  
139 *glauca*), Sitka spruce (*Picea sitchensis*), Lutz spruce (*Picea x lutzii*, hybrid of white and Sitka  
140 spruce), paper birch (*Betula kenaica*), alder (*Alnus sp.*), black cottonwood (*Populus trichocarpa*),  
141 and aspen (*Populus tremuloides*) all occur within various forest stand types. Herbaceous and  
142 woody wetland complexes intermingle with these forests in low-lying areas and river corridors.  
143 Within wetland complexes, elevated forested plateaus, primarily black spruce but with some paper  
144 birch and cottonwood and an understory of dwarf shrubs, exist where the ground surface has been

145 elevated above the regional water table. We suspected these features were associated with a  
146 volumetric expansion of freezing peat, forming a permafrost plateau, an elevated permafrost  
147 feature associated with frost heave (Zoltai, 1972, Zoltai, 1993). Characterization of degrading  
148 permafrost plateaus is the focus of our studies on the Kenai Peninsula.

### 149 **3 Methods**

150 In September 2012, we conducted field studies at a number of black spruce plateaus located within  
151 herbaceous wetland complexes (Fig. 1, Fig. 3). These studies documented frozen ground below  
152 an unfrozen layer with thicknesses ranging from 0.49m to >1.00 m. The plateau features tended  
153 to have sharply defined scalloped edges, marginal thermokarst moats, and collapse-scar  
154 depressions on their summits (Fig. 3). These traits were characteristic of the permafrost features  
155 described by Hopkins et al. (1955) on the Kenai Peninsula and similar to permafrost plateaus across  
156 colder boreal regions (Zoltai, 1972; Thie, 1974; Jorgenson et al., 2001; Camill, 2005; Sannel et al.,  
157 2015). To answer whether the frozen deposits encountered at the black spruce plateaus were  
158 indeed permafrost, we collected continuous ground temperature measurements for three years,  
159 measured late-summer thaw depths, mechanically drilled and cored for the base of the frozen  
160 ground, imaged the subsurface with ground-penetrating radar (GPR), and analysed a time series  
161 of high-resolution remotely sensed imagery. These research efforts are described in more detail  
162 below.

#### 163 **3.1 Field Instrumentation and Surveys**

164 To confirm the presence or absence of permafrost, data loggers were installed on 12 September  
165 2012 at one ground temperature monitoring site in the Browns Lake and at three sites in the Watson  
166 Lake area (Fig. 1). A 5 cm diameter Kovacs Enterprise ice auger was used to drill the boreholes

167 and cased the holes with a 4.5 cm outer-diameter polyvinyl chloride (PVC) tube from the base of  
168 the borehole to within 10 cm of the surface. Each site was instrumented with a four channel Hobo  
169 data logger (Onset U12-008) buried below the ground surface (bgs). The data loggers recorded  
170 hourly ground temperature at four depths from 0.10–3.00 m bgs using Hobo TMC1-HD and  
171 TMC2-HD thermistors (Table 1). The manufacturer-specified accuracy of the thermistors is  $\pm 0.25$   
172  $^{\circ}\text{C}$ . Prior to deployment, we placed the data logger thermistors in a  $0^{\circ}\text{C}$  ice bath for up to 45  
173 minutes to estimate a calibration factor for post-processing of the data following download in the  
174 field. After calibration in a  $0^{\circ}\text{C}$  ice bath, the precision of temperature measurements near  $0^{\circ}\text{C}$  is  
175 limited only by the sensor-logger system sensitivity, which is  $0.031^{\circ}\text{C}$  in this case. Therefore, the  
176 temperatures in our case were measured with the precision better than  $\pm 0.02^{\circ}\text{C}$  and changes in  
177 soil temperature exceeding  $0.031^{\circ}\text{C}$  can be recorded properly using this measuring system. This  
178 fact was established and demonstrated many times during our measurements in deeper boreholes  
179 using similar measuring systems when the annually measured temperature in some boreholes at  
180 deeper depths (50 m and deeper) will remain constant. These calibration techniques and  
181 measurement sensitivities are similar to improvements recorded for other measurement systems  
182 (Sannel et al., 2015; Cable et al., 2016). All data were post-processed prior to summarizing the  
183 hourly ground temperature data into daily, monthly, and annual means.

184 Additional field surveys at each study site provided information on the geometry of the  
185 frozen ground distribution and deposit types. A tile probe was used to measure the depth to frozen  
186 sediments at each ground temperature monitoring location in mid-September 2015 (limited to 2.2  
187 m bgs). At the two forested plateaus in the Watson Lake wetland complex, tile probing locations  
188 were selected randomly and split between hummock and depression microtopography. At the  
189 Browns Lake site, this depth was recorded at three points every meter along a 100 m transect across



190 the plateau feature. In addition, a topographic profile of the primary Browns Lake plateau was  
191 collected using a Leica survey-grade differential GPS (dGPS) system ( $\pm 0.02$  m vertical accuracy)  
192 on 09 October 2015 to adjust the probing measurements relative to the local topography. An  
193 additional dGPS profile was acquired on 19 February 2016 at an adjacent plateau to provide more  
194 relative feature height information in the wetland complex. At both the Browns Lake and Watson  
195 Lake locations, the frozen ground thickness was measured using the Kovacs Enterprise ice auger  
196 system powered by an 18V portable drill. At Browns Lake (site PF-BL-6), a borehole was also  
197 core-drilled to a depth of 5.38 m using a SIPRE permafrost corer (5 cm diameter) with an engine  
198 auger head for analysis of the frozen ground deposit (Fig. 1). The frozen cores were described  
199 according to the cryofacies method (French and Shur, 2010) using cryostructure classification  
200 systems inspired from Murton and French (1994) and Kanevskiy et al. (2014). Gravimetric ice  
201 contents of eleven samples were measured by oven drying at 60 °C for 168 hours. Volumetric ice  
202 contents were measured from ten well-preserved samples.

203 Implementation of GPR allowed to image certain characteristics of the frozen ground along  
204 the primary Browns Lake plateau feature. A shielded 100 MHz Mala antenna was used in July  
205 2014 and Sensors & Software 100 MHz unshielded bi-static antennas in common-offset  
206 configuration in February 2016. The data were processed using commercially available Reflex-  
207 W processing software (Sandmeier, 2008). Basic processing steps included dewow, time-zero  
208 correction, removing bad traces, and bandpass filtering (40–67.2–128–369 MHz for Mala; 25–50–  
209 200–400 MHz for Sensors & Software). Additional processing steps included an average  
210 background subtraction with a running window of 20–100 traces to reduce noise from surface  
211 multiples, where applicable, and variable gain for viewing purposes. Care was taken during  
212 processing to preserve any flat-lying reflectors. Finally, the radargrams were corrected using the

213 dGPS surface topography and converted two-way travel time to depth using an estimated average  
214 subsurface velocity of  $0.038 \text{ m ns}^{-1}$  calibrated to average direct probe depths.

### 215 **3.2 Remotely Sensed Imagery and Change Detection**

216 Historic aerial photography and contemporary high resolution satellite imagery acquired between  
217 1950–ca. 2010 provided an estimated extent of forested plateaus centred on four wetland  
218 complexes on the western Kenai Peninsula lowlands. Four change detection study areas (Fig. 1)  
219 were selected based on the presence of forested-plateau features surrounded by herbaceous  
220 wetland vegetation that likely indicated permafrost presence in the boreal wetlands on the Kenai  
221 Peninsula (Hopkins et al., 1955). Arranged from north to south, these included portions of the  
222 Mystery Creek, Watson Lake, Browns Lake, and Tustumena Lake wetland complexes (Fig. 1).  
223 Mapping forested plateau features and their change over time is a common method for detection  
224 of permafrost thaw in boreal wetlands. The land cover change associated with conversion of a  
225 forested permafrost plateau to a lake or herbaceous wetland (i.e. bog or fen) is readily detectable  
226 in high-resolution remotely sensed imagery (Thie, 1974; Camill and Clark, 1998; Osterkamp et  
227 al., 2000; Jorgenson et al., 2001; Jorgenson et al., 2008; Payette et al., 2004; Quinton et al., 2011;  
228 Lara et al., 2016).

229 A  $25 \text{ km}^2$  square study area was overlaid at each of the potential permafrost areas and  
230 clipped the wetland extent as defined by the 2001 National Land Cover Dataset for Alaska  
231 (<http://www.mrlc.gov/nlcd2011.php>) to define the mapping area. Panchromatic, Digital  
232 Orthophoto Quadrangle (DOQs) images were produced at a spatial resolution of 1.0 m for the  
233 entire Kenai Peninsula between July–August 1996. The DOQs provided the base upon which to  
234 georegister the other remotely sensed image datasets that consisted of panchromatic aerial photos  
235 collected in August 1950 (1:40,000 scale), color-infrared aerial photos acquired in 1984 (1:62,500

236 scale), and panchromatic high-resolution satellite images (<1 m spatial resolution) acquired in ca.  
237 2010. The mean RMS error associated with image georegistration was 1.82 m and ranged from  
238 1.32–2.61 m. All images were sampled to a ground resolution of 1 m. Following image  
239 registration, forested plateaus were manually digitized in a Geographic Information System  
240 (ArcGIS v. 10.1) at a mapping scale of 1:1,000 (Fig. 4). The high-spatial resolution, georegistered  
241 remotely sensed datasets allowed for the assessment of residual permafrost plateau extent in four  
242 time slices (1950, 1984, 1996, ca. 2010) and change rates across three decadal-scale time periods:  
243 (1) 1950–1984 (34 years), (2) 1984–1996 (12 years), and (3) 1996–ca. 2010 (14 years).

### 244 **3.3 Climate and Weather Data**

245 Climate and weather data were compiled from two regional stations to provide context for  
246 interpreting the ground thermal regime data and changes mapped in the remotely sensed data.  
247 Hourly air temperature data were compiled from Kenai Municipal Airport (KMA) (WBAN:  
248 26523) for 1948–1971 and 1973–Present and sub-hourly air temperature data from the Kenai 29  
249 ENE station (WBAN:26563) located at the Alaska Department of Fish and Game Moose Research  
250 Center (MRC) from September 2010–Present. Since the MRC station is more representative of  
251 the field study sites, the temperature record for MRC were reconstructed back to 1948 using a  
252 linear regression function found between KMA and MRC daily mean temperatures as summarized  
253 from hourly and sub-hourly measurements. The regression equation was calculated by comparing  
254 daily mean temperature for 1 January 2012 to 31 December 2015, and validated against daily mean  
255 temperatures at the MRC for 1 September 2010 to 31 December 2011. Lastly, daily snow depth  
256 totals were acquired from September 2012–September 2015 from MRC records  
257 (<http://wcc.sc.egov.usda.gov/nwcc/site?sitenum=966>).

## 258 **4 Results**

### 259 **4.1 Ground thermal regime of southcentral Alaska permafrost**

260 Calibrated ground temperature records collected between 16 September 2012 and 15 September  
261 2015 at one forested plateau near Browns Lake and two forested plateaus near Watson Lake  
262 confirmed the presence of near-surface permafrost on the western Kenai Peninsula lowlands (Fig.  
263 5a–5c). Over this time period, the MAGT of permafrost at 1.0 m bgs ranged from  $-0.04\text{ }^{\circ}\text{C}$  to -  
264  $0.08\text{ }^{\circ}\text{C}$  (Table 1). At the Browns Lake PF1 and the Watson Lake PF2 sites, permafrost at 2.0 m  
265 bgs had a MAGT between  $-0.06\text{ }^{\circ}\text{C}$  and  $-0.08\text{ }^{\circ}\text{C}$ . At the Browns Lake PF1 site, permafrost at 3.0  
266 m bgs had a MAGT between  $-0.07\text{ }^{\circ}\text{C}$  and  $-0.08\text{ }^{\circ}\text{C}$  (Table 1). No permafrost was detected at a  
267 black spruce forested, non-plateau site near Watson Lake between September 2012 and August  
268 2014 (Fig. 5d).

269 During the three-year observation period, an increase in near-surface ground temperatures  
270 was recorded at all three permafrost sites in response to increases in air temperature (Table 1, Fig.  
271 5). The ground temperature at 0.5 m depth was substantially below  $0\text{ }^{\circ}\text{C}$  at all three sites during  
272 the 2012–2013 winter with minimum temperatures between  $-1.33\text{ }^{\circ}\text{C}$  (Browns Lake) and  $-2.50\text{ }^{\circ}\text{C}$   
273 (Watson Lake PF2). In the 2013–2014 winter, the ground at 0.5 m depth was barely frozen at the  
274 Browns Lake and Watson Lake PF1 sites (Fig. 5a and 5b), with minimum winter temperatures at  
275  $-0.32\text{ }^{\circ}\text{C}$  and  $-0.20\text{ }^{\circ}\text{C}$ , respectively. The increase in summer ground temperatures at 0.5 m depth  
276 was also substantial. By the end of the 2012 warm period, this temperature was above  $0\text{ }^{\circ}\text{C}$  only  
277 at the Browns Lake site (the maximum was at  $0.40\text{ }^{\circ}\text{C}$ ). At the Watson Lake PF1 and PF2 sites  
278 the temperature at 0.5 m depth was just below  $0\text{ }^{\circ}\text{C}$  and never exceeded the thawing threshold,  
279 indicating that the maximum summer thaw (the active layer thickness) was just below 0.5 m during  
280 2012. However, during the summer of 2013 and 2014, the active layer thickness was more than

281 0.5 m at both of these sites and the maximum temperatures in 2014 exceeded 1 °C at the Watson  
282 Lake sites (Fig. 5b and 5c). At the Browns Lake site the temperature at 0.5 m depth reached almost  
283 2 °C before the thermistor malfunction. The ground temperature warming at 0.5 m depth continued  
284 in 2015 (Fig. 5b and 5c).

285 The increase in the shallow ground temperatures triggered warming in the near-surface  
286 permafrost at all three permafrost sites (Fig. 6). This warming was strong enough to initiate top-  
287 down permafrost thaw at the Watson Lake PF1 site in the fall of 2014 (Fig. 6b). Sensor failure  
288 during the winter of 2014–2015 prevented further observations of ground temperature at this site  
289 following thaw that winter. At the Watson Lake PF2 site bottom-up permafrost thaw was detected  
290 at a depth of 2 m during the fall of 2015 and likely associated with groundwater flow or degradation  
291 of the permafrost in the thermokarst moat that borders the plateau. At the Browns Lake site  
292 permafrost persisted at the depths between 1.0–3.0 m bgs over the three-year observation period  
293 (Fig. 6a). However, MAGT warmed by 0.02 °C to 0.01 °C at all three depths during the  
294 observation period. The temperature at 1.0 m bgs is only -0.04 °C now.

295 **4.2 Depth to permafrost table and permafrost thickness**

296 The thaw depth at the data logger observation sites, as measured with the tile probe on 16  
297 September 2015, was 0.64 m for the Watson Lake PF1 site (n=3), 0.53 m for the Watson Lake PF2  
298 site (n=6), and 0.57 m for the Browns Lake PF1 site (n=6). More systematic probing at all three  
299 sites on 16 September 2015 showed that the average depth to the permafrost table where detectable  
300 (max probe length=2.20 m) was 1.48 m (n=222). However, probing did not encounter frozen  
301 ground in the upper 2.20 m of the ground surface at an additional 140 measurement points, mostly  
302 associated with collapse-scar features and thermokarst moats. In general, depth to the permafrost  
303 table depended on the local topographic conditions at each site. Hummocks (n=164) tended to

304 have a shallower depth to the permafrost table where measureable (average of 1.12 m), while depth  
305 to the permafrost table measurements in depressions (n=58) was larger (average of 1.53 m).

306         The measurements of the depth to permafrost table were complemented with mechanical  
307 augering, coring, and GPR surveys in July 2014, September 2015, and February 2016 to constrain  
308 permafrost thickness at the field observation sites. The most detailed measurements were collected  
309 at the Browns Lake PF1 plateau feature (Fig. 7a). At this site, a topographic survey of the plateau  
310 feature was conducted to plot depth to permafrost table along with seasonally frozen depth and  
311 constraints on permafrost thickness in relation to the relative ground surface elevation along a 100  
312 m transect (Fig. 7b). The relative mean elevation of the plateau above the surrounding wetland  
313 area and the collapse-scar bog in the center was 0.49 m, with a maximum along the transect of  
314 0.95 m, and a maximum across the feature of 1.30 m. A topographic survey on an adjacent plateau  
315 feature produced a mean relative height of 0.59 m and a maximum of 1.81 m. We measured  
316 permafrost thickness at five locations and minimum-limiting permafrost thicknesses at another  
317 five locations along the Browns Lake primary plateau feature, with one limiting thickness  
318 measurement at an adjacent plateau feature using the Kovacs auger. The base of the permafrost at  
319 the two marginal plateau measurement sites at the primary plateau feature indicated a permafrost  
320 thickness of 0.45 m and 0.33 m (Fig. 7b). At the three interior plateau measurements points,  
321 permafrost was 5.57–5.65 m thick. At one of these locations (0.98 m relative height), a core was  
322 acquired. It consisted of frozen peat from 0.48–5.69 m bgs, overlying 0.25 m of unfrozen peat,  
323 with unfrozen mineral sediment at the base. At the other five locations where the bottom of  
324 permafrost was not reached, drilling operations documented permafrost at least down to between  
325 3.5–4.0 m bgs (Fig. 7b), and contained frozen peat as well.

326 The permafrost core at the PF-BL-6 site was described as poorly decomposed peat with  
327 well-developed organic-matrix cryostructures, except between depths of 3.32–3.65 m where a  
328 layer of silt and peat with mainly microlenticular cryostructure was observed (Fig. 8 and 9). The  
329 gravimetric and volumetric ice contents of the peat varied between 883–1873 % and 80–96 %,  
330 respectively, while they were 379–447 % and 81 %, respectively, in the previously mentioned silt  
331 and peat layer. The upper two meters of peat were characterized mainly by an organic-matrix  
332 porphyritic cryostructure transitioning to an organic-matrix microlenticular cryostructure with  
333 some layered ice lenses. Below two meters, the peat was characterized mainly by an organic-  
334 matrix microlenticular cryostructure with some belt-like and suspended cryostructures.

335 GPR surveys conducted in July 2014 and February 2016 provided more continuous  
336 information on the geometry associated with the permafrost table in the residual plateaus on the  
337 primary Browns Lake plateau feature (Fig. 10). The topography-corrected radargrams show a  
338 prominent reflector between 1–3 m depth that coincides with the permafrost table in both the  
339 summer (Fig. 10a) and winter (Fig. 10b) survey. The center portion of both images is characterized  
340 by moderately continuous and chaotic reflectors (Neal, 2004), as expected for records in unfrozen  
341 peat sequences (Parsekian et al., 2010) associated with the collapse-scar bog. The areas underlain  
342 by permafrost (i.e. 0–30 m, 60–90 m) show subdued reflection events deeper than the permafrost  
343 table; however, we were unable to image the permafrost base. The interpretation of these  
344 radargrams provides lateral subsurface information on the presence of a talik overlying the  
345 permafrost table.

### 346 **4.3 Remote identification of permafrost plateaus**

347 In 1950, residual permafrost plateau extent accounted for 920 ha of the 4,810 ha (19.1 %) of  
348 wetlands mapped within four change detection areas (Fig. 1, Table 2). Between 1950 and 1984,

349 permafrost plateau extent decreased to 750 ha, at an average rate of 5.1 ha yr<sup>-1</sup> (Table 3). Between  
350 1984–1996, permafrost extent dropped to 520 ha, at an average rate of 18.8 ha yr<sup>-1</sup>, the greatest  
351 rate documented in the study periods. Between 1996–2010, permafrost features continued to  
352 degrade at a rate of 9.5 ha yr<sup>-1</sup> so that by 2010, only 370 ha of the permafrost features remained.  
353 Thus, between 1950–ca. 2010, 60 % of the residual permafrost plateaus disappeared in the mapped  
354 study areas (Fig. 11 and Table 2).

355         Assessment of change in the four wetland complexes showed differences in the extent and  
356 change rate of residual permafrost plateaus overtime. The Mystery Creek study area had the most  
357 extensive permafrost plateau coverage (32.8 % of the wetland area analysed) in the 1950s relative  
358 to the Watson Lake (9.8 %), Browns Lake (11.1 %), and Tustumena Lake (15.8 %) study areas  
359 (Table 2). By ca. 2010, permafrost plateau extent in each of the study areas diminished to a cover  
360 of 14.8 %, 3.5 %, 3.8 %, and 5.2 %, respectively. Thus, there was a loss of 54.8 % of the plateau  
361 extent in the Mystery Creek study area, 64.7 % in the Watson Lake study area, 65.5 % in the  
362 Browns Lake study area, and 66.9 % in the Tustumena Lake study area between 1950–ca. 2010.  
363 These changes equate to loss rates of 0.9 % yr<sup>-1</sup> for Mystery Creek and 1.1 % yr<sup>-1</sup> for the Watson,  
364 Browns, and Tustumena Lake study areas (Table 3). Mean area loss for all four sites was 0.8 %  
365 yr<sup>-1</sup> between 1950 and 1984. During this time, loss rate was greatest for Watson Lake and Brown  
366 Lake and least for Mystery Creek. Mean loss rate for all four sites increased to 2.3 % yr<sup>-1</sup> between  
367 1984 and 1996. During this time, loss rates were greatest in the north and least in the south with  
368 Mystery Creek and Tustumena Lake losing 3.0 % yr<sup>-1</sup> and 1.2 % yr<sup>-1</sup>, respectively. Average loss  
369 rates decreased to 1.8 % yr<sup>-1</sup> between 1996 and 2010, with the three most northern sites losing  
370 approximately 1.2 % yr<sup>-1</sup>, while the Tustumena Lake study area lost 3.2 % yr<sup>-1</sup>. In terms of plateau  
371 area lost per year within the three time periods, Mystery Creek (13.8 ha yr<sup>-1</sup>), Watson Lake (1.6 ha



372 yr<sup>-1</sup>), and Browns Lake (1.3 ha yr<sup>-1</sup>) experienced the greatest areal loss rate during the 1984–1996  
373 time period. At the Tustumena Lake study area, the greatest rate of plateau extent loss (4.6 ha yr<sup>-1</sup>)  
374 occurred between 1996–ca. 2010 (Table 3).

375 This study also assessed whether the permafrost degradation occurred along the perimeter  
376 of the plateau (marginal), whether degradation was internal to the plateau, or if complete  
377 degradation of a plateau occurred. Between 1950–2010, 85.0 % of the degradation occurred as  
378 lateral thaw along the plateau margins, while internal thaw and complete loss of features accounted  
379 for 1.5 % and 13.4 %, respectively. Lateral loss of permafrost was greatest in the Watson Lake  
380 study area (90.9 %) and least (77.0 %) in the Browns Lake study area. Both Mystery Creek and  
381 Tustumena Lake shared a lateral loss of 86.0 %. Mystery Creek saw the greatest percent of internal  
382 collapse loss (3.3 %) compared to Tustumena (1.7 %) and Watson and Browns Lake (both <1.0  
383 %). The complete loss of permafrost features was greatest in Browns Lake (22.4 %) and least in  
384 Watson Lake (8.3 %). Mystery Creek and Tustumena Lake had 10.5 % and 12.3 %, respectively,  
385 of their permafrost plateaus disappear in the form of complete feature loss. During the period of  
386 remotely sensed observations complete feature loss increased from 6.7 % (1950–1984) to 21.0 %  
387 (1996–ca. 2010) of the detected change, while lateral feature loss decreased from 91.0 % (1950–  
388 1984) to 78.1 % (1996–ca. 2010) of the detected change, likely highlighting the role of  
389 fragmentation promoting complete feature degradation.

#### 390 **4.4 Climate and Weather Data**

391 The MAAT of the western Kenai Peninsula lowlands between 1981–2010 was 2.22 °C for the  
392 KMA station and estimated to be 1.79 °C for the MRC station. There was significant correlation  
393 between the KMA and MRC daily mean air temperatures for the 2012–2015 period ( $r^2=0.97$ ). The  
394 regression equation performed well during validation tests ( $r^2=0.95$ ) and was therefore used to

395 estimate daily temperature data for the MRC station back to July 1948. Mean annual air  
396 temperature has increased by 0.4 °C since 1950, with a step increase occurring in 1976 associated  
397 with the Pacific Decadal Oscillation (PDO) (Hartmann and Wendler, 2005) (Fig. 2). Between July  
398 1948 and December 1976, MAAT was 0.83 °C and 0.29°C for KMA and MRC, respectively.  
399 Following the PDO shift MAAT increased to 1.97 °C and 1.51 °C for KMA and the MRC,  
400 respectively (Fig. 2). Prior to the PDO shift, 18 (MRC) and 6 (KMA) out of 27 years had a MAAT  
401 below freezing and after the PDO shift, only 10 (MRC) and 0 (KMA) out of 39 years had a MAAT  
402 below freezing. MAAT at the MRC station was 0.88 °C (2012), 2.58 °C (2013), and 3.24 °C  
403 (2014) during our three-year ground temperature observation period of 16 September 2012 to 15  
404 September 2015. Therefore, the observations during 2014–2015 occurred during a period with  
405 anomalously high MAAT relative to the previous climate normal period, with more warming in  
406 the winter than the summer months (Table 1). Additionally, between 1948–2015, warm season  
407 (May-September) air temperatures increased by 0.02 °C yr<sup>-1</sup> for both the Kenai and MRC station,  
408 while winter season (October-April) air temperature increased by 0.04 °C yr<sup>-1</sup> (Table 4).

409

## 410 **5 Discussion**

### 411 **5.1 Presence of ecosystem-protected permafrost in southcentral Alaska**

412 These permafrost data for the residual permafrost plateaus on the Kenai Peninsula are the first such  
413 observations for isolated permafrost bodies in southcentral Alaska (Osterkamp, 2007). Based on  
414 the five classes of permafrost proposed by Shur and Jorgenson (2007), the permafrost present in  
415 wetland complexes of the western Kenai Peninsula lowlands is ecosystem-protected. The  
416 permafrost on the Kenai Peninsula is extremely warm, with a MAGT that ranges from -0.04 to -  
417 0.08 °C (Table 1; Fig. 6). Permafrost ground temperatures at all monitoring sites were near the

418 phase-equilibrium temperature at depths from 1.0–3.0 m. Latent-heat effects associated with  
419 unfrozen water content in permafrost and with seasonal phase changes in the active layer can buffer  
420 the ground thermal regime from changes in air temperature at warm permafrost sites (Romanovsky  
421 and Osterkamp, 2000) and in part can explain the persistence of ecosystem-protected permafrost  
422 on the Kenai Peninsula (Shur and Jorgenson, 2007; Jorgenson et al., 2010). Even though all  
423 thermistors were calibrated prior to installation, the ability to resolve such warm permafrost  
424 temperatures and their change over time using temperature alone is somewhat limiting. Thus,  
425 future measurements at the residual permafrost plateau sites in southcentral Alaska will be  
426 accompanied by the addition of soil moisture probes as well as borehole, nuclear magnetic  
427 resonance (NMR) which provides a direct measure of liquid water content (Parsekian et al., 2013).

428         Field surveys that included probing, augering, coring and GPR provided additional  
429 information on the vertical and spatial distribution of the warm permafrost on the western Kenai  
430 Peninsula lowlands. The average active layer thickness at the permafrost plateau ground  
431 temperature observation sites was 0.58 m. These sites were chosen for initial instrumentation in  
432 September 2012 based in part on the relatively shallow depth to the frost table. More  
433 comprehensive probing in September 2015 revealed that the average depth to the permafrost table  
434 was 1.48 m (n=222) as averaged across three plateaus. At the Brown Lake plateau, a talik  
435 overlying the permafrost table was present in February 2016. Average permafrost thickness at this  
436 feature was 5.61 m thick, whereas at an adjacent feature it was more than 6.90 m, the maximum  
437 depth of the auger flights. GPR survey data confirmed the presence of a continuous surface talik  
438 at the Browns Lake site (Fig. 10); however, we were unable to image the base of the permafrost  
439 using solely GPR, as similarly described by Lewkowicz et al. (2011). Based on visual  
440 interpretation of the permafrost peat core acquired at the PF-BL-6 site in February 2016, the

441 permafrost deposit consists mainly of ice-rich frozen peat with well-developed organic-matrix  
442 porphyritic and microlenticular cryostructures and some layered ice lenses, belt-like, and  
443 suspended cryostructures (Fig. 8 and Fig. 9). Laboratory analysis also revealed gravimetric and  
444 volumetric ice contents up to 1873 % and 96 %, respectively (Fig. 8).

## 445 **5.2 Extent and change in residual permafrost plateaus since the 1950s**

446 While previous reports of permafrost on the Kenai Peninsula exist (Hopkins et al., 1955; Jorgenson  
447 et al., 2008), they were restricted to the wetland complex (Mystery Creek) north of Sterling (Berg  
448 et al., 2009). Information on its dynamics here and elsewhere was lacking. The analysis of  
449 remotely sensed imagery and field surveys identified residual permafrost plateaus in three  
450 additional wetland complexes where it had not been previously identified (Fig. 1 and Fig. 11) and  
451 indicated that the state of permafrost within the Kenai lowlands is highly dynamic. In 1950,  
452 forested-permafrost plateau extent accounted for 19.0 % of the land cover in the 4,810 ha of  
453 wetland complexes analysed in the four change detection study areas. In each of the wetland areas  
454 analysed, permafrost plateaus accounted for more than 10.0 % of the area in 1950. However,  
455 inferred permafrost extent decreased by 60.0 % between 1950–ca. 2010, and its lateral coverage  
456 dropped below 5.0 % in three of the four study areas (Table 2).

457 The residual permafrost plateaus documented in this study share similar attributes to  
458 features elsewhere in boreal peatlands for which permafrost degradation has been inferred due to  
459 the ease of remotely detecting the conversion from forested permafrost plateau to non-permafrost  
460 herbaceous wetland or waterbody (Jorgenson et al., 2001, 2008, 2012). Thie (1974) inferred a  
461 permafrost plateau loss rate of 0.47 % yr<sup>-1</sup> between 1800–1960 for a 130,000 ha area of southern  
462 Manitoba. In Québec, Canada, a 13 ha peat bog lost 1.80 % yr<sup>-1</sup> between 1957–2003 (Payette et  
463 al., 2004). In the Northwest Territories, Canada, Quinton et al. (2011) reported a loss rate of 0.62

464 % yr<sup>-1</sup> between 1947–2008 across a 100 ha study area. In Interior Alaska (Tanana Flats), Jorgenson  
465 et al. (2001) reported a loss rate of 0.76 % yr<sup>-1</sup> for birch forested permafrost plateaus between  
466 1949–1995 using a point sampling method within a 260,000 ha wetland area. Lara et al. (2016)  
467 recently updated these numbers for the Tanana Flats by manually digitizing features with methods  
468 similar to ours and demonstrated that birch forest plateaus decreased at a much slower rate of 0.12  
469 % yr<sup>-1</sup>, and that black spruce forested permafrost plateau features appeared to be stable. Thus, the  
470 loss rate of 1.0 % yr<sup>-1</sup> that we report for the 4,810 ha mapped on the western Kenai Peninsula  
471 Lowlands between 1950–ca. 2010 are the second fastest change rates reported thus far in boreal  
472 peatlands.

### 473 **5.3 Drivers of permafrost loss**

474 Permafrost on the Kenai Peninsula is likely degrading as a result of warming air temperatures (+0.4  
475 °C decade<sup>-1</sup> since 1950), especially where warming during the winter season likely exacerbates  
476 these effects (Table 4). During the three-year observation period as well as since the 1950s,  
477 warming in the winter has been more pronounced than in the summer (Table 1 and Fig. 2) and  
478 2014–2015 had a MAAT roughly double the 1981–2010 climate normal period. Storm systems  
479 regularly bring warm air masses (>4 °C) to the region during the winter. Air temperature warming  
480 during the winter months has decreased the number of freezing degree-days which means that the  
481 ground freezes to a much lesser degree in the winter (Fig. 2 and Table 1). Therefore, ground  
482 temperatures decreased less over the winter period (Fig. 5 and Table 1), potentially leading to talik  
483 development. Previous research on permafrost plateaus in colder regions indicate that preferential  
484 warming in the winter and increased snow accumulation leads to enhanced permafrost thaw in  
485 boreal peatlands (Camill, 2005; Osterkamp, 2007). Since the Kenai Peninsula lowlands experience  
486 a semi-continental climate due to the rain shadow produced by the Kenai Mountains, a lack of

487 winter snowfall may have contributed to permafrost persistence in this region by allowing  
488 relatively cold winter air temperatures to propagate into the sub-surface. Thus, talik formation and  
489 permafrost degradation at the study sites in southcentral Alaska are likely being driven for the most  
490 part by winter air temperature warming (Fig. 2).

491         The increase in permafrost loss rate in southcentral Alaska following the 1980s is likely  
492 due to the combined effects of forest fires and a shift in the PDO after 1976. The respective pulse  
493 and press disturbances may have promoted large areas of permafrost already close to thawing, to  
494 quickly thaw, leaving only colder permafrost and permafrost with intact peat and forest cover. Fire  
495 can be an important driver of permafrost thaw (Yoshikawa et al., 2002) and thermokarst  
496 development (Jones et al., 2015). The Kenai Fire of 1947 burned the majority of the Mystery  
497 Creek study area, all of the Watson Lake study area, and the majority of the Browns Lake study  
498 area. Evidence of this fire was seen at numerous sites in the Watson Lake and Browns Lake study  
499 areas. Watson Lake and Browns Lake subsequently had the two greatest loss rates between 1950–  
500 1984 which may be related to the 1947 fire. However, the presence of black spruce burn poles  
501 were not found on all permafrost plateaus visited indicating that the burning was likely relatively  
502 patchy in the wetlands. At Browns Lake, permafrost islands that did not burn in 1947 exhibited  
503 less degradation, had thicker permafrost, denser tree cover, and larger trees than the islands that  
504 burned. Large portions of the Tustumena Lake study area burned in the 1996 Crooked Creek Fire  
505 and 2005 Fox Creek Fire. These fires likely damaged, and partially removed the protective  
506 ecosystem cover (black spruce forest and peat), and degraded several permafrost plateau features.  
507 This resulted in the Tustumena study area having the highest change rate for the latter time period  
508 and 77.0 % of the plateau loss that occurred between 1996–ca. 2010 did so in areas that burned in  
509 the 1996 and 2005 fires.

510 Bottom-up permafrost degradation was documented over the short period of direct  
511 measurements between 2012–2015. The bottom-up permafrost thaw observed at the Watson Lake  
512 PF2 site indicates that the flow of groundwater below the permafrost plateaus could be responsible  
513 for degradation (Walters et al., 1998). In addition, analysis of the remotely sensed imagery for the  
514 four select wetland complexes primarily documented lateral permafrost degradation since the  
515 1950s, as inferred by the conversion of forested plateau margins to herbaceous wetland vegetation.  
516 This type of feature loss accounted for 85.0% of the change between 1950–ca. 2010. This pattern  
517 of loss was further observed in the field through the presence of thermokarst moats and drowning  
518 black spruce trees along the margins of the permafrost plateaus (Fig. 3). This is similar to the  
519 dominant processes documented in more northerly boreal peatlands with permafrost plateaus  
520 (Thie, 1974; Camill and Clark, 1998; Osterkamp et al., 2000; Jorgenson et al., 2001; Payette et al.,  
521 2004; Quinton et al., 2011; Jorgenson et al., 2012; O’Donnell et al., 2012; Lara et al., 2015). These  
522 findings highlight the importance of groundwater flow and also the impact of saturated herbaceous  
523 wetlands that absorb heat during the summer that likely degrades permafrost along the peat plateau  
524 margins (Walters et al., 1998). It is possible that lateral permafrost degradation caused by these  
525 processes is overwhelming the protection provided by the ecosystem cover for permafrost stability  
526 on the Kenai Peninsula lowlands. Future research is required to more fully understand the role of  
527 groundwater movement on permafrost instability in the study region.

#### 528 **5.4 Proposed history of permafrost on the Kenai Peninsula**

529 During the Last Glacial Maximum (LGM), northern hemisphere permafrost extended much further  
530 south than present day (Lindgren et al., 2015). However, permafrost history in southcentral Alaska  
531 is poorly constrained. Even though the western Kenai Peninsula lowlands were almost completely  
532 glaciated during the LGM (Reger et al., 2007), the permafrost features identified in this study occur

533 in glaciolacustrine or glaciofluvial wetland complexes that were either not glaciated during the  
534 LGM (Mystery Creek) or became deglaciated before 16,000 cal yrs BP (Reger et al., 2007).  
535 Perhaps permafrost formed on the Kenai Peninsula during deglaciation or shortly thereafter during  
536 the Younger Dryas 12,900–11,700 years ago (Jones et al., 2009). However, this permafrost would  
537 have likely thawed during the Holocene Thermal Maximum (Zoltai, 1972; Kaufman et al., 2004).  
538 As the regional climate became cooler and wetter, between 8,000–5,000 years ago, *Sphagnum*  
539 accumulation and preservation on the western Kenai Peninsula lowlands may have promoted more  
540 widespread permafrost aggradation (Jones et al., 2009). Following this period, the peatlands may  
541 have progressively froze, heaving the permafrost plateaus above the water table, drying the peat-  
542 rich soils, promoting growth of black spruce, and creating a buffer layer protecting the underlying  
543 permafrost (ecosystem-protected) from the unfavourable climate for permafrost that currently  
544 exists today (Zoltai, 1972, 1995; Payette et al., 2004; Camill, 2005; Shur and Jorgenson, 2007).  
545 Growth of permafrost and heaving the peatland surface above the water table could explain low  
546 peat accumulation rates calculated in many Kenai Peninsula peatlands between 3,300–2,000 years  
547 ago (Jones and Yu, 2010; Jones et al., 2014). This also coincides with widespread neoglaciation  
548 on the Kenai Peninsula 3,000–1,500 years ago (Wiles and Calkin, 1994, Barclay et al., 2009).  
549 Alternatively, the Little Ice Age (365–165 years ago), promoted shallow permafrost formation in  
550 areas that were predominantly unfrozen throughout the Holocene (Romanovsky et al., 1992;  
551 Jorgenson et al., 2001), and thus, could account for the presence of residual permafrost on the  
552 Kenai Peninsula. The widespread loss of permafrost plateaus in central Alaska may be a result of  
553 degradation of Little Ice Age permafrost (Jorgenson et al., 2001). The age, history, and future  
554 trajectory of permafrost on the western Kenai Peninsula lowlands require further study.



## 555 **5.5 Landscape dynamics and permafrost thaw on the western Kenai Peninsula lowlands**

556 Previous and ongoing land cover change on the western Kenai Peninsula lowlands are primarily  
557 in response to the interaction of climate change and human development. Increases in summer air  
558 temperature and late-summer droughts, along with human disturbance, have been linked to the  
559 massive spruce bark beetle (*Dendroctonus rufipennis*) outbreak of the late 1990s (Berg et al., 2006;  
560 Sherriff et al., 2011), which led to subsequent timber salvage (Jones, 2008). Berg and Anderson  
561 (2006) caution that overall drier conditions on the western Kenai Peninsula, combined with  
562 standing dead spruce stands, may alter the future fire regime of this region. Wetland drying (Klein  
563 et al., 2005) and establishment of woody vegetation in wetlands (Berg et al., 2009) may be  
564 attributed to warmer air temperatures and decreases in precipitation. Furthermore, tectonic activity  
565 associated with the Great Alaska Earthquake of 1964 caused the western Kenai Peninsula to lower  
566 in elevation by 0.7–2.3 m (Plafker, 1969), while the northern portion of the peninsula subsequently  
567 uplifted 0.8–0.9m (Cohen and Freymueller, 1997), potentially altering groundwater flow paths  
568 (Gracz, 2011).

569 In this study, the loss of ecosystem-protected permafrost in the overall understanding of  
570 landscape dynamics on the western Kenai Peninsula lowlands was documented and incorporated.  
571 The degradation of permafrost can impact terrestrial and aquatic ecosystems, hydrology,  
572 infrastructure, and carbon cycling on the Kenai Peninsula (Schuur et al., 2008; Grosse et al., 2011;  
573 Jorgenson et al., 2013; Kokelj et al., 2015; Vonk et al., 2015). Permafrost degradation within the  
574 wetlands is responsible for a shift from black spruce forest plateaus to fen and bog wetland  
575 ecosystems at a mean rate of 9.2ha yr<sup>-1</sup> since the 1950s in the four change detection study areas.  
576 Permafrost plateaus redirect surface and near-surface drainage in boreal wetlands (Quinton et al.,  
577 2011), and the thaw subsidence of these features increases drainage network connectivity (Beilman  
578 and Robinson, 2003), and alters the local hydrological cycle (Hayashi et al., 2007). Thus, the loss

579 of permafrost and/or changes in seasonally frozen ground phenology could in part be aiding in  
580 observations of terrestrial and aquatic changes that have occurred on the Kenai Peninsula during  
581 the past several decades. Further work is required to better understand the past influence of  
582 permafrost on the Kenai Peninsula as well as the future loss of these warm permafrost deposits.

## 583 **6 Conclusions**

584 Based on the ground data and remotely sensed observations, it was found that peatland permafrost  
585 is currently more extensive than previously reported in southcentral Alaska, a region with a MAAT  
586 of 1.5 °C. Warm permafrost (-0.04 °C to -0.08 °C) persists on the western Kenai Peninsula  
587 lowlands in forested (black spruce), peat plateaus found in glaciolacustrine and glaciofluvial  
588 wetland complexes. At the field study sites, the depth to permafrost table on the peat plateaus  
589 averaged 1.48 m in September 2015, but was as shallow as 0.53 m at some locations. Permafrost  
590 thickness ranged from 0.33 m to greater than 6.90 m. Field surveys conducted in February 2016  
591 documented the presence of a surface talik overlying the permafrost table. In 1950, residual  
592 permafrost plateaus covered 19.0 % of the 4,810 ha wetland area mapped in the study. Within the  
593 changed detection study areas, 60.0 % of the permafrost plateaus present in 1950 had degraded by  
594 ca. 2010. In most cases, permafrost degradation equated to the loss of forest and its replacement  
595 by bog or fen vegetation, preferentially occurring along permafrost plateau margins. Permafrost  
596 loss on the Kenai Peninsula is likely associated with a warming climate, particularly during the  
597 winter season, wildfires that remove the protective ecosystem cover, groundwater flow at depth,  
598 and lateral heat transfer from wetland surface waters in the summer. Future studies on the residual  
599 permafrost plateaus on the Kenai Peninsula will provide further insight for mapping and predicting  
600 permafrost extent across Boreal permafrost regions that are currently warming.

601 **7 Data availability**

602 All data available upon request to the corresponding author.

603 **8 Author contribution**

604 B.M. Jones devised the study design and prepared the manuscript with contributions from all co-  
605 authors. B.M. Jones, C.A. Baughman, V.E. Romanovsky, E.L. Babcock, A.D. Parsekian, M.C.  
606 Jones, and E.E. Berg contributed to field instrumentation and field studies. B.M. Jones, C.A.  
607 Baughman, and G. Grosse conducted and contributed to remote sensing analysis. C.A. Baughman  
608 compiled and interpolated regional weather and climate station data. E.Stephani conducted the  
609 cryofacies analysis and laboratory testing. All co-authors contributed substantially to this research.

610 **9 Acknowledgements**

611 Funding for this research was provided by the U.S. Geological Survey Land Change Science and  
612 Land Remote Sensing programs. Support also was provided by the Russian Science Foundation  
613 (project RNF 16-17-00102). We thank the Kenai National Wildlife Refuge for granting  
614 permission to access field sites. We thank Kelly Harrell, Kobuk, Kashi, Lydia Zeglin, Josefine  
615 Lenz, Emiline Ostlind, and Callie Zuck for help with fieldwork. We thank David Swanson, Eric  
616 Klein, Neal Pastick, Mikhail Kanevskiy, and Andreas Käab for providing useful feedback and  
617 edits on an earlier version of this paper. Any use of trade, product, or firm names is for descriptive  
618 purposes only and does not imply endorsement by the U.S. Government.

619 **10 References**

- 620 Allard M., Seguin M. K., and Levesque R.: Palsas and mineral permafrost mounds in northern  
621 Quebec, In *International Geomorphology, Part II*, Gardiner V (ed). John Wiley and Sons Ltd:  
622 Chichester; 285–309, 1986.
- 623 Anderson, R. S., Hallett, D. J., Berg, E., Jass, R. B., Toney, J. L., Fontaine, C. S. de and DeVolder,  
624 A.: Holocene development of Boreal forests and fire regimes on the Kenai Lowlands of  
625 Alaska, *The Holocene*, 16(6), 791–803, doi:10.1191/0959683606hol966rp, 2006.
- 626 Barclay, D.J., Wiles, G.C. and Calkin, P.E.: Holocene glacier fluctuations in Alaska. *Quaternary*  
627 *Science Reviews*, 28, 2034-2048, 2009.
- 628 Beilman, D. W. and Robinson, S. D.: Peatland permafrost thaw and landform type along a climatic  
629 gradient, *Proc 8th Int Conf Permafr. Zurich Switz.* 21–25 July 2003, 61 – 65, 2003.
- 630 Beilman, D. W., Vitt, D. H. and Halsey, L. A.: Localized Permafrost Peatlands in Western Canada:  
631 Definition, Distributions, and Degradation, *Arct. Antarct. Alp. Res.*, 33(1), 70–77,  
632 doi:10.2307/1552279, 2001.
- 633 Berg, E. E. and Anderson, R. S.: Fire history of white and Lutz spruce forests on the Kenai  
634 Peninsula, Alaska, over the last two millennia as determined from soil charcoal, *For. Ecol.*  
635 *Manag.*, 227(3), 275–283, doi:10.1016/j.foreco.2006.02.042, 2006.
- 636 Berg, E. E., David Henry, J., Fastie, C. L., De Volder, A. D. and Matsuoka, S. M.: Spruce beetle  
637 outbreaks on the Kenai Peninsula, Alaska, and Kluane National Park and Reserve, Yukon  
638 Territory: Relationship to summer temperatures and regional differences in disturbance  
639 regimes, *For. Ecol. Manag.*, 227(3), 219–232, doi:10.1016/j.foreco.2006.02.038, 2006.
- 640 Berg, E. E., Hillman, K. M., Dial, R. and DeRuwe, A.: Recent woody invasion of wetlands on the  
641 Kenai Peninsula Lowlands, south-central Alaska: a major regime shift after 18 000 years of  
642 wet *Sphagnum*–sedge peat recruitment, *Can. J. For. Res.*, 39(11), 2033–2046,  
643 doi:10.1139/X09-121, 2009.
- 644 Brown, J., Ferrians, O. J., Heginbottom, J. A. and Melnikov, E. S.: *Circum-Arctic Map of*  
645 *Permafrost and Ground Ice Conditions*, 1998.
- 646 Brown, R. J. E.: The Distribution of Permafrost and Its Relation to Air Temperature in Canada and  
647 the U.S.S.R., *ARCTIC*, 13(3), 163–177, doi:10.14430/arctic3697, 1960.
- 648 Brown, R. J. E.: *Permafrost in Canada: its influence on Northern development*, University of  
649 Toronto Press., 1970.

650 Cable, W. L., Romanovsky, V. E., and Jorgenson, M. T.: Scaling-up Permafrost Thermal  
651 Measurements in Western Alaska using an Ecotype Approach, *The Cryosphere Discussions*,  
652 doi:10.5194/tc-2016-30, 2016.

653 Camill, P.: Patterns of boreal permafrost peatland vegetation across environmental gradients  
654 sensitive to climate warming, *Can. J. Bot.*, 77(5), 721–733, doi:10.1139/b99-008, 1999.

655 Camill, P.: Permafrost Thaw Accelerates in Boreal Peatlands During Late-20th Century Climate  
656 Warming, *Clim. Change*, 68(1–2), 135–152, doi:10.1007/s10584-005-4785-y, 2005.

657 Camill, P. and Clark, J. S.: Climate Change Disequilibrium of Boreal Permafrost Peatlands Caused  
658 by Local Processes, *Am. Nat.*, 151(3), 207–222, doi: 10.1086/286112, 1998.

659 Cohen, S.C. and Freymueller, J.T.: Deformation of the Kenai Peninsula, Alaska, *Journal of*  
660 *Geophysical Research: Solid Earth*, 102, 20479–20487. 1997.

661 Ferrians, O. J.: Permafrost Map of Alaska, U.S. Geological Survey Miscellaneous Geologic  
662 Investigations Map I-445, scale 1:2,500,000, 1965.

663 Gooseff, M. N., Balsler, A., Bowden, W. B. and Jones, J. B.: Effects of Hillslope Thermokarst in  
664 Northern Alaska, *Eos Trans. Am. Geophys. Union*, 90(4), 29–30,  
665 doi:10.1029/2009EO040001, 2009.

666 Gracz, M. B.: Comment on “Wetland drying and succession across the Kenai Peninsula Lowlands,  
667 south-central Alaska” Appears in *Can. J. For. Res.* 35: 1931–1941 (2005)., *Can. J. For. Res.*,  
668 41(2), 425–428, doi:10.1139/X10-147, 2011.

669 Grosse, G., Harden, J., Turetsky, M., McGuire, A. D., Camill, P., Tarnocai, C., Frohling, S.,  
670 Schuur, E. A. G., Jorgenson, T., Marchenko, S., Romanovsky, V., Wickland, K. P., French,  
671 N., Waldrop, M., Bourgeau-Chavez, L. and Striegl, R. G.: Vulnerability of high-latitude soil  
672 organic carbon in North America to disturbance, *J. Geophys. Res. Biogeosciences*, 116(G4),  
673 G00K06, doi:10.1029/2010JG001507, 2011.

674 Hartmann, B., and Wendler, G.: The significance of the 1976 Pacific climate shift in the  
675 climatology of Alaska, *Journal of Climate*, 18(22), 4824–4839, 2005.

676 Hayashi, M., Goeller, N., Quinton, W. L. and Wright, N.: A simple heat-conduction method for  
677 simulating the frost-table depth in hydrological models, *Hydrol. Process.*, 21(19), 2610–2622,  
678 doi:10.1002/hyp.6792, 2007.

679 Hopkins, D. M., Karlstrom, T. N. V., others, others and others: Permafrost and Ground Water in  
680 Alaska, Geological Survey Professional Paper, U.S. Geological Survey, United States  
681 Government Printing Office., 1955.

682 Jafarov, E. E., Romanovsky, V. E., Genet, H., McGuire, A. D. and Marchenko, S. S.: The effects  
683 of fire on the thermal stability of permafrost in lowland and upland black spruce forests of  
684 interior Alaska in a changing climate, *Environ. Res. Lett.*, 8(3), 035030, doi:10.1088/1748-  
685 9326/8/3/035030, 2013.

686 Jones, B. M.: Land-cover change on the southern Kenai Peninsula lowlands, Alaska using USGS  
687 land cover trends methodology, *J. Geogr. Reg. Plan.*, 1(4), 068–071, 2008.

688 Jones, B. M., Grosse, G., Arp, C. D., Miller, E., Liu, L., Hayes, D. J. and Larsen, C. F.: Recent  
689 Arctic tundra fire initiates widespread thermokarst development, *Sci. Rep.*, 5,  
690 doi:10.1038/srep15865, 2015.

691 Jones, M. C., Peteet, D. M., Kurdyla, D. and Guilderson, T.: Climate and vegetation history from  
692 a 14,000-year peatland record, Kenai Peninsula, Alaska, *Quat. Res.*, 72(2), 207–217,  
693 doi:10.1016/j.yqres.2009.04.002, 2009.

694 Jones, M.C. and Yu, Z.: Rapid deglacial and early Holocene expansion of peatlands in Alaska,  
695 *Proceedings of the National Academy of Sciences*, 107, 7347-7352, 2010.

696 Jones, M.C., Wooller, M. and Peteet, D.M.: A deglacial and Holocene record of climate variability  
697 in south-central Alaska from stable oxygen isotopes and plant macrofossils in peat,  
698 *Quaternary Science Reviews*, 87, 1–11, 2014.

699 Jorgenson, M. T., Racine, C. H., Walters, J. C. and Osterkamp, T. E.: Permafrost Degradation and  
700 Ecological Changes Associated with a Warming Climate in Central Alaska, *Clim. Change*,  
701 48(4), 551–579, doi:10.1023/A:1005667424292, 2001.

702 Jorgenson, M. T. and Osterkamp, T. E.: Response of boreal ecosystems to varying modes of  
703 permafrost degradation, *Can. J. For. Res.*, 35(9), 2100–2111, doi:10.1139/x05-153, 2005.

704 Jorgenson, M.T., Roth, J.E., Schlentner, S.F., Pullman, E.R. and Macander, M.: An ecological land  
705 survey for Fort Richardson, Alaska (No. ERDC/CRREL-TR-03-19). Engineer Research And  
706 Development Center Hanover Cold Regions Research And Engineering Lab, 2003.

707 Jorgenson, M. T., Shur, Y. L. and Pullman, E. R.: Abrupt increase in permafrost degradation in  
708 Arctic Alaska, *Geophys. Res. Lett.*, 33(2), L02503, doi:10.1029/2005GL024960, 2006.

709 Jorgenson, T., Shur, Y.L., Osterkamp, T.E.: Thermokarst in Alaska. In Proceedings of the Ninth  
710 International Conference on Permafrost, Vol. 1, June 29–July 3, 2008, Fairbanks, Alaska,  
711 Kane DL, Hinkel KM (eds). Institute of Northern Engineering, University of Alaska  
712 Fairbanks; 869–876, 2008.

713 Jorgenson, M. T., Yoshikawa, K., Kanevskiy, M. Z., Shur, Y., Romanovsky, V. E., Marchenko,  
714 S., Grosse, G., Brown, J. and Jones, B. M.: Permafrost characteristics of Alaska, in 9th  
715 International Conference on Permafrost, 2008.

716 Jorgenson, M. T., Romanovsky, V., Harden, J., Shur, Y., O'Donnell, J., Schuur, E. A. G.,  
717 Kanevskiy, M. and Marchenko, S.: Resilience and vulnerability of permafrost to climate  
718 change, *Can. J. For. Res.*, 40(7), 1219–1236, doi:10.1139/X10-060, 2010.

719 Jorgenson, M.T., Kanevskiy, M., Shur, Y., Osterkamp, T., Fortier, D., Cater, T., Miller, P.:  
720 Thermokarst lake and shore fen development in boreal Alaska. In Proceedings of the Tenth  
721 International Conference on Permafrost, Vol. 1 International contributions, June 25–29, 2012,  
722 Salekhard, Russia, Hinkel KM (ed.). The Northern Publisher: Salekhard, Russia; 179–184,  
723 2012.

724 Jorgenson, M. T., Harden, J., Kanevskiy, M., O'Donnell, J., Wickland, K., Stephanie Ewing,  
725 Manies, K., Zhuang, Q., Shur, Y., Striegl, R. and Koch, J.: Reorganization of vegetation,  
726 hydrology and soil carbon after permafrost degradation across heterogeneous boreal  
727 landscapes, *Environ. Res. Lett.*, 8(3), 035017, doi:10.1088/1748-9326/8/3/035017, 2013.

728 Kanevskiy, M., Jorgenson, T., Shur, Y., O'Donnell, J.A., Harden, J.W., Zhuang, Q., and Fortier,  
729 D.: Cryostratigraphy and Permafrost Evolution in the Lacustrine Lowlands of West-Central  
730 Alaska, *Permafrost and Periglacial Processes*, 25(1), 14-34, 2014.

731 Kanevskiy, M., Shur, Y., Krzewinski, T. and Dillon, M.: Structure and properties of ice-rich  
732 permafrost near Anchorage, Alaska, *Cold Reg. Sci. Technol.*, 93, 1–11,  
733 doi:10.1016/j.coldregions.2013.05.001, 2013.

734 Kaufman, D.S., Ager, T.A., Anderson, N.J., Anderson, P.M., Andrews, J.T., Bartlein, P.J.,  
735 Brubaker, L.B., Coats, L.L., Cwynar, L.C., Duvall, M.L. and Dyke, A.S.: Holocene thermal  
736 maximum in the western Arctic (0–180 W). *Quaternary Science Reviews*, 23, 529–560, 2004.

737 Klein, E., Berg, E. E. and Dial, R.: Wetland drying and succession across the Kenai Peninsula  
738 Lowlands, south-central Alaska, *Can. J. For. Res.*, 35(8), 1931–1941, doi:10.1139/x05-129,  
739 2005.

740 Kokelj, S. V., Tunnicliffe, J., Lacelle, D., Lantz, T. C., Chin, K. S. and Fraser, R.: Increased  
741 precipitation drives mega slump development and destabilization of ice-rich permafrost  
742 terrain, northwestern Canada, *Glob. Planet. Change*, 129, 56–68,  
743 doi:10.1016/j.gloplacha.2015.02.008, 2015.

744 Lachenbruch, A. H. and Marshall, B. V.: Changing Climate: Geothermal Evidence from  
745 Permafrost in the Alaskan Arctic, *Science*, 234(4777), 689–696,  
746 doi:10.1126/science.234.4777.689, 1986.

747 Lantz, T. C. and Kokelj, S. V.: Increasing rates of retrogressive thaw slump activity in the  
748 Mackenzie Delta region, N.W.T., Canada, *Geophys. Res. Lett.*, 35(6), L06502,  
749 doi:10.1029/2007GL032433, 2008.

750 Lara, M. J., Genet, H., McGuire, A. D., Euskirchen, E. S., Zhang, Y., Brown, D. R. N., Jorgenson,  
751 M. T., Romanovsky, V., Breen, A. and Bolton, W. R.: Thermokarst rates intensify due to  
752 climate change and forest fragmentation in an Alaskan boreal forest lowland, *Glob. Change*  
753 *Biol.*, doi:10.1111/gcb.13124, 2015.

754 Lewkowicz, A.G., Etzelmüller, B. and Smith, S.L.: Characteristics of discontinuous permafrost  
755 based on ground temperature measurements and electrical resistivity tomography, southern  
756 Yukon, Canada, *Permafrost and Periglacial Processes*, 22(4), pp.320-342, 2011.

757 Liljedahl, A.K., Boike, J., Daanen, R.P., Fedorov, A.N., Frost, G.V., Grosse, G., Hinzman, L.D.,  
758 Iijma, Y., Jorgenson, J.C., Matveyeva, N., Necsoiu, M., , Raynolds, M. K., Romanovsky, V.  
759 E., Schulla, J., Tape, K., Walker, D. A., and H. Yabuki: Pan-Arctic ice-wedge degradation in  
760 warming permafrost and its influence on tundra hydrology, *Nature Geoscience*, 2016.

761 Lindgren, A., Hugelius, G., Kuhry, P., Christensen, T. R. and Vandenberghe, J.: GIS-based Maps  
762 and Area Estimates of Northern Hemisphere Permafrost Extent during the Last Glacial  
763 Maximum, *Permafr. Periglac. Process.*, n/a–n/a, doi:10.1002/ppp.1851, 2015.

764 Lunardini, V.J.: Climatic warming and the degradation of warm permafrost. *Permafrost and*  
765 *Periglacial Processes*, 7(4), 311-320, 1996.

766 Lynch, J. A., Clark, J. S., Bigelow, N. H., Edwards, M. E. and Finney, B. P.: Geographic and  
767 temporal variations in fire history in boreal ecosystems of Alaska, *J. Geophys. Res.*  
768 *Atmospheres*, 107(D1), 8152, doi:10.1029/2001JD000332, 2002.



769 Morse, P. D., Wolfe, S. A., Kokelj, S. V. and Gaanderse, A. J. R.: The Occurrence and Thermal  
770 Disequilibrium State of Permafrost in Forest Ecotopes of the Great Slave Region, Northwest  
771 Territories, Canada, *Permafrost Periglacial Processes*, doi:10.1002/ppp.1858, 2015.

772 Murton, J.B, and French, H.M.: Cryostructures in permafrost, Tuktoyaktuk coastlands, western  
773 Arctic Canada. *Canadian Journal of Earth Sciences*, 31, pp.737–747, 1994.

774 Neal, A.: Ground-penetrating radar and its use in sedimentology: principles, problems and  
775 progress, *Earth-science reviews*, 66, 261-330, 2004.

776 Nelson, F. E., Anisimov, O. A. and Shiklomanov, N. I.: Subsidence risk from thawing permafrost,  
777 *Nature*, 410(6831), 889–890, doi:10.1038/35073746, 2001.

778 O'Donnell, J.A., Harden, J.W., McGuire, A.D., Kanevskiy, M.Z., Jorgenson, M.T., Xu, X. The  
779 effect of fire and permafrost interactions on soil carbon accumulation in an upland black  
780 spruce ecosystem of interior Alaska: implications for post-thaw carbon loss. *Global Change*  
781 *Biology* 17: 1461–1474. DOI: 10.1111/j.1365-2486. 2010.02358, 2011.

782 O'Donnell, J. A., Jorgenson, M. T., Harden, J. W., McGuire, A. D., Kanevskiy, M. Z. and  
783 Wickland, K. P.: The Effects of Permafrost Thaw on Soil Hydrologic, Thermal, and Carbon  
784 Dynamics in an Alaskan Peatland, *Ecosystems*, 15(2), 213–229, doi:10.1007/s10021-011-  
785 9504-0, 2012.

786 Osterkamp, T. E.: Characteristics of the recent warming of permafrost in Alaska, *J. Geophys. Res.*  
787 *Earth Surf.*, 112(F2), F02S02, doi:10.1029/2006JF000578, 2007.

788 Osterkamp, T. E., Viereck, L., Shur, Y., Jorgenson, M. T., Racine, C., Doyle, A. and Boone, R.  
789 D.: Observations of Thermokarst and Its Impact on Boreal Forests in Alaska, U.S.A., *Arct.*  
790 *Antarct. Alp. Res.*, 32(3), 303–315, doi:10.2307/1552529, 2000.

791 Parsekian, A. D., Slater, L., Comas, X., and Glaser, P. H.: Variations in free-phase gases in peat  
792 landforms determined by ground-penetrating radar, *Journal of Geophysical Research:*  
793 *Biogeosciences*, 115, no. G2, 2010.

794 Parsekian, A.D., Grosse, G., Walbrecker, J.O., Müller-Petke, M., Keating, K., Liu, L., Jones, B.M.  
795 and Knight, R.: Detecting unfrozen sediments below thermokarst lakes with surface nuclear  
796 magnetic resonance, *Geophysical Research Letters*, 40(3), 535-540, 2013.

797 Pastick, N. J., Jorgenson, M. T., Wylie, B. K., Nield, S. J., Johnson, K. D. and Finley, A. O.:  
798 Distribution of near-surface permafrost in Alaska: Estimates of present and future conditions,  
799 *Remote Sens. Environ.*, 168, 301–315, doi:10.1016/j.rse.2015.07.019, 2015.

800 Payette, S., Delwaide, A., Caccianiga, M. and Beauchemin, M.: Accelerated thawing of subarctic  
801 peatland permafrost over the last 50 years, *Geophys. Res. Lett.*, 31(18), L18208,  
802 doi:10.1029/2004GL020358, 2004.

803 Plafker, G.: Tectonics of the March 27, 1964 Alaska earthquake: U.S. Geological Survey  
804 Professional Paper. [online] Available from: <http://pubs.usgs.gov/pp/0543i/>, 1969.

805 Quinton, W. l., Hayashi, M. and Chasmer, L. e.: Permafrost-thaw-induced land-cover change in  
806 the Canadian subarctic: implications for water resources, *Hydrol. Process.*, 25(1), 152–158,  
807 doi:10.1002/hyp.7894, 2011.

808 Reger, R. D., Sturm, A. G., Berg, E. E. and Burns, P. A. C.: A guide to the late Quaternary  
809 history of northern and western Kenai Peninsula, Alaska: Alaska Division of Geological &  
810 Geophysical Surveys Guidebook 8, 2007.

811 Riddle, C.H., Rooney, J.W.: Encounters with relict permafrost in the Anchor- age, Alaska, area.  
812 Proceedings of the Tenth International Conference on Permafrost, Salekhard, Yamal-Nenets  
813 Autonomous District, Russia, June 25–29, 2012, 1, pp. 323–328, 2012.

814 Romanovsky, V. E., Garagula, L. S., and Seregina, N. V.: Freezing and thawing of soils under the  
815 influence of 300- and 90-year periods of temperature fluctuation. Pages 543–548 in  
816 Proceedings of the International Conference on the Role of Polar Regions in Global Change.  
817 Fairbanks (AK): Geophysical Institute, University of Alaska. 1992.

818 Romanovsky, V. E. and Osterkamp, T. E.: Effects of unfrozen water on heat and mass transport  
819 processes in the active layer and permafrost, *Permafr. Periglac. Process.*, 11(3), 219–239,  
820 doi:10.1002/1099-1530(200007/09)11:3<219::AID-PPP352>3.0.CO;2-7, 2000.

821 Romanovsky, V. E. and Osterkamp, T. E. (1995), Interannual variations of the thermal regime of  
822 the active layer and near-surface permafrost in northern Alaska. *Permafrost Periglac. Process.*,  
823 6: 313–335. doi: 10.1002/ppp.3430060404

824 Romanovsky, V., Burgess, M., Smith, S., Yoshikawa, K. and Brown, J.: Permafrost temperature  
825 records: Indicators of climate change, *Eos Trans. Am. Geophys. Union*, 83(50), 589–594,  
826 doi:10.1029/2002EO000402, 2002.

827 Romanovsky, V. E., Smith, S. L. and Christiansen, H. H.: Permafrost thermal state in the polar  
828 Northern Hemisphere during the international polar year 2007–2009: a synthesis, *Permafr.*  
829 *Periglac. Process.*, 21(2), 106–116, doi:10.1002/ppp.689, 2010.

830 Sandmeier, K. J.: REFLEXW – Windows™ 9x/NT/2000/XP-program for the processing of  
831 seismic, acoustic or electromagnetic reflection, refraction and transmission data, 2008.

832 Sannel, A.B.K., Kuhry, P. : Long-term stability of permafrost in subarctic peat plateaus, west-  
833 central Canada. *The Holocene* 18, 589–601, 2008.

834 Sannel, A.B.K., Kuhry, P.: Warming induced destabilization of peat plateau/thermokarst lake  
835 complexes. *Journal of Geophysical Research* 116, G03035, doi:10.1029/2010JG001635,  
836 2011.

837 Sannel, A. B. K., Hugelius, G., Jansson, P. and Kuhry, P.: Permafrost Warming in a Subarctic  
838 Peatland – Which Meteorological Controls are Most Important?, *Permafr. Periglac. Process.*,  
839 n/a–n/a, doi:10.1002/ppp.1862, 2015.

840 Schuur, E. A. G., Bockheim, J., Canadell, J. G., Euskirchen, E., Field, C. B., Goryachkin, S. V.,  
841 Hagemann, S., Kuhry, P., Lafleur, P. M., Lee, H., Mazhitova, G., Nelson, F. E., Rinke, A.,  
842 Romanovsky, V. E., Shiklomanov, N., Tarnocai, C., Venevsky, S., Vogel, J. G. and Zimov,  
843 S. A.: Vulnerability of Permafrost Carbon to Climate Change: Implications for the Global  
844 Carbon Cycle, *BioScience*, 58(8), 701–714, doi:10.1641/B580807, 2008.

845 Sherriff, R. L., Berg, E. E. and Miller, A. E.: Climate variability and spruce beetle (*Dendroctonus*  
846 *rufipennis*) outbreaks in south-central and southwest Alaska, *Ecology*, 92(7), 1459–1470,  
847 doi:10.1890/10-1118.1, 2011.

848 Shur, Y. L. and Jorgenson, M. T.: Patterns of permafrost formation and degradation in relation to  
849 climate and ecosystems, *Permafr. Periglac. Process.*, 18(1), 7–19, doi:10.1002/ppp.582, 2007.

850 Smith, M. W. and Riseborough, D. W.: Climate and the limits of permafrost: a zonal analysis,  
851 *Permafr. Periglac. Process.*, 13(1), 1–15, doi:10.1002/ppp.410, 2002.

852 Swindles, G.T., Morris, P.J., Mullan, D., Watson, E.J., Turner, T.E., Roland, T.P., Amesbury, M.J.,  
853 Kokfelt, U., Schoning, K., Pratte, S. and Gallego-Sala, A.: The long-term fate of permafrost  
854 peatlands under rapid climate warming, *Scientific Reports*, 5, 17951, 10.1038/srep17951,  
855 2015.

856 Thie, J.: Distribution and Thawing of Permafrost in the Southern Part of the Discontinuous  
857 Permafrost Zone in Manitoba, *ARCTIC*, 27(3), 1974.

858 Turetsky, M. R., Wieder, R. K., Vitt, D. H., Evans, R. J. and Scott, K. D.: The disappearance of  
859 relict permafrost in boreal North America: Effects on peatland carbon storage and fluxes,  
860 *Glob. Change Biol.*, 13(9), 1922–1934, doi:10.1111/j.1365-2486.2007.01381.x, 2007.

861 Van Everdingen, R. O.: Multi-Language Glossary of Permafrost and Related Ground-Ice Terms  
862 in Chinese, English, French, German, Icelandic, Italian, Norwegian, Polish, Romanian,  
863 Russian, Spanish, and Swedish. International Permafrost Association, Terminology Working  
864 Group, 1998.

865 Vonk, J. E., Tank, S. E., Bowden, W. B., Laurion, I., Vincent, W. F., Alekseychik, P., Amyot, M.,  
866 Billet, M. F., Canário, J., Cory, R. M., Deshpande, B. N., Helbig, M., Jammet, M., Karlsson,  
867 J., Larouche, J., MacMillan, G., Rautio, M., Walter Anthony, K. M. and Wickland, K. P.:  
868 Reviews and syntheses: Effects of permafrost thaw on Arctic aquatic ecosystems,  
869 *Biogeosciences*, 12(23), 7129–7167, doi:10.5194/bg-12-7129-2015, 2015.

870 Walters J.C., Racine C.H., and Jorgenson M.T.: Characteristics of permafrost in the Tanana Flats,  
871 interior Alaska, In: *Proceedings of the Seventh International Conference on Permafrost Vol*  
872 *57* (eds. Lewkowicz AG, Allard M), Université Laval, Québec. *Collection Nordicana*, 1109–  
873 1116, 1998.

874 Wiles, G. C. and Calkin, P. E.: Late Holocene, high-resolution glacial chronologies and climate,  
875 Kenai Mountains, Alaska, *Geol. Soc. Am. Bull.*, 106(2), 281–303, doi:10.1130/0016-  
876 7606(1994)106<0281:LHHRGC>2.3.CO;2, 1994.

877 Yoshikawa, K., Bolton, W. R., Romanovsky, V. E., Fukuda, M. and Hinzman, L. D.: Impacts of  
878 wildfire on the permafrost in the boreal forests of Interior Alaska, *J. Geophys. Res.*  
879 *Atmospheres*, 107(D1), 8148, doi:10.1029/2001JD000438, 2002.

880 Zoltai, S. C.: Palsas and Peat Plateaus in Central Manitoba and Saskatchewan, *Can. J. For. Res.*,  
881 2(3), 291–302, doi:10.1139/x72-046, 1972.

882 Zoltai, S.C.: Cyclic development of permafrost in the peatlands of northwestern Alberta, Canada.  
883 *Arctic and Alpine Research* 25, 240–246, 1993.

884 Zoltai, S. C.: Permafrost Distribution in Peatlands of West-Central Canada During the Holocene  
885 Warm Period 6000 Years BP, *Géographie Phys. Quat.*, 49(1), 45, doi:10.7202/033029ar,  
886 1995.

887 **Tables**

888 Table 1. Mean annual ground temperature (MAGT) data for four observation sites on the Kenai Peninsula lowlands. Browns Lake PF1,  
 889 Watson Lake PF1, and Watson Lake PF2 represent permafrost plateaus and the Watson Lake non-PF site a black spruce forested non-  
 890 plateau site. Sensor depths that were perennially frozen in a given year are in bold. Mean annual air temperature (MAAT), thawing  
 891 and freezing degree days (TDD and FDD), and average winter snow depth (MASD) are from the he MRC station (Kenai 29 ENE - AWS  
 892 702590).

893

9/16/2012 - 9/15/2013											
Browns Lake PF1		Watson Lake PF1		Watson Lake PF2		Watson Lake non-PF		KENAI 29 ENE AWS 702590 Met Station Data			
Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	MAAT (°C)	TDD Sums	FDD Sums	MASD (cm)
50	-0.02	10	0.34	10	0.05	25	0.94				
<b>100</b>	<b>-0.06</b>	25	-0.09	50	-0.30	50	0.42	0.88	1865.9	1544.3	19.3
<b>200</b>	<b>-0.08</b>	50	-0.20	<b>100</b>	<b>-0.08</b>	100	0.14				
<b>300</b>	<b>-0.08</b>	<b>100</b>	<b>-0.08</b>	<b>200</b>	<b>-0.06</b>	130	0.16				

9/16/2013 - 9/15/2014											
Browns Lake PF1		Watson Lake PF1		Watson Lake PF2		Watson Lake non-PF		KENAI 29 ENE AWS 702590 Met Station Data			
Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	MAAT (°C)	TDD Sums	FDD Sums	MASD (cm)
50*	0.17	10	0.93	10	0.86	25	0.57				
<b>100</b>	<b>-0.06</b>	25	0.40	50	-0.07	50	0.32	2.58	2066.6	1123.4	8.3
<b>200</b>	<b>-0.06</b>	50	-0.02	<b>100</b>	<b>-0.08</b>	100	0.14				
<b>300</b>	<b>-0.08</b>	<b>100</b>	<b>-0.06</b>	<b>200</b>	<b>-0.08</b>	130	0.14				

\*Thermistor failed on 24 August 2014

9/16/2014 - 9/15/2015											
Browns Lake PF1		Watson Lake PF1		Watson Lake PF2		Watson Lake non-PF		KENAI 29 ENE AWS 702590 Met Station Data			
Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	MAAT (°C)	TDD Sums	FDD Sums	MASD (cm)
50*	---	10*	---	10	1.53	25*	---				
<b>100</b>	<b>-0.04</b>	25*	---	50	0.14	50*	---	3.24	2009.8	829.1	2.7
<b>200</b>	<b>-0.06</b>	50*	---	<b>100</b>	<b>-0.07</b>	100*	---				
<b>300</b>	<b>-0.07</b>	100 <sup>#</sup> *	---	200 <sup>#</sup>	-0.07	130*	---				

\*Thermistor or data logger failure

<sup>#</sup>Permafrost thaw during observation period

894

895 Table 2. Permafrost plateau extent mapped in each study region in 1950, 1984, 1996, and ca. 2010. Analyzed wetland area for each  
 896 study region is given along with the number of features, total plateau area, mean plateau area, and plateau extent for each image  
 897 observation year. In ca. 2010, images were acquired in 2011 (Mystery Creek and Watson Lake), 2012 (Tustumena Lake), and 2013  
 898 (Browns Lake).  
 899

Study Region	Wetland Area (ha)	1950				1984				1996				ca. 2010			
		Number of Features	Total Plateau Area (ha)	Mean Plateau Area (ha)	Plateau Extent (%)	Number of Features	Total Plateau Area (ha)	Mean Plateau Area (ha)	Plateau Extent (%)	Number of Features	Total Plateau Area (ha)	Mean Plateau Area (ha)	Plateau Extent (%)	Number of Features	Total Plateau Area (ha)	Mean Plateau Area (ha)	Plateau Extent (%)
Mystery Creek	1562.0	212	511.5	2.4	32.7	237	457.7	1.9	29.3	335	292.6	0.9	18.7	321	232.3	0.7	14.9
Watson Lake	904.2	44	86.6	2.0	9.6	55	54.0	1.0	6.0	68	35.4	0.5	3.9	67	29.8	0.4	3.3
Browns Lake	1013.0	102	111.9	1.1	11.0	117	67.2	0.6	6.6	107	51.2	0.5	5.1	89	38.6	0.4	3.8
Tustumena Lake	1333.4	92	210.2	2.2	15.8	150	168.6	1.1	12.6	183	143.5	0.8	10.8	206	69.9	0.3	5.2
All Sites	4812.7	450	920.2	2.0	19.1	559	747.5	1.3	15.5	693	522.6	0.8	10.9	683	370.6	0.5	7.7

900

901

902 Table 3. Change in the extent of permafrost plateaus for each of the study regions between 1950  
 903 and ca. 2010, 1950 and 1984, 1984 and 1996, and 1996 and ca. 2010. Change is reported in aerial  
 904 units per year, proportional area change, percent change per year, and by the type of change.  
 905 Change type refers to whether the plateau loss occurred along the periphery of a feature (lateral),  
 906 in the centre of a feature (internal), or whether complete loss of a feature occurred. In ca. 2010,  
 907 images were acquired in 2011 (Mystery Creek and Watson Lake), 2012 (Tustumena Lake), and  
 908 2013 (Browns Lake).  
 909

1950 to ca. 2010						
Study Area	Area Change (ha yr <sup>-1</sup> )	Proportional Area Change (ha yr <sup>-1</sup> 100 ha <sup>-1</sup> )	Percent Change (% yr <sup>-1</sup> )	Change Type		
				Lateral (%)	Internal (%)	Complete (%)
Mystery Creek	-4.6	-0.3	-0.9	86.2	3.3	10.5
Watson Lake	-0.9	-0.1	-1.1	90.9	0.8	8.3
Browns Lake	-1.2	-0.1	-1.0	77.2	0.3	22.4
Tustumena Lake	-2.3	-0.2	-1.1	86.0	1.7	12.3
All Sites	-9.2	-0.2	-1.0	85.1	1.5	13.4

1950 to 1984						
Study Area	Area Change (ha yr <sup>-1</sup> )	Proportional Area Change (ha yr <sup>-1</sup> 100 ha <sup>-1</sup> )	Percent Change (% yr <sup>-1</sup> )	Change Type		
				Lateral (%)	Internal (%)	Complete (%)
Mystery Creek	-1.6	-0.1	-0.3	88.8	5.2	5.9
Watson Lake	-1.0	-0.1	-1.1	91.7	1.4	6.9
Browns Lake	-1.3	-0.1	-1.2	89.0	0.6	10.1
Tustumena Lake	-1.2	-0.1	-0.6	94.1	2.1	3.8
All Sites	-5.1	-0.1	-0.6	91.0	2.3	6.7

1984 to 1996						
Study Area	Area Change (ha yr <sup>-1</sup> )	Proportional Area Change (ha yr <sup>-1</sup> 100 ha <sup>-1</sup> )	Percent Change (% yr <sup>-1</sup> )	Change Type		
				Lateral (%)	Internal (%)	Complete (%)
Mystery Creek	-13.8	-0.9	-3.0	87.1	1.8	11.2
Watson Lake	-1.6	-0.2	-2.9	88.7	0.6	10.7
Browns Lake	-1.3	-0.1	-2.0	84.0	0.1	16.0
Tustumena Lake	-2.1	-0.2	-1.2	85.1	2.9	12.0
All Sites	-18.7	-0.4	-2.5	86.2	1.3	12.5

1996 to ca. 2010						
Study Area	Area Change (ha yr <sup>-1</sup> )	Proportional Area Change (ha yr <sup>-1</sup> 100 ha <sup>-1</sup> )	Percent Change (% yr <sup>-1</sup> )	Change Type		
				Lateral (%)	Internal (%)	Complete (%)
Mystery Creek	-4.0	-0.3	-1.4	82.7	3.0	14.3
Watson Lake	-0.4	-0.1	-1.1	92.2	0.5	7.3
Browns Lake	-0.7	-0.1	-1.4	58.7	0.1	41.2
Tustumena Lake	-4.6	-0.3	-3.2	78.7	0.2	21.1
All Sites	-9.5	-0.2	-1.8	78.1	1.0	21.0

910


911 Table 4. Mean annual, mean summer (May to September), and mean winter (October to April) air  
 912 temperature for the three remotely sensed image observation periods compiled from the Kenai  
 913 Municipal Airport (WBAN 26523) and estimated from the MRC station (Kenai 29 ENE - AWS  
 914 702590).  
 915

**Mean Annual Air    Mean Summer Air    Mean Winter Air  
 Temperature (°C)    Temperature (°C)    Temperature (°C)**

Remote Sensing Time Period	Mean Annual Air Temperature (°C)		Mean Summer Air Temperature (°C)		Mean Winter Air Temperature (°C)	
	Kenai Airport	MRC	Kenai Airport	MRC	Kenai Airport	MRC
1950 to 1984	1.12	0.59	9.92	10.39	-5.29	-6.54
1984 to 1996	1.77	1.31	10.28	10.78	-4.37	-5.52
1996 to 2015	2.34	1.86	10.81	11.31	-3.77	-4.95


916





918

919 Figure 1: Study area figure. (a) Recent permafrost map of Alaska (Pastick et al., 2015) indicating  
 920 permafrost presence (purple) and absence (white) in the upper one meter of the ground surface.  
 921 (b) Hillshade relief image showing a portion of southcentral Alaska. The study region on the Kenai  
 922 Peninsula lowlands is shown with the black box outline. (c) The portion of the Kenai Peninsula  
 923 lowlands where field studies and remotely sensed observations were conducted. Ground  
 924 temperature observations were collected at the Browns Lake and Watson Lake sites. The remote  
 925 sensing change detection areas are shown with a white box: (A) Mystery Creek, (B) Watson Lake,  
 926 (C) Browns Lake, and (D) Tustumena Lake wetland complexes.



927

928 Figure 2: a) Historical (1948-2015) mean annual air temperature compiled from Kenai Municipal  
 929 Airport (WBAN 26523) hourly surface data and interpolated (broken) and measured (solid) mean  
 930 annual air temperature for the MRC station (Kenai 29 ENE AWS 702590). b) Thawing degree  
 931 day (TDD) and freezing degree day (FDD) sums for 1948-2015 derived from historical and  
 932 interpolated daily mean temperature. c) Cumulative annual precipitation data from the Kenai  
 933 Municipal Airport (WBAN 26523) between 1948 and 2015.



934


935

936 Figure 3: Field photos of residual permafrost plateau landforms and thermokarst on the western  
937 Kenai Peninsula lowlands. (a) A forested permafrost plateau in the Browns Lake wetland  
938 complex. A thermokarst moat and drowning black spruce trees in the (b) Browns Lake and (c)  
939 Watson Lake wetland complexes.

940


941






942

943 Figure 4: Time series documenting the extent of a subset of permafrost features in the Browns  
 944 Lake wetland complex in (a) 1950, (b) 1984, (c) 1996, and (d) 2013. Permafrost plateau extent in  
 945 1950 is shown as a yellow polygon in each frame and other time slices outlined as green (1984),  
 946 blue (1996), and red (2013).



948

949 Figure 5: Mean daily ground temperature data plots for the four shallow boreholes on the western  
 950 Kenai Peninsula lowlands for the period of 16 September 2012 to 15 September 2015: (a) Browns  
 951 Lake PF1 site, (b) Watson Lake PF1 site, (c) Watson Lake PF2 site, and (d) Watson Lake non-PF  
 952 site. All axes scales are the same but sensor depths vary among sites based on site characteristics.  
 953 Missing data indicates sensor or thermistor failure.  
 954




955

956 Figure 6: Mean daily ground temperature data plots indicating the presence of permafrost on the  
 957 western Kenai Peninsula lowlands for the period of 16 September 2012 to 15 September 2015: (a)  
 958 Browns Lake PF1 site, (b) Watson Lake PF1 site, and (c) Watson Lake PF2 site. Top-down  
 959 permafrost thaw occurred at Watson Lake PF1 during the fall of 2014 and bottom-up permafrost  
 960 thaw occurred at Watson Lake PF2 during the fall of 2015.


961





962

963 Figure 7: (a) High-resolution satellite image showing the permafrost plateau in the Browns Lake  
 964 wetland complex where detailed field surveys were conducted as well as the location of the Browns  
 965 Lake PF1 data logger (green dot). (b) A ~100 m transect across the Browns Lake PF1 permafrost  
 966 plateau site showing ground surface height above the wetland (green line), depth to the permafrost  
 967 table (blue line and blue arrows), permafrost thickness constraints (grey arrows), seasonally frozen  
 968 ground depth (maroon arrows), and lack of permafrost (yellow stars) as measured by probing,  
 969 drilling, and coring. Locations where the permafrost table exceeded 2.2 m from the ground surface  
 970 (limiting depth for September surveys) are indicated with a non-existent blue line. Locations  
 971 where the base of the permafrost was encountered are indicated with an upward looking grey  
 972 triangle and those locations where it was not encountered, a downward looking grey triangle. The  
 973 black dashed vertical line represents the location of the PF-BL-6 permafrost core.



974  
 975

976 Figure 8: Cryostratigraphy and ice contents from borehole PF-BL-6 (inspired by Kanevskiy et  
 977 al., 2014). The gravimetric and volumetric ice contents of the peat varied between 883–1873%  
 978 and 80–96%, respectively, while they were 379–447% and 81%, respectively, in the silt and peat  
 979 layer located between 3.32 m and 3.65 m.






980  
981

982 Figure 9: (a) Stratigraphic contact between the base of the frozen silt and peat layer occurring at  
983 3.65 m and the underlying frozen peat. (b) Example of organic-matrix microlenticular to  
984 suspended cryostructures developed in the peat below the stratigraphic contact shown in (a).



985



986

987 Figure 10: GPR profiles at the intensive Browns Lake permafrost plateau (Fig. 7a) from a) late-  
988 July 2014 with Mala shielded 100-MHz antennas and b) mid-February 2016 with Sensors &  
989 Software unshielded bi-static 100-MHz antennas. Processed radargrams are on the left and  
990 processed, interpreted radargrams are on the right. Both summer and winter profiles clearly show  
991 reflectors associated with the permafrost table and in the case of (b) show the presence of a talik.  
992 However, we were unable to image the permafrost base using GPR. Note that the two GPR  
993 transects differ slightly in their orientation across the feature.  
994





995

996 Figure 11: Spatial and temporal pattern of permafrost loss within four change detection areas: a)  
 997 Mystery Creek, b) Watson Lake, c) Browns Lake, and d) Tustumena Lake. Red indicates feature  
 998 loss between 1950 and 1980, orange is feature loss between 1984 and 1996, yellow is feature loss  
 999 between 1996 and ca. 2010, and green is ca. 2010 permafrost plateau extent. The white dots  
 1000 indicate the location of field verified permafrost between 2009 and 2016. The hatched white  
 1001 polygons indicate the wetland extent where plateau features were mapped in each study area.  
 1002 Background imagery is the 1996 orthophotography.