



1 **Presence of rapidly degrading permafrost plateaus in**  
2 **southcentral Alaska**

3  
4 Benjamin M. Jones<sup>1</sup>, Carson A. Baughman<sup>1</sup>, Vladimir E. Romanovsky<sup>2</sup>, Andrew D. Parsekian<sup>3</sup>,  
5 Esther L. Babcock<sup>1</sup>, Miriam C. Jones<sup>4</sup>, Guido Grosse<sup>5</sup>, and Edward E. Berg<sup>6</sup>

6  
7  
8 <sup>1</sup> Alaska Science Center, U.S. Geological Survey, Anchorage, AK 99508, USA

9 <sup>2</sup> Geophysical Institute, University of Alaska Fairbanks, Fairbanks, AK 99775, USA

10 <sup>3</sup> Department of Geology and Geophysics, University of Wyoming, Laramie, WY 82070, USA

11 <sup>4</sup> Eastern Geology and Paleoclimate Science Center, U.S. Geological Survey, Reston, VA 20192, USA

12 <sup>5</sup> Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Potsdam, 14473, Germany

13 <sup>6</sup> Kenai National Wildlife Refuge, U.S. Fish and Wildlife Service, Soldotna, AK 99669, USA (retired)

14  
15 *Correspondence to:* Benjamin M. Jones (bjones@usgs.gov)

16 **Abstract.** Permafrost presence is determined by a complex interaction of climatic, topographic,  
17 and ecological conditions operating over long time scales. In particular, vegetation and organic  
18 layer characteristics may act to protect permafrost in regions with a mean annual air temperature  
19 (MAAT) above 0 °C. In this study, we document the presence of residual permafrost plateaus on  
20 the western Kenai Peninsula lowlands of southcentral Alaska, a region with a MAAT of 1.5±1  
21 °C (1981 to 2010). Continuous ground temperature measurements between 16 September 2012  
22 and 15 September 2015, using calibrated thermistor strings, documented the presence of warm  
23 permafrost (-0.04 to -0.08 °C). Field measurements (probing) on several plateau features during  
24 the fall of 2015 showed that the depth to the permafrost table averaged 1.48 m but was as  
25 shallow as 0.53 m. Late winter surveys (drilling, coring, and GPR) in 2016 showed that the  
26 average seasonally frozen ground thickness was 0.45 m, overlying a talik above the permafrost  
27 table. Measured permafrost thickness ranged from 0.33 to > 6.90 m. Manual interpretation of  
28 historic aerial photography acquired in 1950 indicates that residual permafrost plateaus covered  
29 920 ha as mapped across portions of four wetland complexes encompassing 4,810 ha. However,



30 between 1950 and ca. 2010, permafrost plateau extent decreased by 60 %, with lateral feature  
31 degradation accounting for 85 % of the reduction in area. Permafrost loss on the Kenai  
32 Peninsula is likely associated with a warming climate, wildfires that remove the protective forest  
33 and organic layer cover, groundwater flow at depth, and lateral heat transfer from wetland  
34 surface waters in the summer. Better understanding the resilience and vulnerability of  
35 ecosystem-protected permafrost is critical for mapping and predicting future permafrost extent  
36 and degradation across all permafrost regions that are currently warming. Further work should  
37 focus on reconstructing permafrost history in southcentral Alaska as well as additional  
38 contemporary observations of these ecosystem-protected permafrost sites lying south of the  
39 regions with relatively stable permafrost.

## 40 **1 Introduction**

41 Permafrost is a major component of the cryosphere in the northern hemisphere, covering ~24%  
42 of the terrestrial landscape (Brown et al., 1998). Permafrost is defined as ground that remains at  
43 or below 0 °C for at least two consecutive years (Van Everdingen, 1998). Four zones describe  
44 the lateral extent of permafrost regions: continuous (90-100%), discontinuous (50-90%),  
45 sporadic discontinuous (10-50%), and isolated discontinuous (< 10%). This zonation typically  
46 represents the north to south changes in spatial distribution for terrestrial permafrost in high  
47 latitudes. Mean annual ground temperatures (MAGT) in the continuous permafrost zone can be  
48 as cold as -15 °C, fall within a narrow range around -2 °C in the discontinuous permafrost zone,  
49 and can be warmer than -1 °C in sporadic and isolated permafrost zones (Smith and  
50 Riseborough, 2002; Romanovsky et al., 2010; Smith et al., 2010). In the absence of extensive  
51 ground temperature data, researchers have estimated the southern limit of permafrost in northern  
52 high latitudes with continental-scale patterns of air temperature isotherms (Brown, 1960, 1970;  
53 Ferrians, 1965; Brown et al., 1998). However, in reality complex interactions between climatic,  
54 topographic, hydrologic, and ecologic conditions operating over long time scales regulate



55 permafrost presence and stability (Shur and Jorgenson, 2007). Due to these interactions,  
56 permafrost may persist in regions with a mean annual air temperature (MAAT) above 0 °C, and  
57 it may degrade in regions with a MAAT below -10 °C (Jorgenson et al., 2010). Thus, the extent  
58 and dynamics of permafrost and permafrost-related landscape features remain poorly mapped  
59 and modelled at sufficiently fine resolution needed for predicting the impact of climate change  
60 on specific local landscapes, which is necessary for many decision makers.

61 Permafrost warming, degradation, and thaw subsidence can have significant implications  
62 for ecosystems, infrastructure, and climate at local, regional, and global scales (Jorgenson et al.,  
63 2001; Nelson et al., 2001; Schuur et al., 2008). In general, permafrost in Alaska has warmed  
64 between 0.3 °C and 6 °C since ground temperature measurements began between the 1950s and  
65 1980s (Lachenbruch and Marshall, 1986; Romanovsky and Osterkamp, 1995; Romanovsky et  
66 al., 2002; Osterkamp, 2007; Romanovsky et al., 2010). Warming and thawing of near-surface  
67 permafrost may lead to widespread terrain instability in ice-rich permafrost in the Arctic  
68 (Jorgenson et al., 2006; Lantz and Kokelj, 2008; Gooseff et al., 2009; Jones et al., 2015;  
69 Liljedahl et al., 2016) and the sub-Arctic (Osterkamp et al., 2000; Jorgenson and Osterkamp,  
70 2005; Lara et al., 2016). Such land surface changes can impact vegetation, hydrology, aquatic  
71 ecosystems, and soil-carbon dynamics (Grosse et al., 2011; Jorgenson et al., 2013; Kokelj et al.,  
72 2015; O'Donnell et al., 2011; Schuur et al., 2008; Vonk et al., 2015). For example, in boreal  
73 peatlands, thaw of ice-rich permafrost often converts forested permafrost plateaus into lake and  
74 wetland bog and fen complexes (Camill, 1999; Jorgenson et al., 2001; Payette et al., 2004;  
75 Quinton et al., 2011; Lara et al., 2016; Swindles et al., 2015). Furthermore, the transition from  
76 permafrost peatlands to thawed or only seasonally frozen peatlands can have a positive or a  
77 negative feedback on regional and global carbon cycles depending on permafrost conditions and



78 differential effects of thaw on net primary productivity and heterotrophic respiration (Turetsky et  
79 al., 2007; Swindles et al., 2015), as well as on the degree of loss of the former deep permafrost  
80 carbon pool (O'Donnell et al., 2012).

81 In Alaska, a variety of permafrost conditions shape roughly 80% of the landscape (Jorgenson  
82 et al., 2008). Shur and Jorgenson (2007) proposed five classes of permafrost that describe the  
83 interaction of climatological and ecological processes. Arranged from coldest to warmest, these  
84 permafrost classes are as follows: climate-driven; climate-driven but ecosystem-modified;  
85 climate-driven but ecosystem-protected; ecosystem-driven; and ecosystem-protected.  
86 Ecosystem-protected permafrost is the warmest and most vulnerable of the five classes of  
87 permafrost and characterizes the sporadic and isolated permafrost zones. It comprises residual  
88 permafrost that persists due to favourable ecosystem factors under a climate that is not conducive  
89 to its formation. Press disturbances, associated with warming air temperatures and increases in  
90 precipitation (especially snow), and pulse disturbances, such as fire or human activities, can  
91 trigger immediate ecosystem modification and permafrost thaw in these regions (Shur and  
92 Jorgenson, 2007).

93 Since permafrost acts as a sentinel, integrator, and regulator of climate change, improved  
94 understanding of its distribution and dynamics is essential, particularly along the southern  
95 permafrost boundary (Lunardini, 1996). Southcentral Alaska, a region with a MAAT  $\sim 2$  °C, is  
96 typically mapped as being within the permafrost-free zone (Ferrians, 1965; Brown et al., 1998;  
97 Pastick et al., 2015). However, ecosystem-protected permafrost persists in southcentral Alaska  
98 in regions with present-day climatic conditions that are no longer conducive to its formation  
99 (Shur and Jorgenson, 2007). Isolated permafrost patches in southcentral Alaska exist on the  
100 western Kenai Peninsula Lowlands (Berg et al., 2009; Hopkins et al., 1955; Jorgenson et al.,



101 2008) and in the vicinity of Anchorage (Jorgenson et al., 2003; Kanevskiy et al., 2013).  
102 Enhanced insight into the resilience and vulnerability of ecosystem-protected permafrost is  
103 important due to its utility as a climate indicator and a forecaster of the environmental  
104 consequences expected to arise from permafrost thaw elsewhere in the boreal forest where  
105 MAAT is expected to warm beyond 0 °C in the coming decades (Beilman et al., 2001).  
106 Nevertheless, to date, detailed studies of these southcentral Alaska ecosystem-protected  
107 permafrost deposits have remained limited (Kanevskiy et al., 2013).

108 In this study, we document the presence of rapidly degrading permafrost plateaus on the  
109 western Kenai Peninsula lowlands of southcentral Alaska (Fig. 1), a region with a MAAT of  
110  $1.5 \pm 1$  °C (Fig. 2). In mid-September 2012, we conducted field studies at several black spruce  
111 plateaus located within herbaceous wetland complexes. Continuous ground temperature  
112 measurements between 16 September 2012 and 15 September 2015 confirmed the presence and  
113 degradation of permafrost. Probing, drilling, coring, and ground-penetrating radar surveys  
114 conducted in the summer, fall, and winter seasons provided additional information on the  
115 geometry of the frozen ground below the forested plateaus. We also used historic aerial  
116 photography and high-resolution satellite imagery from 1950, 1984, 1996, and ca. 2010 to map  
117 decadal-scale changes in the aerial extent of the residual permafrost plateaus in portions of four  
118 wetland complexes on the western Kenai Peninsula. This study aims to document and  
119 incorporate the loss of ecosystem-protected permafrost into the overall understanding of  
120 landscape dynamics on the western Kenai Peninsula lowlands. More importantly, insights into  
121 its stability will enhance mapping and predicting current and future permafrost extent along the  
122 southern fringe of the circumpolar permafrost region.



## 123 **2 Study Area**

124 The western Kenai Peninsula lowlands are located in southcentral Alaska, between 59.6 and 61.0  
125 °N, and are generally less than 100 m asl (Fig. 1). The lowlands experience a semi-continental  
126 climate due to a rain shadow produced by the Kenai Mountains to the east and the presence of  
127 Cook Inlet to the west and north, and Kachemak Bay to the south (Jones et al., 2009). Regional  
128 MAAT for 1981–2010 is 1.5 °C, with a mean annual precipitation of 441 mm  
129 (<http://www.ncdc.noaa.gov/crn/observations.htm>). The lowlands represent a unique landscape  
130 where two major glacial ice fields converged during the Late Wisconsin, 25,000–21,000 kya  
131 (Reger et al., 2007). The modern topography, composed of moraines, outwash fans, kettle lakes,  
132 kames, and eskers, is indicative of this glacial history (Hopkins et al., 1955). During the  
133 Holocene, the Kenai Peninsula lowlands have succeeded to boreal forest, muskeg, and wetlands  
134 laced with rivers and creeks and dotted with lakes (Anderson et al., 2006; Reger et al., 2007).  
135 Pastick et al. (2015) recently mapped this region as being permafrost-free in the upper one meter  
136 of the ground surface.

137 The Kenai lowlands are situated in an ecotone between the coastal temperate rainforest  
138 and interior boreal forest. Species assemblages depend on topography and disturbance history,  
139 as well as their location relative to the rain shadow. Black spruce (*Picea mariana*), white spruce  
140 (*Picea glauca*), Sitka spruce (*Picea sitchensis*), Lutz spruce (*Picea x lutzii*, hybrid of white and  
141 Sitka spruce), paper birch (*Betula kenaica*), alder (*Alnus sp.*), black cottonwood (*Populus*  
142 *trichocarpa*), and aspen (*Populus tremuloides*) all occur within various forest stand types.  
143 Herbaceous and woody wetland complexes intermingle with these forests in low-lying areas and  
144 river corridors. Within wetland complexes, elevated forested plateaus, primarily black spruce  
145 but with some paper birch and cottonwood and an understory of dwarf shrubs, exist where the



146 ground surface has been elevated above the regional water table. We suspected these features  
147 were associated with a volumetric expansion of freezing peat, forming a permafrost plateau, an  
148 elevated permafrost feature associated with frost heave (Zoltai, 1972). These features are the  
149 focus of our studies on the Kenai Peninsula.

### 150 **3 Methods**

151 In September 2012, we conducted field studies at a number of black spruce plateaus located  
152 within herbaceous wetland complexes (Fig. 3). These studies documented frozen ground below  
153 an unfrozen layer with thicknesses ranging from 0.49 to >1.00 m. The plateau features tended to  
154 have sharply defined scalloped edges, marginal thermokarst moats, and collapse-scar depressions  
155 on their summits (Fig. 3). These traits were characteristic of the permafrost features described  
156 by Hopkins et al. (1955) on the Kenai Peninsula and similar to permafrost plateaus across colder  
157 boreal regions (Zoltai, 1972; Thie, 1974; Jorgenson et al., 2001; Camill, 2005; Sannel et al.,  
158 2015). To answer whether the frozen ground deposits encountered at the black spruce plateaus  
159 were indeed permafrost, we collected continuous ground temperature measurements for three  
160 years, measured late-summer thaw depths, mechanically drilled and cored for the base of the  
161 frozen ground, imaged the subsurface with ground-penetrating radar (GPR), and analysed a time  
162 series of high-resolution remotely sensed imagery. We describe these research efforts in more  
163 detail below.

#### 164 **3.1 Field Instrumentation and Surveys**

165 To confirm the presence or absence of permafrost, we installed data loggers on 12 September  
166 2012 at one ground temperature monitoring site in the Browns Lake and three in the Watson  
167 Lake area (Fig. 1). We used a 5-cm diameter Kovacs Enterprise ice auger to drill the boreholes



168 and cased the holes with a 4.5 cm outer-diameter PVC tube from the base of the borehole to  
169 within 10 cm of the surface. We instrumented each site with a 4-channel Hobo data logger  
170 (Onset U12-008) buried below the ground surface (bgs). The data loggers recorded hourly  
171 ground temperature at four depths from 0.10 m to 3.00 m bgs using Hobo TMC1-HD and  
172 TMC2-HD thermistors (Table 1). The manufacturer-specified accuracy of the thermistors is +/-  
173 0.25 °C. Prior to deployment, we placed the data logger thermistors in a 0 °C ice bath for up to  
174 45 minutes to estimate a calibration factor for post-processing of the data following download in  
175 the field. This calibration increased the accuracy of the ground temperature data to better than  
176 +/- 0.05 °C on average and is similar to improvements recorded for other measurement systems  
177 (Sannel et al., 2015; Cable et al., 2016). We post-processed all data prior to summarizing the  
178 hourly ground temperature data into daily, monthly, and annual means.

179 Additional field surveys at each study site provided information on the geometry of the  
180 frozen ground distribution and deposit types. We used a tile probe to measure the depth to  
181 frozen sediments at each ground temperature monitoring location in mid-September 2015  
182 (limited to 2.2 m bgs). At the two forested plateaus in the Watson Lake wetland complex, we  
183 selected tile probing locations randomly and split between hummock and depression  
184 microtopography. At the Browns Lake site, we recorded this depth at three points every meter  
185 along a 100 m transect across the plateau feature. In addition, we collected a topographic profile  
186 of the primary Browns Lake plateau using a Leica survey-grade differential GPS (dGPS) system  
187 (+/- 0.02 m vertical accuracy) on 09 October 2015 to adjust the probing measurements relative to  
188 the local topography. An additional dGPS profile was acquired on 19 February 2016 at an  
189 adjacent plateau to provide more relative feature height information in the wetland complex. At  
190 both the Browns Lake and Watson Lake locations, we measured the frozen ground thickness





191 using the Kovacs Enterprise ice auger system powered by an 18V portable drill. At Browns  
192 Lake, we also collected a core for visual analysis of the frozen ground deposit using a SIPRE  
193 permafrost corer with an engine auger head. We calculated the excess ice fraction (EIF) for  
194 three sites at the Browns Lake plateau for which we had detailed height, depth to permafrost  
195 table, and permafrost base information following Lewkowicz et al. (2011) to enable comparison  
196 of EIF with previously studied permafrost plateaus.

197 Implementation of GPR allowed us to image certain characteristics of the frozen ground  
198 along the primary Browns Lake plateau feature. We used a shielded 100-MHz Mala antenna in  
199 July 2014 and Sensors & Software 100-MHz unshielded bi-static antennas in common-offset  
200 configuration in February 2016. We processed the data using commercially available Reflex-W  
201 processing software (Sandmeier, 2008). Basic processing steps included dewow, time-zero  
202 correction, removing bad traces, and bandpass filtering (40-67.2-128-369 MHz for Mala; 25-50-  
203 200-400 MHz for Sensors & Software). Additional processing steps included an average  
204 background subtraction with a running window of 20 to 100 traces to reduce noise from surface  
205 multiples, where applicable, and variable gain for viewing purposes. Care was taken during  
206 processing to preserve any flat-lying reflectors. Finally, we corrected the radargrams using the  
207 dGPS surface topography and converted two-way travel time to depth using an estimated  
208 average subsurface velocity of  $0.038 \text{ m ns}^{-1}$  calibrated to average direct probe depths.

### 209 **3.2 Remotely Sensed Imagery and Change Detection**

210 Historic aerial photography and contemporary high resolution satellite imagery acquired between  
211 1950 and ca. 2010 provided an estimated extent of forested plateaus centred on four wetland  
212 complexes on the western Kenai Peninsula lowlands. We selected four change detection study  
213 areas (Fig. 1) based on the presence of forested, plateau features surrounded by herbaceous



214 wetland vegetation that likely indicated permafrost presence in the boreal wetlands on the Kenai  
215 Peninsula (Hopkins et al., 1955). Arranged from north to south, these included portions of the  
216 Mystery Creek, Watson Lake, Browns Lake, and Tustumena Lake wetland complexes (Fig. 1).  
217 Mapping forested plateau features and their change over time is a common method for detection  
218 of permafrost thaw in boreal wetlands. The land cover change associated with conversion of a  
219 forested permafrost plateau to a lake or herbaceous wetland (i.e. bog or fen) is readily detectable  
220 in high-resolution remotely sensed imagery (Thie, 1974; Camill and Clark, 1998; Osterkamp et  
221 al., 2000; Jorgenson et al., 2001; Payette et al., 2004; Quinton et al., 2011; Lara et al., 2016).

222 We overlaid a 25-km<sup>2</sup> square study area at each of the potential permafrost areas and  
223 clipped the wetland extent as defined by the 2001 National Land Cover Dataset for Alaska  
224 (<http://www.mrlc.gov/nlcd2011.php>) to define the mapping area. Panchromatic, Digital  
225 Orthophoto Quadrangle (DOQs) images were produced at a spatial resolution of 1.0 m for the  
226 entire Kenai Peninsula between July and August 1996. The DOQs provided the base upon which  
227 to georegister the other remotely sensed image datasets that consisted of panchromatic aerial  
228 photos collected in August 1950 (1:40,000 scale), color-infrared aerial photos acquired in 1984  
229 (1:62,500 scale), and panchromatic high-resolution satellite images (< 1 m spatial resolution)  
230 acquired in ca. 2010. The mean RMS error associated with image georegistration was 1.82 m  
231 and ranged from 1.32 m to 2.61 m and all images were sampled to a ground resolution of 1 m.  
232 Following image registration, we manually digitized forested plateaus in a Geographic  
233 Information System (ArcGIS v. 10.1) at a mapping scale of 1:1,000 (Fig. 4). The high-spatial  
234 resolution, georegistered remotely sensed datasets allowed for the assessment of residual  
235 permafrost plateau extent in four time slices (1950, 1984, 1996, ca. 2010) and change rates



236 across three decadal-scale time periods: (1) 1950 to 1984 (34 years), (2) 1984 to 1996 (12 years),  
237 and (3) 1996 to ca. 2010 (14 years).

### 238 **3.3 Climate and Weather Data**

239 We compiled climate and weather data from two regional stations to provide context for  
240 interpreting the ground thermal regime data and changes mapped in the remotely sensed data.  
241 We compiled hourly air temperature data from Kenai Municipal Airport (WBAN: 26523) for  
242 1948 - 1971 and 1973 – Present and sub-hourly air temperature data from the Kenai 29 ENE  
243 station (WBAN:26563) located at the Alaska Department of Fish and Game Moose Research  
244 Center (MRC) from September 2010 - Present. Since the MRC station is more representative of  
245 the field study sites, we reconstructed the temperature record for MRC back to 1948 using a  
246 linear regression function found between Kenai and MRC daily mean temperatures as  
247 summarized from hourly and sub-hourly measurements. The regression equation was calculated  
248 by comparing daily mean temperature for 1 January 2012 to 31 December 2015, and validated  
249 against daily mean temperatures at the MRC for 1 September 2010 to 31 December 2011.  
250 Lastly, we acquired daily snow depth totals recorded at the MRC from September 2012 –  
251 September 2015 (<http://wcc.sc.egov.usda.gov/nwcc/site?sitenum=966>).

## 252 **4 Results**

### 253 **4.1 Ground thermal regime of southcentral Alaska permafrost**

254 Calibrated ground temperature records collected between 16 September 2012 and 15 September  
255 2015 at one forested plateau near Browns Lake and two forested plateaus near Watson Lake  
256 confirmed the presence of near-surface permafrost on the western Kenai Peninsula lowlands  
257 (Fig. 5a-5c). Over this time period, the MAGT of permafrost at 1 m bgs ranged from -0.04 °C to



258 -0.08 °C (Table 1). At the Browns Lake PF1 and the Watson Lake PF2 sites, permafrost at 2.0 m  
259 bgs had a MAGT between -0.06 °C and -0.08 °C. At the Browns Lake PF1 site, permafrost at  
260 3.0 m bgs had a MAGT between -0.07 °C and -0.08 °C (Table 1). We detected no permafrost at  
261 a black spruce forested, non-plateau site near Watson Lake between September 2012 and August  
262 2014 (Fig. 5d).

263 During the three-year observation period, an increase in near-surface ground  
264 temperatures was recorded at all three permafrost sites in response to increases in air temperature  
265 (Table 1, Fig. 5). The ground temperature at 0.5 m depth was substantially below 0 °C at all  
266 three sites during the 2012-2013 winter with minimum temperatures between -1.33 °C (Browns  
267 Lake) and -2.5°C (Watson Lake PF2). In the 2013-2014 winter, the ground at 0.5 m depth was  
268 barely frozen at the Browns Lake and Watson Lake PF1 sites (Fig. 5a and 5b), with minimum  
269 winter temperatures at -0.32 °C and -0.2 °C, respectively. The increase in summer ground  
270 temperatures at 0.5 m depth was also substantial. By the end of the 2012 warm period, this  
271 temperature was above 0 °C only at the Browns Lake site (the maximum was at 0.4 °C). At the  
272 Watson Lake PF1 and PF2 sites the temperature at 0.5 m depth was just below 0 °C and never  
273 exceeded the thawing threshold, indicating that the maximum summer thaw (the active layer  
274 thickness) was just below 0.5 m during 2012. However, during the summer of 2013 and 2014,  
275 the active layer thickness was more than 0.5 m at both of these sites and the maximum  
276 temperatures in 2014 exceeded 1°C at the Watson Lake sites (Fig. 5b and 5c). At the Browns  
277 Lake site the temperature at a 0.5 m reached almost 2 °C before the thermistor malfunction. The  
278 ground temperature warming at 0.5 m depth continued in 2015 (Fig. 5b and 5c).

279 The increase in the shallow ground temperatures triggered warming in the near-surface  
280 permafrost at all three permafrost sites (Fig. 6). This warming was strong enough to initiate top-



281 down permafrost thaw at the Watson Lake PF1 site in the fall of 2014 (Fig. 6b). Sensor failure  
282 during the winter of 2014/2015 prevented further observations of ground temperature at this site  
283 following thaw that winter. At the Watson Lake PF2 site bottom up permafrost thaw was  
284 detected during the fall of 2015 and likely associated with groundwater flow or degradation of  
285 the permafrost in the thermokarst moat that borders the plateau. At the Browns Lake site  
286 permafrost persisted at the depths between 1 and 3 m bgs over the three-year observation period  
287 (Fig. 6a). However, MAGT warmed by 0.02 to 0.01 °C at all three depths during the observation  
288 period. The temperature at 1 m bgs is only -0.04 °C now.

#### 289 **4.2 Depth to permafrost table and permafrost thickness**

290 The thaw depth at our data logger observation sites as measured with the tile probe on 16  
291 September 2015 was 0.64 m for the Watson Lake PF1 site (n = 3), 0.53 m for the Watson Lake  
292 PF2 site (n = 6), and 0.57 m for the Browns Lake PF1 site (n = 6). More systematic probing at  
293 all three sites on 16 September 2015 showed that the average depth to the permafrost table where  
294 detectable (max probe length = 2.20 m) was 1.48 m (n = 222). However, probing did not  
295 encounter frozen ground in the upper 2.20 m of the ground surface at an additional 140  
296 measurement points, mostly associated with collapse-scar features and thermokarst moats. In  
297 general, depth to the permafrost table depended on the local topographic conditions at each site.  
298 Hummocks (n = 164) tended to have a shallower depth to the permafrost table where  
299 measureable (average of 1.12 m), while depth to the permafrost table measurements in  
300 depressions (n = 58) was larger (average of 1.53 m).

301 The measurements of the depth to permafrost table were complemented with mechanical  
302 drilling, coring, and GPR surveys in July 2014, September 2015, and February 2016 to constrain  
303 permafrost thickness at the field observation sites. The most detailed measurements were



304 collected at the Browns Lake PF1 plateau feature (Fig. 7a). At this site, we conducted a  
305 topographic survey of the plateau feature to plot depth to permafrost table along with seasonally  
306 frozen depth and constraints on permafrost thickness in relation to the relative ground surface  
307 elevation along a 100 m transect (Fig. 7b). The relative mean elevation of the plateau above the  
308 surrounding wetland area was 0.49 m (not including the collapse-scar bog in the center), with a  
309 maximum along the transect of 0.95 m, and a maximum across the feature of 1.3 m. A  
310 topographic survey on an adjacent plateau feature produced a mean relative height of 0.59 m and  
311 a maximum of 1.81 m. We measured permafrost thickness at five locations and minimum-  
312 limiting permafrost thicknesses at another five locations along the Browns Lake primary plateau  
313 feature, with one limiting thickness measurement at an adjacent plateau feature using the Kovacs  
314 auger. The base of the permafrost at the two marginal plateau measurement sites at the primary  
315 plateau feature indicated a permafrost thickness of 0.45 and 0.33 m (Fig. 7b). At the three  
316 interior plateau measurements points, permafrost was 5.57 to 5.65 m thick. At one of these  
317 locations (0.98 m relative height), we acquired a core that consisted of frozen peat from 0.48 m  
318 bgs down to 5.69 m bgs, overlying 0.25 m of unfrozen peat, with unfrozen mineral sediment at  
319 the base. At the other five locations where the bottom of permafrost was not reached, drilling  
320 operations documented permafrost at least down to between 3.5 and 4.0 bgs (Fig. 7b), and  
321 contained frozen peat as well. The EIF for the three interior measurements points on the Browns  
322 Lake plateau, where we had information on relative height, depth to permafrost table, and depth  
323 of permafrost base, ranged from 0.09 to 0.13. At an adjacent plateau (not shown) the minimum  
324 permafrost thickness was 6.90 m bgs, at which point we ran out of auger flight extensions. At  
325 Watson Lake PF1, drilling efforts detected permafrost base between 1.30 and 1.50 meters bgs.  
326 At the Watson Lake PF2 site, the permafrost base was between 1.96 and 2.04 m bgs.



327 GPR surveys conducted in July 2014 and February 2016 provided more continuous  
328 information on the geometry associated with the permafrost table in the residual plateaus on the  
329 primary Browns Lake plateau feature (Fig. 8). The topography-corrected radargrams show a  
330 prominent reflector between 1 – 3 m depth that coincides with the permafrost table in both the  
331 summer (Fig. 8a) and winter (Fig. 8b) survey. The center portion of both images is characterized  
332 by moderately continuous and chaotic reflectors (Neal, 2004) as expected for records in unfrozen  
333 peat sequences (Parsekian et al., 2010) associated with the collapse-scar bog. The areas  
334 underlain by permafrost (i.e. 0 – 30 m, 60 – 90 m) show subdued reflection events deeper than  
335 the permafrost table; however, we were unable to image the permafrost base. Our interpretation  
336 of these radargrams provides lateral subsurface information on the presence of a talik overlying  
337 the permafrost table.

#### 338 **4.3 Remote identification of permafrost plateaus**

339 In 1950, residual permafrost plateau extent accounted for 920 ha of the 4,810 ha (19.1%) of  
340 wetlands mapped within four change detection areas (Fig. 1, Table 2). Between 1950 and 1984,  
341 permafrost plateau extent decreased to 750 ha, at an average rate of 5.1 ha yr<sup>-1</sup> (Table 3).  
342 Between 1984 and 1996, permafrost extent dropped to 520 ha, at an average rate of 18.8 ha yr<sup>-1</sup>,  
343 the greatest rate documented in our study periods. Between 1996 and 2010, permafrost features  
344 continued to degrade at a rate of 9.5 ha yr<sup>-1</sup> so that by 2010, only 370 ha of the permafrost  
345 features remained. Thus, between 1950 and ca. 2010, 60% of the residual permafrost plateaus  
346 disappeared in our mapped study areas (Fig. 9).

347 Assessment of change in the four wetland complexes showed differences in the extent  
348 and change rate of residual permafrost plateaus overtime. The Mystery Creek study area had the  
349 most extensive permafrost plateau coverage (32.8 % of the wetland area analysed) in the 1950s



350 relative to the Watson Lake (9.8 %), Browns Lake (11.1 %), and Tustumena Lake (15.8 %) study  
351 areas (Table 2). By ca. 2010, permafrost plateau extent in each of the study areas diminished to  
352 a cover of 14.8 %, 3.5 %, 3.8 %, and 5.2 %, respectively. Thus, there was a loss of 54.8 % of the  
353 plateau extent in the Mystery Creek study area, 64.7 % in the Watson Lake study area, 65.5 % in  
354 the Browns Lake study area, and 66.9 % in the Tustumena Lake study area between 1950 and ca.  
355 2010. These changes equate to loss rates of 0.9 % yr<sup>-1</sup> for Mystery Creek and 1.1 % yr<sup>-1</sup> for the  
356 Watson, Browns, and Tustumena Lake study areas (Table 3). Mean area loss for all four sites  
357 was 0.8 % yr<sup>-1</sup> between 1950 and 1984. During this time, loss rate was greatest for Watson Lake  
358 and Brown Lake and least for Mystery Creek. Mean loss rate for all four sites increased to 2.3 %  
359 yr<sup>-1</sup> between 1984 and 1996. During this time, loss rates were greatest in the north and least in  
360 the south with Mystery Creek and Tustumena Lake losing 3.0 % yr<sup>-1</sup> and 1.2 % yr<sup>-1</sup>, respectively.  
361 Average loss rates decreased to 1.8 % yr<sup>-1</sup> between 1996 and 2010, with the three most northern  
362 sites losing approximately 1.2 % yr<sup>-1</sup>, while the Tustumena Lake study area lost 3.2 % yr<sup>-1</sup>. In  
363 terms of plateau area lost per year within the three time periods, Mystery Creek (13.8 ha yr<sup>-1</sup>),  
364 Watson Lake (1.6 ha yr<sup>-1</sup>), and Browns Lake (1.3 ha yr<sup>-1</sup>) experienced the greatest areal loss rate  
365 during the 1984 to 1996 time period. At the Tustumena Lake study area, the greatest rate of  
366 plateau extent loss (4.6 ha yr<sup>-1</sup>) occurred between 1996 and ca. 2010 (Table 3).

367 We also assessed whether the permafrost degradation occurred along the perimeter of the  
368 plateau (marginal), whether degradation was internal to the plateau, or if complete degradation of  
369 a plateau occurred. Between 1950 and 2010, 85 % of the degradation occurred as lateral thaw  
370 along the plateau margins, while internal thaw and complete loss of features accounted for 1.5 %  
371 and 13.4 %, respectively. Lateral loss of permafrost was greatest in the Watson Lake study area  
372 (90.9 %) and least (77 %) in the Browns Lake study area. Both Mystery Creek and Tustumena





373 Lake shared a lateral loss of 86 %. Mystery Creek saw the greatest percent of internal collapse  
374 loss (3.3 %) compared to Tustumena (1.7 %) and Watson and Browns Lake (both <1 %). The  
375 complete loss of permafrost features was greatest in Browns Lake (22.4 %) and least in Watson  
376 Lake (8.3 %). Mystery Creek and Tustumena Lake had 10.5 % and 12.3 %, respectively, of their  
377 permafrost plateaus disappear in the form of complete feature loss. During the period of  
378 remotely sensed observations complete feature loss increased from 6.7 % (1950 to 1984) to 21.0  
379 % (1996 to ca. 2010) of the detected change, while lateral feature loss decreased from 91.0 %  
380 (1950 to 1984) to 78.1 % (1996 to ca. 2010) of the detected change, likely highlighting the role  
381 of fragmentation promoting complete feature degradation.

#### 382 **4.4 Climate and Weather Data**

383 The MAAT of the western Kenai Peninsula lowlands between 1981 and 2010 was 2.22 °C for  
384 the Kenai Municipal Airport and estimated to be 1.79 °C for the MRC station. There was  
385 significant correlation between Kenai daily mean air temperature and the MRC daily mean air  
386 temperature for the 2012-2015 period ( $r^2 = 0.97$ ). The regression equation performed well during  
387 validation tests ( $r^2 = 0.95$ ) and was therefore used to estimate daily temperature data for the MRC  
388 station back to July 1948. Mean annual air temperature has increased by 0.4 °C since 1950, with  
389 a step increase occurring in 1976 associated with the Pacific Decadal Oscillation (PDO)  
390 (Hartmann and Wendler, 2005) (Fig. 2). Between July 1948 and December 1976, MAAT was  
391 0.83 °C and 0.29 °C for Kenai and MRC, respectively. Following the PDO shift MAAT  
392 increased to 1.97 °C and 1.51 °C for Kenai and the MRC, respectively (Fig. 2). Prior to the PDO  
393 shift, 18 (MRC) and 6 (Kenai) out of 27 years had a MAAT below freezing and after the PDO  
394 shift, only 10 (MRC) and 0 (Kenai) out of 39 years had a MAAT below freezing. MAAT at the  
395 MRC station was 0.88 °C (2012), 2.58 °C (2013), and 3.24 °C (2014) during our three-year



396 ground temperature observation period of 16 Sept 2012 to 15 Sept 2015. Therefore, our  
397 observations during 2014 and 2015 occurred during a period with anomalously high MAAT  
398 relative to the previous climate normal period, with more warming in the winter than the summer  
399 months (Table 1). Additionally, between 1948 and 2015, warm season (May-Sept) air  
400 temperatures increased by  $0.02\text{ °C yr}^{-1}$  for both the Kenai and MRC station, while winter season  
401 (Oct-April) air temperature increased by  $0.04\text{ °C yr}^{-1}$  (Table 4).

402

## 403 **5 Discussion**

### 404 **5.1 Presence of ecosystem-protected permafrost in southcentral Alaska**

405 Our permafrost data for the residual permafrost plateaus on the Kenai Peninsula are the first such  
406 observations for isolated permafrost bodies in southcentral Alaska (Osterkamp, 2007). Based on  
407 the five classes of permafrost proposed by Shur and Jorgenson (2007), the permafrost present in  
408 wetland complexes of the western Kenai Peninsula lowlands is ecosystem-protected. The  
409 permafrost on the Kenai Peninsula is extremely warm, with a MAGT that ranges from  $-0.04$  to -  
410  $0.08\text{ °C}$  (Table 1; Fig. 6). Permafrost at all ground temperature monitoring sites and depths from  
411 1.0 to 3.0 m were near the phase-equilibrium temperature. Latent-heat effects associated with  
412 unfrozen water content in permafrost and with seasonal phase changes in the active layer can  
413 buffer the ground thermal regime from changes in air temperature at warm permafrost sites  
414 (Romanovsky and Osterkamp, 2000) and in part can explain the persistence of ecosystem-  
415 protected permafrost on the Kenai Peninsula (Shur and Jorgenson, 2007; Jorgenson et al., 2010).  
416 Even though we calibrated all thermistors prior to installation, the ability to resolve such warm  
417 permafrost temperatures and their change over time using temperature alone is somewhat  
418 limiting. Thus, future measurements at the residual permafrost plateau sites in southcentral



419 Alaska will be accompanied by the addition of soil moisture probes as well as borehole, nuclear  
420 magnetic resonance (NMR) which provides a direct measure of liquid water content (Parsekian  
421 et al., 2013).

422 Field surveys that included probing, drilling, coring and GPR provided additional  
423 information on the vertical and spatial distribution of the warm permafrost on the western Kenai  
424 Peninsula lowlands. The average active layer thickness at our permafrost plateau ground  
425 temperature observation sites was 0.58 m. We chose these sites for initial instrumentation in  
426 September 2012 based in part on the relatively shallow depth to the frost table. More  
427 comprehensive probing in September 2015 revealed that the average depth to the permafrost  
428 table was 1.48 m (n=222) as averaged across three plateaus. At the Brown Lake plateau, a talik  
429 overlying the permafrost table was present in February 2016. Average permafrost thickness at  
430 this feature was 5.61 m thick, whereas at an adjacent feature it was more than 6.90 m, the  
431 maximum depth of our auger flights. GPR survey data confirmed the presence of a continuous  
432 surface talik at the Browns Lake site (Fig. 8); however, we were unable to image the base of the  
433 permafrost using solely GPR, as similarly described by Lewkowicz et al. (2011). EIF was 0.09  
434 to 0.13 for three measurement sites on the primary Browns Lake plateau feature. In comparison,  
435 Allard et al. (1986) studied similar peat plateau features in Canada which typically were as high  
436 as one-third the thickness of permafrost or an EIF of 0.33. Lewkowicz et al. (2011)  
437 demonstrated that features with EIF values below 0.33 likely results from ice-poor permafrost  
438 and/or a high unfrozen water content of the permafrost. Based on visual interpretation of the  
439 permafrost peat core acquired in February 2016, the permafrost deposit consists entirely of  
440 frozen peat that appears to be ice-rich, with a number of ice bands, ice lenses, and ice inclusions.  
441 This evidence combined with the low EIF values and the flat-line ground temperature data



442 suggest high unfrozen water content associated with degrading permafrost on the Kenai  
443 Peninsula.

#### 444 **5.2 Extent and change in residual permafrost plateaus since the 1950s**

445 While previous reports of permafrost on the Kenai Peninsula exist (Hopkins et al., 1955;  
446 Jorgenson et al., 2008), they were restricted to the wetland complex (Mystery Creek) north of  
447 Sterling (Berg et al., 2009). Information on its dynamics here and elsewhere was lacking. Our  
448 analysis of remotely sensed imagery and field surveys identified residual permafrost plateaus in  
449 three additional wetland complexes where it had not been previously identified (Fig. 1 and Fig.  
450 9) and indicates that the state of permafrost within the Kenai lowlands is highly dynamic. In  
451 1950, forested, permafrost plateau extent accounted for 19 % of the land cover in the 4,810 ha of  
452 wetland complexes analysed in the four change detection study areas. In each of the wetland  
453 areas analysed, permafrost plateaus accounted for more than 10 % of the area in 1950. However,  
454 inferred permafrost extent decreased by 60 % between 1950 and ca. 2010, and its lateral  
455 coverage dropped below 5 % in three of the four study areas (Table 2).

456 The residual permafrost plateaus documented in this study share similar attributes to  
457 features elsewhere in boreal peatlands for which permafrost degradation has been inferred due to  
458 the ease of remotely detecting the conversion from forested permafrost plateau to non-permafrost  
459 herbaceous wetland or waterbody (Jorgenson et al., 2001). Thie (1974) inferred a permafrost  
460 plateau loss rate of 0.47 % yr<sup>-1</sup> between 1800 and 1960 for a 130,000 ha area of southern  
461 Manitoba. In Québec, Canada, a 13 ha peat bog lost 1.80 % yr<sup>-1</sup> between 1957 and 2003 (Payette  
462 et al., 2004). In the Northwest Territories, Canada, Quinton et al. (2011) reported a loss rate of  
463 0.62 % yr<sup>-1</sup> between 1947 and 2008 across a 100 ha study area. In Interior Alaska (Tanana  
464 Flats), Jorgenson et al. (2001) reported a loss rate of 0.76 % yr<sup>-1</sup> for birch forested permafrost



465 plateaus between 1949 and 1995 using a point sampling method within a 260,000 ha wetland  
466 area. Lara et al. (2016) recently updated these numbers for the Tanana Flats by manually  
467 digitizing features with methods similar to ours and demonstrated that birch forest plateaus  
468 decreased at a much slower rate of  $0.12 \text{ \% yr}^{-1}$ , and that black spruce forested permafrost plateau  
469 features appeared to be stable. Thus, the loss rate of  $1.0 \text{ \% yr}^{-1}$  that we report for the 4,810 ha  
470 mapped on the western Kenai Peninsula Lowlands between 1950 and ca. 2010 are the second  
471 fastest change rates reported thus far in boreal peatlands.

### 472 **5.3 Drivers of permafrost loss**

473 Permafrost on the Kenai Peninsula is likely degrading as a result of warming air temperatures  
474 ( $+0.4 \text{ }^\circ\text{C decade}^{-1}$  since 1950), especially where warming during the winter season likely  
475 exacerbates these effects (Table 4). During our three-year observation period as well as since the  
476 1950s, warming in the winter has been more pronounced than in the summer (Table 1 and Fig. 2)  
477 and 2014 and 2015 had a MAAT roughly double the 1981 to 2010 climate normal period. Storm  
478 systems regularly bring warm air masses ( $> 4 \text{ }^\circ\text{C}$ ) to the region during the winter. Air  
479 temperature warming during the winter months has decreased the number of freezing degree  
480 days which means that the ground freezes to a much lesser degree in the winter (Fig. 2, Table 1).  
481 Therefore ground temperatures decreased less over the winter period (Table 1 and Fig. 5),  
482 potentially leading to talik development. Previous research on permafrost plateaus in colder  
483 regions indicate that preferential warming in the winter and increased snow accumulation leads  
484 to enhanced permafrost thaw in boreal peatlands (Camill, 2005; Osterkamp, 2007). Since the  
485 Kenai Peninsula lowlands experience a semi-continental climate due to the rain shadow  
486 produced by the Kenai Mountains, a lack of winter snow fall may have contributed to permafrost  
487 persistence in this region by allowing relatively cold winter air temperatures to propagate into the



488 sub-surface. Thus, talik formation and permafrost degradation at our study sites in southcentral  
489 Alaska are likely being driven for the most part by winter fire  
490 air temperature warming (Fig. 2).

491 The increase in permafrost loss rate in southcentral Alaska following the 1980s is likely  
492 due to the combined effects of forest fires and a shift in the PDO after 1976. The respective  
493 pulse and press disturbances may have promoted large areas of permafrost already close to  
494 thawing, to quickly thaw, leaving only colder permafrost and permafrost with intact peat and  
495 forest cover. Fire can be an important driver of permafrost thaw (Yoshikawa et al., 2002) and  
496 thermokarst development (Jones et al., 2015). The Kenai Fire of 1947 burned the majority of the  
497 Mystery Creek study area, all of the Watson Lake study area, and the majority of the Browns  
498 Lake study area. We saw evidence of this fire at numerous sites within the Watson Lake and  
499 Browns Lake study areas. Watson Lake and Browns Lake subsequently had the two greatest loss  
500 rates between 1950 and 1984 and may be related to the 1947 fire. However, the presence of  
501 black spruce burn poles were not found on all permafrost plateaus visited indicating that the  
502 burning was likely relatively patchy in the wetlands. At Browns Lake, permafrost islands that  
503 did not burn in 1947 exhibited less degradation, had thicker permafrost, denser tree cover, and  
504 larger trees than the islands that burned. Large portions of the Tustumena Lake study area  
505 burned in the 1996 Crooked Creek Fire and 2005 Fox Creek Fire. These fires likely damaged,  
506 and partially removed the protective ecosystem cover (black spruce forest and peat), and  
507 degraded several permafrost plateau features. This resulted in the Tustumena study area having  
508 the highest change rate for the latter time period and 77 % of the plateau loss that occurred  
509 between 1996 and ca. 2010 study area did so in areas that burned in the 1996 and 2005 fires.



510 We documented bottom-up permafrost degradation over the short period of direct  
511 measurements between 2012 and 2015. The bottom-up permafrost thaw observed at the Watson  
512 Lake PF2 site indicates that the flow of groundwater below the permafrost plateaus could be  
513 responsible for degradation (Walters et al., 1998). In addition, analysis of the remotely sensed  
514 imagery for the four select wetland complexes primarily documented lateral permafrost  
515 degradation since the 1950s as inferred by the conversion of forested plateau margins to  
516 herbaceous wetland vegetation. This type of feature loss accounted for 85% of the change  
517 between 1950 and ca. 2010. This pattern of loss was further observed in the field through the  
518 presence of thermokarst moats and drowning black spruce trees along the margins of the  
519 permafrost plateaus (Fig. 3). This is similar to the dominant processes documented in more  
520 northerly boreal peatlands with permafrost plateaus (Thie, 1974; Camill and Clark, 1998;  
521 Osterkamp et al., 2000; Jorgenson et al., 2001; Payette et al., 2004; Quinton et al., 2011; Lara et  
522 al., 2015). These findings highlight the importance of groundwater flow and also the impact of  
523 saturated herbaceous wetlands that absorb heat during the summer that likely degrades  
524 permafrost along the peat plateau margins (Walters et al., 1998). It is possible that lateral  
525 permafrost degradation caused by these processes are overwhelming the protection provided by  
526 the ecosystem cover for permafrost stability on the Kenai Peninsula lowlands. Future research is  
527 required to more fully understand the role of groundwater movement on permafrost instability in  
528 the study region.

#### 529 **5.4 Proposed history of permafrost on the Kenai Peninsula**

530 During the Last Glacial Maximum (LGM), northern hemisphere permafrost extended much  
531 further south than present day (Lindgren et al., 2015). However, permafrost history in  
532 southcentral Alaska is poorly constrained. Even though the western Kenai Peninsula lowlands



533 were almost completely glaciated during the LGM (Reger et al., 2007), the permafrost features  
534 identified in this study occur in glaciolacustrine or glaciofluvial wetland complexes that were  
535 either not glaciated during the LGM (Mystery Creek) or became deglaciated before 16,000 cal  
536 yrs BP (Reger et al., 2007). Perhaps permafrost formed on the Kenai Peninsula during  
537 deglaciation or shortly thereafter during the Younger Dryas 12,900 to 11,700 years ago (Jones et  
538 al., 2009). However, this permafrost would have likely thawed during the Holocene Thermal  
539 Maximum (Zoltai, 1972; Kaufman et al., 2004). As the regional climate became cooler and  
540 wetter, between 8,000 and 5,000 years ago, *Sphagnum* accumulation and preservation on the  
541 western Kenai Peninsula lowlands may have promoted more widespread permafrost aggradation  
542 (Jones et al., 2009). Following this period, the peatlands may have progressively froze, heaving  
543 the permafrost plateaus above the water table, drying the peat-rich soils, promoting growth of  
544 black spruce, and creating a buffer layer protecting the underlying permafrost (ecosystem-  
545 protected) from the unfavourable climate for permafrost that currently exists today (Zoltai, 1972,  
546 1995; Payette et al., 2004; Camill, 2005). Growth of permafrost and heaving the peatland  
547 surface above the water table could explain low peat accumulation rates calculated in many  
548 Kenai Peninsula peatlands between 3,300-2,000 years ago (Jones and Yu, 2010; Jones et al.,  
549 2014). This also coincides with widespread neoglaciation on the Kenai Peninsula 3,000 to 1,500  
550 years ago (Wiles and Calkin, 1994, Barclay et al., 2009). Alternatively, the Little Ice Age (365 -  
551 165 years ago), promoted shallow permafrost formation in areas that were predominantly  
552 unfrozen throughout the Holocene (Romanovsky et al., 1992; Jorgenson et al., 2001), and thus,  
553 could account for the presence of residual permafrost on the Kenai Peninsula. The widespread  
554 loss of permafrost plateaus in central Alaska may be a result of degradation of Little Ice Age





555 permafrost (Jorgenson et al., 2001). The age, history, and future trajectory of permafrost on the  
556 western Kenai Peninsula lowlands require further study.

### 557 **5.5 Landscape dynamics and permafrost thaw on the western Kenai Peninsula lowlands**

558 Previous and ongoing land cover change on the western Kenai Peninsula lowlands are primarily  
559 in response to the interaction of climate change and human development. Increases in summer  
560 air temperature and late-summer droughts, along with human disturbance, have been linked to  
561 the massive spruce bark beetle (*Dendroctonus rufipennis*) outbreak of the late 1990s (Berg et al.,  
562 2006; Sherriff et al., 2011), which led to subsequent timber salvage (Jones, 2008). Berg and  
563 Anderson (2006) caution that overall drier conditions on the western Kenai Peninsula, combined  
564 with standing dead spruce stands, may alter the future fire regime of this region. Wetland drying  
565 (Klein et al., 2005) and establishment of woody vegetation in wetlands (Berg et al., 2009) may  
566 be attributed to warmer air temperatures and decreases in precipitation. Furthermore, tectonic  
567 activity associated with the Great Alaska Earthquake of 1964 caused the western Kenai  
568 Peninsula to lower in elevation by 0.7 to 2.3 m (Plafker, 1969), while the northern portion of the  
569 peninsula subsequently uplifted 0.8 – 0.9 m (Cohen and Freymueller, 1997), potentially altering  
570 groundwater flow paths (Gracz, 2011).

571 In our study, we document and incorporate the loss of ecosystem-protected permafrost in  
572 the overall understanding of landscape dynamics on the western Kenai Peninsula lowlands. The  
573 degradation of permafrost can impact terrestrial and aquatic ecosystems, hydrology,  
574 infrastructure, and carbon cycling on the Kenai Peninsula (Schuur et al., 2008; Grosse et al.,  
575 2011; Jorgenson et al., 2013; Kokelj et al., 2015; Vonk et al., 2015). Permafrost degradation  
576 within the wetlands is responsible for a shift from black spruce forest plateaus to fen and bog  
577 wetland ecosystems at a mean rate of 9.2 ha yr<sup>-1</sup> since the 1950s in the four change detection



578 study areas. Permafrost plateaus redirect surface and near-surface drainage in boreal wetlands  
579 (Quinton et al., 2011), and the thaw subsidence of these features increases drainage network  
580 connectivity (Beilman and Robinson, 2003), and alters the local hydrological cycle (Hayashi et  
581 al., 2007). Thus, the loss of permafrost and/or changes in seasonally frozen ground phenology  
582 could in part be aiding in observations of terrestrial and aquatic changes that have occurred on  
583 the Kenai Peninsula during the past several decades. Further work is required to better  
584 understand the past influence of permafrost on the Kenai Peninsula as well as the future loss of  
585 these warm permafrost deposits.

## 586 **6 Conclusions**

587 Based on our ground data and remotely sensed observations, we found that peatland permafrost  
588 is currently more extensive than previously reported in southcentral Alaska, a region with a  
589 MAAT of 1.5 °C. Warm permafrost (-0.04 to -0.08 °C) persists on the western Kenai Peninsula  
590 lowlands in forested (black spruce), peat plateaus found in glaciolacustrine and glaciofluvial  
591 wetland complexes. At our field study sites, the depth to permafrost table on the peat plateaus  
592 averaged 1.48 m in September 2015, but was as shallow as 0.53 m. Permafrost thickness ranged  
593 from 0.33 m to greater than 6.90 m. Field surveys conducted in February 2016 documented the  
594 presence of a surface talik overlying the permafrost table. In 1950, residual permafrost plateaus  
595 covered 19 % of the 4,810 ha wetland area mapped in our study. Within our changed detection  
596 study areas, 60 % of the permafrost plateaus present in 1950 had degraded by ca. 2010. In most  
597 cases, permafrost degradation equated to the loss of forest and its replacement by bog or fen  
598 vegetation, preferentially occurring along permafrost plateau margins. Permafrost loss on the  
599 Kenai Peninsula is likely associated with a warming climate, particularly during the winter



600 season, wildfires that remove the protective ecosystem cover, groundwater flow at depth, and  
601 lateral heat transfer from wetland surface waters in the summer. Future studies on the residual  
602 permafrost plateaus on the Kenai Peninsula will provide further insight for mapping and  
603 predicting permafrost extent across Boreal permafrost regions that are currently warming.

#### 604 **7 Data availability**

605 All data available upon request to the corresponding author.

#### 606 **8 Author contribution**

607 B.M. Jones devised the study design and prepared the manuscript with contributions from all co-  
608 authors. B.M. Jones, C.A. Baughman, V.E. Romanovsky, E.L. Babcock, A.D. Parsekian, M.C.  
609 Jones, and E.E. Berg contributed to field instrumentation and field studies. B.M. Jones, C.A.  
610 Baughman, and G. Grosse conducted and contributed to remote sensing analysis. C.A.  
611 Baughman compiled and interpolated regional weather and climate station data. All co-authors  
612 contributed substantially to this research.

#### 613 **9 Acknowledgements**

614 Funding for this research was provided by the U.S. Geological Survey Land Change Science and  
615 Land Remote Sensing programs. We thank the Kenai National Wildlife Refuge for granting  
616 permission to access field sites. We thank Kelly Harrell, Kobuk, Kashi, Lydia Zeglin, Josefina  
617 Lenz, Emiline Ostlind, and Callie Zuck for help with fieldwork. We thank David Swanson, Eric  
618 Klein, and Neal Pastick for providing useful feedback on an earlier version of this paper. Any



619 use of trade, product, or firm names is for descriptive purposes only and does not imply  
620 endorsement by the U.S. Government.

## 621 **10 References**

- 622 Allard M., Seguin M. K., and Levesque R.: Palsas and mineral permafrost mounds in northern  
623 Quebec, In *International Geomorphology, Part II*, Gardiner V (ed). John Wiley and Sons  
624 Ltd: Chichester; 285–309, 1986.
- 625 Anderson, R. S., Hallett, D. J., Berg, E., Jass, R. B., Toney, J. L., Fontaine, C. S. de and  
626 DeVolder, A.: Holocene development of Boreal forests and fire regimes on the Kenai  
627 Lowlands of Alaska, *The Holocene*, 16(6), 791–803, doi:10.1191/0959683606hol966rp,  
628 2006.
- 629 Barclay, D.J., Wiles, G.C. and Calkin, P.E.: Holocene glacier fluctuations in Alaska. *Quaternary*  
630 *Science Reviews*, 28, 2034-2048, 2009.
- 631 Beilman, D. W. and Robinson, S. D.: Peatland permafrost thaw and landform type along a  
632 climatic gradient, *Proc 8th Int Conf Permafr. Zurich Switz.* 21–25 July 2003, 61 – 65, 2003.
- 633 Beilman, D. W., Vitt, D. H. and Halsey, L. A.: Localized Permafrost Peatlands in Western  
634 Canada: Definition, Distributions, and Degradation, *Arct. Antarct. Alp. Res.*, 33(1), 70–77,  
635 doi:10.2307/1552279, 2001.
- 636 Berg, E. E. and Anderson, R. S.: Fire history of white and Lutz spruce forests on the Kenai  
637 Peninsula, Alaska, over the last two millennia as determined from soil charcoal, *For. Ecol.*  
638 *Manag.*, 227(3), 275–283, doi:10.1016/j.foreco.2006.02.042, 2006.
- 639 Berg, E. E., David Henry, J., Fastie, C. L., De Volder, A. D. and Matsuoka, S. M.: Spruce beetle  
640 outbreaks on the Kenai Peninsula, Alaska, and Kluane National Park and Reserve, Yukon  
641 Territory: Relationship to summer temperatures and regional differences in disturbance  
642 regimes, *For. Ecol. Manag.*, 227(3), 219–232, doi:10.1016/j.foreco.2006.02.038, 2006.
- 643 Berg, E. E., Hillman, K. M., Dial, R. and DeRuwe, A.: Recent woody invasion of wetlands on  
644 the Kenai Peninsula Lowlands, south-central Alaska: a major regime shift after 18 000 years  
645 of wet Sphagnum–sedge peat recruitment, *Can. J. For. Res.*, 39(11), 2033–2046,  
646 doi:10.1139/X09-121, 2009.



- 647 Brown, J., Ferrians, O. J., Heginbottom, J. A. and Melnikov, E. S.: Circum-Arctic Map of  
648 Permafrost and Ground Ice Conditions, 1998.
- 649 Brown, R. J. E.: The Distribution of Permafrost and Its Relation to Air Temperature in Canada  
650 and the U.S.S.R., *ARCTIC*, 13(3), 163–177, doi:10.14430/arctic3697, 1960.
- 651 Brown, R. J. E.: Permafrost in Canada: its influence on Northern development, University of  
652 Toronto Press., 1970.
- 653 Cable, W. L., Romanovsky, V. E., and Jorgenson, M. T.: Scaling-up Permafrost Thermal  
654 Measurements in Western Alaska using an Ecotype Approach, *The Cryosphere*  
655 *Discussions*, doi:10.5194/tc-2016-30, 2016.
- 656 Camill, P.: Patterns of boreal permafrost peatland vegetation across environmental gradients  
657 sensitive to climate warming, *Can. J. Bot.*, 77(5), 721–733, doi:10.1139/b99-008, 1999.
- 658 Camill, P.: Permafrost Thaw Accelerates in Boreal Peatlands During Late-20th Century Climate  
659 Warming, *Clim. Change*, 68(1-2), 135–152, doi:10.1007/s10584-005-4785-y, 2005.
- 660 Camill, P. and Clark, J. S.: Climate Change Disequilibrium of Boreal Permafrost Peatlands  
661 Caused by Local Processes, *Am. Nat.*, 151(3), 207–222, doi: 10.1086/286112, 1998.
- 662 Cohen, S.C. and Freymueller, J.T.: Deformation of the Kenai Peninsula, Alaska, *Journal of*  
663 *Geophysical Research: Solid Earth*, 102, 20479-20487. 1997.
- 664 Ferrians, O. J.: Permafrost Map of Alaska, U.S. Geological Survey Miscellaneous Geologic  
665 Investigations Map I-445, scale 1:2,500,000, 1965.
- 666 Gooseff, M. N., Balsler, A., Bowden, W. B. and Jones, J. B.: Effects of Hillslope Thermokarst in  
667 Northern Alaska, *Eos Trans. Am. Geophys. Union*, 90(4), 29–30,  
668 doi:10.1029/2009EO040001, 2009.
- 669 Gracz, M. B.: Comment on “Wetland drying and succession across the Kenai Peninsula  
670 Lowlands, south-central Alaska” Appears in *Can. J. For. Res.* 35: 1931-1941 (2005)., *Can.*  
671 *J. For. Res.*, 41(2), 425–428, doi:10.1139/X10-147, 2011.
- 672 Grosse, G., Harden, J., Turetsky, M., McGuire, A. D., Camill, P., Tarnocai, C., Frohling, S.,  
673 Schuur, E. A. G., Jorgenson, T., Marchenko, S., Romanovsky, V., Wickland, K. P., French,  
674 N., Waldrop, M., Bourgeau-Chavez, L. and Striegl, R. G.: Vulnerability of high-latitude soil  
675 organic carbon in North America to disturbance, *J. Geophys. Res. Biogeosciences*, 116(G4),  
676 G00K06, doi:10.1029/2010JG001507, 2011.



- 677 Hartmann, B., and Wendler, G.: The significance of the 1976 Pacific climate shift in the  
678 climatology of Alaska, *Journal of Climate*, 18(22), 4824-4839, 2005.
- 679 Hayashi, M., Goeller, N., Quinton, W. L. and Wright, N.: A simple heat-conduction method for  
680 simulating the frost-table depth in hydrological models, *Hydrol. Process.*, 21(19), 2610–  
681 2622, doi:10.1002/hyp.6792, 2007.
- 682 Hopkins, D. M., Karlstrom, T. N. V., others, others and others: Permafrost and Ground Water in  
683 Alaska, Geological Survey Professional Paper, U.S. Geological Survey, United States  
684 Government Printing Office., 1955.
- 685 Jafarov, E. E., Romanovsky, V. E., Genet, H., McGuire, A. D. and Marchenko, S. S.: The effects  
686 of fire on the thermal stability of permafrost in lowland and upland black spruce forests of  
687 interior Alaska in a changing climate, *Environ. Res. Lett.*, 8(3), 035030, doi:10.1088/1748-  
688 9326/8/3/035030, 2013.
- 689 Jones, B. M.: Land-cover change on the southern Kenai Peninsula lowlands, Alaska using USGS  
690 land cover trends methodology, *J. Geogr. Reg. Plan.*, 1(4), 068–071, 2008.
- 691 Jones, B. M., Grosse, G., Arp, C. D., Miller, E., Liu, L., Hayes, D. J. and Larsen, C. F.: Recent  
692 Arctic tundra fire initiates widespread thermokarst development, *Sci. Rep.*, 5,  
693 doi:10.1038/srep15865, 2015.
- 694 Jones, M. C., Peteet, D. M., Kurdyla, D. and Guilderson, T.: Climate and vegetation history from  
695 a 14,000-year peatland record, Kenai Peninsula, Alaska, *Quat. Res.*, 72(2), 207–217,  
696 doi:10.1016/j.yqres.2009.04.002, 2009.
- 697 Jones, M.C. and Yu, Z.: Rapid deglacial and early Holocene expansion of peatlands in Alaska,  
698 *Proceedings of the National Academy of Sciences*, 107, 7347-7352, 2010.
- 699 Jones, M.C., Wooller, M. and Peteet, D.M.: A deglacial and Holocene record of climate  
700 variability in south-central Alaska from stable oxygen isotopes and plant macrofossils in  
701 peat, *Quaternary Science Reviews*, 87, 1-11, 2014.
- 702 Jorgenson, M. T., Racine, C. H., Walters, J. C. and Osterkamp, T. E.: Permafrost Degradation  
703 and Ecological Changes Associated with a Warming Climate in Central Alaska, *Clim.*  
704 *Change*, 48(4), 551–579, doi:10.1023/A:1005667424292, 2001.
- 705 Jorgenson, M. T. and Osterkamp, T. E.: Response of boreal ecosystems to varying modes of  
706 permafrost degradation, *Can. J. For. Res.*, 35(9), 2100–2111, doi:10.1139/x05-153, 2005.



- 707 Jorgenson, M.T., Roth, J.E., Schlentner, S.F., Pullman, E.R. and Macander, M.: An ecological  
708 land survey for Fort Richardson, Alaska (No. ERDC/CRREL-TR-03-19). Engineer  
709 Research And Development Center Hanover Cold Regions Research And Engineering Lab,  
710 2003.
- 711 Jorgenson, M. T., Shur, Y. L. and Pullman, E. R.: Abrupt increase in permafrost degradation in  
712 Arctic Alaska, *Geophys. Res. Lett.*, 33(2), L02503, doi:10.1029/2005GL024960, 2006.
- 713 Jorgenson, M. T., Yoshikawa, K., Kanevskiy, M. Z., Shur, Y., Romanovsky, V. E., Marchenko,  
714 S., Grosse, G., Brown, J. and Jones, B. M.: Permafrost characteristics of Alaska, in 9th  
715 International Conference on Permafrost, 2008.
- 716 Jorgenson, M. T., Romanovsky, V., Harden, J., Shur, Y., O'Donnell, J., Schuur, E. A. G.,  
717 Kanevskiy, M. and Marchenko, S.: Resilience and vulnerability of permafrost to climate  
718 change, *Can. J. For. Res.*, 40(7), 1219–1236, doi:10.1139/X10-060, 2010.
- 719 Jorgenson, M. T., Harden, J., Kanevskiy, M., O'Donnell, J., Wickland, K., Stephanie Ewing,  
720 Manies, K., Zhuang, Q., Shur, Y., Striegl, R. and Koch, J.: Reorganization of vegetation,  
721 hydrology and soil carbon after permafrost degradation across heterogeneous boreal  
722 landscapes, *Environ. Res. Lett.*, 8(3), 035017, doi:10.1088/1748-9326/8/3/035017, 2013.
- 723 Kanevskiy, M., Shur, Y., Krzewinski, T. and Dillon, M.: Structure and properties of ice-rich  
724 permafrost near Anchorage, Alaska, *Cold Reg. Sci. Technol.*, 93, 1–11,  
725 doi:10.1016/j.coldregions.2013.05.001, 2013.
- 726 Kaufman, D.S., Ager, T.A., Anderson, N.J., Anderson, P.M., Andrews, J.T., Bartlein, P.J.,  
727 Brubaker, L.B., Coats, L.L., Cwynar, L.C., Duvall, M.L. and Dyke, A.S.: Holocene thermal  
728 maximum in the western Arctic (0–180 W). *Quaternary Science Reviews*, 23, 529-560,  
729 2004.
- 730 Klein, E., Berg, E. E. and Dial, R.: Wetland drying and succession across the Kenai Peninsula  
731 Lowlands, south-central Alaska, *Can. J. For. Res.*, 35(8), 1931–1941, doi:10.1139/x05-129,  
732 2005.
- 733 Kokelj, S. V., Tunnicliffe, J., Lacelle, D., Lantz, T. C., Chin, K. S. and Fraser, R.: Increased  
734 precipitation drives mega slump development and destabilization of ice-rich permafrost  
735 terrain, northwestern Canada, *Glob. Planet. Change*, 129, 56–68,  
736 doi:10.1016/j.gloplacha.2015.02.008, 2015.



- 737 Lachenbruch, A. H. and Marshall, B. V.: Changing Climate: Geothermal Evidence from  
738 Permafrost in the Alaskan Arctic, *Science*, 234(4777), 689–696,  
739 doi:10.1126/science.234.4777.689, 1986.
- 740 Lantz, T. C. and Kokelj, S. V.: Increasing rates of retrogressive thaw slump activity in the  
741 Mackenzie Delta region, N.W.T., Canada, *Geophys. Res. Lett.*, 35(6), L06502,  
742 doi:10.1029/2007GL032433, 2008.
- 743 Lara, M. J., Genet, H., McGuire, A. D., Euskirchen, E. S., Zhang, Y., Brown, D. R. N.,  
744 Jorgenson, M. T., Romanovsky, V., Breen, A. and Bolton, W. R.: Thermokarst rates  
745 intensify due to climate change and forest fragmentation in an Alaskan boreal forest  
746 lowland, *Glob. Change Biol.*, doi:10.1111/gcb.13124, 2015.
- 747 Lewkowicz, A.G., Etzelmüller, B. and Smith, S.L.: Characteristics of discontinuous permafrost  
748 based on ground temperature measurements and electrical resistivity tomography, southern  
749 Yukon, Canada, *Permafrost and Periglacial Processes*, 22(4), pp.320-342, 2011.
- 750 Liljedahl, A.K., Boike, J., Daanen, R.P., Fedorov, A.N., Frost, G.V., Grosse, G., Hinzman, L.D.,  
751 Iijma, Y., Jorgenson, J.C., Matveyeva, N., Necsoiu, M., , Raynolds, M. K., Romanovsky, V.  
752 E., Schulla, J., Tape, K., Walker, D. A., and H. Yabuki: Pan-Arctic ice-wedge degradation  
753 in warming permafrost and its influence on tundra hydrology, *Nature Geoscience*, 2016.
- 754 Lindgren, A., Hugelius, G., Kuhry, P., Christensen, T. R. and Vandenberghe, J.: GIS-based Maps  
755 and Area Estimates of Northern Hemisphere Permafrost Extent during the Last Glacial  
756 Maximum, *Permafr. Periglac. Process.*, n/a–n/a, doi:10.1002/ppp.1851, 2015.
- 757 Lunardini, V.J.: Climatic warming and the degradation of warm permafrost. *Permafrost and*  
758 *Periglacial Processes*, 7(4), 311-320, 1996.
- 759 Lynch, J. A., Clark, J. S., Bigelow, N. H., Edwards, M. E. and Finney, B. P.: Geographic and  
760 temporal variations in fire history in boreal ecosystems of Alaska, *J. Geophys. Res.*  
761 *Atmospheres*, 107(D1), 8152, doi:10.1029/2001JD000332, 2002.
- 762 Morse, P. D., Wolfe, S. A., Kokelj, S. V. and Gaanderse, A. J. R.: The Occurrence and Thermal  
763 Disequilibrium State of Permafrost in Forest Ecotopes of the Great Slave Region, Northwest  
764 Territories, Canada, *Permafr. Periglac. Process.*, doi:10.1002/ppp.1858, 2015.
- 765 Neal, A.: Ground-penetrating radar and its use in sedimentology: principles, problems and  
766 progress, *Earth-science reviews*, 66, 261-330, 2004.





- 767 Nelson, F. E., Anisimov, O. A. and Shiklomanov, N. I.: Subsidence risk from thawing  
768 permafrost, *Nature*, 410(6831), 889–890, doi:10.1038/35073746, 2001.
- 769 O'Donnell, J. A., Jorgenson, M. T., Harden, J. W., McGuire, A. D., Kanevskiy, M. Z. and  
770 Wickland, K. P.: The Effects of Permafrost Thaw on Soil Hydrologic, Thermal, and Carbon  
771 Dynamics in an Alaskan Peatland, *Ecosystems*, 15(2), 213–229, doi:10.1007/s10021-011-  
772 9504-0, 2012.
- 773 Osterkamp, T. E.: Characteristics of the recent warming of permafrost in Alaska, *J. Geophys.*  
774 *Res. Earth Surf.*, 112(F2), F02S02, doi:10.1029/2006JF000578, 2007.
- 775 Osterkamp, T. E., Viereck, L., Shur, Y., Jorgenson, M. T., Racine, C., Doyle, A. and Boone, R.  
776 D.: Observations of Thermokarst and Its Impact on Boreal Forests in Alaska, U.S.A., *Arct.*  
777 *Antarct. Alp. Res.*, 32(3), 303–315, doi:10.2307/1552529, 2000.
- 778 Parsekian, A. D., Slater, L., Comas, X., and Glaser, P. H.: Variations in free-phase gases in peat  
779 landforms determined by ground-penetrating radar, *Journal of Geophysical Research:*  
780 *Biogeosciences*, 115, no. G2, 2010.
- 781 Parsekian, A.D., Grosse, G., Walbrecker, J.O., Müller-Petke, M., Keating, K., Liu, L., Jones,  
782 B.M. and Knight, R.: Detecting unfrozen sediments below thermokarst lakes with surface  
783 nuclear magnetic resonance, *Geophysical Research Letters*, 40(3), 535-540, 2013.
- 784 Pastick, N. J., Jorgenson, M. T., Wylie, B. K., Nield, S. J., Johnson, K. D. and Finley, A. O.:  
785 Distribution of near-surface permafrost in Alaska: Estimates of present and future  
786 conditions, *Remote Sens. Environ.*, 168, 301–315, doi:10.1016/j.rse.2015.07.019, 2015.
- 787 Payette, S., Delwaide, A., Caccianiga, M. and Beauchemin, M.: Accelerated thawing of subarctic  
788 peatland permafrost over the last 50 years, *Geophys. Res. Lett.*, 31(18), L18208,  
789 doi:10.1029/2004GL020358, 2004.
- 790 Plafker, G.: Tectonics of the March 27, 1964 Alaska earthquake: U.S. Geological Survey  
791 Professional Paper. [online] Available from: <http://pubs.usgs.gov/pp/0543i/>, 1969.
- 792 Quinton, W. I., Hayashi, M. and Chasmer, L. e.: Permafrost-thaw-induced land-cover change in  
793 the Canadian subarctic: implications for water resources, *Hydrol. Process.*, 25(1), 152–158,  
794 doi:10.1002/hyp.7894, 2011.
- 795 Reger, R. D., Sturm, A. G., Berg, E. E. and Burns, P. A. C.: A guide to the late Quaternary  
796 history of northern and western Kenai Peninsula, Alaska: Alaska Division of Geological &  
797 Geophysical Surveys Guidebook 8, 2007.



- 798 Romanovsky, V. E., Garagula, L. S., and Seregina, N. V.: Freezing and thawing of soils under  
799 the influence of 300- and 90-year periods of temperature fluctuation. Pages 543–548 in  
800 Proceedings of the International Conference on the Role of Polar Regions in Global Change.  
801 Fairbanks (AK): Geophysical Institute, University of Alaska. 1992.
- 802 Romanovsky, V. E. and Osterkamp, T. E.: Effects of unfrozen water on heat and mass transport  
803 processes in the active layer and permafrost, *Permafr. Periglac. Process.*, 11(3), 219–239,  
804 doi:10.1002/1099-1530(200007/09)11:3<219::AID-PPP352>3.0.CO;2-7, 2000.
- 805 Romanovsky, V. E. and Osterkamp, T. E. (1995), Interannual variations of the thermal regime of  
806 the active layer and near-surface permafrost in northern Alaska. *Permafrost Periglac.*  
807 *Process.*, 6: 313–335. doi: 10.1002/ppp.3430060404
- 808 Romanovsky, V., Burgess, M., Smith, S., Yoshikawa, K. and Brown, J.: Permafrost temperature  
809 records: Indicators of climate change, *Eos Trans. Am. Geophys. Union*, 83(50), 589–594,  
810 doi:10.1029/2002EO000402, 2002.
- 811 Romanovsky, V. E., Smith, S. L. and Christiansen, H. H.: Permafrost thermal state in the polar  
812 Northern Hemisphere during the international polar year 2007–2009: a synthesis, *Permafr.*  
813 *Periglac. Process.*, 21(2), 106–116, doi:10.1002/ppp.689, 2010.
- 814 Sandmeier, K. J.: REFLEXW – Windows™ 9x/NT/2000/XP-program for the processing of  
815 seismic, acoustic or electromagnetic reflection, refraction and transmission data, 2008.
- 816 Sannel, A. B. K., Hugelius, G., Jansson, P. and Kuhry, P.: Permafrost Warming in a Subarctic  
817 Peatland – Which Meteorological Controls are Most Important?, *Permafr. Periglac.*  
818 *Process.*, n/a–n/a, doi:10.1002/ppp.1862, 2015.
- 819 Schuur, E. A. G., Bockheim, J., Canadell, J. G., Euskirchen, E., Field, C. B., Goryachkin, S. V.,  
820 Hagemann, S., Kuhry, P., Lafleur, P. M., Lee, H., Mazhitova, G., Nelson, F. E., Rinke, A.,  
821 Romanovsky, V. E., Shiklomanov, N., Tarnocai, C., Venevsky, S., Vogel, J. G. and Zimov,  
822 S. A.: Vulnerability of Permafrost Carbon to Climate Change: Implications for the Global  
823 Carbon Cycle, *BioScience*, 58(8), 701–714, doi:10.1641/B580807, 2008.
- 824 Sherriff, R. L., Berg, E. E. and Miller, A. E.: Climate variability and spruce beetle  
825 (*Dendroctonus rufipennis*) outbreaks in south-central and southwest Alaska, *Ecology*, 92(7),  
826 1459–1470, doi:10.1890/10-1118.1, 2011.



- 827 Shur, Y. L. and Jorgenson, M. T.: Patterns of permafrost formation and degradation in relation to  
828 climate and ecosystems, *Permafr. Periglac. Process.*, 18(1), 7–19, doi:10.1002/ppp.582,  
829 2007.
- 830 Smith, M. W. and Riseborough, D. W.: Climate and the limits of permafrost: a zonal analysis,  
831 *Permafr. Periglac. Process.*, 13(1), 1–15, doi:10.1002/ppp.410, 2002.
- 832 Swindles, G.T., Morris, P.J., Mullan, D., Watson, E.J., Turner, T.E., Roland, T.P., Amesbury,  
833 M.J., Kokfelt, U., Schoning, K., Pratte, S. and Gallego-Sala, A.: The long-term fate of  
834 permafrost peatlands under rapid climate warming, *Scientific Reports*, 5, 17951,  
835 10.1038/srep17951, 2015.
- 836 Thie, J.: Distribution and Thawing of Permafrost in the Southern Part of the Discontinuous  
837 Permafrost Zone in Manitoba, *ARCTIC*, 27(3), 1974.
- 838 Turetsky, M. R., Wieder, R. K., Vitt, D. H., Evans, R. J. and Scott, K. D.: The disappearance of  
839 relict permafrost in boreal North America: Effects on peatland carbon storage and fluxes,  
840 *Glob. Change Biol.*, 13(9), 1922–1934, doi:10.1111/j.1365-2486.2007.01381.x, 2007.
- 841 Van Everdingen, R. O.: Multi-Language Glossary of Permafrost and Related Ground-Ice Terms  
842 in Chinese, English, French, German, Icelandic, Italian, Norwegian, Polish, Romanian,  
843 Russian, Spanish, and Swedish. International Permafrost Association, Terminology  
844 Working Group, 1998.
- 845 Vonk, J. E., Tank, S. E., Bowden, W. B., Laurion, I., Vincent, W. F., Alekseychik, P., Amyot,  
846 M., Billet, M. F., Canário, J., Cory, R. M., Deshpande, B. N., Helbig, M., Jammet, M.,  
847 Karlsson, J., Larouche, J., MacMillan, G., Rautio, M., Walter Anthony, K. M. and  
848 Wickland, K. P.: Reviews and syntheses: Effects of permafrost thaw on Arctic aquatic  
849 ecosystems, *Biogeosciences*, 12(23), 7129–7167, doi:10.5194/bg-12-7129-2015, 2015.
- 850 Walters J.C., Racine C.H., and Jorgenson M.T.: Characteristics of permafrost in the Tanana  
851 Flats, interior Alaska, In: *Proceedings of the Seventh International Conference on*  
852 *Permafrost Vol 57* (eds. Lewkowicz AG, Allard M), Université Laval, Québec. Collection  
853 *Nordicana*, 1109-1116, 1998.
- 854 Wiles, G. C. and Calkin, P. E.: Late Holocene, high-resolution glacial chronologies and climate,  
855 Kenai Mountains, Alaska, *Geol. Soc. Am. Bull.*, 106(2), 281–303, doi:10.1130/0016-  
856 7606(1994)106<0281:LHHRGC>2.3.CO;2, 1994.



- 857 Yoshikawa, K., Bolton, W. R., Romanovsky, V. E., Fukuda, M. and Hinzman, L. D.: Impacts of  
858 wildfire on the permafrost in the boreal forests of Interior Alaska, *J. Geophys. Res.*  
859 *Atmospheres*, 107(D1), 8148, doi:10.1029/2001JD000438, 2002.
- 860 Zoltai, S. C.: Palsas and Peat Plateaus in Central Manitoba and Saskatchewan, *Can. J. For. Res.*,  
861 2(3), 291–302, doi:10.1139/x72-046, 1972.
- 862 Zoltai, S. C.: Permafrost Distribution in Peatlands of West-Central Canada During the Holocene  
863 Warm Period 6000 Years BP, *Géographie Phys. Quat.*, 49(1), 45, doi:10.7202/033029ar,  
864 1995.



865 **Tables**

866 Table 1. Mean annual ground temperature data for four observation sites on the Kenai Peninsula lowlands. Browns Lake PF1,  
 867 Watson Lake PF1, and Watson Lake PF2 represent permafrost plateaus and the Watson Lake non-PF site a black spruce forested non-  
 868 plateau site. Sensor depths that were perennially frozen in a given year are in bold. Mean annual air temperature (MAAT), thawing  
 869 and freezing degree days (TDD and FDD), and average winter snow depth (MASD) are from the Kenai 29 ENE AWS 702590 weather  
 870 station.  
 871

9/16/2012 - 9/15/2013											
Browns Lake PF1		Watson Lake PF1		Watson Lake PF2		Watson Lake non-PF		KENAI 29 ENE AWS 702590 Met Station Data			
Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	MAAT (°C)	TDD Sums	FDD Sums	MASD (cm)
50	-0.02	10	0.34	10	0.05	25	0.94				
<b>100</b>	<b>-0.06</b>	25	-0.09	50	-0.30	50	0.42	0.88	1865.9	1544.3	19.3
<b>200</b>	<b>-0.08</b>	50	-0.20	<b>100</b>	<b>-0.08</b>	100	0.14				
<b>300</b>	<b>-0.08</b>	<b>100</b>	<b>-0.08</b>	<b>200</b>	<b>-0.06</b>	130	0.16				

9/16/2013 - 9/15/2014											
Browns Lake PF1		Watson Lake PF1		Watson Lake PF2		Watson Lake non-PF		KENAI 29 ENE AWS 702590 Met Station Data			
Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	MAAT (°C)	TDD Sums	FDD Sums	MASD (cm)
50*	0.17	10	0.93	10	0.86	25	0.57				
<b>100</b>	<b>-0.06</b>	25	0.40	50	-0.07	50	0.32	2.58	2066.6	1123.4	8.3
<b>200</b>	<b>-0.06</b>	50	-0.02	<b>100</b>	<b>-0.08</b>	100	0.14				
<b>300</b>	<b>-0.08</b>	<b>100</b>	<b>-0.06</b>	<b>200</b>	<b>-0.08</b>	130	0.14				

\*Thermistor failed on 24 August 2014

9/16/2014 - 9/15/2015											
Browns Lake PF1		Watson Lake PF1		Watson Lake PF2		Watson Lake non-PF		KENAI 29 ENE AWS 702590 Met Station Data			
Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	Sensor depth (cm)	MAGT (°C)	MAAT (°C)	TDD Sums	FDD Sums	MASD (cm)
50*	---	10*	---	10	1.53	25*	---				
<b>100</b>	<b>-0.04</b>	25*	---	50	0.14	50*	---	3.24	2009.8	829.1	2.7
<b>200</b>	<b>-0.06</b>	50*	---	<b>100</b>	<b>-0.07</b>	100*	---				
<b>300</b>	<b>-0.07</b>	100**	---	200 <sup>#</sup>	-0.07	130*	---				

\*Thermistor or data logger failure

<sup>#</sup>Permafrost thaw during observation period

872



873 Table 2. Permafrost plateau extent mapped in each study region in 1950, 1984, 1996, and ca. 2010. Analyzed wetland area for each  
 874 study region is given along with the number of features, total plateau area, mean plateau area, and plateau extent for each image  
 875 observation year. In ca. 2010, images were acquired in 2011 (Mystery Creek and Watson Lake), 2012 (Tustumena Lake), and 2013  
 876 (Browns Lake).  
 877

Study Region	Wetland Area (ha)	1950				1984				1996				ca. 2010			
		Number of Features	Total Plateau Area (ha)	Mean Plateau Area (ha)	Plateau Extent (%)	Number of Features	Total Plateau Area (ha)	Mean Plateau Area (ha)	Plateau Extent (%)	Number of Features	Total Plateau Area (ha)	Mean Plateau Area (ha)	Plateau Extent (%)	Number of Features	Total Plateau Area (ha)	Mean Plateau Area (ha)	Plateau Extent (%)
Mystery Creek	1562.0	212	511.5	2.4	32.7	237	457.7	1.9	29.3	335	292.6	0.9	18.7	321	232.3	0.7	14.9
Watson Lake	904.2	44	86.6	2.0	9.6	55	54.0	1.0	6.0	68	35.4	0.5	3.9	67	29.8	0.4	3.3
Browns Lake	1013.0	102	111.9	1.1	11.0	117	67.2	0.6	6.6	107	51.2	0.5	5.1	89	38.6	0.4	3.8
Tustumena Lake	1333.4	92	210.2	2.2	15.8	150	168.6	1.1	12.6	183	143.5	0.8	10.8	206	69.9	0.3	5.2
All Sites	4812.7	450	920.2	2.0	19.1	559	747.5	1.3	15.5	693	522.6	0.8	10.9	683	370.6	0.5	7.7

878  
 879



880 Table 3. Change in the extent of permafrost plateaus for each of the study regions between 1950  
 881 and ca. 2010, 1950 and 1984, 1984 and 1996, and 1996 and ca. 2010. Change is reported in  
 882 aerial units per year, proportional area change, percent change per year, and by the type of  
 883 change. Change type refers to whether the plateau loss occurred along the periphery of a feature  
 884 (lateral), in the centre of a feature (internal), or whether complete loss of a feature occurred. In  
 885 ca. 2010, images were acquired in 2011 (Mystery Creek and Watson Lake), 2012 (Tustumena  
 886 Lake), and 2013 (Browns Lake).  
 887

Study Area	1950 to ca. 2010			Change Type		
	Area Change	Proportional Area	Percent Change	Lateral (%)	Internal (%)	Complete (%)
	(ha yr <sup>-1</sup> )	Change (ha yr <sup>-1</sup> 100 ha <sup>-1</sup> )	(% yr <sup>-1</sup> )			
Mystery Creek	-4.6	-0.3	-0.9	86.2	3.3	10.5
Watson Lake	-0.9	-0.1	-1.1	90.9	0.8	8.3
Browns Lake	-1.2	-0.1	-1.0	77.2	0.3	22.4
Tustumena Lake	-2.3	-0.2	-1.1	86.0	1.7	12.3
All Sites	-9.2	-0.2	-1.0	85.1	1.5	13.4

Study Area	1950 to 1984			Change Type		
	Area Change	Proportional Area	Percent Change	Lateral (%)	Internal (%)	Complete (%)
	(ha yr <sup>-1</sup> )	Change (ha yr <sup>-1</sup> 100 ha <sup>-1</sup> )	(% yr <sup>-1</sup> )			
Mystery Creek	-1.6	-0.1	-0.3	88.8	5.2	5.9
Watson Lake	-1.0	-0.1	-1.1	91.7	1.4	6.9
Browns Lake	-1.3	-0.1	-1.2	89.0	0.6	10.1
Tustumena Lake	-1.2	-0.1	-0.6	94.1	2.1	3.8
All Sites	-5.1	-0.1	-0.6	91.0	2.3	6.7

Study Area	1984 to 1996			Change Type		
	Area Change	Proportional Area	Percent Change	Lateral (%)	Internal (%)	Complete (%)
	(ha yr <sup>-1</sup> )	Change (ha yr <sup>-1</sup> 100 ha <sup>-1</sup> )	(% yr <sup>-1</sup> )			
Mystery Creek	-13.8	-0.9	-3.0	87.1	1.8	11.2
Watson Lake	-1.6	-0.2	-2.9	88.7	0.6	10.7
Browns Lake	-1.3	-0.1	-2.0	84.0	0.1	16.0
Tustumena Lake	-2.1	-0.2	-1.2	85.1	2.9	12.0
All Sites	-18.7	-0.4	-2.5	86.2	1.3	12.5

Study Area	1996 to ca. 2010			Change Type		
	Area Change	Proportional Area	Percent Change	Lateral (%)	Internal (%)	Complete (%)
	(ha yr <sup>-1</sup> )	Change (ha yr <sup>-1</sup> 100 ha <sup>-1</sup> )	(% yr <sup>-1</sup> )			
Mystery Creek	-4.0	-0.3	-1.4	82.7	3.0	14.3
Watson Lake	-0.4	-0.1	-1.1	92.2	0.5	7.3
Browns Lake	-0.7	-0.1	-1.4	58.7	0.1	41.2
Tustumena Lake	-4.6	-0.3	-3.2	78.7	0.2	21.1
All Sites	-9.5	-0.2	-1.8	78.1	1.0	21.0

888



889 Table 4. Mean annual, mean summer (May to September), and mean winter (October to April)  
 890 air temperature for the three remotely sensed image observation periods compiled from the  
 891 Kenai Municipal Airport (WBAN 26523) and estimated from the MRC station (Kenai 29 ENE -  
 892 AWS 702590).  
 893


Remote Sensing Time Period	Mean Annual Air Temperature (°C)		Mean Summer Air Temperature (°C)		Mean Winter Air Temperature (°C)	
	Kenai Airport	MRC	Kenai Airport	MRC	Kenai Airport	MRC
1950 to 1984	1.12	0.59	9.92	10.39	-5.29	-6.54
1984 to 1996	1.77	1.31	10.28	10.78	-4.37	-5.52
1996 to 2015	2.34	1.86	10.81	11.31	-3.77	-4.95

894






895 **Figures**



896

897 Figure 1: Study area figure. (a) Recent permafrost map of Alaska (Pastick et al., 2015)  
898 indicating permafrost presence (purple) and absence (white) in the upper one meter of the ground  
899 surface. (b) Hillshade relief image showing a portion of southcentral Alaska. The study region  
900 on the Kenai Peninsula lowlands is shown with the black box outline. (c) The portion of the  
901 Kenai Peninsula lowlands where field studies and remotely sensed observations were conducted.  
902 Ground temperature observations were collected at the Browns Lake and Watson Lake sites.  
903 The remote sensing change detection areas are shown with a white box: (A) Mystery Creek, (B)  
904 Watson Lake, (C) Browns Lake, and (D) Tustumena Lake wetland complexes.



905

906 Figure 2: a) Historical (1948-2015) mean annual air temperature compiled from Kenai  
907 Municipal Airport (WBAN 26523) hourly surface data and interpolated (broken) and measured  
908 (solid) mean annual air temperature for the MRC station (Kenai 29 ENE AWS 702590). b)  
909 Thawing degree day (TDD) and freezing degree day (FDD) sums for 1948-2015 derived from  
910 historical and interpolated daily mean temperature. c) Cumulative annual precipitation data from  
911 the Kenai Municipal Airport (WBAN 26523) between 1948 and 2015.



912


913

914 Figure 3: Field photos of residual permafrost plateau landforms and thermokarst on the western  
915 Kenai Peninsula lowlands. (a) A forested permafrost plateau in the Browns Lake wetland  
916 complex. A thermokarst moat and drowning black spruce trees in the (b) Browns Lake and (c)  
917 Watson Lake wetland complexes.

918

919






920

921 Figure 4: Time series documenting the extent of a subset of permafrost features in the Browns  
922 Lake wetland complex in (a) 1950, (b) 1984, (c) 1996, and (d) 2013. Permafrost plateau extent  
923 in 1950 is shown as a yellow polygon in each frame and other time slices outlined as green  
924 (1984), blue (1996), and red (2013).




925




926

927 Figure 5: Mean daily ground temperature data plots for the four shallow boreholes on the  
928 western Kenai Peninsula lowlands for the period of 16 September 2012 to 15 September 2015:  
929 (a) Browns Lake PF1 site, (b) Watson Lake PF1 site, (c) Watson Lake PF2 site, and (d) Watson  
930 Lake non-PF site. All axes scales are the same but sensor depths vary among sites based on site  
931 characteristics. Missing data indicates sensor or thermistor failure.  
932



933

934 Figure 6: Mean daily ground temperature data plots indicating the presence of permafrost on the  
935 western Kenai Peninsula lowlands for the period of 16 September 2012 to 15 September 2015:  
936 (a) Browns Lake PF1 site, (b) Watson Lake PF1 site, and (c) Watson Lake PF2 site. Top-down  
937 permafrost thaw occurred at Watson Lake PF1 during the fall of 2014 and bottom-up permafrost  
938 thaw occurred at Watson Lake PF2 during the fall of 2015.



939


940 Figure 7: (a) High-resolution satellite image showing the permafrost plateau in the Browns Lake  
941 wetland complex where detailed field surveys were conducted as well as the location of the  
942 Browns Lake PF1 data logger (green dot). (b) A ~100 m transect across the Browns Lake PF1  
943 permafrost plateau site showing ground surface height above the wetland (green line), depth to  
944 the permafrost table (blue line and blue arrows), permafrost thickness constraints (grey arrows),  
945 seasonally frozen ground depth (maroon arrows), and lack of permafrost (yellow stars) as  
946 measured by probing, drilling, and coring. Locations where the permafrost table exceeded 2.2 m  
947 from the ground surface (limiting depth for September surveys) are indicated with a non-existent  
948 blue line. Locations where the base of the permafrost was encountered are indicated with an  
949 upward looking grey triangle and those locations where it was not encountered, a downward  
950 looking grey triangle.

951






952



953

954 Figure 8. GPR profiles at the intensive Browns Lake permafrost plateau (Fig. 7a) from a) late-  
955 July 2014 with Mala shielded 100-MHz antennas and b) mid-February 2016 with Sensors &  
956 Software unshielded bi-static 100-MHz antennas. Processed radargrams are on the left and  
957 processed, interpreted radargrams are on the right. Both summer and winter profiles clearly  
958 show reflectors associated with the permafrost table and in the case of (b) show the presence of a  
959 talik. However, we were unable to image the permafrost base using GPR. Note that the two  
960 GPR transects differ slightly in their orientation across the feature.  
961





962

963 Figure 9. Spatial and temporal pattern of permafrost loss within four change detection areas: a)  
964 Mystery Creek, b) Watson Lake, c) Browns Lake, and d) Tustumena Lake. Red indicates feature  
965 loss between 1950 and 1980, orange is feature loss between 1984 and 1996, yellow is feature  
966 loss between 1996 and ca. 2010, and green is ca. 2010 permafrost plateau extent. The white dots  
967 indicate the location of field verified permafrost between 2009 and 2016. The hatched white  
968 polygons indicate the wetland extent where plateau features were mapped in each study area.  
969 Background imagery is the 1996 orthophotography.