

# A Short Proof that Lebesgue Outer Measure of an Interval is Its Length

Jitender Singh

Department of Mathematics, Guru Nanak Dev University Amritsar, India

[sonumaths@gmail.com](mailto:sonumaths@gmail.com); [jitender.math@gndu.ac.in](mailto:jitender.math@gndu.ac.in)

MSC: Primary 28A12. The Lebesgue outer measure  $m^*(E)$  of a subset  $E$  of real line is defined as  $m^*(E) := \inf\{\sum_{k=1}^{\infty} \ell(I_k) \mid E \subseteq \cup_{k=1}^{\infty} I_k\}$ , where each  $I_k$  is an open interval and  $\ell(I_k)$  is its length. Establishing one of the inequalities in the standard proof of the fact in the title above turns out to be tedious in [1, p. 31]. Using the connectedness of the interval shortens the proof as follows.

*Proof.* Given two real numbers  $a$  and  $b$  with  $a < b$ , it is enough to prove that  $m^*([a, b]) = b - a$ . Clearly,  $m^*([a, b]) \leq b - a$ . Now let  $[a, b] \subset \cup_{k=1}^n I_k$  for some positive integer  $n$ , which is always possible since  $[a, b]$  is compact. Without loss of generality, assume that the set  $[a, b] \cap I_k$  is nonempty for each  $k$ . Observe that the set  $\cup_{k=1}^n I_k$  is connected. (Otherwise, if  $(P, Q)$  is its separation, then for each  $k$ , by connectedness of  $I_k$ , either  $I_k \subset P$  or  $I_k \subset Q$ . Thus each of  $P$  and  $Q$  is equal to union of sets from the list  $\{I_1, \dots, I_n\}$ . So the pair  $(P \cap [a, b], Q \cap [a, b])$  determines a separation of  $[a, b]$ , which contradicts connectedness of  $[a, b]$ .) So  $\cup_{k=1}^n I_k$  is an open interval containing  $[a, b]$ . Thus,  $b - a \leq \ell(\cup_{k=1}^n I_k) \leq \sum_{k=1}^n \ell(I_k)$ , where the last inequality holds since some intervals overlap<sup>†</sup>. Hence,  $b - a \leq m^*([a, b])$ .  $\square$

<sup>†</sup>The inequality  $\ell(\cup_{k=1}^n I_k) \leq \sum_{k=1}^n \ell(I_k)$  can be justified as follows. Given a bounded interval  $I$ , by definition,  $\ell(I) = \sup I - \inf I$ . Observe that if two distinct bounded intervals  $I_1$  and  $I_2$  overlap, then  $\ell(I_1 \cup I_2) \leq \ell(I_1) + \ell(I_2)$ .

Given  $n > 1$  bounded open intervals  $I_1, \dots, I_n$  with their union being connected implies that for each  $k = 1, \dots, n - 1$  we may choose  $I_k$  after re-indexing these intervals, such that  $\cup_{j=1}^k I_j$  is connected and it overlaps at least one interval among rest of the  $(n - k)$  intervals, which we denote by  $I_{k+1}$ . So, we have  $\ell(\cup_{k=1}^n I_k) = \ell(\cup_{k=1}^{n-1} I_k \cup I_n) \leq \ell(\cup_{k=1}^{n-1} I_k) + \ell(I_n) \leq \ell(\cup_{k=1}^{n-2} I_k) + \ell(I_{n-1}) + \ell(I_n) \leq \dots \leq \ell(I_1) + \dots + \ell(I_n)$ .

## References

- [1] H. L. Royden, P. M. Fitzpatrick, *Real Analysis*. Fourth ed. Pearson, Boston, 2010.