



European Red List of Birds 2021

Compiled by BirdLife International



Manuscript completed in May 2022

The European Commission is not liable for any consequence stemming from the reuse of this publication.

Luxembourg: Publications Office of the European Union, 2022



© European Union, 2022

The reuse policy of European Commission documents is implemented based on Commission Decision 2011/833/EU of 12 December 2011 on the reuse of Commission documents (OJ L 330, 14.12.2011, p. 39).

Except otherwise noted, the reuse of this document is authorised under a Creative Commons Attribution 4.0 International (CC-BY 4.0) licence (<https://creativecommons.org/licenses/by/4.0/>). This means that reuse is allowed provided appropriate credit is given and any changes are indicated.

For any use or reproduction of elements that are not owned by the European Union, permission may need to be sought directly from the respective rightholders.

Print ISBN 978-92-76-45973-6
doi:10.2779/967570
KH-07-21-130-EN-C

PDF ISBN 978-92-76-45974-3
doi:10.2779/959320
KH-07-21-130-EN-N

Getting in touch with the EU

In person

All over the European Union there are hundreds of Europe Direct information centres. You can find the address of the centre nearest you at: https://europa.eu/european-union/contact_en

On the phone or by email

Europe Direct is a service that answers your questions about the European Union. You can contact this service:

- by freephone: 00 800 6 7 8 9 10 11 (certain operators may charge for these calls),
- at the following standard number: +32 22999696 or
- by email via: https://europa.eu/european-union/contact_en

Finding information about the EU

Online

Information about the European Union in all the official languages of the EU is available on the Europa website at: https://europa.eu/european-union/index_en

EU publications

You can download or order free and priced EU publications at: <https://op.europa.eu/en/publications>. Multiple copies of free publications may be obtained by contacting Europe Direct or your local information centre (see https://europa.eu/european-union/contact_en).

EU law and related documents

For access to legal information from the EU, including all EU law since 1952 in all the official language versions, go to EUR-Lex at: <http://eur-lex.europa.eu>

Open data from the EU

The EU Open Data Portal (<http://data.europa.eu/euodp/en>) provides access to datasets from the EU. Data can be downloaded and reused for free, for both commercial and non-commercial purposes.

Published by Stichting BirdLife Europe.

This publication has been prepared by BirdLife International.

The designation of geographical entities in this book, and the presentation of the material, do not imply the expression of any opinion whatsoever on the part of the European Commission or BirdLife International concerning the legal status of any country, territory, or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries. The views expressed in this publication do not necessarily reflect those of the European Commission or BirdLife International.

Citation: BirdLife International (2021) European Red List of Birds. Luxembourg: Publications Office of the European Union.

Table of contents

Introduction	4
Foreword	5
Acknowledgements	6
Executive summary	8
1. Background	10
1.1 The European context	10
1.2 Red Lists	10
2. Methodology	12
2.1 Red List assessment process	12
2.2 Geographic scope	14
2.3 Taxonomic scope	14
3. Results and discussion	16
3.1 Overview	16
3.1.1 Red List status of European birds	16
3.1.2 Population trend of European birds	17
3.2 European bird species changing Red List category	18
3.3 Status and trends of European birds by taxonomic and functional group	24
3.4 Status and trends of European birds by main type of habitat	26
3.5 Gaps in knowledge	29
4. Biodiversity protection in Europe and the EU	30
5. Recommendations and future work	32
5.1 Recommendations for policy and practice	32
5.2 Conclusions	34
5.3 Future work	34
References	35
Appendixes	38
A.1 Red List status, trend and population size estimate, as well as endemism of birds in Europe	38
A.2 Red List status, trend and population size estimate, as well as endemism of birds in the EU	48
A.3 Red List status of birds in Europe and the EU	58
A.4 Countries and territories covered by the European Red List of Birds	67

Introduction



Martin Harper

Regional Director, BirdLife Europe and Central Asia

The goal of nature conservation is to keep common species common, to prevent threatened species from becoming extinct and to ensure that humans continue to benefit from the services that nature provides; such as pollination, clean water, flood protection, a stable climate, recreation and inspiration.

By assessing the extinction risk of all 544 wild bird species in Europe, this update to the European Red List shows how we are doing in meeting this ambition by using birds as the most visible and best studied indicators of biodiversity.

Like so many recent reports about the state of nature, the headlines do not make for comfortable reading. Nearly one in eight European bird species are threatened. The groups of species most at risk are seabirds, wildfowl, waders and raptors; but our analysis shows renewed concern about declining trends in the populations of migratory birds, especially long-distance migrants, farm/grassland species and potentially forest birds. It is alarming to see the status of iconic species like Demoiselle Crane, Common Swift and even Rook deteriorate alongside declines in the population of most larks, shrikes and buntings.

The results demonstrate that we are failing to manage our land, freshwater and seas sustainably. We want and need Europe to lead the world in restoring nature, but for that to happen, nothing short of a transformation to our economies is needed.

This transformation must begin NOW, in what the UN has called the Decade of Ecological Restoration. This is essential for our region to fulfil its new international obligations under the Convention on Biological Diversity, which (in its current draft of the global biodiversity framework) is setting a new milestone for extinction risk to be "reduced by at least 10 percent, with a decrease in the proportion of species that are threatened, and the abundance and distribution of populations of species enhanced or at least maintained."

While the headlines in the new Red List are stark, we should feel optimistic that we can make things better, because within the report, there is some good news. The improved status of Red Kite and Azores Bullfinch – two remarkable conservation success stories – shows that targeted approaches to species recovery can work; while interventions such as well-designed agri-environment schemes can provide a lifeline to threatened species like Corncrake. Yet, this optimism must be conditional on governments, businesses and civil society playing their part to the full.

Governments across Europe need to translate the new global ambition to restore nature into legal targets, backed up by the right policies and funding. Businesses must find new ways to be nature positive, ensuring that their commercial activity improves the state of biodiversity. Civil Society, including BirdLife International Partners, must continue to champion urgent action but also be prepared to take practical steps to help recover threatened species.

The results from this report (and other resources such as the new European Breeding Bird Atlas or the Common Bird Monitoring Scheme) should help decision-makers and will inform the development of a new list of Species of European Conservation Concern to help prioritise future conservation effort. More than that, we hope that it serves as a catalyst for more people and organisations to take action to help Europe's birds.

In future European Red Lists, we want to be able to report a reduction in the number of species at risk of extinction, but to do that, we all have to change the way we think and act.

Foreword



Micheal O'Briain

Deputy Head of the Nature Unit, DG Environment, European Commission

Ever since Rachel Carson's 'Silent Spring' the role of wild birds as key indicators of the state of our environment has become increasingly recognised. Monitoring bird populations is essential to review the progress made towards achieving conservation goals, address gaps in knowledge, and define priorities for conservation action.

BirdLife International's latest updated European Red List of Birds provides a critical assessment of the health of Europe's birds. Based on data gathered in over 50 countries and territories, this major compilation and synthesis of knowledge, much of it undertaken by volunteers, reveals that we are still experiencing the loss of nature across our continent. It is particularly important in determining the risk of extinction, revealing that 13% of European birds are threatened with extinction with another 6% Near Threatened. Of 544 species assessed 30% of them are showing a decreasing trend.

How can we still be facing such a continued erosion of nature given that we have strong legislation for the conservation of birds and other wildlife in Europe since 1979? The EU Birds and Habitats Directives, the Bern Convention on European Wildlife, the Convention of Migratory Species and the Agreement on the Conservation of African-Eurasian Migratory Waterbirds all provide a valuable framework for co-operation and action to conserve species, sites and habitats.

However, despite some significant achievements in European conservation, especially in setting up the Natura 2000 network of protected areas, there are still major shortcomings in the implementation of nature legislation. Conservation action is frequently lacking or incomplete and not taking place at the necessary scale to address the key pressures and threats that wild birds are facing. These include land use changes, particularly linked to agriculture, as well as urbanization, forestry, alterations to freshwater habitats, pollution, overexploitation and illegal killing and invasive alien species, all of which negatively affect nature. On top of all these problems, climate change is emerging as a key threat to the future of our avifauna.

Is there any basis for optimism? We know that targeted action does halt and reverse the decline of species. There are inspiring success stories in Europe with emblematic species such as the Bittern, Griffons Vulture and Red Kite all showing recovery

in areas where they have benefitted from conservation action, frequently supported by initiatives such as the EU LIFE programme. There is now also a much greater public appreciation of the fact that the climate and biodiversity crises are closely interlinked and that restoring our natural capital is essential for economic prosperity and human wellbeing.

This is reflected in renewed commitments for the recovery of nature, including the UN declaring the 2020's as the decade on ecosystem restoration to heal our planet. The EU Biodiversity Strategy for 2030 has set ambitious targets, including extending legal site protection to at least 30% of the EU's land and sea area, 10% of which should be strictly protected, with all of the sites to be under effective management. It also commits to ensuring that by 2030 there is no further deterioration in any habitats and species listed in the Nature Directives and that there is a positive strong trend for at least 30% of those in poor or bad state. A new EU Nature Restoration legal instrument is under preparation, which should provide a major impetus to the restoration agenda in the coming years. Hopefully, these initiatives will inspire similar commitments for the recovery of nature elsewhere in Europe and at global level.

Translating ambitious targets for the recovery of nature into effective action will require very significant human and financial resources over the coming years. Realising the full potential of existing nature and other environmental legislation will be central to success. However, as recognised in the EU Green Deal there is also an urgent need for a fundamental shift in the way we live, particularly in our production and consumption patterns. The attainment of biodiversity and climate objectives will need to be embedded into all relevant policy sectors. There are great opportunities to transition to a just and fair green economy but this will require fully engaging business and other key stakeholder groups such as farmers, foresters and fishers, which affect land and water use across Europe.

As revealed by Eurobarometer surveys that gauge public opinion, European citizens want and expect us to succeed in halting and reversing the loss of biodiversity. This new Red List of Birds provides a vital baseline to assess future progress towards restoring the health of our shared natural heritage.

Acknowledgements

The 2021 update of the European Red List of Birds was made possible thanks to financing from the European Commission (service contract No 07.0202/2018/790700/SER/ENV.D.3).

For their fundamental support to the project, we would like to give special thanks to Angelika Rubin, Micheal O'Briain and Sofia Pachini from the Nature Unit of the European Commission's Directorate General for the Environment (DG ENV); Carlos Romao and Eleni Tryfon from the European Environment Agency; as well as Zelmira Šípková-Gaudillat, Nirmala Seon-Massin, Jérôme Bailly Maitre, Pablo Bolanos, Sabine Roscher and Michelle Watson from the European Topic Centre on Biological Diversity.

This project was implemented in collaboration between BirdLife Europe and Central Asia, BirdLife International and the International Union for the Conservation of Nature (IUCN).

The coordination of the project, Red List assessments and writing of the report was carried out by Anna Staneva and Claire Rutherford from BirdLife Europe and Central Asia. Red List assessments at EU level were also carried out by Alexa Piggott, Hannah Grice and Eresha Fernando. Expert advice on applying the IUCN Red List categories, criteria and regional guidelines; as well as invaluable support throughout the project, was provided by Ian Burfield and Rob Martin from the Global Red List Team at BirdLife International. We are grateful to Rob Pople for his time, expertise and knowledge sharing during the initial months of the project. This project would not have been possible without the encouragement and supervision of Iván Ramírez, former Head of Conservation of BirdLife Europe and Central Asia. Mark Balman, Mike Evans and Sarah Mogg from the BirdLife International Global Information Management Team have contributed with producing species range maps and supplementary materials, which are integral parts of the Red List species profiles. Martin Harper, Ariel Brunner, Justine Guiny,

Harriet Bradley, Anouk Puymartin, Sofie Ruyschaert, Antonio Vulcano, Willem Van den Bossche and other colleagues from BirdLife Europe and Central Asia provided expert advice and feedback during the preparation of this report. Their support and encouragement made a world of a difference for the coordinating team during the intense last months of the project. Emma Brown and Annie Huang, the BirdLife International Conservation Team interns, gave valuable external insights and proofreading feedback. Alison Morris and Kate Williamson from the BirdLife International Human Resources Team provided support with recruitment for the project, while Georgie Godby has provided legal and contracting advice.

At IUCN, Caroline Pollock and Janet Scott have provided their expert advice in assessing and validating the Red List status of birds. Their understanding and readiness to support the project in its very crucial moments were essential to the successful delivery of this work.

Important parts of the data used for the European Red List of Birds assessment were officially reported by EU Member States to the European Commission under Article 12 of the Birds Directive in 2013-2018. To complete the dataset for the pan-European Red List assessments, similar data were sourced from BirdLife Partners and other collaborating experts in most other European countries. The European Red List of Birds was heavily dependent on the collaboration of hundreds of experts and thousands of volunteers in 54 countries and territories, who generously gave their time, data and knowledge. The enthusiasm and commitment of these people has enabled us to generate a comprehensive and detailed picture of bird status and trends in Europe. We record below our particular thanks to the following people, who have contributed and provided species data included within this report, asking for forgiveness from anyone whose name is inadvertently omitted or misspelled:

- **All EU Member States:** all those that contributed to the official reporting under Article 12 of the EU's Birds Directive to the European Commission
- **Albania:** Taulant Bino and Erald Xeka (Albanian Ornithological Society, AOS)
- **Andorra:** Clara Pladevall (Institut d'Estudis Andorrans, IEA)
- **Armenia:** Karen Aghababyan (BirdLinks Armenia)
- **Azerbaijan:** Elchin Sultanov (Azerbaijan Ornithological Society, AOS)
- **Belarus:** Anastasiya Kuzmiankova (independent expert)
- **Bosnia and Herzegovina:** Dražen Kotrošan (Ornithological Society Naše Ptice)
- **Faroe Islands:** Jóhannis Danielsen (Faroese Marine Research Institute) and Janus Hansen (Faroe Islands National Museum)
- **Georgia:** Khatia Basilashvili, Zurab Gurgenidze (SABUKO), with the help and input of Zurab Javakhishvili and Nika Paposhvili
- **Greenland:** David Boertmann (Aarhus University)
- **Iceland:** Hólmfríður Arnardóttir and Kristinn Haukur Skarphéðinsson (Fuglavernd)
- **Kosovo:** Qenan Maxhuni (Institute for Nature Protection)
- **Liechtenstein:** Georg Willi (Botanisch-Zoologische Gesellschaft, BZG)
- **Moldova:** Silvia Ursul and Vitalie Ajder (Societatea Pentru Protecția Păsărilor și a Naturii, SPPN)
- **Montenegro:** Bojan Zeković (Centre for Protection and Research of Birds of Montenegro, CZIP)
- **Norway and Svalbard:** Paul Shimmings (Norwegian Ornithological Society, NOF), with input from Arne Follestad (Norwegian Institute for Nature Research) and Jon Bekken
- **Republic of North Macedonia:** Metodija Veleviski, Danka Uzunova, Ksenija Putilin Stamkoska, Slave Nakev, Nenad Petrovski (Macedonian Ecological Society, MES), Henk Sierdsema, Wouter Plomp, Vlasta Škorpíková, Václav Prášek, Emanuel Lisitchanets, Nicky Petkov and Mihail Iliev
- **Russia:** Alexander Mischenko, Olga Sukhanova (BirdsRussia), Victor Belik (Southern Federal University), Yuri Krasnov (Murmansk Marine Biological Institute Russian Academy of Sciences), Vladimir Sarychev, Ekaterina Preobrazhenskaya, Olga Voltzit, Alexander Sharikov, Gennady Grishanov and Maria Gavriilo
- **Serbia:** Marko Šćiban (Bird Protection and Study Society of Serbia, BPSST), Rajkov, S., Rajković, D., Mirić, R., Medenica, I., Radaković, M., Stanojević, N., Puzović, S., Vučanović, M., Agošton, A., Rudić, B., Sekereš, O., Gergelj, A., Radišić, D., Đorđević, I., Čuturilov, S., Đapić, D., Grabovac, D., Spremo, N., Dožai, J., Pantović, U., Nikolić-Antonijević, J., Vračarić, M., Tucakov, M., Simić, D., Grujić, D., Buzoganj, T., Vasiljević, A., Ružić, M., Dajović, M., Balog, I. and Medved, A.
- **Switzerland:** Peter Knaus, Nicolas Strebel, Hans Schmid, Thomas Sattler, Claudia Müller, Bernard Volet, Samuel Wechsler, Sylvain Antoniazza, Jérôme Guélat and Marc Kéry (Swiss Ornithological Institute)
- **Turkey:** Serdar Özuslu (Doğa), Ömer Döndüren, Soner Bekir, Özge Balkız, Süreyya İsfendiyaroğlu, Kaan Özgencil, Cansu Özcan, Dilek Şahin, Burak Tatar, Kerem Ali Boyla, Lider Sınay, Arzu Gürsoy, Kiraz Erciyas, Ferdi Akarsu and Şafak Arslan
- **Ukraine:** Tatiana Kuzmenko, Olexandr Bronskov, Mykhaylo Banik, Ihor Shydlovskiy, Yuriy Strus, Vitaliy Gryshchenko, Dr. Maksym Gavryliuk and Yuriy Kuzmenko (Ukrainian Society for the Protection of Birds, USPB)

In addition, Danae Portolou (Hellenic Ornithological Society, HOS), Christina Ieronymidou (BirdLife Cyprus), Stoycho Stoychev, Irina Mateeva and Nicky Petkov (Bulgarian Society for the Protection of Birds, BSPB), Aleksii Lehtikoinen (Finnish Museum of Natural History), Juan Carlos del Moral, Juan Antonio Lorenzo (SEO/BirdLife Spain), Hany Alonso (SPEA) and Szabolcs Nagy (Wetlands International) have provided the team with important feedback on the data and data quality for several species, which contributed to the final assessments.

Executive summary

The European Red List of Birds is a review of the regional extinction risk of all 544 species of birds occurring regularly and naturally in Europe. The assessment, performed by BirdLife International for the fourth time (1994, 2004, 2015 and 2021), follows the IUCN Red List Categories and Criteria applied at regional level. Evaluating the extinction risk of each species – i.e., Least Concern, Near Threatened or threatened, and if the latter, to what level – helps to inform decision making, shaping national and international environmental policies and on-the-ground conservation action. The current update of the Red List assessment was (similarly to 2015) carried out at two regional levels – EU and Europe.

The results presented in this report¹ are based on data collected by thousands of experts and volunteers from 54 countries and territories across Europe, extending from Greenland, Iceland and Svalbard in the North to the Canary Islands, Malta and Cyprus in the South, and from the Azores in the West to the Caucasus and Ural Mountains in the East. These include data reported by the 28 EU Member States (including the UK) under Article 12 of the Birds Directive. Additional sources such as scientific reports, national atlases and Red Data Books, as well as peer-reviewed literature were also used to make sure the most recent data available were used. The data, compiled in 2019, encompass the 1980–2018 period.

At the European regional level, 13% (71 species) of bird species are threatened, with approximately 2% (8 species) Critically Endangered, nearly 3% (15 species) Endangered, and nearly 9% (48 species) Vulnerable. A further 6% (34 species) are considered Near Threatened, while five species are considered Regionally Extinct. Changes between Red List categories since the previous assessment in 2015 were recorded for 83 species, of which 36 species were classified in a higher extinction risk category and 47 in a lower category. Most of these changes are due to genuine deterioration or improvement in the species populations, while a smaller proportion are related to artefacts, e.g., due to changes in the methodology of field data collection, calculation of trend values, or previous under- or overestimation of population size. Another 13 species have been assessed for the first time due mainly to taxonomic changes.

At the EU regional level, the results show that nearly 19% of bird species are threatened, with 2% Critically Endangered, nearly 5% Endangered, 12% Vulnerable, and a further 8% are Near Threatened. When looking at Red List trends, 34% of species in the EU are in decline, another 22% have stable or fluctuating trend, 28% are increasing, while 16% of species have unknown trend. A total of 42 species were assessed under higher risk Red List category and 39 species placed in a lower risk Red List category, compared to 2015. Fifteen species were assessed for first time in the EU.

At both European and EU level, seabirds, wildfowl, waders and raptors each have the highest proportion of threatened and Near Threatened species per taxonomic group, along with smaller groups such as kingfishers, bustards and grebes. When looking at trends in Europe, over 40% of wildfowl and wader species, over 30% of seabirds, as well as 25% of raptors, are experiencing population declines. Within the EU, the proportion increases to over 50% for waders and wildfowl, and more than 30% for raptors. Relatively high proportions of declining trends are also observed among migratory birds, with 121 species considered sub-Saharan migrants (of which 33% have declining trends), compared to species that are migratory but do not migrate as far as sub-Saharan Africa (of which 31.5% have declining trends).

A comparison between main habitat types in Europe and the EU shows that farmland and grassland, as well as marine habitats, have the highest number of associated threatened and Near Threatened species. This is supported by the fact that approximately 30% of species associated with these habitats have a decreasing trend in Europe, while in the EU, the proportion is even higher, reaching over 40%. Among species associated with forest habitats, those with decreasing, stable, increasing and unknown trends share a quarter of the total each. Rocky habitats like inland cliffs and mountain peaks have the highest proportion of species with unknown trends (more than 50%), and a third of all species associated with marine habitats and shrubland also have unknown trends. Within the EU, forest and shrubland habitats seem to fare better, with 60% of species having stable or increasing populations.

Despite having an exceptional history of data collection and environmental surveillance, Europe's knowledge base on birds still has a lot of room for improvement. Good cooperation between governments, organisations and institutions, as well as coordinated international efforts in research and bird monitoring, supported by the availability of dedicated resources, are essential to successfully inform and address conservation issues across the continent. The models of the legal mechanisms (e.g. reporting under art. 12 of the Birds Directive) and financial mechanisms (e.g. LIFE Programme) of the European Union play a crucial role in this process.

The results confirm the overall conclusions found in other studies. Large scale land use change, the intensification of agricultural practices, the development of infrastructure, the over-exploitation of marine resources, the pollution of inland waters and unsustainable yet commonly used forestry practices are the main drivers of bird population declines observed in European habitats. Ongoing population declines and range contractions in widespread species of open habitats – like larks, shrikes and buntings – clearly demonstrate the impact of disappearing mosaic landscapes and the increased use of agricultural chem-

¹ Also available on <http://datazone.birdlife.org> and <https://www.iucnredlist.org>.



icals. Although some raptor species have made recent population recoveries due to legal protection and targeted conservation activities, many raptor species dependent on open habitats (e.g. grassland and shrubland) for prey are still in decline. In the marine environment, birds suffer from bycatch and incidental killing (due to fishing and hunting activities), as well as invasive non-native species, disturbance and water pollution (originating from both land and sea). Many species of birds are legally hunted across Europe, while millions die every year from illegal killing.

In addition to these threats come the unpredictable effects of climate change on species populations. Current knowledge suggests that for certain species climate change is or will be a major drive for range shifts, while for others it may significantly increase their extinction risk.

Examples of successful species and habitat recovery can be found across Europe. The decline of many widespread and common species (bird species, but also species in other taxonomic classes such as butterflies and other pollinating insects) signals, however, a problem of a much larger scale: a systemic issue in the functioning of ecosystems caused by a diverse array of human-induced changes.

Future conservation efforts in Europe should not only focus on action locally, but also seek to deliver more fundamental change at a bigger scale. Collaboration between decision-makers and conservation practitioners needs to be expanded while securing greater involvement from consumers, producers, landowners and other businesses.

The European Union commits to fight the intrinsically linked biodiversity and climate crises with its 2030 Biodiversity Strategy, as part of the Green Deal, for the broader European continent and the world. At the heart of the Strategy lie ambitious plans to protect nature through a coherent and extended network of well-managed protected areas, the restoration of biodiversity and the reduction of the environmental footprint of production (e.g. of food, timber or energy). To avoid actions that are only implemented on paper, these policies need to be legally binding, have clear guidance criteria and be actively enforced. In addition, urgent action is needed to reform agricultural, fisheries and forestry policies, mainly through legal obligations but also through incentives, to safeguard biodiversity. International agreements and coordinated efforts remain crucial in all aspects of the protection and restoration of the natural environment, but particularly in the sustainable use of marine resources, the conservation of migratory birds at a flyway scale, and to tackle the climate crisis.

Ultimately, as humans, we depend on a healthy natural environment (including irreplaceable services and functions such as the provision of clean water, pollination of crops, flood protection, and a stable climate; which are also necessary for our recreation and well-being). Without the services provided by the natural environment, our health, society and economy would not hold up. Therefore, it is important to understand that investing in nature pays off, not just to the habitats and species that are being protected, but also to us as individuals, to governments and to society as a whole.

1.

Background

1.1 The European context

Europe, with its 10.18 million km² (approximately 7% of Earth's land surface), offers suitable habitats for over 70 bird taxonomic families comprising over 540 species (approximately 5% of the global avifauna). A fifth of these species are endemic or near-endemic to the region. Birds are an important part of European biodiversity. They fulfil specific roles within the diversity of ecosystems, and their status has been recognised as a good indicator of the state of the natural environment. Therefore, knowledge about birds can help us identify, predict and understand changes, and take adequate action to prevent the loss of biodiversity and valuable resources derived from nature.

The European landscape has been continually shaped and modified by humans over thousands of years. With the historical expansion of people, habitats have gradually transformed from natural wild areas to settlements, roads, managed woodland, rivers and lakes, farmed land and pasture. With these alterations, there has come a change in the bird communities present (e.g., species that would have naturally favoured grassland and woodland edge habitats as new opportunities emerged have become relatively more abundant; whereas species dependent on habitats and features that were not present in towns and farmland (such as forest- or wetland-dependent species) have been gradually displaced).

However, the rate of global change in nature during the past 50 years is unprecedented in human history². Inevitably, many species populations have declined dramatically. For example, birds of prey like the Red-footed Falcon (*Falco vespertinus*) have suffered from significant deterioration in their food supply due to the massive use of pesticides in agriculture³, adding to other threat factors such as persecution and infrastructure intensification.

Species monitoring has been indispensable in understanding these drivers of decline over time, and have contributed to the assessment of species' risk of extinction. As a result, informed decisions to undertake conservation actions like legal protection, ceasing the use of certain chemicals, or creating new nesting opportunities for birds have resulted in halting declines in certain species (e.g., Peregrine Falcon (*Falco peregrinus*) or Red Kite (*Milvus milvus*)).

1.2 Red Lists

The European Red List of Birds is a review of the regional extinction risk of all species of birds occurring regularly and naturally in Europe. The current publication presents the Red List status of birds at two regional levels - the EU and Europe. The assessment is part of the IUCN Red List of Threatened Species, which aims to evaluate the extinction risk of each speciesTM, which then helps to inform decision making, the shaping of national and international environmental policies and on-the-ground conservation action. The IUCN Red List is used as a critical indicator of the health of the world's biodiversity. It is often used as a guide to revise the annexes of some important international agreements, such as the Convention on International Trade in Endangered Species (CITES) and the Convention on Migratory Species (CMS), or to measure the progress towards achieving international biodiversity and sustainable development goals (SDGs). Data from the IUCN Red List are also used to calculate the Red List Index (RLI), which is one of the biodiversity indicators used by the Convention on Biological Diversity (CBD). For instance, Milestone 2b, aims to ensure that "the increase in the extinction rate is halted or reversed, and the extinction risk is reduced by at least 10 percent, with a decrease in the proportion of species that are threatened, and the abundance and distribution of populations of species is enhanced or at least maintained".


² <https://ipbes.net/global-assessment>

³ Ferguson-Lees, J.; Christie, D.A. 2001. Raptors of the World. Christopher Helm, London.

Along with assessing the risk of species becoming globally extinct, it is also important to assess species' extinction risk at regional and national levels, where conservation policy is more often established and implemented. The European Red List of Birds is a regional Red List assessment initiated by BirdLife International in 1994. Since then, updates have been published in 2004 and 2015, with the 2021 update being the fourth one. In 2015, Red List assessment of birds was carried out also at EU level, following the obligations of Member States to report under Art. 12 of the Birds Directive (repeated at 6-yearly cycle).

The current assessment is the second one performed at EU regional level.

The results of the European Red List of Birds are used to inform decision-making across the region, set up biodiversity protection targets (e.g., at national level, or as part of the EU Biodiversity Strategies to 2020 and 2030), and identify priorities for further research and conservation (e.g., for threatened species). The European Red List of Birds is the baseline for assessing their conservation status as European Species of Conservation Concern (SPECs)⁴.



Red-necked Grebe (*Podiceps grisegena*)
Photo: Becky Matsubara

⁴ https://www.birdlife.org/sites/default/files/attachments/European%20Birds%20of%20Conservation%20Concern_Low.pdf

2.

Methodology

2.1 Red List assessment process

The European Red List of Birds assessment follows the IUCN Red List Categories and Criteria (version 3.1, second edition)⁵ as well as the Guidelines for Using the IUCN Red List Categories and Criteria (version 14, August 2019)⁶. The extinction risk of a species can be assessed at global, regional or national levels. It

is quite common for a species to be listed in different Red List categories on the global IUCN Red List and a Regional Red List. For example, a species that is common worldwide and classified as Least Concern on the IUCN Red List may face a high level of threat in a specific region and therefore be listed as Endangered in that region (see Figure 1 for the explanation of the IUCN Red List Categories). To avoid under or overestimating a particular species, the Guidelines for Application of IUCN Red List Criteria at Regional and National Levels (version 4.0)⁷ are applied.



Common Redshank (*Tringa totanus*)
Photo: Britt-Marie Sohlström

⁵ <https://portals.iucn.org/library/node/10315>

⁶ <https://www.iucnredlist.org/resources/redlistguidelines>

⁷ <https://www.iucn.org/content/guidelines-application-iucn-red-list-criteria-regional-and-national-levels-version-40>




Figure 1. IUCN Red List Categories at a regional scale

The assessment is based on data reported by the 28 Member States of the European Union (EU)⁸ under Article 12 of the EU Birds Directive, as described in the State of Nature in the EU –Methodological paper⁹, as well as national reports from 22 countries and territories in Europe which are not part of the EU. The latter were collated by BirdLife International through national experts, coordinators of bird monitoring programmes and organisations, including BirdLife International Partners and many others. The reports were submitted in 2019-2020 and included information on the size and trend of populations and distributions of individual bird species. Additional sources such as scientific reports, national atlases and Red Data Books, as well as peer-reviewed literature were also used to make sure the most recent data available were considered. Due to delays in reporting under Article 12, the 2013-2018 data from Romania were not used for the EU level assessment, nonetheless they were included at European level.

The national data were combined to produce overall European and EU population sizes and trends for each species. For popu-

lation sizes, the reported minimum and maximum population size data across countries were summed to calculate the overall minimum and maximum value. In addition, the best estimate was used if provided, and if not, the geometric mean of each population size was calculated and used as a best estimate of population size where appropriate. All European countries were requested to use the same population unit, which was breeding pairs for most breeding birds, (except for a minority of taxa with unusual or complex breeding biology or cryptic behaviour, for which other units, such as calling or lekking males, were used), and individuals for birds in winter (non-breeding season). For population trends, data from all countries were combined, weighting each country's contribution according to the size of its population. Weightings were based on the best estimate provided or the geometric mean of the countries' minimum and maximum population size compared to the equivalent total for the overall European/EU population. This analysis was carried out using a dedicated tool developed by IUCN to estimate overall trends based on data from multiple (national) sub-populations¹⁰.

⁸ As it stood in 2019, and therefore at the time, still comprising the UK (hereafter referred to as “the EU28 or just “EU”, a level for which regional Red List assessments have also been previously undertaken in 2020 for native species that regularly occur within the region).

⁹ Röschel, L., Noebel, R., Stein, U., Naumann, S., Romão, C., Tryfon, E., Gaudillat, Z., Roscher, S., Moser, D., Ellmauer, T., Löhnertz, M., Halada, L., Staneva, A. and Rutherford, C. A. (2020). State of Nature in the EU - Methodological paper. Methodologies under the Nature Directives reporting 2013-2018 and analysis for the State of Nature 2000. ETC/BD report to the EEA.

¹⁰ <https://www.iucnredlist.org/resources/criterion-a>

A species' population size trend is measured over the length of time of three generations of that species and described as 'increasing' where the population size has expanded over three generations, 'decreasing' where the population size has undergone reduction, 'stable' where the population size has not experienced any significant changes or 'fluctuating' if the species experiences strong inter-annual changes in its population size. Where it was not possible to allocate a trend category with confidence, either because trend directions were reported as unknown for a large proportion of the total European/EU population, in the case of conflicting trend information, or lack of trend magnitudes, the overall European/EU trend was classified as 'unknown'.

Where possible, the robustness of trend categories to the effects of any missing data was tested using plausible 'good' and 'bad' scenarios, based on other sources of information, such as any other reported trend information, recent national Red Lists, scientific literature, and other publications and consultations with experts. In addition, attention was given to the quality of the reported trend information, e.g., where trends were derived from 'complete surveys or robust estimates' in the majority of a species' range countries; the overall calculated trend was considered of higher quality compared to trends based on 'extrapolation from limited amount of data', the latter being of higher quality than 'expert opinion'.

For most species, assessments were based on data from the breeding season, but in a small number of cases, winter data were also used. Winter data were only reported for a subset of species, mainly wintering waterbirds (especially migratory wildfowl and waders), whose populations are often best monitored in the winter when they congregate in large numbers. The assessments of species that do not breed (regularly) within Europe were based solely on winter data, while for species that occur in both seasons and for which the reported datasets were both representative of the regional population, the assessment process was carried out independently on data for both the breeding and wintering populations. For some species in winter, underlying population trends can be obscured by demographic factors, often related to inter-annual variation in weather conditions. In some years, for example, birds that usually winter in the region may be forced to move elsewhere by harsh winter conditions, whilst in others, birds that usually winter outside the region may show marked influxes into the region. Consequently, assessments were carried out principally based on breeding data, provided that the resulting status

category was the same as or higher (i.e., more threatened) than that obtained using winter data. In cases where status categories were the same in both seasons, the final assessment was based on the season for which better quality data was available.

The species' trends in themselves at European/EU level are also considered in the analyses in this report as they can reveal subtle changes in population sizes, such as slow deteriorations or stabilisations, before meeting a Red List category threshold. When used to study species in groups, such as species dependent on similar habitats, they can indicate a general deterioration of that habitat, implying the degradation of an ecosystem. Trends also provide an additional level or meaning to the Red List assessment of a species. For example, if a species is assessed as Vulnerable due to its rapidly declining trend, this can mean that although it has existed for millennia in this region, in a period of just three generations, at least a third of the species' population has been lost.


2.2 Geographic scope

The geographical scope of the Red List assessments presented in this publication comprises two regional levels - Europe and the EU.

For Europe, the geographical scope is continent-wide and comprises 54 countries and territories across Europe, extending from Greenland (Denmark), Iceland and Svalbard (Norway) in the north to the Canary Islands (Spain), Malta and Cyprus in the south, and from the Azores (Portugal) in the west to the Caucasus and Ural Mountains in the east. For the EU, the geographical scope is limited to the territory of the 28 Member States, including the UK (as the EU stood in 2019). Figure 2 shows the geographic area covered by the European Red List assessment.

2.3 Taxonomic scope

All terrestrial and marine bird species native to and/or naturally (and regularly) occurring in Europe are included in the assessment. Assessments were carried out at the species level only, following BirdLife International's current taxonomy (BirdLife International 2021), and population size and trend data reported by countries at subspecific or flyway population levels were aggregated to the species level.



The complete list of countries and territories covered by the European Red List of Birds is presented in Appendix 4.

Figure 2. Geographic area covered by the Red List of Birds assessment - European level shown in green and EU28 shown in green with blue stripes.

3.

Results and discussion

3.1 Overview

3.1.1. Red List status of European birds

At the European regional level, 13% (71 species) of all 544 assessed bird species are threatened, with approximately 2% (8 species) Critically Endangered (CR), nearly 3% (15 species) Endangered (EN), and nearly 9% (48 species) Vulnerable (VU). A further 6% (34 species) are considered Near Threatened (NT) (Figure 3 shows the percentage of species in each IUCN Red List Category). This translates as one in five bird species in Europe being threatened or Near Threatened with extinction. By comparison, within the EU28 region, nearly 19% of bird species (86 species) are threatened, with 2% (9 species) Critically Endangered, 4.5% (21 species) Endangered, 12% (56 species) Vulnerable, and a further 8% (37 species) are Near Threatened. Therefore, in the EU28 region, over a quarter of bird species are threatened or Near Threatened.

Compared to the European Red List assessment in 2015¹¹ (when 533 species were assessed), the results appear very similar with 13% threatened species, 2% Critically Endangered, 3% Endangered species, and 7% Vulnerable species (see Section 3.2 for more details). The proportion of Near Threatened species in Europe has remained unchanged (6%). Analogous comparison at EU level shows that the proportion of threatened species has increased by almost 1% (was 18%), while that of Near Threatened species has increased by 2% (was 6%).

Two species which used to occur in Europe are now globally Extinct; they were not assessed or counted in the statistics below as they disappeared much earlier than the current time period – these are the Canary Oystercatcher (*Haematopus meadewaldoi*) and Great Auk (*Pinguinus impennis*). Five species are Regionally Extinct in Europe: Pallas's Sandgrouse (*Syrhaptes*

paradoxus), Northern Bald Ibis¹² (*Geronticus eremita*), African Darter (*Anhinga rufa*), Common Buttonquail (*Turnix sylvaticus*) and Pine Bunting (*Emberiza leucocephalos*). New data has contributed to the re-assessment of two species previously classified as Regionally Extinct: the Caspian Plover (*Charadrius asiaticus*) and Desert Warbler (*Curruca nana*), which are now considered Endangered and Vulnerable, respectively. Another three species are Critically Endangered (Possibly Extinct): the Asian Houbara (*Chlamydotis macqueenii*), Slender-billed Curlew (*Numenius tenuirostris*) and Pechora Pipit (*Anthus gustavi*), with the latter being the latest addition to the list.

Amongst the most threatened (Critically Endangered) European birds is one wader species: the Sociable Lapwing (*Vanellus gregarius*); one seabird: the Balearic Shearwater (*Puffinus mauretanicus*); as well as the Steppe Eagle (*Aquila nipalensis*), Yellow-breasted Bunting (*Emberiza aureola*) and Red-knobbed Coot (*Fulica cristata*).

A fifth of Europe's birds (103 species) are endemic or near endemic to the region. Of these, almost a quarter (24 species) are threatened or Near Threatened, and over a quarter (28%) are declining, increasing and unknown, respectively, with the remaining species being stable or fluctuating. The most threatened endemic European species is the Balearic Shearwater (*Puffinus mauretanicus*) (Critically Endangered), followed by the Endangered endemic or near endemic species: the Atlantic Puffin (*Fratercula arctica*), the Gran Canaria Blue Chaffinch (*Fringilla polatzeki*), and Zino's Petrel (*Pterodroma madeira*).

The number of endemic and near endemic species in the EU is 52. Among these, the Balearic Shearwater (Critically Endangered), Zino's Petrel (Endangered) and Gran Canaria Chaffinch (Endangered) are the most threatened.

¹¹ http://datazone.birdlife.org/userfiles/file/Species/erlob/EuropeanRedListOfBirds_June2015.pdf

¹² The Northern Bald Ibis does exist in Europe but only as partially captive, captive-bred and / or reintroduced populations, none of which are yet considered self-sustainable, hence its continued evaluation as RE.




Figure 3. IUCN Red List status of birds in Europe (left) and the EU28 (right).

3.1.2. Population trend of European birds

Overall, 30% of European birds have a decreasing population trend, 23% are stable or fluctuating, 21% are increasing, while 26% have an unknown trend (Fig. 4). By comparison, at EU level, the proportion of species with decreasing trend is higher

(34%), the proportion of species with stable or fluctuating trend is similar (22%), those with an increasing trend is higher (28%), while the proportion of species with unknown trend is significantly lower (16%).




Figure 4. Population trends of birds in Europe (left) and the EU28 (right).

3.2 European bird species changing Red List category

In total, 83 species have changed Red List status at European level since the previous assessment in 2015¹³ (Table 1), mostly by one category but occasionally by more. Among these, 36 species are now classified in a higher extinction risk category. Most of these cases are due to genuine declines in species populations. An example is the Demoiselle Crane (*Anthropoides virgo*), where new data from across its European range (mainly Russia and Ukraine) show genuine declines at a rate of over 50%, warranting a reclassification from Least Concern in 2015 to Endangered. The remaining 47 species have been reclassified in lower Red List categories. Although new data suggest they face a lower level of extinction risk compared to 2015, half of these species are still undergoing (slow) population declines. An example is the Common Kingfisher (*Alcedo atthis*), which is now considered Least Concern (it was Vulnerable in 2015) (see Box 1).

The change between categories in roughly one out of ten (of the 83) species is considered to result from changes in the methodology of field data collection, calculation of trend values, or previous under- or overestimation of population size (i.e., non-genuine changes). Updated estimates of species' generation lengths¹⁴ have also affected the final assessment of certain species; e.g. long-lived species for which trends were calculated over a significantly shorter or longer 3-generation period. The re-assessment of the Northern Fulmar (*Fulmarus glacialis*) is an example of such a non-genuine change in category, where the use of the new generation length value has resulted in its status now being assessed as Vulnerable (rather than Endangered).

The improvement of status in some species reflects a genuine population increase due to targeted conservation action, for instance, with the Red Kite (*Milvus milvus*) being downlisted from Near Threatened to Least Concern, and the Bearded Vulture (*Gypaetus barbatus*) being downlisted from Vulnerable to Near Threatened. It also underlines the importance of such actions being continued in the long term, until the threats to the species have ceased and the populations in the region are no longer conservation dependent. For example, the White-headed Duck (*Oxyura leucocephala*) was previously Endangered in Europe. Thanks to long-term conservation action taking place and the elaboration and implementation of an international Species Action Plan (SAP) limiting the reduction and degradation of its habitat, the pressure of hunting, and the risk of hybridisation with the non-native Ruddy Duck (*Oxyura jamaicensis*); its rate of decline has now reduced to the point where it no longer meets the thresholds for qualifying as Endangered, and is now classified as Vulnerable. In the case of the Egyptian Vulture (*Neophron percnopterus*) though, reclassified as Vulnerable from Endangered (based on the lower rate of decline mainly in its Iberian population), there is still a long way to go; legal protection and on-the-ground conservation action need to be further implemented at national level and across borders, both within its breeding and wintering grounds and along migration routes. Underlying this further is the Saker Falcon (*Falco cherrug*) which, after decades of successful population restoration in its core European range, is again experiencing recent overall population declines that have resulted in its status being reassessed as Endangered (from Vulnerable).

At EU level, the analysis shows very similar results, with 42 species placed in a higher risk Red List category and 39 species placed in a lower risk Red List category, compared to 2015.

¹³ <http://datazone.birdlife.org/info/euroredlist>

¹⁴ Bird, J. P., Martin, R., Akçakaya, H. R., Gilroy, J., Burfield, I. J., Garnett, S. G., Symes, A., Taylor, J., Şekercioğlu, Ç. H. and Butchart, S. H. M. (2020). Generation lengths of the world's birds and their implications for extinction risk. *Conservation Biology* 34(5): 1252-1261. DOI: 10.1111/cobi.13486.

Table 1. Species whose European Red List categories have changed in 2021 (shown in taxonomic order). On page 19, species whose reassessment has placed them in a higher risk Red List category; on pages 20 and 21, species whose reassessment has placed them in a lower risk Red List category. Species shown in bold have changed Red List category at EU as well as European level.

SCIENTIFIC NAME BY TAXONOMIC ORDER	ENGLISH NAME	2021 EUROPEAN RED LIST CATEGORY	2015 EUROPEAN RED LIST CATEGORY
<i>Branta ruficollis</i>	Red-breasted Goose	VU	NT
<i>Anas acuta</i>	Northern Pintail	VU	LC
<i>Aythya fuligula</i>	Tufted Duck	NT	LC
<i>Somateria mollissima</i>	Common Eider	EN	VU
<i>Lyrurus mlokosiewiczi</i>	Caucasian Grouse	NT	LC
<i>Alectoris rufa</i>	Red-legged Partridge	NT	LC
<i>Ammoperdix griseogularis</i>	See-see Partridge	VU	LC
<i>Coturnix coturnix</i>	Common Quail	NT	LC
<i>Turnix sylvaticus</i>	Common Buttonquail	RE	CR
<i>Podiceps grisegena</i>	Red-necked Grebe	VU	LC
<i>Podiceps nigricollis</i>	Black-necked Grebe	VU	LC
<i>Hydrobates leucorhous</i>	Leach's Storm-petrel	NT	LC
<i>Puffinus yelkouan</i>	Yelkouan Shearwater	VU	LC
<i>Falco cherrug</i>	Saker Falcon	EN	VU
<i>Falco columbarius</i>	Merlin	VU	LC
<i>Falco vespertinus</i>	Red-footed Falcon	VU	NT
<i>Fulica cristata</i>	Red-knobbed Coot	CR	EN
<i>Anthropoides virgo</i>	Demoiselle Crane	EN	LC
<i>Chlamydotis undulata</i>	African Houbara	VU	NT
<i>Calidris falcinellus</i>	Broad-billed Sandpiper	VU	LC
<i>Calidris pugnax</i>	Ruff	NT	LC
<i>Gallinago gallinago</i>	Common Snipe	VU	LC
<i>Tringa totanus</i>	Common Redshank	VU	LC
<i>Stercorarius parasiticus</i>	Arctic Jaeger	EN	LC
<i>Larus audouinii</i>	Audouin's Gull	VU	LC
<i>Larus genei</i>	Slender-billed Gull	VU	LC
<i>Pagophila eburnea</i>	Ivory Gull	VU	LC
<i>Syrhaptes paradoxus</i>	Pallas's Sandgrouse	RE	EN
<i>Clamator glandarius</i>	Great Spotted Cuckoo	VU	LC
<i>Caprimulgus ruficollis</i>	Red-necked Nightjar	NT	LC
<i>Apus apus</i>	Common Swift	NT	LC
<i>Anthus gustavi</i>	Pechora Pipit	CR	VU
<i>Phoenicurus erythrogastrus</i>	White-winged Redstart	NT	LC
<i>Cyanistes cyanus</i>	Azure Tit	NT	LC
<i>Lanius senator</i>	Woodchat Shrike	NT	LC
<i>Corvus frugilegus</i>	Rook	VU	LC
<i>Emberiza leucocephalos</i>	Pine Bunting	RE	VU

SCIENTIFIC NAME BY TAXONOMIC ORDER	ENGLISH NAME	2021 EUROPEAN RED LIST CATEGORY	2015 EUROPEAN RED LIST CATEGORY
<i>Cygnus columbianus</i>	Tundra Swan	VU	EN
<i>Anser erythropus</i>	Lesser White-fronted Goose	VU	EN
<i>Aythya marila</i>	Greater Scaup	LC	VU
<i>Clangula hyemalis</i>	Long-tailed Duck	LC	VU
<i>Oxyura leucocephala</i>	White-headed Duck	VU	EN
<i>Lagopus lagopus</i>	Willow Grouse	LC	VU
<i>Lagopus muta</i>	Rock Ptarmigan	LC	NT
<i>Alectoris chukar</i>	Chukar	LC	NT
<i>Gavia immer</i>	Common Loon	LC	VU
<i>Fulmarus glacialis</i>	Northern Fulmar	VU	EN
<i>Gypaetus barbatus</i>	Bearded Vulture	NT	VU
<i>Neophron percnopterus</i>	Egyptian Vulture	VU	EN
<i>Aquila fasciata</i>	Bonelli's Eagle	LC	NT
<i>Clanga clanga</i>	Greater Spotted Eagle	VU	EN
<i>Milvus milvus</i>	Red Kite	LC	NT
<i>Circus cyaneus</i>	Hen Harrier	LC	NT
<i>Circus macrourus</i>	Pallid Harrier	LC	NT
<i>Accipiter badius</i>	Shikra	NT	VU
<i>Charadrius asiaticus</i>	Caspian Plover	EN	RE
<i>Charadrius leschenaultii</i>	Greater Sandplover	NT	VU
<i>Glareola nordmanni</i>	Black-winged Pratincole	LC	VU
<i>Limosa limosa</i>	Black-tailed Godwit	NT	VU
<i>Numenius arquata</i>	Eurasian Curlew	NT	VU
<i>Hydrocoloeus minutus</i>	Little Gull	LC	NT
<i>Larus argentatus</i>	European Herring Gull	LC	NT
<i>Larus armenicus</i>	Armenian Gull	LC	NT
<i>Alca torda</i>	Razorbill	LC	NT
<i>Uria aalge</i>	Common Murre	LC	NT
<i>Ketupa zeylonensis</i>	Brown Fish-owl	EN	CR
<i>Apus affinis</i>	Little Swift	NT	VU
<i>Alcedo atthis</i>	Common Kingfisher	LC	VU
<i>Ceryle rudis</i>	Pied Kingfisher	VU	EN
<i>Picus sharpei</i>	Iberian Green Woodpecker	LC	NT
<i>Ammomanes deserti</i>	Desert Lark	EN	CR
<i>Melanocorypha yeltoniensis</i>	Black Lark	EN	CR
<i>Anthus pratensis</i>	Meadow Pipit	LC	NT

SCIENTIFIC NAME BY TAXONOMIC ORDER	ENGLISH NAME	2021 EUROPEAN RED LIST CATEGORY	2015 EUROPEAN RED LIST CATEGORY
<i>Prunella montanella</i>	Siberian Accentor	LC	NT
<i>Oenanthe chrysopygia</i>	Red-tailed Wheatear	VU	EN
<i>Oenanthe leucura</i>	Black Wheatear	LC	VU
<i>Turdus iliacus</i>	Redwing	LC	NT
<i>Curruca nana</i>	Asian Desert Warbler	VU	RE
<i>Lanius excubitor</i>	Great Grey Shrike	LC	VU
<i>Passer moabiticus</i>	Dead Sea Sparrow	LC	VU
<i>Pyrrhula murina</i>	Azores Bullfinch	VU	EN
<i>Emberiza cineracea</i>	Cinereous Bunting	NT	VU
<i>Emberiza rustica</i>	Rustic Bunting	LC	VU

Box 1. Species with significant change in their European Red List status



Kingfisher *Alcedo atthis*

VU → LC

This species' trend varies between countries in its range, with many reporting an unknown direction. Based on the latest data available, this species' trend appears to still be decreasing, but at a lower rate than in 2015, and not sufficiently rapidly to meet a higher category than Least Concern. This highlights the importance of regular species monitoring, as if new data were gathered on the trends which are currently unknown, this could paint a more accurate picture of the state of the species within the region. The same change in Red List category is valid at EU level.



Greater Spotted Eagle *Clanga clanga*

EN → VU

A non-genuine change in the European Red List status due to change in generation length from 50 to 28 years (for 3 generations). Furthermore, the overall declining trend is dictated by a 30–49% population reduction within the European part of Russia (which holds 82% of the European population).



Black-tailed Godwit *Limosa limosa*

VU → NT

A reportedly genuine increase in the breeding population in Iceland (which holds approximately 47% of the European population), likely due to increasing spring temperature¹⁵, dictates the overall stable European trend. However, the species is undergoing rapid declines in other key range countries like the Netherlands and Russia (together holding another 45% of the European population), despite recent reports of northward expansion of the species in Northwest Russia¹⁶. This species depends on wet grasslands during the breeding season, and therefore habitat loss and degradation in breeding grounds is a key threat to it, according to the breeding waders Multi-Species Action Plan (MSAP)¹⁷. Despite the stabilisation of the overall population, this threat has not yet been halted.

¹⁵ Effects of spring temperature and volcanic eruptions on wider productivity. Tómas Grétar Gunnarsson, Lilja Jóhannesdóttir, José A Alves, Böðvar Þórisson & Jennifer A Gill. IBIS (2017). DOI: 10.1111/ibi.12449

¹⁶ Popov I. & Starikov D. 2015. Recent northward expansion of breeding Black-tailed Godwits *Limosa limosain* NW Russia. Wader Study 122(3): 173–183.

¹⁷ Leyrer, J., Brown, D., Gerritsen, G., Hötter, H. and Ottvall, R. (compilers). (2018). International Multispecies Action Plan for the Conservation of Breeding Waders in Wet Grassland Habitats in Europe (2018–2028). Report of Action A13 under the framework of Project LIFE EuroSAP (LIFE14 PRE/UK/002). NABU, RSPB, VBN and SOF.



VU

NT

Eurasian Curlew *Numenius arquata*

Although the species has been re-classified under a lower extinction risk category- at both European and EU level. Its overall trend is still decreasing (by 25-29%). The species seems to be doing well in Finland (which holds 37% of the European population), however, in the European part of Russia (holding 29% of the population) and in the UK (holding 25% of the population), its population is still decreasing rapidly. Overall, in Europe, this species is thought to be mostly suffering from the impact of habitat loss and degradation



LC

VU

Common Snipe *Gallinago gallinago*

A genuine change in the main range country in Europe, Russia (holding 65% of the population) drives an overall decrease in the population trend by 36%. This species is declining globally too. The highest impacting threat known for this species is habitat loss and degradation in breeding grounds, which is likely to be driving the declining trend. This species is dependent on wet grassland in the breeding season, and therefore their trends are very much determined by the extent and condition of these habitats in Europe.



LC

VU

Common Redshank *Tringa totanus*

This species is undergoing an overall decline (of over 30% in 3 generations), dictated by decreasing trends in the 3 main range countries (holding circa 50% of the European population): Iceland¹⁸, Russia and Belarus. The overall trend for this species is also supported by the Pan-European Common Bird Monitoring Scheme (PECBMS)¹⁹. As with the Common Snipe, and as per the breeding waders MSAP, this species is highly dependent on wet grasslands and consequently its highest impacting threat is habitat loss and degradation, which is likely to be driving the declining trend.



LC

VU

Northern Pintail *Anas acuta*

A reportedly genuine decline in the European Russian population, as well as in the Finnish and Swedish populations (representing practically the entire European population) result in an overall decline of 38% in three generations. This species is threatened by habitat loss and modification (wetland and coastal habitats) in both its breeding and wintering grounds²⁰, which is likely to be the main driver of its current decline. The species was placed in a higher risk Red List category also in the EU (from VU to EN).

¹⁸ <https://en.ni.is/biota/animalia/chordata/aves/charadriiformes/stelkur-tringa-totanus>

¹⁹ <https://pecbms.info/>

²⁰ Scott, D. A.; Rose, P. M. (1996). Atlas of Anatidae populations in Africa and western Eurasia. Wetlands International, Wageningen, Netherlands.



VU

EN

Common Eider *Somateria mollissima*

It is expected this species will have declined by over 50% in three generations (by 2033) based on reported decreasing trends from across its European range, i.e. Iceland, Finland, Norway and Greenland (which together hold almost three quarters of the European population). The rapid population declines in the Baltic-Wadden Sea after the 2000s are considered to be due a combination of factors, including a decreased survival due to natural predation, diseases and food shortage.²¹



LC

VU

Rook *Corvus frugilegus*

Approximately half of the species' population is experiencing declines. These include countries like Russia, Poland, France and the Netherlands. Some countries have explicitly reported a genuine change in the species' population size, including the European part of Russia (which holds almost a third of the European population). Overall, the population size of the Rook is declining by over 30% in Europe, likely due to the persecution and destruction of nesting sites.



LC

NT

Common Quail *Coturnix coturnix*

The species population in Europe is decreasing by at least 25%, with genuine declines reported by two key range countries – Russia and Spain. The main reasons for decline are considered to be the loss of rough grassland and uncultivated land due to agricultural intensification, which has diminished food availability such as wild plants, seeds and insects; with hunting also playing a significant role.²²



LC

VU

Red-necked Grebe *Podiceps grisegena*

The overall population decline by 38% is considered genuine and is observed in most of the key range countries, including Russia and Finland (which hold together over half the European population). Despite improved knowledge about the species population in some range countries, little is known about the reasons for change.



LC

VU

Black-necked Grebe *Podiceps nigricollis*

An overall genuine decline by 35% was calculated based on declining trends from Russia and Ukraine (which together hold about 70% of the European population). Little is known about the main reasons of the decline, although water pollution from agricultural activities and urban run-off, decreases in precipitation due to climate change, the construction of flood protection infrastructure, disturbance from human recreational activities and bycatch may all be contributing factors.²³

²¹ Keller, V., Herrando, S., Voříšek, P., Franch, M., Kipson, M., Milanese, P., Martí, D., Anton, M., Klvaňová, A., Kalyakin, M.V., Bauer, H.-G. and Foppen, R.P.B. (2020). *European Breeding Bird Atlas 2: Distribution, Abundance and Change*. European Bird Census Council & Lynx Edicions, Barcelona.

²² Heath, M. and Tucker, G. (1994). *Birds in Europe*. World Birdwatch 16: 9-13.

²³ Pressures and threats data reported as part of the EU Birds Directive Article 12 reporting exercise 2013-2018 <https://cdr.eionet.europa.eu/>

In addition, 13 species have been assessed for first time at European level (see Table 2), mainly due to taxonomic changes since 2015. Examples are the newly recognised Cyprus Scops-owl (*Otus cypricus*) (Least Concern) and the Italian Sparrow (*Passer italiae*) (Vulnerable). At EU level, 15 species were assessed for first time.

Table 2. Species assessed at European and EU level for the first time (shown in taxonomic order).

SCIENTIFIC NAME	ENGLISH NAME	2021 RED LIST STATUS
<i>Gallinago megala</i>	Swinhoe's Snipe	NT
<i>Onychoprion fuscatus</i>	Sooty Tern	VU
<i>Otus cypricus</i>	Cyprus Scops-owl	LC
<i>Phylloscopus nitidus</i>	Green Warbler	LC
<i>Phylloscopus orientalis</i>	Eastern Orphean Warbler	LC
<i>Curruca balearica</i>	Balearic Warbler	LC
<i>Curruca crassirostris</i>	Eastern Bonelli's Warbler	LC
<i>Curruca subalpina</i>	Moltoni's Warbler	LC
<i>Cyanistes teneriffae</i>	African Blue Tit	LC
<i>Poecile hyrcanus</i>	Caspian Tit	LC
<i>Lanius meridionalis</i>	Iberian Grey Shrike	VU
<i>Passer italiae</i>	Italian Sparrow	VU
<i>Fringilla polatzeki</i>	Gran Canaria Blue Chaffinch	EN

3.3 Status and trends of European birds by taxonomic and functional group

There are 73 bird families occurring in Europe and there are considerable differences amongst them, both in the number of species they include and their threatened status. When merging families into taxonomic or functional groups, amongst those comprising the most species; seabirds, wildfowl, waders, and raptors have highest proportion of threatened and Near Threatened species. In slightly different order - wildfowl, waders, raptors and seabirds, these are the most threatened groups of birds also in the EU. When looking at trends in Europe, over 40% of wildfowl and wader species, over 30% of seabirds, as well as 25% of raptors, are experiencing population declines. Within the EU, the proportion increases to over 50% for waders and wildfowl, and more than 30% for raptors. Worryingly,

deterioration in trend is observed amongst the most widespread and numerous group of birds, the passerines, with one in four species in this group having a decreasing population trend in Europe, while in the EU the ratio increases to one in three. Among these, larks, shrikes and buntings, as well as pipits, chats, flycatchers, finches, tits and sparrows, contain relatively high numbers of species with decreasing trends (Fig. 5).

Less speciose taxonomic groups also show significant population declines in Europe, e.g. kingfishers (all 3 species), bustards (3 out of 4 species), rails and crakes (5 out of 9 species), grebes (3 out of 5 species) and grouse, partridges and pheasants (7 out of 17 species).

A comparison between passerines, terrestrial non-passerines and species dependent on the marine habitat during most of their life cycle shows that passerines and marine species have a higher proportion of species with unknown trends (Fig. 6). Similar results are obtained at EU level.




Figure 5. Population trend of passerine birds by family (in brackets, total number of species per family, in graph, number of species per trend direction). Results at European level.




Figure 6. Differences in proportion of species trends between passerines, non-passerines (excl. seabirds, diving ducks, loons and cormorants) and non-passerines associated with the marine environment (seabirds, diving ducks, loons and cormorants) in Europe. In brackets, number of species per group, in graph, proportion of trend direction per group.

A relatively higher proportion of declining trends are observed among migratory birds (more than 30%) compared to resident species (20%) (see Fig. 7). Moreover, among 121 species considered sub-Saharan migrants²⁴, 33% have declining trends, while among species that are migratory but do not migrate as far as to sub-Saharan Africa; 31.5% have declining trends.




Figure 7. Trend directions in relation to migrating behaviour (in brackets, total number of species per group, in graph, proportion of species per trend direction). Results at European level.

3.4 Status and trends of European birds by main type of habitat

A comparison between main habitat types in Europe and the EU shows that farmland and grassland habitats have the highest number of associated threatened and Near Threatened species (33 and 40 species, respectively in Europe and the EU) of all habitat types, and are closely followed by marine habitats (28 and 32 species, respectively), and inland wetlands next, with 13 and 17 species, respectively (Fig. 8).




Figure 8. Number of threatened and Near Threatened species per main type of habitat in Europe (brown) and the EU (grey).

²⁴ Vickery, J.A., Ewing, S.R., Smith, K.W., Pain, D.J., Bairlein, F., Škorpirová, J. and Gregory, R.D. (2014). The decline of Afro-Palaeartic migrants and an assessment of potential causes. *Ibis*, 156(1), pp.1-22. <https://onlinelibrary.wiley.com/doi/full/10.1111/ibi.12118>

The picture looks similar when comparing the proportion of species with different European population trend directions by habitat type (Fig. 9), where at least a third of species associated with farmland and grassland (55 out of 158 species), wetlands (24 out of 70 species) and marine habitats (30 out of 92 species) have a decreasing trend. Within the EU, the number of species with decreasing trend in each of these three habitat types is even higher, reaching over 40%.

In Europe marine and freshwater aquaculture, artificial ponds, irrigated land and wastewater areas appear to have the highest proportion of species with a decreasing trend (over 45%), however a relatively small number of species (19) are associated with this type of habitat.

At the same time, another circa 40% of all species associated with marine habitats and wetlands have a stable or increasing trend. However, the proportion of species with an increasing trend is the smallest compared to all other habitat types (excluding rocky areas where only one species falls within this category). The results are valid also at EU level.

Over a quarter of all species associated with forest habitats have decreasing trends (22 out of 80 species), although this habitat type also has the highest proportion of stable (25%) and increasing (25%) trends, with the rest being unknown. Similarly, approximately 50% of all species associated with shrubland have either stable or increasing trends. Interestingly, forest and shrubland habitats in the EU have significantly higher number of species with either stable or increasing trend (equal to 60%), compared to those in Europe and to other habitat types in the EU.

Rocky habitats like inland cliffs and mountain peaks have the highest proportion of species with unknown trends (more than 50%), both in Europe and the EU. A third of all species associated with marine habitats and shrubland also have an unknown trend. This proportion is much smaller in the EU (less than 20%). The smallest proportion of species with unknown trends is associated with artificial aquatic habitats (less than 20%), farmland and grassland, and forest habitats (circa 20% each). In the EU, the lowest number of species with unknown trend is related to inland wetlands, farmland and grassland, and forest habitats (c. 10% each).




Figure 9. Population trend by main type of habitat in Europe (in brackets, total number of species per group, in graph, number of species per trend direction).

The results confirm the overall conclusions observed in other studies, such as the State of Nature in the EU 2013-2018 or the IPBES 2019. Large scale land use changes and the intensification of agricultural practices, the development of infrastructure, the overexploitation of marine resources, the pollution of inland waters and unsustainable yet commonly used forestry practices are the main drivers of bird population declines observed in European habitats. The Pan-European Common Bird Monitoring Scheme indicates a 57% decline in common farmland birds between 1980 and 2018²⁵. Ongoing population declines and range contractions in widespread species of open

habitats – like larks, shrikes and buntings – clearly demonstrate the impact of disappearing mosaic landscapes and the increased use of agricultural chemicals (see Box 2 and Fig. 6). Although some raptor species have made recent population recoveries due to legal protection and targeted conservation activities²⁶, many raptor species dependent on open habitats (e.g., grassland and shrubland) for prey are still in decline. In the marine environment, birds suffer from bycatch and incidental killing (due to fishing and hunting activities), as well as from the impact of invasive non-native species, disturbance and water pollution (originating from both land and sea).

²⁵ https://pecbms.info/trends-and-indicators/indicators/indicators/E_C_Fa/

²⁶ <https://www.eea.europa.eu/publications/state-of-nature-in-the-eu-2020>

Many species of birds are legally hunted across Europe. While hunting is prohibited in the EU during the breeding season, this method of exploitation is a threat to at least 82 species listed in Annex II of the Birds Directive²⁷ during their migration or in winter. Additionally, some species fall under the so-called EU derogations, where countries are permitted to cull part of the population under defined conditions. Species like the Rook

(*Corvus frugilegus*) have suffered massive population declines, likely due to permitted persecution²⁸ (killing, destruction of nests and roosting sites) across most of its range, which has resulted in its current re-assessment as Vulnerable. Illegal killing is another major threat to European birds, with more than 25 million birds illegally slaughtered in the Mediterranean region each year²⁹.

Box 2. Specialists of open habitats: larks, shrikes and buntings



Woodchat Shrike (*Lanius senator*)
Photo: Juan Emilio

The 2021 Red List assessment of European birds shows that more species of open habitats, like larks, shrikes and buntings, experience population decline compared to the previous update (2015).

Of the 6 species of shrikes in Europe, 5 have a decreasing Red List trend, which has resulted in one species classified as Vulnerable (Iberian Grey Shrike (*Lanius meridionalis*)) and one as Near Threatened (Woodchat Shrike (*Lanius senator*)).

Shrikes prefer open small-scale (mosaic) landscapes, usually with scattered trees or shrubs, and have adapted (to a different degree) to low-intensity cultivated areas like farmland and grassland. Land use changes and the intensification of agricultural practices have led to the loss of areas of natural vegetation and field margins, as well as a massive decline in the abundance of large insects³⁰. Land abandonment resulting in natural succession processes have been identified as another driver for disappearing breeding habitats for shrikes in some parts of the European continent. Climatic factors are expected to significantly influence shifts in population distribution and abundance in Red-backed Shrike (*Lanius collurio*) and Masked Shrike (*Lanius nubicus*).

Two thirds of all 13 lark species occurring in Europe and more than half of the 16 species of buntings also have decreasing population trends. The populations of two widespread lark species, the Eurasian Skylark (*Alauda arvensis*) and Calandra Lark (*Melanocorypha calandra*), as well as the Yellowhammer (*Emberiza citrinella*), Corn (*Emberiza calandra*), Ortolan (*Emberiza hortulana*) and Rustic Buntings (*Emberiza rustica*) have undergone significant declines in the past decades. Similarly, these species depend on a mosaic of open habitats, and their declines are linked to habitat loss and/or degradation generally related to agricultural intensification.³¹

In addition, the compounding effects of drivers such as climate change, land-/sea-use change, overexploitation of resources, pollution and invasive alien species are likely to exacerbate the negative impacts on nature, as seen in different ecosystems³². Climate change, which can alter the habitats or seasons within a species natural range, or cause the displacement or disappearance of prey species, is a rather unpredictable variable when considering species population changes. For example, sedentary species confined to mountain areas, like the Caucasian Grouse (*Tetraogallus caucasicus*), may be driven to near extinction in the upcoming 30-50 years, as a result of shrinking alpine and sub-alpine habitats³³. Migratory species in

their breeding season, particularly Arctic breeding waders are among the species with the smallest proportion of their current range projected to be climatically suitable by 2050³⁴. On the other hand, a recent study using the comprehensive European Breeding Birds Atlas 2 (EBBA2) data to understand birds' range dynamics in the last decades in relation to environmental drivers and species' traits states that climate change is not (yet) a major driver behind the observed changes in bird distributions in Europe in recent years³⁵. This means that the conservation actions needed for species sensitive to its effects are very difficult to determine without further research and understanding of the issue.

²⁷ https://ec.europa.eu/environment/nature/legislation/birdsdirective/index_en.htm

²⁸ Krüger, T., Heckenroth, H., Prior, N. et al. Persecution and statutory protection have driven Rook *Corvus frugilegus* population dynamics over the past 120 years in NW-Germany. *J Ornithol* 161, 569–584 (2020). <https://doi.org/10.1007/s10336-020-01750-3>

²⁹ Brochet, A., Van den Bossche, W., Jbour, S., Ndang'ang'a, P., Jones, V., Abdou, W., ... Butchart, S. (2016). Preliminary assessment of the scope and scale of illegal killing and taking of birds in the Mediterranean. *Bird Conservation International*, 26(1), 1-28. doi:10.1017/S0959270915000416.

³⁰ Keller, V., Herrando, S., Voříšek, P., Franch, M., Kipson, M., Milanese, P., Martí, D., Anton, M., Klvaňová, A., Kalyakin, M.V., Bauer, H.-G. and Foppen, R.P.B. (2020) European Breeding Bird Atlas 2: Distribution, Abundance and Change. European Bird Census Council & Lynx Edicions, Barcelona.

³¹ Suarez et al. (1997) Newton 2017 after EBBA2.

³² https://ipbes.net/sites/default/files/2020-02/ipbes_global_assessment_report_summary_for_policymakers_en.pdf

³³ Anouschka R. Hof, Andrew M. Allen, An uncertain future for the endemic Galliformes of the Caucasus, *Science of The Total Environment*, Volume 651, Part 1, 2019, Pages 725-735, ISSN 0048-9697, <https://doi.org/10.1016/j.scitotenv.2018.09.227>.

³⁴ Nagy, S., Breiner, F., Anand, M., Butchart, S., Flörke, M., Fluet-Chouinard, E., ... Voltz, O. (2021). Climate change exposure of waterbird species in the African-Eurasian flyways. *Bird Conservation International*, 1-26. doi:10.1017/S0959270921000150.

³⁵ Aleksí Lehikoinen et al. (2021). Local colonisations and extinctions of European birds are poorly explained by changes in climate suitability (unpublished).

3.5 Gaps in knowledge

Red Lists are a dynamic tool that evolves with time as species are re-assessed according to new information or situations. They are aimed at promoting and supporting research, monitoring and conservation actions at local, regional and international levels; especially for threatened, Near Threatened and Data Deficient species. Through the process of compiling data for the European Red List of Birds, a number of knowledge gaps have been identified.

Europe has a long history of data collection and environmental surveillance. Many countries have established surveying and monitoring programmes (e.g. PECBMS³⁶ and the International Waterbird Census³⁷), and contribute to international data collection projects (e.g. European Bird Portal³⁸), leading to the region having a comparatively good biodiversity knowledge database. This is an advantage, as recent and good quality data are essential to successfully inform and address conservation issues across the continent.

Despite this, Europe's knowledge base about birds still has room for improvement. Across Europe there are significant geographic, geopolitical and taxonomic biases in the quality of data available on the distribution and status of species. Unless they are part of a specific monitoring or conservation programme, surveys are often irregularly undertaken or species are selectively surveyed for, often depending on their conservation priority – a factor which is often based on the existing knowledge of a species. The population size and trend data reported during the time of this project were found to sometimes be old and re-used, or were based solely on expert opinion, again, due to lack of recent data availability, sometimes with the most recent sources using data dating back to the 1990s or early 2000s. This means that there exists a certain proportion of species for which the information is too old or uncertain, or for which there is not enough information at all to be able to calculate a realistic trend. In these cases, the trend is labelled as unknown. In Europe, although the data may be old or of poor quality, there still is enough information to allow the assessment of all bird species, even those with unknown trends,

without the use of the Data Deficient category. However, in a few circumstances, this can result in an assessment in which the lack of trend information leads to a default Least Concern assessment, whereas if the trends were known, the species may actually be Near Threatened or threatened.

When data is lacking, it is often evident that for some countries this is because of a shortage of resources for regular monitoring of bird populations (e.g. time, personnel and/or financial resources) which would allow for better coverage of species and areas. The lack of information could also be due to poor cooperation between governments, institutions and organisations. In some rare cases, data may be scarce due to the species being present in inaccessible places, although with today's technology a solution is likely possible. Context is an important factor in understanding the reasons for changes in a species' population size trends or range. Understanding the causes of a species' decline or recovery can help analyse the key threats to birds across Europe. Nonetheless, currently this tends to be a secondary priority in terms of data reporting, and more research into this subject is needed to fully understand the various impacts of human activity on nature.

To counter these issues, there is a need for more coordinated national and international efforts in bird monitoring, to ensure the monitoring cycles in different countries concur with each other and, ideally, with the timings of the European Red List of Birds updates. These survey programmes need to be coordinated and communicated across organisations, institutions and governmental bodies in order to ensure that information is transferred and that everyone is working together towards the same goals. To support these surveys, dedicated resources should be available at local, national and international levels, with policy tools created enabling the easy access to these resources and encouraging the collection of data. The EU legal and financial mechanisms, for example, already play a crucial role in this process, such as the reporting of national data under the Birds and Habitats Directives, and the development of the LIFE Programme, which enables the access to funding for conservation-based projects.

³⁶ <https://pecbms.info/>

³⁷ <https://www.wetlands.org/knowledge-base/international-waterbird-census/>

³⁸ <https://eurobirdportal.org/ebp/en/#home/HIRRUS/r52weeks/CUCCAN/r52weeks/>

4.

Biodiversity protection in Europe

Europe has a long history of nature protection. From decrees to protect sovereigns' own hunting grounds during the Middle Ages to the modern-day democratic international conventions and agreements; environmental law has always aimed at preserving nature for its socio-economic and intrinsic value.

In Europe, environmental legislation is created and implemented at different geographic and administrative levels. National and regional biodiversity laws set up the extent and means for protection of species, habitats and other natural resources. Different systems of protected areas, e.g., national parks, nature reserves or protected landscape monuments exist in each country. To put the protection of these areas in practice, site-specific management or action plans are often developed in collaboration with local authorities and stakeholders. Similarly, action plans for the conservation of priority (e.g., nationally, European or globally threatened) species may be developed. Other types of national or local law usually define the protection and management of particular parts of the environment, such as rivers and lakes, marine or coastal areas, forests etc.

Nature is a continuum, which political borders cross and often fragment, as do different human activities. Therefore, international cooperation is crucial for the successful protection of biodiversity at an appropriate scale. Most European countries and all EU member states are signatories to a number of relevant international conventions, including the Bern Convention on the Conservation of European Wildlife and Natural Habitats (1979), the Convention on Biological Diversity (1992) and the African-Eurasian Migratory Waterbird Agreement (AEWA). The international trade of a small number of European bird species is regulated under the Convention on International Trade in Endangered Species (CITES).

The Bern Convention is a binding international legal instrument that aims to conserve wild flora and fauna and their natural habitats. It covers all European countries and some African states. It aims to increase cooperation between contracting parties and to regulate the exploitation of species (including migratory species). Key provisions of the Convention include

the establishment of protected areas (defined as the Emerald network), the protection of breeding and resting sites and the regulation of disturbance, capture, killing and trade of wild species. The Bern Convention has played a fundamental role in promoting the adoption of International Species Action Plans on European threatened or Near Threatened species.

The African-Eurasian Migratory Waterbird Agreement (AEWA) is an international treaty for the conservation of migratory waterbirds, developed under the auspices of the Convention on Migratory Species (CMS, or Bonn Convention). Among other obligations, Parties to the Agreement are required to adopt so-called International Single Species Action Plans (SSAPs) for species of particular concern (i.e., species/populations with an unfavourable conservation status). These SSAPs cover all Range States relevant for the species to ensure a coordinated conservation approach along the entire flyway.

The Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES), adopted in 1973, has 178 countries that are Parties to the Convention. The Convention aims to protect species from the detrimental effects of international trade by establishing an international legal framework for preventing or controlling trade. Species listed on Appendix I of the Convention are considered to be threatened with extinction and are not allowed to be traded commercially, while those on Appendix II are only allowed to enter international trade under specific controlled circumstances (including a licensing system). Parties are obliged to develop national legislation effectively implementing the obligations of the Convention, including setting sustainable quotas for Appendix II species. Currently, 687 species of fauna, including 155 bird species are listed on Appendix I and more than 5,000 species of fauna, including 1,279 bird species, on Appendix II³⁹. CITES is implemented in the EU through a set of Regulations known as the EU Wildlife Trade Regulations⁴⁰.

The Convention on Biological Diversity is an international legal instrument for the conservation and sustainable use of biological diversity. Parties have to develop national biodiversity strategies or action plans (NBSAPs). They also need to agree on

³⁹ <https://cites.org/>

⁴⁰ https://ec.europa.eu/environment/cites/legislation_en.htm

a ten-year global strategic plan with concrete targets in order to achieve the overall objectives of the Convention. At the time of publication of this report, world leaders are developing the new Post-2020 Global Biodiversity Framework with a package of commitments to address and reverse global biodiversity loss by 2030⁴¹.

In the EU, nature conservation policy is based on two main pieces of legislation: the Birds Directive of 1979 and the Habitats Directive of 1992, both often referred to as the EU Nature Directives. The Birds Directive provides a binding, legal framework for the protection of all wild birds in the EU, including their eggs, nests and habitats. EU Member States need to ensure that a sufficient area and diversity of habitats is available for all wild bird species, including those that live on farmland and in urban environments. The most important areas for threatened birds on Annex I of the Birds Directive and for regularly occurring migratory birds should be designated as Special Protection Areas (SPAs). The Birds Directive also regulates the hunting of birds in the EU, restricting the hunting seasons and methods, as well as the species that can be hunted (listed in Annex II).

The Habitats Directive is Europe's most powerful tool to address habitat protection, listing the priority habitats (e.g. specific types of wetlands, meadows and marine habitats) on Annex I and species on Annex II. As with the Birds Directive, important sites under the Habitats Directive should be designated as Special Areas of Conservation (SACs). These form, together with the SPAs, the Natura 2000 network: the largest coordinated network of protected areas in the world. It stretches across all Member States and covers over 18% of the EU's land area and more than 6% of its sea territories⁴². Each Member State is required to identify sites of European importance for Natura 2000 and is encouraged to put in place special management plans, combining long-term conservation objectives with economic and social activities.

The EU Marine Strategy Framework Directive (MSFD)⁴³ aims to protect the marine environment across Europe more effectively. It requires Member States to achieve Good Environmental Status (GES), following an ecosystem-based approach, focused on 11 descriptors related to ecosystem features, human drivers and pressures⁴⁴. The EU Maritime Spatial Planning (MSP) Directive aims to manage the use of our seas and oceans coherently, and to ensure that human activities take place in an efficient, safe and sustainable way, guaranteeing the protection of nature at sea. In addition, Europe's marine environment is protected

under regional conventions like the Baltic Marine Environment Protection Commission⁴⁵ (HELCOM), the OSPAR Commission⁴⁶, Conservation of Arctic Flora and Fauna⁴⁷ (CAFF), the Commission on the Protection of the Black Sea Against Pollution⁴⁸.

The majority of Europe's threatened bird species are listed on the Birds Directive Annexes and the Bern Convention Appendices.

Besides the above-mentioned nature protection legal frameworks, several other sectoral policies exist in the EU with certain measures and practices that aim to protect nature. Most relevant for the agricultural environment is the EU's Common Agricultural Policy⁴⁹ (CAP), with its 'agri-environment and climate measures' under Pillar II. However, these are of very varying quality and a review by the European Court of Auditors in 2020 found that 'most CAP funding has little positive impact on biodiversity'⁵⁰. For marine species and habitats, together with the Marine Strategy Framework Directive, the EU's Common Fisheries Policy (CFP) aims at setting rules for sustainably managing European fishing fleets and conserving fish stocks⁵¹. Inland surface waters, coastal waters and groundwater are protected under the framework of the EU Water Framework Directive. It aims to prevent and reduce pollution, promote sustainable water use, protect and improve the aquatic environment and mitigate the effects of floods and droughts.

Within a renewed political effort to address biodiversity loss, the European Union responded to its global commitments by developing a 10-year Biodiversity Strategy. In May 2020, a new comprehensive and ambitious EU 2030 Biodiversity Strategy was put forward by the European Commission and largely endorsed by EU legislators. Its aim is to put Europe's biodiversity on a path to recovery with specific actions and commitments⁵². These include the development of a new nature restoration law, a Forest Strategy (published in July 2021), an action plan to conserve fisheries resources and protect marine ecosystem, improved financing for biodiversity measures in the EU, and an update of the Directive of Sustainable Use of Pesticides. The strategy sets targets for the non-deterioration of all species and habitats, as well as significant improvement of at least 30% of species and habitats. It also urges Member States to assure the protection of 30% of land and 30% of sea by 2030, and strictly protect at least 10%. The agreement on the long-term EU Budget from 2021 to 2027 includes a commitment to spend at least 7.5% of the EU Budget on biodiversity as of 2025, and 10% as of 2026.

⁴¹ www.cbd.int

⁴² https://ec.europa.eu/environment/nature/index_en.htm

⁴³ <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex%3A32008L0056>

⁴⁴ https://ec.europa.eu/environment/nature/index_en.htm

⁴⁵ <https://helcom.fi/>

⁴⁶ <https://www.ospar.org/>

⁴⁷ <https://www.caff.is/>

⁴⁸ http://www.blacksea-commission.org/_convention.asp

⁴⁹ https://ec.europa.eu/info/food-farming-fisheries/key-policies/common-agricultural-policy/cap-glance_en

⁵⁰ https://www.eca.europa.eu/Lists/ECADocuments/SR20_13/SR_Biodiversity_on_farmland_EN.pdf

⁵¹ https://ec.europa.eu/oceans-and-fisheries/policy/common-fisheries-policy-cfp_en

⁵² https://ec.europa.eu/environment/strategy/biodiversity-strategy-2030_en

5.

Recommendations and future work

5.1 Recommendations for policy and practice

The existing biodiversity protection framework in Europe (see Section 4) has contributed, directly or indirectly, to the successful recovery of many threatened species and protection and restoration of key habitats. These have arisen thanks to the establishment of a network of protected areas, collaboration to conserve sites and to take action for species often across borders. Yet, this collective effort, while essential, has to date been insufficient to reverse the overall decline in our continent's biodiversity. Sadly, many success stories have been limited to the local level. For certain species, such actions may be enough to maintain stable or even improving populations. The decline of many widespread and common species (e.g., birds and butterflies) signals, however, a problem of a much larger scale, a systemic issue in the functioning of ecosystems caused by a diverse array of human-induced changes. Therefore, future conservation efforts in Europe should not only focus on action locally but also seek to deliver more fundamental change at a bigger scale. Collaboration between decision-makers and conservation practitioners needs to be expanded while securing greater involvement from consumers, producers, landowners and other businesses.

As intergovernmental reports on climate change and biodiversity have recommended^{53,54}, we need to rethink the way we produce, trade, consume and reuse our vital resources. Ultimately, as humans, we depend on a healthy natural environment (including irreplaceable services and functions such as the provision of clean water, pollination of crops, flood protection, and a stable climate; which are also fundamental for our recreation and well-being). Without the services provided by the natural environment, our health, society and economy would not hold up. Thus, a revision of the way we develop and implement sectoral policies will be

key to future success. For example, a review by the European Court of Auditors in 2020 found that "most CAP funding has little positive impact on biodiversity"⁵⁵, while another report from 2020 published by the same institution⁵⁶ stated that "overall, while a framework was in place to protect the marine environment, the EU's actions had not restored seas to good environmental status, nor fishing to sustainable levels in all seas". This was also supported by the European Environment Agency, which identified in 2020 that "Marine biodiversity remains under threat in Europe's seas. A high proportion of marine species and habitats' assessments continue to show an 'unfavourable conservation status' or a status that is 'unknown'⁵⁷. Similar critiques of progress can be found outside of the EU, for example in the UK State of Nature Report⁵⁸.

Through its Green Deal, the EU has expressed the ambition to tackle these current environmental and socio-economic challenges in a transformative way. It aims to transform the EU into a modern, resource-efficient and competitive economy, leading by example towards being the first climate-neutral continent⁵⁹. The EU commits to combat the intrinsically linked biodiversity and climate crises with its 2030 Biodiversity Strategy, as part of the Green Deal, for the broader European continent and the world. In addition, 88 Heads of State (including 14 outside the EU) have signed up to the Leaders' Pledge for Nature⁶⁰ to step up global ambition for biodiversity. This provides momentum for securing an ambitious new global biodiversity framework for 2030, set to be approved by a Conference of the Parties to the UN Convention on Biological Diversity in Kunming, China in April 2022.

To create a different future, where nature is thriving in harmony with people, where common species remain common and fewer species are at risk of extinction, governments across Europe need to develop coherent regional and national biodiversity strategies including the actions outlined in the next pages:

⁵³ <https://www.ipcc.ch/2019/>

⁵⁴ <https://ipbes.net/global-assessment>

⁵⁵ https://www.eca.europa.eu/Lists/ECADocuments/SR20_13/SR_Biodiversity_on_farmland_EN.pdf

⁵⁶ <https://www.eca.europa.eu/en/Pages/DocItem.aspx?did=57066>

⁵⁷ <https://www.eea.europa.eu/publications/marine-messages-2>

⁵⁸ <https://nbn.org.uk/wp-content/uploads/2019/09/State-of-Nature-2019-UK-full-report.pdf>

⁵⁹ https://ec.europa.eu/info/strategy/priorities-2019-2024/european-green-deal_en

⁶⁰ <https://www.leaderspledgefornature.org/>

Conservation ambition and action

- Translate global targets for nature's recovery into EU and domestic law.
- Deliver an expanded and better managed network of protected areas (bolstering the existing Emerald and Natura 2000 networks) consistent with global ambition for 30% of land and sea to be protected and well managed by 2030. Protected areas (PAs) must include substantial areas under strict protection such as 'no take' marine protected areas and 'no logging' forests allowed to fully recover to near pristine state, as proposed by the EU biodiversity strategy target of putting a third of PAs under strict protection.
- Improve and expand targeted conservation actions for threatened and declining species, populations and habitats.
- Prioritise the large-scale restoration of carbon-rich ecosystems to deliver benefits for both wildlife and the climate. There should be a particular focus on grassland, peat, forest and coastal habitats.
- Ensure that carbon sequestration always contributes to ecosystem and biodiversity health and resilience, while adaptation policies to climate change prioritise nature-based solutions and are synergistic with ecosystem restoration.
- Enhance the collaboration across borders (including nations in sub-Saharan Africa) to support the conservation of long-distance migratory birds.

Funding

- Secure adequate funding from national, EU and international levels to support conservation objectives fulfilling, for example, the EU commitment to spend 10% of the EU Budget on biodiversity.
- Provide specific funding to developing countries, especially in sub-Saharan Africa to support conservation of migratory birds on their wintering grounds.
- End perverse subsidies that harm nature (for example by overhauling agriculture policy and reforming agriculture incentives to reward wildlife-friendly farming to support the recovery of farmland birds), while ensuring that nature objectives are embedded in other funding schemes (for example European Maritime Fisheries and Aquaculture Fund and Neighbourhood Development and International Cooperation Instrument).

Regulation and enforcement for sustainable management of land and sea

- To support the recovery of farmland birds, introduce better regulation and law enforcement concerning agricultural practices that are harmful to the environment. In particular, policies should massively reduce the use of pesticides, nitrogen pollution, water over abstraction and simplification of landscapes. Farm policies must support a transition towards agro-ecological practices, and crucially, the provision of a minimum of 10% of landscape features, native vegetation and ecological infrastructure at farm level (as foreseen by the EU Biodiversity Strategy, for example).
- To support the recovery of threatened forest bird species, enable the full and strict protection of vulnerable forests (e.g. primary and old-growth forests), support a transition of forestry towards "close to nature forest management" and champion ecologically compatible afforestation policies (applying the principle of "the right trees in the right places for the right reasons", as indicated in the new EU Forest Strategy).
- Roll out effective policies addressing unsustainable consumption that drives agricultural forestry and fisheries expansion and intensification. Subsidies that boost consumption (e.g. for burning biomass or promoting of meat and dairy) should be removed, waste prevention and recycling must be promoted through circular economy legislation, and socially sensitive policies should support a dietary shift away from animal consumption.
- To support seabird conservation, evolve fisheries policies to eliminate bycatch and restore food chains, while also improving biosecurity measures and taking action to reduce the threat posed by invasive non-native species especially on seabird breeding grounds.
- To support both migratory bird and raptor conservation, tackle the illegal and legal, unsustainable hunting and persecution (e.g. poisoning) through improved enforcement of national legislation within and beyond the European continent.
- To reduce future pressures on species, ensure energy (including renewable energy) and other infrastructure development are designed, sited and built in harmony with nature, without adversely impacting birds and their habitats.

Monitoring and reporting

- Continue standardised regular monitoring of birds and other taxa, while addressing knowledge gaps (see Section 3.5), to ensure that changes are well understood and measures are taken in a timely manner to prevent further deterioration.
- Continue monitoring bird species with observed or projected population declines, including common species associated with farmland or forests and considered key indicators of the state of the environment (see Sections 3.3 and 3.4).
- Ensure transparent reporting of progress of EU and national action to encourage scrutiny, accountability and to adapt policies as necessary to improve future conservation success.

5.2 Conclusions

The results of this report show that while there is progress made in the conservation of some European birds, it is clear that if we want to achieve the global targets to 2030, we need an immediate action at a continental scale. While certain threats, e.g., persecution can be tackled relatively easily through legal regulations and implementation, the systemic threats observed in the natural environment (e.g. unsustainable agriculture, forestry or fisheries practices) will require a change that:

1. Is transformative, starting from the way we source prime materials, through transportation, production, discarding and recovering resources.
2. Involves all stakeholders – from consumers to producers, landowners and other businesses.
3. Ensures the protection of ecosystems which are still in a good status and the restoration of degraded ones, through adequate implementation, funding and research.

5.3 Future work

The data used for the European Red List of Birds gives a general overview of the threats to birds in Europe. Furthermore, considering that threat analysis can be very complex and difficult to interpret, a more detailed analysis should be the purpose of separate research, perhaps addressed by taxonomic group, type of habitat or geographic region.

Within the scope of further analyses should be the link between the extinction risk and various threat factors for European species breeding in the Arctic, migratory birds, seabirds, species inhabiting the alpine and sub-alpine areas, as well as forest specialists.

The differentiation between the direct and indirect impact of agricultural practices on bird populations should help direct practical measures to benefit nature at systemic level.

In this regard, assessing the differences between the main driving factors of species declines within the territory of the

European Union and beyond (as a potential consequence of the different legal and enforcement processes) could provide valuable insights into practices that benefit biodiversity as opposed to those that harm it.

The impact of climate change or invasive non-native species as threats with likely significant impact on species in the future, should be better understood and monitored.

These and many other research topics can be delivered through the data gathered under the European Red List of Birds, while they could largely benefit from a joint approach with other data sources, including spatial analysis, statistical modelling and citizen science.

The European Red List of Birds is part of a wider initiative aimed at assessing the status of European species. Together with other sources like the European Breeding Bird Atlas 2 or the Common Bird Monitoring Scheme, it provides a key resource for decision-makers, policymakers, resources managers, environmental planners and NGOs. It has gathered large amounts of data on the population, ecology, habitats and threats of each bird species. These data are freely available on the IUCN Red List website (<https://www.iucnredlist.org/regions/europe>), on the European Commission's website (<http://ec.europa.eu/environment/nature/conservation/species/redlist>) and through paper publications.

The European Red List of Birds will also form the basis for the identification of Species of European Conservation Concern (SPEC), following the methodology developed in the previous assessments (Tucker & Heath 1994, BirdLife International 2004, BirdLife International 2017). It will also be used (depending on the availability of resources) to calculate the Red List Index for European Birds which measures the projected overall regional extinction risk of sets of species, and tracks changes in this risk⁶¹.

The results presented in this report can be applied to inform policy and identify priority species to include in research and monitoring programmes. It can also serve in the process of measuring progress towards national and international biodiversity targets.

⁶¹ <https://www.eea.europa.eu/data-and-maps/indicators/red-list-index-for-european-species/red-list-index-for-european/?searchterm=None>

References

REPORTS

- BirdLife International (2015). *European Red List of Birds*. Luxembourg: Office for Official Publications of the European Communities. <http://datazone.birdlife.org/info/euroredlist>
- BirdLife International (2017). *European birds of conservation concern: populations, trends and national responsibilities* Cambridge, UK: BirdLife International.
- European Environment Agency (2020). *State of nature in the EU - Results from reporting under the nature directives 2013-2018*. EEA. 142 pp. <https://www.eea.europa.eu/publications/state-of-nature-in-the-eu-2020>
- Hayhow D.B., Eaton M.A., Stanbury A.J., Burns F., Kirby W.B., Bailey N., Beckmann B., Bedford J., Boersch-Supan P.H., Coomber F., Dennis E.B., Dolman S.J., Dunn E., Hall J., Harrower C., Hatfield J.H., Hawley J., Haysom K., Hughes J., Johns D.G., Mathews F., McQuatters-Gollop A., Noble D.G., Outhwaite C.L., Pearce-Higgins J.W., Pescott O.L., Powney G.D. and Symes N. (2019). *The State of Nature 2019*. The State of Nature partnership <https://nbn.org.uk/wp-content/uploads/2019/09/State-of-Nature-2019-UK-full-report.pdf>
- Heath, M. and Tucker, G. (1994). *Birds in Europe*. World Birdwatch 16: 9-13.
- IPBES (2019). *Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. E. S. Brondizio, J. Settele, S. Díaz, and H. T. Ngo (editors). Bonn, Germany: IPBES secretariat. 1148 pp. <https://doi.org/10.5281/zenodo.3831673>
- IPBES (2019). *Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. S. Díaz, J. Settele, E. S. Brondizio, H. T. Ngo, M. Guèze, J. Agard, A. Arneeth, P. Balvanera, K. A. Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M. Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S. Polasky, A. Purvis, J. Razzaque, B. Reyers, R. Roy Chowdhury, Y. J. Shin, I. J. Visseren-Hamakers, K. J. Willis, and C. N. Zayas (eds.). Bonn, Germany: IPBES secretariat. 56 pages. <https://doi.org/10.5281/zenodo.3553579>
- Leyrer, J., Brown, D., Gerritsen, G., Hötter, H. and Ottvall, R. (compilers). (2018) *International Multispecies Action Plan for the Conservation of Breeding Waders in Wet Grassland Habitats in Europe (2018-2028)*. Report of Action A13 under the framework of Project LIFE EuroSAP (LIFE14 PRE/UK/002). NABU, RSPB, VBN and SOF.

GUIDELINES

- IUCN (2012). *Guidelines for Application of IUCN Red List Criteria at Regional and National Levels: Version 4.0*. Gland, Switzerland and Cambridge, UK: IUCN. iii + 41 pp
- IUCN (2012). *IUCN Red List Categories and Criteria: Version 3.1*. Second edition. Gland, Switzerland and Cambridge, UK: IUCN. iv + 32pp.
- IUCN Standards and Petitions Committee (2019). *Guidelines for Using the IUCN Red List Categories and Criteria. Version 14*. Prepared by the Standards and Petitions Committee. Downloadable from <http://www.iucnredlist.org/documents/RedList-Guidelines.pdf>
- Röschel, L., Noebel, R., Stein, U., Naumann, S., Romão, C., Tryfon, E., Gaudillat, Z., Roscher, S., Moser, D., Ellmauer, T., Löhnertz, M., Halada, L., Staneva, A., Rutherford, C.A. (2020). *State of Nature in the EU - Methodological paper. Methodologies under the Nature Directives reporting 2013-2018 and analysis for the State of Nature 2000*. ETC/BD report to the EEA.

ONLINE SOURCES

- Baltic Marine Environment Protection Commission. *HELCOM*. <https://helcom.fi/> Last accessed: 28.08.2021
- BirdLife International (2008) Red List Index of European Birds. <https://www.eea.europa.eu/data-and-maps/indicators/red-list-index-for-european-species/red-list-index-for-european/?searchterm=None>

- Conservation of Arctic Fauna and Flora (CAFF). <https://www.caff.is/> Last accessed: 28.08.2021
- Convention on International Trade in Endangered Species of Wild Fauna and Flora. <https://cites.org/> Last accessed: 28.08.2021
- EBCC (2018). *Pan-European Common Bird Monitoring Scheme - European Indicators*. Available at: https://pecbms.info/trends-and-indicators/indicators/indicators/E_C_Fa/
- EBCC (2018). *Pan-European Common Bird Monitoring Scheme*. Available at: <https://pecbms.info/>
- EBCC (2021). *EuroBird Portal*. Available at: <https://eurobirdportal.org/ebp/en/#home/HIRRUS/r52weeks/CUCCAN/r52weeks/>
- EC (2021). The European Union and Trade in Wild Fauna and Flora https://ec.europa.eu/environment/cites/legislation_en.htm Last accessed: 28.08.2021
- EEA (2019). *Pressures and threats data reported as part of the EU Birds Directive Article 12 reporting exercise 2013-2018* <https://cdr.eionet.europa.eu/>
- EEA (2020). Article 12 webtool. <https://nature-art12.eionet.europa.eu/article12/> Last accessed: 28.08.2021
- EU (2008). *Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive)* https://ec.europa.eu/environment/marine/eu-coast-and-marine-policy/marine-strategy-framework-directive/index_en.htm - Last accessed: 28.08.2021
- EU (2009). *Directive 2009/147/EC of the European Parliament and of the Council of 30 November 2009 on the conservation of wild birds*, OJ L 20, 26.1.2018, pp. 7-25. https://ec.europa.eu/environment/nature/legislation/birdsdirective/index_en.htm
- European Commission (2021). Nature and biodiversity. https://ec.europa.eu/environment/nature/index_en.htm Last accessed: 28.08.2021
- European Commission (2021). *Biodiversity strategy for 2030* https://ec.europa.eu/environment/strategy/biodiversity-strategy-2030_en Last accessed: 28.08.2021
- European Commission (2021). *Common fisheries policy (CFP)* https://ec.europa.eu/oceans-and-fisheries/policy/common-fisheries-policy-cfp_en Last accessed: 28.08.2021
- European Commission (2021). *Our oceans, seas and coasts*. https://ec.europa.eu/environment/nature/index_en.htm https://ec.europa.eu/environment/marine/good-environmental-status/descriptor-1/index_en.htm Last accessed: 28.08.2021
- European Commission (2021). *The common agricultural policy at a glance*. https://ec.europa.eu/info/food-farming-fisheries/key-policies/common-agricultural-policy/cap-glance_en Last accessed: 28.08.2021
- European Court of Auditors (2020). *Biodiversity on farmland: CAP contribution has not halted the decline*. 58pp. https://www.eca.europa.eu/Lists/ECADocuments/SR20_13/SR_Biodiversity_on_farmland_EN.pdf
- Intergovernmental Panel on Climate Change (IPCC). <https://www.ipcc.ch/2019/> Last accessed: 28.08.2021
- Ospar Commission. <https://www.ospar.org/> Last accessed: 28.08.2021
- Skarphéðinsson, K.H. (2018). Stelkur (*Tringa totanus*) In: Náttúrufræðistofnun Íslands (2018) *Válisti fugla*. <https://en.ni.is/biota/animalia/chordata/aves/charadriiformes/stelkur-tringa-totanus>
- The Commission on the Protection of the Black Sea Against Pollution (1992). *Convention on the Protection of the Black Sea Against Pollution*. http://www.blacksea-commission.org/_convention.asp
- *The Leaders' Pledge for Nature - United to Reverse Biodiversity Loss by 2030 for Sustainable Development* (2020). <https://www.leaderspledgefornature.org/>
- United Nations (1992). Convention on Biological Diversity. Available at: www.cbd.int
- Wetlands International. International Waterbird Census (2020). Available at: <https://www.wetlands.org/knowledge-base/international-waterbird-census/> Last accessed 28.08.2021

SCIENTIFIC PAPERS

- Bird, J.P., Martin, R., Akçakaya, H.R., Gilroy, J., Burfield, I.J., Garnett, S.G., Symes, A., Taylor, J., Şekercioğlu, Ç.H. and Butchart, S.H.M. (2020). *Generation lengths of the world's birds and their implications for extinction risk*. Conservation Biology 34(5): 1252-1261. Doi: 10.1111/cobi.13486.
- Brochet, A., Van den Bossche, W., Jbour, S., Ndang'ang'a, P., Jones, V., Abdou, W., ... Butchart, S. (2016). *Preliminary assessment of the scope and scale of illegal killing and taking of birds in the Mediterranean*. Bird Conservation International, 26(1), 1-28. doi:10.1017/S0959270915000416
- Gunnarsson, T.G., Jóhannesdóttir, L., Alves, J.A., Þórisson, B., & Gill, J.A. (2017) *Effects of spring temperature and volcanic eruptions on wider productivity*. IBIS. DOI: 10.1111/ibi.12449
- Hof, Anouschka R. and Allen, Andrew M.(2019). *An uncertain future for the endemic Galliformes of the Caucasus*. Science of The Total Environment. 651(1) 725-735.
- Krüger, T., Heckenroth, H., Prior, N., Seitz, J., Zang, H. (2020). *Persecution and statutory protection have driven Rook *Corvus frugilegus* population dynamics over the past 120 years in NW-Germany*. Journal of Ornithology 161, 569–584. <https://doi.org/10.1007/s10336-020-01750-3>
- Lehtikoinen, Aleksu *et al.* (2021). Local colonisations and extinctions of European birds are poorly explained by changes in climate suitability (in prep.)
- Nagy, S., Breiner, F., Anand, M., Butchart, S., Flörke, M., Fluet-Chouinard, E., ... Voltz, O. (2021). *Climate change exposure of waterbird species in the African-Eurasian flyways*. Bird Conservation International, 1-26. doi:10.1017/S0959270921000150
- Popov I. & Starikov D. (2015) *Recent northward expansion of breeding Black-tailed Godwits *Limosa limosa* in NW Russia*. Wader Study 122(3): 173–183.
- Scott, D. A.; Rose, P. M. (1996). *Atlas of Anatidae populations in Africa and western Eurasia*. Wageningen, Netherlands: Wetlands International.
- Suárez, F. & Naveso, M.A. & de Juana, E. (1997a). *Farming in the drylands of Spain: Birds of the pseudosteppes*. Farming and Birds in Europe. 297-330.
- Vickery, J.A., Ewing, S.R., Smith, K.W., Pain, D.J., Bairlein, F., Škorpilová, J. and Gregory, R.D. (2014). *The decline of Afro-Palaearctic migrants and an assessment of potential causes*. Ibis, 156(1), pp.1-22. <https://onlinelibrary.wiley.com/doi/full/10.1111/ibi.12118>

BOOKS

- Ferguson-Lees, J.; Christie, D.A. (2001). *Raptors of the World*. Christopher Helm, London.
- Keller, V., Herrando, S., Voříšek, P., Franch, M., Kipson, M., Milanese, P., Martí, D., Anton, M., Klvaňová, A., Kalyakin, M.V., Bauer, H.-G. and Foppen, R.P.B. (2020). *European Breeding Bird Atlas 2: Distribution, Abundance and Change*. Barcelona: European Bird Census Council & Lynx Edicions.
- Newton, I. (2017). *Farming and birds*. London: William Collins

Appendix

Appendix 1. Red List status, trend and population size estimate, as well as endemism of birds in Europe⁶²

Scientific species name	Common species name	Season of assessment	IUCN Red List category (Europe)	IUCN Red List criteria (Europe)	Population trend	Population size [mature individuals] min-max, best estimate	(Near*) endemic to Europe?
<i>Cygnus columbianus</i>	Tundra Swan	W	VU	A4abcde	Decreasing	11700-23500,14600	
<i>Cygnus cygnus</i>	Whooper Swan	B	LC		Increasing	66800-102000,82000	
<i>Cygnus olor</i>	Mute Swan	B	LC		Increasing	169000-236000,199000	
<i>Anser albifrons</i>	Greater White-fronted Goose	W	LC		Stable	1180000-1970000,1420000	
<i>Anser anser</i>	Greylag Goose	W	LC		Increasing	797000-975000,866000	
<i>Anser brachyrhynchus</i>	Pink-footed Goose	W	LC		Increasing	370000-377000,373000	Yes
<i>Anser caerulescens</i>	Snow Goose	B	LC		Increasing	2000-10000,4500	
<i>Anser erythropus</i>	Lesser White-fronted Goose	B	VU	D1	Decreasing	340-530,420	
<i>Anser fabalis</i>	Bean Goose	W	LC		Increasing	468000-647000,534000	
<i>Branta bernicla</i>	Brent Goose	W	LC		Stable	198000-225000,211000	
<i>Branta canadensis</i>	Canada Goose	B	LC		Increasing	20000-100000,44700	
<i>Branta leucopsis</i>	Barnacle Goose	W	LC		Increasing	877000-1010000,939000	Yes
<i>Branta ruficollis</i>	Red-breasted Goose	W	VU	A2bcd+3bcd+4bcd	Decreasing	10000-49400,18800	Yes*
<i>Tadorna ferruginea</i>	Ruddy Shelduck	B	LC		Unknown	35500-64100,46500	
<i>Tadorna tadorna</i>	Common Shelduck	B	LC		Stable	104000-154000,124000	
<i>Anas acuta</i>	Northern Pintail	B	VU	A2bcde+A3bcde+A4bcde	Decreasing	310000-401000,353000	
<i>Anas crecca</i>	Common Teal	W	LC		Increasing	1040000-1640000,1240000	
<i>Anas platyrhynchos</i>	Mallard	W	LC		Decreasing	2830000-3770000,3200000	
<i>Mareca penelope</i>	Eurasian Wigeon	B	LC		Decreasing	451000-733000,574000	
<i>Mareca strepera</i>	Gadwall	B	LC		Increasing	164000-262000,202000	
<i>Marmaronetta angustirostris</i>	Marbled Teal	B	VU	D1	Stable	760-2200	
<i>Spatula clypeata</i>	Northern Shoveler	B	LC		Decreasing	295000-391000,339000	
<i>Spatula querquedula</i>	Garganey	B	LC		Decreasing	681000-920000,789000	
<i>Aythya ferina</i>	Common Pochard	W	VU	A2abcde+3bcde+4abcde	Decreasing	373000-679000,500000	
<i>Aythya fuligula</i>	Tufted Duck	B	NT	A2bcde	Decreasing	879000-1240000,1060000	
<i>Aythya marila</i>	Greater Scaup	B	LC		Decreasing	96400-170000,128000	
<i>Aythya nyroca</i>	Ferruginous Duck	B	LC		Unknown	18000-47000,28600	
<i>Bucephala clangula</i>	Common Goldeneye	B	LC		Decreasing	770000-990000,870000	
<i>Bucephala islandica</i>	Barrow's Goldeneye	B	NT	D2	Stable	1600-1800,1700	
<i>Clangula hyemalis</i>	Long-tailed Duck	W	LC		Decreasing	865000-1610000,1120000	
<i>Histrionicus histrionicus</i>	Harlequin Duck	B	LC		Stable	14000-18000,15700	
<i>Melanitta fusca</i>	Velvet Scoter	B	VU	A2abcde	Decreasing	36800-80300,57100	Yes*
<i>Melanitta nigra</i>	Common Scoter	B	LC		Unknown	45200-71100,56500	Yes*
<i>Mergellus albellus</i>	Smew	B	LC		Stable	21700-34000,27200	
<i>Mergus merganser</i>	Goosander	B	LC		Increasing	162000-267000,208000	
<i>Mergus serrator</i>	Red-breasted Merganser	B	NT	A2bcde+4bcde	Decreasing	141000-282000,198000	
<i>Netta rufina</i>	Red-crested Pochard	B	LC		Decreasing	69300-104000,82000	
<i>Oxyura leucocephala</i>	White-headed Duck	B	VU	C1+2a(i)	Decreasing	1000-1600,1300	
<i>Polysticta stelleri</i>	Steller's Eider	W	LC		Unknown	15800-18400,17000	
<i>Somateria mollissima</i>	Common Eider	B	EN	A4abcde	Decreasing	1220000-1630000,1410000	
<i>Somateria spectabilis</i>	King Eider	W	LC		Unknown	682000-683000,683000	
<i>Bonasa bonasia</i>	Hazel Grouse	B	LC		Unknown	4010000-6960000,5240000	
<i>Lagopus lagopus</i>	Willow Grouse	B	LC		Stable	2560000-4050000,3230000	
<i>Lagopus muta</i>	Rock Ptarmigan	B	LC		Fluctuating	619000-1700000,963000	
<i>Lyrurus mlokosiewiczzi</i>	Caucasian Grouse	B	NT	A3cde	Unknown	28300-57600,40300	Yes
<i>Lyrurus tetrix</i>	Black Grouse	B	LC		Unknown	2430000-3750000,3000000	
<i>Tetrao urogallus</i>	Western Capercaillie	B	LC		Decreasing	1590000-3420000,2110000	
<i>Alectoris barbara</i>	Barbary Partridge	B	LC		Unknown	15500-41000,25100	
<i>Alectoris chukar</i>	Chukar	B	LC		Unknown	1350000-3900000,2240000	
<i>Alectoris graeca</i>	Rock Partridge	B	NT	A2bcd+3bcd+4bcd	Decreasing	75300-123000,95400	Yes
<i>Alectoris rufa</i>	Red-legged Partridge	B	NT	A2bcde+3bcde+4bcde	Decreasing	9950000-13700000,11700000	Yes
<i>Ammoperdix griseogularis</i>	See-see Partridge	B	VU	C2a(ii)	Decreasing	7900-13600,10200	
<i>Coturnix coturnix</i>	Common Quail	B	NT	A2bcde+3bcde+4bcde	Decreasing	5000000-9030000,6560000	
<i>Francolinus francolinus</i>	Black Francolin	B	LC		Fluctuating	21400-37700,28000	

⁶² Red List Categories adjusted down by one/two/three Categories because of rescue effect from outside the region are denoted with °/°°/°°°, respectively.

Scientific species name	Common species name	Season of assessment	IUCN Red List category (Europe)	IUCN Red List criteria (Europe)	Population trend	Population size [mature individuals] min-max, best estimate	(Near*) endemic to Europe?
<i>Perdix perdix</i>	Grey Partridge	B	LC		Decreasing	2290000-3750000,2820000	
<i>Phasianus colchicus</i>	Common Pheasant	B	LC		Stable	9700-16300,12000	
<i>Tetraogallus caspius</i>	Caspian Snowcock	B	LC		Decreasing	8400-20800,13200	
<i>Tetraogallus caucasicus</i>	Caucasian Snowcock	B	LC		Stable	10000-32400,16300	Yes
<i>Turnix sylvaticus</i>	Common Buttonquail	B	RE		N/A	0-0,0	
<i>Gavia adamsii</i>	Yellow-billed Loon	W	VU	D1	Unknown	660-670,670	
<i>Gavia arctica</i>	Arctic Loon	B	LC		Decreasing	142000-194000,166000	
<i>Gavia immer</i>	Common Loon	W	LC		Stable	5700-7200,6300	
<i>Gavia stellata</i>	Red-throated Loon	B	LC		Unknown	91100-146000,115000	
<i>Podiceps auritus</i>	Horned Grebe	B	NT	C1	Decreasing	14200-19200,16100	
<i>Podiceps cristatus</i>	Great Crested Grebe	B	LC		Stable	772000-1060000,903000	
<i>Podiceps grisegena</i>	Red-necked Grebe	B	VU	A2bcde+4bcde	Decreasing	31200-58500,42100	
<i>Podiceps nigricollis</i>	Black-necked Grebe	B	VU	A2bce+3bce+4bce	Decreasing	71100-116000,88500	
<i>Tachybaptus ruficollis</i>	Little Grebe	B	LC		Stable	209000-390000,275000	
<i>Bulweria bulwerii</i>	Bulwer's Petrel	B	LC		Unknown	92100-103000,97000	
<i>Calonectris borealis</i>	Cory's Shearwater	B	LC		Unknown	503000-506000,505000	Yes*
<i>Calonectris diomedea</i>	Scopoli's Shearwater	B	LC		Unknown	57000-94300,72800	
<i>Fulmarus glacialis</i>	Northern Fulmar	B	VU	A4abcde	Decreasing	6350000-7660000,6790000	
<i>Hydrobates castro</i>	Band-rumped Storm-petrel	B	LC		Unknown	13200-14300,13700	
<i>Hydrobates leucorhous</i>	Leach's Storm-petrel	B	NT	B2ab(v)	Decreasing	117000-176000,141000	
<i>Hydrobates monteiroi</i>	Monteiro's Storm-petrel	B	VU	D1+2	Unknown	720-770,740	Yes
<i>Hydrobates pelagicus</i>	European Storm-petrel	B	LC		Unknown	810000-848000,825000	Yes
<i>Pelagodroma marina</i>	White-faced Storm-petrel	B	EN	B2ab(iii,v)	Decreasing	155000-222000,186000	
<i>Pterodroma deserta</i>	Desertas Petrel	B	VU	D1+2	Stable	320-360,340	Yes
<i>Pterodroma madeira</i>	Zino's Petrel	B	EN	D	Increasing	130-160,140	Yes
<i>Puffinus lherminieri</i>	Audubon's Shearwater	B	NT	B2ab(v); C1+2a(i)	Decreasing	5000-11800,7600	
<i>Puffinus mauretanicus</i>	Balearic Shearwater	B	CR	A4abcde	Decreasing	5800-5900,5800	Yes
<i>Puffinus puffinus</i>	Manx Shearwater	B	LC		Unknown	693000-779000,733000	Yes
<i>Puffinus yelkouan</i>	Yelkouan Shearwater	B	VU	A2abcde	Unknown	47000-81800,61900	Yes
<i>Morus bassanus</i>	Northern Gannet	B	LC		Increasing	821000-823000,822000	Yes
<i>Pelecanus crispus</i>	Dalmatian Pelican	B	LC		Increasing	7500-9400,8300	
<i>Pelecanus onocrotalus</i>	Great White Pelican	B	LC		Increasing	18700-40700,27400	
<i>Anhinga rufa</i>	African Darter	B	RE		N/A	0-0,0	
<i>Gulosus aristotelis</i>	European Shag	B	LC		Decreasing	142000-162000,152000	Yes
<i>Microcarbo pygmaeus</i>	Pygmy Cormorant	B	LC		Increasing	84600-111000,96600	
<i>Phalacrocorax carbo</i>	Great Cormorant	B	LC		Increasing	828000-1030000,926000	
<i>Ardea alba</i>	Great White Egret	B	LC		Increasing	79800-132000,102000	
<i>Ardea cinerea</i>	Grey Heron	B	LC		Decreasing	492000-700000,583000	
<i>Ardea purpurea</i>	Purple Heron	B	LC		Increasing	57700-97300,73500	
<i>Ardeola ralloides</i>	Squacco Heron	B	LC		Stable	36600-67000,49300	
<i>Botaurus stellaris</i>	Eurasian Bittern	B	LC		Stable	147000-206000,174000	
<i>Bubulcus ibis</i>	Cattle Egret	B	LC		Increasing	147000-184000,164000	
<i>Ciconia ciconia</i>	White Stork	B	LC		Increasing	502000-563000,530000	Yes
<i>Ciconia nigra</i>	Black Stork	B	LC		Unknown	20200-32400,25400	
<i>Egretta garzetta</i>	Little Egret	B	LC		Decreasing	121000-190000,146000	
<i>Geronticus eremita</i>	Northern Bald Ibis	B	RE		N/A	0-0,0	
<i>Ixobrychus minutus</i>	Common Little Bittern	B	LC		Stable	171000-301000,230000	
<i>Nycticorax nycticorax</i>	Black-crowned Night-heron	B	LC		Stable	114000-169000,137000	
<i>Platalea leucorodia</i>	Eurasian Spoonbill	B	LC		Increasing	23800-36300,29000	
<i>Plegadis falcinellus</i>	Glossy Ibis	B	LC		Increasing	60200-120000,80500	
<i>Phoenicopterus roseus</i>	Greater Flamingo	B	LC		Increasing	71800-265000,144000	
<i>Aegyptius monachus</i>	Cinereous Vulture	B	LC		Increasing	5800-6700,6100	
<i>Gypaetus barbatus</i>	Bearded Vulture	B	NT	D1	Increasing	1200-2000,1500	
<i>Gyps fulvus</i>	Griffon Vulture	B	LC		Increasing	69600-89400,75700	
<i>Neophron percnopterus</i>	Egyptian Vulture	B	VU	C1	Decreasing	6100-9000,7100	
<i>Aquila adalberti</i>	Spanish Imperial Eagle	B	VU	D1	Increasing	1000-1100,1100	Yes
<i>Aquila chrysaetos</i>	Golden Eagle	B	LC		Increasing	19200-25600,22100	
<i>Aquila fasciata</i>	Bonelli's Eagle	B	LC		Stable	2100-2500,2300	
<i>Aquila heliaca</i>	Eastern Imperial Eagle	B	LC		Increasing	3900-6000,4800	
<i>Aquila nipalensis</i>	Steppe Eagle	B	CR	A2abcd+3bcd+4abcd	Decreasing	1500-2300,1800	
<i>Circus gallicus</i>	Short-toed Snake-eagle	B	LC		Increasing	19800-31900,24600	
<i>Clanga clanga</i>	Greater Spotted Eagle	B	VU	A2abcde+3bcde+4abcde	Decreasing	1900-2500,2200	
<i>Clanga pomarina</i>	Lesser Spotted Eagle	B	LC		Increasing	34200-46200,39700	Yes

Scientific species name	Common species name	Season of assessment	IUCN Red List category (Europe)	IUCN Red List criteria (Europe)	Population trend	Population size [mature individuals] min-max, best estimate	(Near*) endemic to Europe?
<i>Haliaeetus albicilla</i>	White-tailed Sea-eagle	B	LC		Increasing	20900-29200,24500	
<i>Hieraetus pennatus</i>	Booted Eagle	B	LC		Increasing	46600-60500,52100	
<i>Pandion haliaetus</i>	Osprey	B	LC		Increasing	19200-27100,23000	
<i>Milvus migrans</i>	Black Kite	B	LC		Increasing	372000-507000,433000	
<i>Milvus milvus</i>	Red Kite	B	LC		Increasing	65100-76600,70300	Yes
<i>Circus aeruginosus</i>	Western Marsh-harrier	B	LC		Stable	303000-485000,380000	
<i>Circus cyaneus</i>	Hen Harrier	B	LC		Decreasing	112000-174000,139000	
<i>Circus macrourus</i>	Pallid Harrier	B	LC		Stable	2000-4300,2900	
<i>Circus pygargus</i>	Montagu's Harrier	B	LC		Decreasing	139000-219000,174000	
<i>Buteo buteo</i>	Eurasian Buzzard	B	LC		Increasing	1760000-2460000,2070000	
<i>Buteo lagopus</i>	Rough-legged Buzzard	B	LC		Stable	57600-117000,81400	
<i>Buteo rufinus</i>	Long-legged Buzzard	B	LC		Increasing	27600-45800,35300	
<i>Pernis apivorus</i>	European Honey-buzzard	B	LC		Stable	241000-350000,289000	Yes
<i>Accipiter badius</i>	Shikra	B	NT ^o	D1	Stable	200-1100,450	
<i>Accipiter brevipes</i>	Levant Sparrowhawk	B	LC		Stable	7700-15300,10800	
<i>Accipiter gentilis</i>	Northern Goshawk	B	LC		Decreasing	234000-380000,291000	
<i>Accipiter nisus</i>	Eurasian Sparrowhawk	B	LC		Stable	728000-1150000,915000	
<i>Elanus caeruleus</i>	Black-winged Kite	B	LC		Stable	2600-5700,3800	
<i>Falco biarmicus</i>	Lanner Falcon	B	NT*	D1	Unknown	840-1700,1200	
<i>Falco cherrug</i>	Saker Falcon	B	EN	C1	Decreasing	860-1300,1100	
<i>Falco columbarius</i>	Merlin	B	VU	A2bce	Decreasing	40100-83400,59200	
<i>Falco eleonorae</i>	Eleonora's Falcon	B	LC		Stable	28400-28900,28600	Yes*
<i>Falco naumanni</i>	Lesser Kestrel	B	LC		Stable	65900-85200,73400	
<i>Falco peregrinus</i>	Peregrine Falcon	B	LC		Increasing	32200-62100,41300	
<i>Falco rusticolus</i>	Gyr Falcon	B	LC		Stable	2000-3500,2600	
<i>Falco subbuteo</i>	Eurasian Hobby	B	LC		Stable	280000-437000,342000	
<i>Falco tinnunculus</i>	Common Kestrel	B	LC		Decreasing	823000-1270000,991000	
<i>Falco vespertinus</i>	Red-footed Falcon	B	VU	A2abcde+4abcde	Decreasing	115000-170000,140000	
<i>Crex crex</i>	Corncrake	B	LC		Decreasing	2310000-3430000,2800000	
<i>Fulica atra</i>	Common Coot	B	NT	A2abcde+4abcde	Decreasing	2030000-3360000,2590000	
<i>Fulica cristata</i>	Red-knobbed Coot	B	CR	C2a(ii)	Decreasing	42-90,60	
<i>Gallinula chloropus</i>	Common Moorhen	B	LC		Decreasing	1790000-2670000,2150000	
<i>Porphyrio porphyrio</i>	Purple Swamphen	B	LC		Fluctuating	30400-114000,56500	
<i>Porzana porzana</i>	Spotted Crake	B	LC		Decreasing	200000-289000,235000	
<i>Rallus aquaticus</i>	Western Water Rail	B	LC		Unknown	315000-653000,438000	
<i>Zapornia parva</i>	Little Crake	B	LC		Unknown	82000-203000,121000	
<i>Zapornia pusilla</i>	Baillon's Crake	B	LC		Unknown	2000-4800,3000	
<i>Anthropoides virgo</i>	Demoiselle Crane	B	EN	A2bcd+3bcd+4bcd	Decreasing	9300-11900,10500	
<i>Grus grus</i>	Common Crane	B	LC		Increasing	309000-423000,362000	
<i>Chlamydotis macqueenii</i>	Asian Houbara	B	CR (PE)	D	Decreasing	0-0,0	
<i>Chlamydotis undulata</i>	African Houbara	B	VU	D1	Increasing	610-620,620	
<i>Otis tarda</i>	Great Bustard	B	LC		Decreasing	35200-41800,38300	
<i>Tetrax tetrax</i>	Little Bustard	B	VU	A2bcde+4bcde	Decreasing	194000-280000,228000	
<i>Actitis hypoleucos</i>	Common Sandpiper	B	LC		Decreasing	2060000-2980000,2460000	
<i>Arenaria interpres</i>	Ruddy Turnstone	W	LC		Stable	57200-67300,61400	
<i>Burhinus oedicephalus</i>	Eurasian Thick-knee	B	LC		Decreasing	123000-193000,152000	
<i>Calidris alba</i>	Sanderling	W	LC		Increasing	50200-66800,57400	
<i>Calidris alpina</i>	Dunlin	W	LC		Stable	865000-1050000,937000	
<i>Calidris bairdii</i>	Baird's Sandpiper	B	LC ^{oo}		Unknown	200-1000,450	
<i>Calidris canutus</i>	Red Knot	W	LC		Decreasing	268000-325000,296000	
<i>Calidris falcinellus</i>	Broad-billed Sandpiper	B	VU	A2bc+3bc+4bc	Decreasing	40500-70700,52500	
<i>Calidris ferruginea</i>	Curlew Sandpiper	W	VU	D1	Increasing	100-1100,420	
<i>Calidris maritima</i>	Purple Sandpiper	W	LC		Unknown	179000-196000,185000	
<i>Calidris minuta</i>	Little Stint	B	LC		Unknown	80200-122000,98400	
<i>Calidris pugnax</i>	Ruff	B	NT	A2abcde+3bcde+4abcde	Decreasing	513000-1380000,841000	
<i>Calidris temminckii</i>	Temminck's Stint	B	LC		Unknown	166000-248000,203000	
<i>Charadrius alexandrinus</i>	Kentish Plover	B	LC		Decreasing	34900-67400,46600	
<i>Charadrius asiaticus</i>	Caspian Plover	B	EN ^o	D	Unknown	6-60,25	
<i>Charadrius dubius</i>	Little Ringed Plover	B	LC		Decreasing	425000-634000,513000	
<i>Charadrius hiaticula</i>	Common Ringed Plover	B	LC		Increasing	261000-399000,323000	
<i>Charadrius leschenaultii</i>	Greater Sandplover	B	NT	D1	Decreasing	1300-2400,1700	
<i>Cursorius cursor</i>	Cream-coloured Courser	B	NT	D1	Unknown	1300-1600,1400	
<i>Eudromias morinellus</i>	Eurasian Dotterel	B	LC		Unknown	21100-43100,31300	

*Erratum: Red List status and trend in Europe updated in July 2022 to EN (for more details see: www.iucnredlist.org). This update is not reflected in the graphs and statistics presented in the report.

Scientific species name	Common species name	Season of assessment	IUCN Red List category (Europe)	IUCN Red List criteria (Europe)	Population trend	Population size [mature individuals] min-max, best estimate	(Near*) endemic to Europe?
<i>Gallinago gallinago</i>	Common Snipe	B	VU	A2bcde+3bcde+4bcde	Decreasing	5270000-7260000,6160000	
<i>Gallinago media</i>	Great Snipe	B	LC		Decreasing	104000-181000,136000	
<i>Gallinago megala</i>	Swinhoe's Snipe	B	NT ⁰⁰⁰	D1	Unknown	20-60,35	
<i>Gallinago stenura</i>	Pintail Snipe	B	LC		Unknown	20000-38000,27600	
<i>Glareola nordmanni</i>	Black-winged Pratincole	B	LC		Stable	12000-24200,17000	
<i>Glareola pratincola</i>	Collared Pratincole	B	LC		Decreasing	17500-35000,23800	
<i>Haematopus ostralegus</i>	Eurasian Oystercatcher	B	VU	A4abcde	Decreasing	526000-692000,598000	
<i>Himantopus himantopus</i>	Black-winged Stilt	B	LC		Increasing	101000-269000,158000	
<i>Limosa lapponica</i>	Bar-tailed Godwit	W	LC		Stable	98600-119000,108000	
<i>Limosa limosa</i>	Black-tailed Godwit	B	NT	A2bcde	Stable	272000-334000,300000	
<i>Lymnocyptes minimus</i>	Jack Snipe	B	LC		Unknown	30500-77700,51100	
<i>Numenius arquata</i>	Eurasian Curlew	B	NT	A2bcde+4bcde	Decreasing	405000-553000,470000	
<i>Numenius phaeopus</i>	Whimbrel	B	LC		Unknown	674000-774000,719000	
<i>Numenius tenuirostris</i>	Slender-billed Curlew	W	CR (PE)	D	N/A	0-0,0	
<i>Phalaropus fulicarius</i>	Red Phalarope	B	LC		Unknown	30400-62200,43400	
<i>Phalaropus lobatus</i>	Red-necked Phalarope	B	LC		Unknown	551000-834000,678000	
<i>Pluvialis apricaria</i>	Eurasian Golden Plover	B	LC		Stable	1660000-2310000,1960000	Yes*
<i>Pluvialis squatarola</i>	Grey Plover	W	LC		Stable	84800-119000,95100	
<i>Recurvirostra avosetta</i>	Pied Avocet	B	LC		Decreasing	81200-155000,95300	
<i>Scolopax rusticola</i>	Eurasian Woodcock	B	LC		Decreasing	9790000-13500000,11500000	
<i>Tringa erythropus</i>	Spotted Redshank	B	LC		Unknown	37900-81300,56300	
<i>Tringa glareola</i>	Wood Sandpiper	B	LC		Unknown	2090000-2920000,2480000	
<i>Tringa nebularia</i>	Common Greenshank	B	LC		Increasing	452000-942000,646000	
<i>Tringa ochropus</i>	Green Sandpiper	B	LC		Unknown	1180000-1850000,1470000	
<i>Tringa stagnatilis</i>	Marsh Sandpiper	B	LC		Decreasing	18200-28800,22900	
<i>Tringa totanus</i>	Common Redshank	B	VU	A2bcde+4bcde	Decreasing	522000-694000,596000	
<i>Vanellus gregarius</i>	Sociable Lapwing	B	CR	A2bcde+3bcde+4bcde; C1+2a(i,ii); D	Decreasing	6-20,11	
<i>Vanellus indicus</i>	Red-wattled Lapwing	B	VU ⁰⁰	D1	Unknown	40-60,49	
<i>Vanellus leucurus</i>	White-tailed Lapwing	B	LC		Stable	2200-20400,6600	
<i>Vanellus spinosus</i>	Spur-winged Lapwing	B	LC		Unknown	2100-3300,2700	
<i>Vanellus vanellus</i>	Northern Lapwing	B	VU	A2bcde	Decreasing	3180000-4590000,3820000	
<i>Xenus cinereus</i>	Terek Sandpiper	B	LC		Decreasing	80300-121000,98400	
<i>Catharacta skua</i>	Great Skua	B	LC		Increasing	25800-30000,27600	Yes
<i>Stercorarius longicaudus</i>	Long-tailed Jaeger	B	LC		Fluctuating	73200-161000,112000	
<i>Stercorarius parasiticus</i>	Arctic Jaeger	B	EN	A4bcde	Decreasing	54200-83000,66000	
<i>Stercorarius pomarinus</i>	Pomarine Jaeger	B	LC		Unknown	5000-8100,6300	
<i>Hydrocoloeus minutus</i>	Little Gull	B	LC		Unknown	64300-125000,88900	
<i>Larus argentatus</i>	European Herring Gull	B	LC		Decreasing	1060000-1220000,1130000	Yes
<i>Larus armenicus</i>	Armenian Gull	B	LC		Unknown	44400-71800,56000	Yes
<i>Larus audouinii</i>	Audouin's Gull	B	VU	A3bce+4abce	Decreasing	31400-42000,36300	Yes
<i>Larus cachinnans</i>	Caspian Gull	B	LC		Increasing	231000-400000,303000	
<i>Larus canus</i>	Mew Gull	B	LC		Unknown	1920000-2600000,2250000	
<i>Larus fuscus</i>	Lesser Black-backed Gull	B	LC		Increasing	738000-828000,781000	
<i>Larus genei</i>	Slender-billed Gull	B	VU	A4bcde	Decreasing	61100-73200,66300	
<i>Larus glaucoides</i>	Iceland Gull	B	LC		Stable	100000-200000,141000	
<i>Larus hyperboreus</i>	Glaucous Gull	B	LC		Stable	69200-246000,125000	
<i>Larus ichthyæetus</i>	Pallas's Gull	B	LC		Increasing	34700-56300,43900	
<i>Larus marinus</i>	Great Black-backed Gull	B	LC		Stable	172000-215000,196000	
<i>Larus melanocephalus</i>	Mediterranean Gull	B	LC		Decreasing	128000-203000,160000	Yes
<i>Larus michahellis</i>	Yellow-legged Gull	B	LC		Unknown	830000-1110000,947000	Yes
<i>Larus ridibundus</i>	Black-headed Gull	B	LC		Decreasing	2810000-4120000,3380000	
<i>Onychoprion fuscatus</i>	Sooty Tern	B	VU ⁰⁰	D1	Unknown	2-2,2	
<i>Pagophila eburnea</i>	Ivory Gull	B	VU	C1	Decreasing	8600-11600,9900	
<i>Rhodostethia rosea</i>	Ross's Gull	B	EN ⁰	D	Fluctuating	10-10,10	
<i>Rissa tridactyla</i>	Black-legged Kittiwake	B	VU	A2abcd+3bcd+4abcd	Decreasing	3250000-3450000,3330000	
<i>Xema sabini</i>	Sabine's Gull	B	LC		Increasing	2000-2100,2000	
<i>Gelochelidon nilotica</i>	Common Gull-billed Tern	B	LC		Stable	26700-37400,30900	
<i>Hydroprogne caspia</i>	Caspian Tern	B	LC		Stable	12700-19900,15700	
<i>Sterna dougallii</i>	Roseate Tern	B	LC		Increasing	4900-6200,5500	
<i>Sterna hirundo</i>	Common Tern	B	LC		Unknown	881000-1430000,1120000	
<i>Sterna paradisaea</i>	Arctic Tern	B	LC		Stable	1060000-1510000,1220000	
<i>Sternula albifrons</i>	Little Tern	B	LC		Decreasing	66800-101000,81600	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (Europe)	IUCN Red List criteria (Europe)	Population trend	Population size [mature individuals] min-max, best estimate	(Near*) endemic to Europe?
<i>Thalasseus sandvicensis</i>	Sandwich Tern	B	LC		Increasing	160000-320000,258000	
<i>Chlidonias hybrida</i>	Whiskered Tern	B	LC		Increasing	117000-294000,177000	
<i>Chlidonias leucopterus</i>	White-winged Tern	B	LC		Unknown	176000-432000,262000	
<i>Chlidonias niger</i>	Black Tern	B	LC		Unknown	225000-355000,280000	
<i>Alca torda</i>	Razorbill	B	LC		Increasing	519000-1070000,763000	Yes
<i>Alle alle</i>	Little Auk	B	LC		Unknown	7600000-77000000,76400000	
<i>Cephus grylle</i>	Black Guillemot	B	LC		Unknown	368000-397000,380000	
<i>Fratercula arctica</i>	Atlantic Puffin	B	EN	A2abcde+4abcde	Decreasing	7400000-8240000,7820000	Yes
<i>Uria aalge</i>	Common Murre	B	LC		Increasing	2990000-4430000,3660000	
<i>Uria lomvia</i>	Thick-billed Murre	B	LC		Decreasing	3410000-4560000,3950000	
<i>Pterocles alchata</i>	Pin-tailed Sandgrouse	B	LC		Stable	15000-18800,16600	
<i>Pterocles orientalis</i>	Black-bellied Sandgrouse	B	EN	A2bcde+4bcde	Decreasing	14900-37700,24300	
<i>Syrhaptes paradoxus</i>	Pallas's Sandgrouse	B	RE		N/A	0-0,0	
<i>Columba bollii</i>	Dark-tailed Laurel-pigeon	B	LC		Unknown	2500-10000,5000	Yes
<i>Columba junoniae</i>	White-tailed Laurel-pigeon	B	NT	B1ab(iii)+2ab(iii)	Unknown	1000-2500,1600	Yes
<i>Columba livia</i>	Rock Dove	B	LC		Unknown	6840000-14200000,9710000	
<i>Columba oenas</i>	Stock Dove	B	LC		Increasing	1360000-2290000,1700000	Yes
<i>Columba palumbus</i>	Common Woodpigeon	B	LC		Increasing	41400000-57600000,48400000	Yes
<i>Columba trocaz</i>	Madeira Laurel-pigeon	B	LC		Increasing	10000-14000,11800	Yes
<i>Spilopelia senegalensis</i>	Laughing Dove	B	LC		Increasing	141000-328000,210000	
<i>Streptopelia decaocto</i>	Eurasian Collared-dove	B	LC		Stable	17900000-44100000,25500000	
<i>Streptopelia turtur</i>	European Turtle-dove	B	VU	A2bcde+4bcde	Decreasing	5020000-9510000,6900000	
<i>Clamator glandarius</i>	Great Spotted Cuckoo	B	VU	A2bcd+3bcd+4bcd	Decreasing	293000-316000,301000	
<i>Cuculus canorus</i>	Common Cuckoo	B	LC		Decreasing	8640000-13600000,10800000	
<i>Cuculus optatus</i>	Oriental Cuckoo	B	LC		Unknown	240000-310000,273000	
<i>Aegolius funereus</i>	Boreal Owl	B	LC		Fluctuating	189000-471000,288000	
<i>Asio flammeus</i>	Short-eared Owl	B	LC		Fluctuating	169000-284000,215000	
<i>Asio otus</i>	Northern Long-eared Owl	B	LC		Unknown	624000-1030000,788000	
<i>Athene noctua</i>	Little Owl	B	LC		Unknown	1150000-2140000,1560000	
<i>Bubo bubo</i>	Eurasian Eagle-owl	B	LC		Increasing	37100-59500,45900	
<i>Bubo scandiacus</i>	Snowy Owl	B	LC		Fluctuating	2400-5400,3500	
<i>Glaucidium passerinum</i>	Eurasian Pygmy-owl	B	LC		Stable	165000-313000,230000	
<i>Ketupa zeylonensis</i>	Brown Fish-owl	B	EN	D	Unknown	100-160,130	
<i>Otus brucei</i>	Pallid Scops-owl	B	EN	D	Stable	100-500,220	
<i>Otus cypricus</i>	Cyprus Scops-owl	B	LC		Stable	6000-10000,7700	Yes
<i>Otus scops</i>	Eurasian Scops-owl	B	LC		Stable	447000-749000,566000	
<i>Strix aluco</i>	Tawny Owl	B	LC		Stable	1260000-1870000,1520000	
<i>Strix nebulosa</i>	Great Grey Owl	B	LC		Stable	12400-19900,15700	
<i>Strix uralensis</i>	Ural Owl	B	LC		Increasing	160000-263000,203000	
<i>Surnia ulula</i>	Northern Hawk-owl	B	LC		Increasing	20800-92400,36800	
<i>Tyto alba</i>	Common Barn-owl	B	LC		Decreasing	164000-356000,239000	
<i>Caprimulgus europaeus</i>	European Nightjar	B	LC		Unknown	1190000-2220000,1620000	
<i>Caprimulgus ruficollis</i>	Red-necked Nightjar	B	NT	A2bc+3bc+4bc	Decreasing	202000-270000,233000	
<i>Apus affinis</i>	Little Swift	B	NT	A2bc; C1	Decreasing	1500-4800,2700	
<i>Apus apus</i>	Common Swift	B	NT	A2bc+3bc+4bc	Decreasing	3220000-56700000,41700000	
<i>Apus caffer</i>	White-rumped Swift	B	NT*	D1	Increasing	500-700,590	
<i>Apus pallidus</i>	Pallid Swift	B	LC		Unknown	152000-424000,230000	
<i>Apus unicolor</i>	Plain Swift	B	LC		Unknown	15000-40000,24100	Yes
<i>Tachymarptis melba</i>	Alpine Swift	B	LC		Unknown	647000-1460000,960000	
<i>Upupa epops</i>	Common Hoopoe	B	LC		Stable	2740000-6410000,4030000	
<i>Alcedo atthis</i>	Common Kingfisher	B	LC		Decreasing	179000-440000,270000	
<i>Ceryle rudis</i>	Pied Kingfisher	B	VU	D1	Decreasing	600-1000,770	
<i>Halcyon smyrnensis</i>	White-breasted Kingfisher	B	VU	D1	Decreasing	300-500,390	
<i>Coracias garrulus</i>	European Roller	B	LC		Decreasing	102000-208000,138000	
<i>Merops apiaster</i>	European Bee-eater	B	LC		Stable	7370000-11200000,8970000	
<i>Merops persicus</i>	Blue-cheeked Bee-eater	B	LC		Increasing	16900-55600,30400	
<i>Dendrocopos leucotos</i>	White-backed Woodpecker	B	LC		Decreasing	420000-822000,585000	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (Europe)	IUCN Red List criteria (Europe)	Population trend	Population size [mature individuals] min-max, best estimate	(Near*) endemic to Europe?
<i>Dendrocopos major</i>	Great Spotted Woodpecker	B	LC		Decreasing	17200000-27300000,21500000	
<i>Dendrocopos syriacus</i>	Syrian Woodpecker	B	LC		Unknown	645000-1540000,988000	
<i>Dryobates minor</i>	Lesser Spotted Woodpecker	B	LC		Increasing	1510000-3020000,2130000	
<i>Dryocopus martius</i>	Black Woodpecker	B	LC		Unknown	1240000-2270000,1670000	
<i>Jynx torquilla</i>	Eurasian Wryneck	B	LC		Stable	1590000-3040000,2190000	
<i>Leiopicus medius</i>	Middle Spotted Woodpecker	B	LC		Increasing	802000-1390000,1050000	Yes
<i>Picoides tridactylus</i>	Three-toed Woodpecker	B	LC		Unknown	603000-1120000,816000	
<i>Picus canus</i>	Grey-faced Woodpecker	B	LC		Increasing	365000-609000,470000	
<i>Picus sharpei</i>	Iberian Green Woodpecker	B	LC		Decreasing	488000-938000,671000	Yes
<i>Picus viridis</i>	Eurasian Green Woodpecker	B	LC		Stable	1200000-2050000,1560000	Yes
<i>Panurus biarmicus</i>	Bearded Reedling	B	LC		Increasing	343000-783000,496000	
<i>Alauda arvensis</i>	Eurasian Skylark	B	LC		Decreasing	87800000-132000000,106000000	
<i>Alauda leucoptera</i>	White-winged Lark	B	LC		Stable	40000-110000,66300	
<i>Alaudala rufescens</i>	Lesser Short-toed Lark	B	LC		Decreasing	1700000-3070000,2250000	
<i>Ammomanes deserti</i>	Desert Lark	B	EN	D	Decreasing	40-200,90	
<i>Calandrella brachydactyla</i>	Greater Short-toed Lark	B	LC		Unknown	9300000-17400000,12700000	
<i>Chersophilus duponti</i>	Dupont's Lark	B	VU	A2bc+3bc+4bc; C1+2a(i)	Decreasing	4600-4700,4700	
<i>Eremophila alpestris</i>	Horned Lark	B	LC		Unknown	4130000-12300000,7090000	
<i>Galerida cristata</i>	Crested Lark	B	LC		Decreasing	34300000-47500000,39900000	
<i>Galerida theklae</i>	Thekla's Lark	B	LC		Increasing	3500000-5690000,4380000	
<i>Lullula arborea</i>	Woodlark	B	LC		Decreasing	4290000-9130000,6010000	Yes
<i>Melanocorypha bimaculata</i>	Bimaculated Lark	B	LC		Unknown	2000000-4020000,2840000	
<i>Melanocorypha calandra</i>	Calandra Lark	B	LC		Decreasing	19800000-38400000,27500000	
<i>Melanocorypha yeltoniensis</i>	Black Lark	B	EN	D	Decreasing	60-150,90	
<i>Cecropis daurica</i>	Red-rumped Swallow	B	LC		Increasing	2860000-6680000,4450000	
<i>Delichon urbicum</i>	Northern House Martin	B	LC		Stable	23000000-48100000,33000000	
<i>Hirundo rustica</i>	Barn Swallow	B	LC		Decreasing	51900000-89500000,66500000	
<i>Ptyonoprogne rupestris</i>	Eurasian Crag Martin	B	LC		Stable	420000-816000,562000	
<i>Riparia riparia</i>	Collared Sand Martin	B	LC		Decreasing	13400000-19800000,15900000	
<i>Anthus berthelotii</i>	Berthelot's Pipit	B	LC		Unknown	60000-300000,134000	Yes
<i>Anthus campestris</i>	Tawny Pipit	B	LC		Unknown	2240000-4050000,2980000	
<i>Anthus cervinus</i>	Red-throated Pipit	B	LC		Unknown	2210000-4240000,3060000	
<i>Anthus gustavi</i>	Pechora Pipit	B	CR (PE)	D	N/A	0-0,0	
<i>Anthus hodgsoni</i>	Olive-backed Pipit	B	LC		Unknown	160000-280000,212000	
<i>Anthus petrosus</i>	Rock Pipit	B	LC		Unknown	334000-571000,433000	Yes
<i>Anthus pratensis</i>	Meadow Pipit	B	LC		Decreasing	22000000-29800000,25100000	Yes
<i>Anthus spinoletta</i>	Water Pipit	B	LC		Decreasing	2100000-5280000,3110000	
<i>Anthus trivialis</i>	Tree Pipit	B	LC		Decreasing	38900000-70100000,52000000	
<i>Motacilla alba</i>	White Wagtail	B	LC		Stable	30300000-51100000,39000000	
<i>Motacilla cinerea</i>	Grey Wagtail	B	LC		Stable	2240000-4750000,3260000	
<i>Motacilla citreola</i>	Citrine Wagtail	B	LC		Increasing	803000-1150000,963000	
<i>Motacilla flava</i>	Western Yellow Wagtail	B	LC		Decreasing	26700000-36000000,30800000	
<i>Cinclus cinclus</i>	White-throated Dipper	B	LC		Decreasing	222000-547000,335000	
<i>Bombycilla garrulus</i>	Bohemian Waxwing	B	LC		Increasing	1300000-2540000,1820000	
<i>Prunella atrogularis</i>	Black-throated Accentor	B	LC		Unknown	3000-4800,3800	
<i>Prunella collaris</i>	Alpine Accentor	B	LC		Stable	203000-379000,271000	
<i>Prunella modularis</i>	Dunnock	B	LC		Decreasing	22100000-37000000,26800000	
<i>Prunella montanella</i>	Siberian Accentor	B	LC		Unknown	24400-46000,33500	
<i>Prunella ocularis</i>	Radde's Accentor	B	LC		Unknown	29800-76300,47200	
<i>Calliope calliope</i>	Siberian Rubythroat	B	LC		Unknown	1800-3200,2400	
<i>Cercotrichas galactotes</i>	Rufous-tailed Scrub-robin	B	LC		Decreasing	270000-775000,456000	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (Europe)	IUCN Red List criteria (Europe)	Population trend	Population size [mature individuals] min-max, best estimate	(Near*) endemic to Europe?
<i>Cyanecula svecica</i>	Bluethroat	B	LC		Stable	7930000-14800000,10900000	
<i>Erithacus rubecula</i>	European Robin	B	LC		Stable	109000000-168000000,133000000	Yes
<i>Irania gutturalis</i>	White-throated Robin	B	LC		Unknown	809000-1820000,1210000	
<i>Luscinia luscinia</i>	Thrush Nightingale	B	LC		Increasing	8790000-14300000,11100000	
<i>Luscinia megarhynchos</i>	Common Nightingale	B	LC		Stable	20400000-30700000,24800000	
<i>Monticola saxatilis</i>	Rufous-tailed Rock-thrush	B	LC		Unknown	179000-572000,310000	
<i>Monticola solitarius</i>	Blue Rock-thrush	B	LC		Unknown	201000-463000,301000	
<i>Oenanthe chrysopygia</i>	Red-tailed Wheatear	B	VU ^o	D1	Stable	90-120,100	
<i>Oenanthe cyprica</i>	Cyprus Wheatear	B	LC		Increasing	100000-200000,141000	Yes
<i>Oenanthe deserti</i>	Desert Wheatear	B	NT ^o	D1	Unknown	220-2200,700	
<i>Oenanthe finschii</i>	Finsch's Wheatear	B	LC		Unknown	204000-617000,354000	
<i>Oenanthe hispanica</i>	Black-eared Wheatear	B	LC		Unknown	2530000-7250000,4280000	
<i>Oenanthe isabellina</i>	Isabelline Wheatear	B	LC		Unknown	4620000-13200000,7740000	
<i>Oenanthe leucura</i>	Black Wheatear	B	LC		Stable	8100-32200,13000	
<i>Oenanthe oenanthe</i>	Northern Wheatear	B	LC		Unknown	10800000-29300000,17500000	
<i>Oenanthe pleschanka</i>	Pied Wheatear	B	LC		Unknown	54500-99700,70400	
<i>Oenanthe xanthopyrna</i>	Kurdish Wheatear	B	LC		Unknown	2000-12000,4600	
<i>Phoenicurus erythrogastrus</i>	White-winged Redstart	B	NT	C1	Decreasing	3400-15000,6600	
<i>Phoenicurus ochruros</i>	Black Redstart	B	LC		Stable	13100000-21300000,16600000	
<i>Phoenicurus phoenicurus</i>	Common Redstart	B	LC		Increasing	10500000-19600000,14400000	
<i>Saxicola dacotiae</i>	Fuerteventura Stonechat	B	NT	B1ab(ii,iii); C2a(ii)	Unknown	13300-15500,14400	Yes
<i>Saxicola rubetra</i>	Whinchat	B	LC		Decreasing	15400000-21100000,18100000	Yes
<i>Saxicola torquatus</i>	Common Stonechat	B	LC		Decreasing	11800000-18400000,14600000	
<i>Tarsiger cyanurus</i>	Orange-flanked Bush-robin	B	LC		Increasing	481000-892000,653000	
<i>Turdus atrogularis</i>	Black-throated Thrush	B	LC		Unknown	50000-80000,63200	
<i>Turdus iliacus</i>	Redwing	B	LC		Decreasing	16200000-28100000,21300000	
<i>Turdus merula</i>	Eurasian Blackbird	B	LC		Increasing	116000000-176000000,140000000	
<i>Turdus philomelos</i>	Song Thrush	B	LC		Stable	47300000-77900000,59700000	
<i>Turdus pilaris</i>	Fieldfare	B	LC		Stable	23000000-44700000,32000000	
<i>Turdus torquatus</i>	Ring Ouzel	B	LC		Stable	552000-1160000,779000	Yes
<i>Turdus viscivorus</i>	Mistle Thrush	B	LC		Stable	7300000-13600000,9720000	
<i>Zoothera aurea</i>	White's Thrush	B	LC		Unknown	20000-34000,26100	
<i>Acrocephalus agricola</i>	Paddyfield Warbler	B	LC		Stable	466000-703000,568000	
<i>Acrocephalus arundinaceus</i>	Great Reed-warbler	B	LC		Stable	4760000-8880000,6410000	
<i>Acrocephalus dumetorum</i>	Blyth's Reed-warbler	B	LC		Increasing	6240000-10900000,8210000	
<i>Acrocephalus melanopogon</i>	Moustached Warbler	B	LC		Stable	73800-188000,110000	
<i>Acrocephalus paludicola</i>	Aquatic Warbler	B	VU	B2ab(i,ii,iii,iv,v)	Decreasing	18300-28500,22700	Yes
<i>Acrocephalus palustris</i>	Marsh Warbler	B	LC		Unknown	11700000-17000000,14100000	Yes
<i>Acrocephalus schoenobaenus</i>	Sedge Warbler	B	LC		Unknown	9170000-13700000,11000000	
<i>Acrocephalus scirpaceus</i>	Common Reed-warbler	B	LC		Stable	3890000-6630000,5060000	
<i>Cettia cetti</i>	Cetti's Warbler	B	LC		Stable	4080000-6550000,5110000	
<i>Cisticola juncidis</i>	Zitting Cisticola	B	LC		Stable	2270000-3680000,2750000	
<i>Hippolais icterina</i>	Icterine Warbler	B	LC		Decreasing	4720000-8430000,6270000	Yes
<i>Hippolais languida</i>	Upcher's Warbler	B	LC		Unknown	36800-90200,55900	
<i>Hippolais olivetorum</i>	Olive-tree Warbler	B	LC		Stable	21500-47700,31800	Yes
<i>Hippolais polyglotta</i>	Melodious Warbler	B	LC		Increasing	4270000-7070000,5210000	Yes
<i>Iduna caligata</i>	Booted Warbler	B	LC		Increasing	800000-1010000,896000	
<i>Iduna opaca</i>	Isabelline Warbler	B	LC		Stable	10000-10000,10000	
<i>Iduna pallida</i>	Olivaceous Warbler	B	LC		Unknown	6510000-13600000,9360000	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (Europe)	IUCN Red List criteria (Europe)	Population trend	Population size [mature individuals] min-max, best estimate	(Near*) endemic to Europe?
<i>Iduna rama</i>	Sykes's Warbler	B	LC		Unknown	8000-14100,10600	
<i>Locustella fluviatilis</i>	River Warbler	B	LC		Decreasing	2780000-4960000,3670000	Yes
<i>Locustella lanceolata</i>	Lanceolated Warbler	B	LC		Unknown	40000-60000,49000	
<i>Locustella luscinioides</i>	Savi's Warbler	B	LC		Unknown	749000-1310000,981000	
<i>Locustella naevia</i>	Common Grasshopper-warbler	B	LC		Decreasing	1270000-2250000,1660000	
<i>Phylloscopus bonelli</i>	Western Bonelli's Warbler	B	LC		Increasing	4800000-6360000,5490000	Yes
<i>Phylloscopus borealis</i>	Arctic Warbler	B	LC		Increasing	6000000-8010000,6930000	
<i>Phylloscopus canariensis</i>	Canary Islands Chiffchaff	B	LC		Unknown	40000-200000,89400	Yes
<i>Phylloscopus collybita</i>	Common Chiffchaff	B	LC		Stable	72100000-114000000,89400000	
<i>Phylloscopus ibericus</i>	Iberian Chiffchaff	B	LC		Stable	924000-1540000,1190000	Yes
<i>Phylloscopus inornatus</i>	Yellow-browed Warbler	B	LC		Unknown	40000-60000,49000	
<i>Phylloscopus nitidus</i>	Green Warbler	B	LC		Increasing	935000-1870000,1320000	
<i>Phylloscopus orientalis</i>	Eastern Bonelli's Warbler	B	LC		Unknown	58500-196000,105000	Yes
<i>Phylloscopus sibilatrix</i>	Wood Warbler	B	LC		Decreasing	16200000-28500000,21400000	Yes
<i>Phylloscopus sindianus</i>	Mountain Chiffchaff	B	LC		Unknown	213000-644000,298000	
<i>Phylloscopus trochiloides</i>	Greenish Warbler	B	LC		Increasing	8080000-15300000,11100000	
<i>Phylloscopus trochilus</i>	Willow Warbler	B	LC		Decreasing	106000000-161000000,130000000	
<i>Prinia gracilis</i>	Graceful Prinia	B	LC		Unknown	2000-8000,4000	
<i>Sylvia atricapilla</i>	Eurasian Blackcap	B	LC		Increasing	88400000-138000000,108000000	Yes
<i>Curruca balearica</i>	Balearic Warbler	B	LC		Stable	28000-50000,37400	Yes
<i>Sylvia borin</i>	Garden Warbler	B	LC		Decreasing	23200000-38900000,30200000	Yes
<i>Curruca cantillans</i>	Subalpine Warbler	B	LC		Increasing	7040000-10700000,8490000	Yes
<i>Curruca communis</i>	Common Whitethroat	B	LC		Increasing	39200000-59200000,47800000	
<i>Curruca conspicillata</i>	Spectacled Warbler	B	LC		Stable	327000-891000,535000	
<i>Curruca crassirostris</i>	Eastern Orphean Warbler	B	LC		Unknown	131000-411000,231000	
<i>Curruca curruca</i>	Lesser Whitethroat	B	LC		Stable	9750000-16700000,12600000	
<i>Curruca hortensis</i>	Western Orphean Warbler	B	LC		Increasing	247000-493000,349000	
<i>Curruca melanocephala</i>	Sardinian Warbler	B	LC		Stable	15400000-32100000,21400000	
<i>Curruca melanothorax</i>	Cyprus Warbler	B	LC		Stable	120000-180000,147000	Yes
<i>Curruca mystacea</i>	Menetries's Warbler	B	LC		Increasing	107000-254000,162000	
<i>Curruca nana</i>	Asian Desert Warbler	B	VU ⁰	D1	Unknown	80-140,110	
<i>Curruca nisoria</i>	Barred Warbler	B	LC		Increasing	1150000-2110000,1530000	
<i>Curruca ruppeli</i>	Rüppel's Warbler	B	LC		Unknown	206000-1030000,458000	Yes
<i>Curruca sarda</i>	Marmora's Warbler	B	LC		Unknown	120000-130000,124000	Yes
<i>Curruca subalpina</i>	Moltoni's Warbler	B	LC		Increasing	102000-265000,164000	Yes
<i>Curruca undata</i>	Dartford Warbler	B	NT	A2bc+3bc+4bc	Decreasing	1190000-2860000,1750000	Yes
<i>Regulus ignicapilla</i>	Common Firecrest	B	LC		Increasing	9120000-16300000,12100000	Yes
<i>Regulus madeirensis</i>	Madeira Firecrest	B	LC		Increasing	100000-200000,141000	Yes
<i>Regulus regulus</i>	Goldcrest	B	LC		Decreasing	29100000-50400000,37400000	
<i>Troglodytes troglodytes</i>	Northern Wren	B	LC		Increasing	66500000-113000000,80200000	
<i>Ficedula albicollis</i>	Collared Flycatcher	B	LC		Stable	3210000-5130000,4010000	Yes
<i>Ficedula hypoleuca</i>	European Pied Flycatcher	B	LC		Decreasing	15500000-24900000,19600000	Yes
<i>Ficedula parva</i>	Red-breasted Flycatcher	B	LC		Increasing	3590000-6870000,4930000	Yes
<i>Ficedula semitorquata</i>	Semi-collared Flycatcher	B	LC		Unknown	60300-297000,123000	Yes
<i>Muscicapa striata</i>	Spotted Flycatcher	B	LC		Decreasing	20600000-37400000,27900000	
<i>Aegithalos caudatus</i>	Long-tailed Tit	B	LC		Stable	16300000-29200000,21500000	
<i>Cyanistes caeruleus</i>	Eurasian Blue Tit	B	LC		Increasing	59700000-95100000,73400000	Yes
<i>Cyanistes cyanus</i>	Azure Tit	B	NT	C1	Decreasing	5000-15700,8800	
<i>Cyanistes teneriffae</i>	African Blue Tit	B	LC		Unknown	40600-201000,90100	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (Europe)	IUCN Red List criteria (Europe)	Population trend	Population size [mature individuals] min-max, best estimate	(Near*) endemic to Europe?
<i>Lophophanes cristatus</i>	Crested Tit	B	LC		Decreasing	9230000-16500000,12100000	Yes
<i>Parus major</i>	Great Tit	B	LC		Stable	127000000-205000000,158000000	
<i>Periparus ater</i>	Coal Tit	B	LC		Stable	31500000-55700000,40600000	
<i>Poecile cinctus</i>	Siberian Tit	B	LC		Decreasing	507000-1100000,755000	
<i>Poecile hyrcanus</i>	Caspian Tit	B	LC		Stable	1000-5000,2200	
<i>Poecile lugubris</i>	Sombre Tit	B	LC		Unknown	902000-2670000,1550000	Yes
<i>Poecile montanus</i>	Willow Tit	B	LC		Decreasing	14000000-25400000,18800000	
<i>Poecile palustris</i>	Marsh Tit	B	LC		Stable	7480000-13300000,9850000	
<i>Remiz pendulinus</i>	Eurasian Penduline-tit	B	LC		Increasing	518000-940000,689000	
<i>Sitta europaea</i>	Eurasian Nuthatch	B	LC		Stable	1900000-34700000,25100000	
<i>Sitta krueperi</i>	Krueper's Nuthatch	B	LC		Unknown	256000-944000,485000	Yes
<i>Sitta neumayer</i>	Western Rock Nuthatch	B	LC		Unknown	2030000-10200000,4540000	Yes
<i>Sitta tephronota</i>	Eastern Rock Nuthatch	B	LC		Unknown	33400-107000,59300	
<i>Sitta whiteheadi</i>	Corsican Nuthatch	B	VU	C2a(ii)	Decreasing	3000-4400,3800	Yes
<i>Tichodroma muraria</i>	Wallcreeper	B	LC		Unknown	26300-96900,48800	
<i>Certhia brachydactyla</i>	Short-toed Treecreeper	B	LC		Increasing	10100000-16800000,12900000	Yes
<i>Certhia familiaris</i>	Eurasian Treecreeper	B	LC		Increasing	10900000-19200000,14400000	
<i>Lanius collurio</i>	Red-backed Shrike	B	LC		Decreasing	16400000-26000000,20400000	
<i>Lanius excubitor</i>	Great Grey Shrike	B	LC		Decreasing	128000-250000,175000	
<i>Lanius meridionalis</i>	Iberian Grey Shrike	B	VU	A2abc+3bc+4abc	Decreasing	721000-1220000,932000	Yes
<i>Lanius minor</i>	Lesser Grey Shrike	B	LC		Decreasing	752000-1610000,1060000	
<i>Lanius nubicus</i>	Masked Shrike	B	LC		Unknown	72400-213000,124000	
<i>Lanius senator</i>	Woodchat Shrike	B	NT	A2bcde+3bcde+4bcde	Decreasing	3860000-6240000,4890000	
<i>Pycnonotus xanthopygos</i>	White-spectacled Bulbul	B	LC		Unknown	12000-36000,20800	
<i>Argya altirostris</i>	Iraq Babbler	B	NT ⁰⁰	D1	Unknown	80-160,110	
<i>Corvus corax</i>	Common Raven	B	LC		Increasing	1170000-3570000,1720000	
<i>Corvus corone</i>	Carrion Crow	B	LC		Stable	16200000-27500000,20800000	
<i>Corvus frugilegus</i>	Rook	B	VU	A2bcde+4bcde	Decreasing	11400000-21300000,15000000	
<i>Corvus monedula</i>	Eurasian Jackdaw	B	LC		Increasing	17800000-36200000,24700000	
<i>Cyanopica cooki</i>	Iberian Azure-winged Magpie	B	LC		Increasing	713000-1800000,1080000	Yes
<i>Garrulus glandarius</i>	Eurasian Jay	B	LC		Stable	12000000-23100000,16400000	
<i>Nucifraga caryocatactes</i>	Northern Nutcracker	B	LC		Stable	596000-1220000,846000	
<i>Perisoreus infaustus</i>	Siberian Jay	B	LC		Unknown	426000-830000,615000	
<i>Pica pica</i>	Eurasian Magpie	B	LC		Stable	19300000-31300000,24100000	
<i>Pyrrhocorax graculus</i>	Yellow-billed Chough	B	LC		Unknown	218000-550000,340000	
<i>Pyrrhocorax pyrrhocorax</i>	Red-billed Chough	B	LC		Unknown	84600-197000,120000	
<i>Pastor roseus</i>	Rosy Starling	B	LC		Fluctuating	101000-403000,187000	
<i>Sturnus unicolor</i>	Spotless Starling	B	LC		Stable	47700000-61400000,54100000	Yes
<i>Sturnus vulgaris</i>	Common Starling	B	LC		Stable	53300000-94400000,69900000	
<i>Oriolus oriolus</i>	Eurasian Golden Oriole	B	LC		Stable	8740000-14600000,11200000	
<i>Carospiza brachydactyla</i>	Pale Sparrow	B	LC		Stable	12700-24100,17300	
<i>Gymnoris xanthocollis</i>	Chestnut-shouldered Bush-sparrow	B	LC		Unknown	1000-4000,2000	
<i>Montifringilla nivalis</i>	White-winged Snowfinch	B	LC		Unknown	255000-1120000,526000	
<i>Passer domesticus</i>	House Sparrow	B	LC		Decreasing	270000000-378000000,316000000	
<i>Passer hispaniolensis</i>	Spanish Sparrow	B	LC		Unknown	7560000-20000000,12000000	
<i>Passer italiae</i>	Italian Sparrow	B	VU	A2bce+3bce+4bce	Decreasing	4240000-6900000,5360000	Yes
<i>Passer moabiticus</i>	Dead Sea Sparrow	B	LC		Unknown	2000-6000,3500	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (Europe)	IUCN Red List criteria (Europe)	Population trend	Population size [mature individuals] min-max, best estimate	(Near*) endemic to Europe?
<i>Passer montanus</i>	Eurasian Tree Sparrow	B	LC		Decreasing	53700000-76000000,63300000	
<i>Petronia petronia</i>	Rock Sparrow	B	LC		Increasing	4320000-9340000,6270000	
<i>Acanthis flammea</i>	Redpoll	B	LC		Increasing	9010000-19100000,13200000	
<i>Bucanetes githagineus</i>	Trumpeter Finch	B	LC		Unknown	21300-44000,30500	
<i>Bucanetes mongolicus</i>	Mongolian Finch	B	LC		Unknown	1100-3200,1800	
<i>Carduelis carduelis</i>	European Goldfinch	B	LC		Increasing	56100000-88700000,68900000	
<i>Carduelis citrinella</i>	Citrl Finch	B	LC		Fluctuating	502000-569000,530000	Yes
<i>Carduelis corsicana</i>	Corsican Finch	B	LC		Unknown	26000-51000,36400	Yes
<i>Carpodacus erythrinus</i>	Common Rosefinch	B	LC		Decreasing	6830000-14000000,9730000	
<i>Carpodacus rubicilla</i>	Great Rosefinch	B	LC		Unknown	5800-23600,10500	
<i>Chloris chloris</i>	European Greenfinch	B	LC		Decreasing	41500000-64400000,50800000	Yes
<i>Coccothraustes coccothraustes</i>	Hawfinch	B	LC		Increasing	7280000-12100000,9290000	
<i>Fringilla coelebs</i>	Common Chaffinch	B	LC		Stable	30800000-46200000,37100000	
<i>Fringilla montifringilla</i>	Brambling	B	LC		Decreasing	14000000-26000000,19500000	
<i>Fringilla polatzeki</i>	Gran Canaria Blue Chaffinch	B	EN	D	Increasing	430-430,430	Yes
<i>Fringilla teydea</i>	Tenerife Blue Chaffinch	B	NT	D2	Increasing	2000-5000,3200	Yes
<i>Linaria cannabina</i>	Common Linnet	B	LC		Increasing	35800000-64400000,46100000	
<i>Linaria flavirostris</i>	Twite	B	LC		Unknown	159000-548000,285000	
<i>Pinicola enucleator</i>	Pine Grosbeak	B	LC		Stable	254000-448000,341000	
<i>Pyrrhula murina</i>	Azores Bullfinch	B	VU	D1+2	Stable	510-1700,970	Yes
<i>Pyrrhula pyrrhula</i>	Eurasian Bullfinch	B	LC		Increasing	9790000-17300000,12900000	
<i>Rhodopechys sanguineus</i>	Eurasian Crimson-winged Finch	B	LC		Unknown	107000-411000,209000	
<i>Rhodospiza obsoleta</i>	Desert Finch	B	LC		Fluctuating	6100-12400,8700	
<i>Serinus canaria</i>	Island Canary	B	LC		Stable	2380000-4970000,3820000	Yes
<i>Serinus pusillus</i>	Red-fronted Serin	B	LC		Unknown	594000-1920000,1050000	
<i>Serinus serinus</i>	European Serin	B	LC		Decreasing	35200000-55900000,43000000	Yes
<i>Spinus spinus</i>	Eurasian Siskin	B	LC		Stable	13700000-25800000,19000000	
<i>Loxia curvirostra</i>	Red Crossbill	B	LC		Stable	5610000-14200000,9020000	
<i>Loxia leucoptera</i>	Two-barred Crossbill	B	LC		Stable	204000-613000,354000	
<i>Loxia pytyopsittacus</i>	Parrot Crossbill	B	LC		Fluctuating	416000-957000,648000	Yes
<i>Loxia scotica</i>	Scottish Crossbill	B	LC		Unknown	8100-22700,13600	Yes
<i>Calcarius lapponicus</i>	Lapland Longspur	B	LC		Unknown	3120000-5170000,4000000	
<i>Emberiza aureola</i>	Yellow-breasted Bunting	B	CR	A2bcd+3bcd+4bcd	Decreasing	1200-2500,1700	
<i>Emberiza bruniceps</i>	Red-headed Bunting	B	LC		Increasing	8400-25000,14500	
<i>Emberiza buchanani</i>	Grey-necked Bunting	B	LC		Unknown	13800-40300,23400	
<i>Emberiza caesia</i>	Cretzschmar's Bunting	B	LC		Unknown	230000-451000,320000	Yes
<i>Emberiza calandra</i>	Corn Bunting	B	LC		Decreasing	42200000-67800000,52100000	
<i>Emberiza cia</i>	Rock Bunting	B	LC		Decreasing	3880000-8580000,5640000	
<i>Emberiza cineracea</i>	Cinereous Bunting	B	NT	C1	Decreasing	5300-10400,7400	Yes
<i>Emberiza cirrus</i>	Cirl Bunting	B	LC		Decreasing	4910000-9150000,6570000	Yes
<i>Emberiza citrinella</i>	Yellowhammer	B	LC		Decreasing	39000000-54000000,45500000	
<i>Emberiza hortulana</i>	Ortolan Bunting	B	LC		Decreasing	7220000-11300000,8940000	
<i>Emberiza leucocephalos</i>	Pine Bunting	B	RE		N/A	0-0,0	
<i>Emberiza melanocephala</i>	Black-headed Bunting	B	LC		Unknown	4940000-17900000,9080000	
<i>Emberiza pallasi</i>	Pallas's Bunting	B	LC		Unknown	20000-34000,26100	
<i>Emberiza pusilla</i>	Little Bunting	B	LC		Unknown	5020000-7070000,5960000	
<i>Emberiza rustica</i>	Rustic Bunting	B	LC		Decreasing	834000-1640000,1220000	
<i>Emberiza schoeniclus</i>	Reed Bunting	B	LC		Decreasing	9500000-16700000,12300000	
<i>Plectrophenax nivalis</i>	Snow Bunting	B	LC		Unknown	2510000-4940000,3480000	

Appendix 2. Red List status, trend and population size estimate, as well as endemism of birds in the EU28⁶³

Scientific species name	Common species name	Season of assessment	IUCN Red List category (EU28)	IUCN Red List criteria (EU28)	Population trend (EU28)	Population size [mature individuals] min-max, best estimate (EU28)	(Near*) endemic to EU28?
<i>Cygnus columbianus</i>	Tundra Swan	W	VU	A4abcde	Decreasing	11400-16100,13300	
<i>Cygnus cygnus</i>	Whooper Swan	B	LC		Increasing	33700-47400,41300	
<i>Cygnus olor</i>	Mute Swan	B	LC		Increasing	134000-190000,159000	
<i>Anser albifrons</i>	Greater White-fronted Goose	W	LC		Increasing	1040000-1570000,1190000	
<i>Anser anser</i>	Greylag Goose	B	LC		Increasing	448000-652000,539000	
<i>Anser brachyrhynchus</i>	Pink-footed Goose	W	LC		Increasing	370000-378000,373000	Yes
<i>Anser erythropus</i>	Lesser White-fronted Goose	B	CR	D	Decreasing	30-60,46	
<i>Anser fabalis</i>	Bean Goose	B	VU	C1+2a(ii)	Decreasing	2500-6200,4200	
<i>Branta bernicla</i>	Brent Goose	W	LC		Increasing	198000-225000,211000	
<i>Branta canadensis</i>	Canada Goose	B	N/A		N/A	182000-217000,199000	
<i>Branta leucopsis</i>	Barnacle Goose	B	LC		Increasing	59400-77700,71200	
<i>Branta ruficollis</i>	Red-breasted Goose	W	VU	A2bcd+3bcd+4bcd	Decreasing	1800-34600,7900	Yes*
<i>Tadorna ferruginea</i>	Ruddy Shelduck	B	NT ^o	D1	Increasing	460-1800,800	
<i>Tadorna tadorna</i>	Common Shelduck	B	LC		Increasing	69500-99600,81100	
<i>Anas acuta</i>	Northern Pintail	B	EN	A2bcde+3bcde+4bcde	Decreasing	9100-37600,22400	
<i>Anas crecca</i>	Common Teal	B	LC		Decreasing	469000-734000,598000	
<i>Anas platyrhynchos</i>	Mallard	B	LC		Stable	3860000-6330000,4950000	
<i>Mareca penelope</i>	Eurasian Wigeon	B	VU	A2bcde+3bcde+4bcde	Decreasing	83300-141000,108000	
<i>Mareca strepera</i>	Gadwall	B	LC		Increasing	100000-161000,122000	
<i>Marmaronetta angustirostris</i>	Marbled Teal	B	EN	D	Stable	140-160,150	
<i>Spatula clypeata</i>	Northern Shoveler	B	NT	A2bcde+3bcde+4bcde	Decreasing	48300-69300,58000	
<i>Spatula querquedula</i>	Garganey	B	VU	A2bcde+3bcde+4bcde	Decreasing	16100-38600,23500	
<i>Aythya ferina</i>	Common Pochard	B	VU	A2abcde+4abcde	Decreasing	57000-110000,77500	
<i>Aythya fuligula</i>	Tufted Duck	B	VU	A2abcde+4abcde	Decreasing	333000-575000,458000	
<i>Aythya marila</i>	Greater Scaup	B	EN	A2bcde+4bcde; C1	Decreasing	2100-3000,2500	
<i>Aythya nyroca</i>	Ferruginous Duck	B	LC		Unknown	9900-30300,17100	
<i>Bucephala clangula</i>	Common Goldeneye	B	LC		Decreasing	374000-481000,421000	
<i>Clangula hyemalis</i>	Long-tailed Duck	W	LC		Decreasing	773000-1420000,993000	
<i>Melanitta fusca</i>	Velvet Scoter	B	VU	A2abcde	Decreasing	22000-47900,35200	Yes*
<i>Melanitta nigra</i>	Common Scoter	B	LC		Increasing	12100-19800,15400	
<i>Mergellus albellus</i>	Smew	B	LC		Stable	5600-13900,9200	
<i>Mergus merganser</i>	Goosander	B	LC		Increasing	126000-209000,163000	
<i>Mergus serrator</i>	Red-breasted Merganser	B	NT	A2bcde+3bcde+4bcde	Decreasing	89300-148000,117000	
<i>Netta rufina</i>	Red-crested Pochard	B	LC		Increasing	22700-30800,25400	
<i>Oxyura leucocephala</i>	White-headed Duck	B	VU	D1	Fluctuating	510-620,620	
<i>Polysticta stelleri</i>	Steller's Eider	W	EN	A2bcd+3bcd+4bcd; C1	Decreasing	530-1100,730	
<i>Somateria mollissima</i>	Common Eider	B	VU	A2abcde+4abcde	Decreasing	353000-592000,466000	
<i>Bonasa bonasia</i>	Hazel Grouse	B	VU	A2bcd+3bcd+4bcd	Decreasing	1110000-1900000,1420000	
<i>Lagopus lagopus</i>	Willow Grouse	B	LC		Decreasing	749000-1530000,1090000	
<i>Lagopus muta</i>	Rock Ptarmigan	B	LC		Increasing	227000-494000,332000	
<i>Lyrurus tetrix</i>	Black Grouse	B	VU	A2bcde+3bcde+4bcde	Decreasing	1040000-1740000,1330000	
<i>Tetrao urogallus</i>	Western Capercaillie	B	LC		Decreasing	937000-2330000,1260000	
<i>Alectoris barbara</i>	Barbary Partridge	B	LC		Unknown	15500-41000,25100	
<i>Alectoris chukar</i>	Chukar	B	LC		Stable	521000-747000,620000	
<i>Alectoris graeca</i>	Rock Partridge	B	NT	A2bcd+3bcd+4bcd	Decreasing	51100-75400,61600	
<i>Alectoris rufa</i>	Red-legged Partridge	B	NT	A2bcde+3bcde+4bcde	Decreasing	9950000-13700000,11700000	Yes*
<i>Coturnix coturnix</i>	Common Quail	B	LC		Unknown	2850000-5020000,3710000	
<i>Francolinus francolinus</i>	Black Francolin	B	LC		Increasing	7200-15200,10500	
<i>Perdix perdix</i>	Grey Partridge	B	VU	A2bcde+3bcde+4bcde	Decreasing	1370000-2470000,1740000	
<i>Phasianus colchicus</i>	Common Pheasant	B	EN	D	Stable	100-200,140	
<i>Phasianus colchicus</i>	Common Pheasant	B	LC		Stable	5700000-7690000,6410000	
<i>Turnix sylvaticus</i>	Common Buttonquail	B	RE		Unknown	0-0,0	
<i>Gavia arctica</i>	Arctic Loon	B	LC		Increasing	29600-38900,34300	
<i>Gavia immer</i>	Common Loon	W	LC		Increasing	4600-5200,4800	
<i>Gavia stellata</i>	Red-throated Loon	B	LC		Increasing	6100-9500,7500	
<i>Podiceps auritus</i>	Horned Grebe	B	VU	C1	Decreasing	8900-11400,9900	
<i>Podiceps cristatus</i>	Great Crested Grebe	B	LC		Decreasing	367000-526000,437000	
<i>Podiceps grisegena</i>	Red-necked Grebe	B	VU	A2bcde+3bcde+4bcde	Decreasing	14600-25100,19000	

⁶³ Red List Categories adjusted down by one/two/three Categories because of rescue effect from outside the region are denoted with °/°°/°°°, respectively.

Scientific species name	Common species name	Season of assessment	IUCN Red List category (EU28)	IUCN Red List criteria (EU28)	Population trend (EU28)	Population size [mature individuals] min-max, best estimate (EU28)	(Near*) endemic to EU28?
<i>Podiceps nigricollis</i>	Black-necked Grebe	B	LC		Decreasing	13800-27400,18700	
<i>Tachybaptus ruficollis</i>	Little Grebe	B	LC		Stable	136000-275000,184000	
<i>Bulweria bulwerii</i>	Bulwer's Petrel	B	LC		Unknown	92100-103000,97000	
<i>Calonectris borealis</i>	Cory's Shearwater	B	LC		Unknown	503000-506000,505000	Yes
<i>Calonectris diomedea</i>	Scopoli's Shearwater	B	LC		Unknown	57000-94200,72800	
<i>Fulmarus glacialis</i>	Northern Fulmar	B	EN	A4abcde	Decreasing	457000-1430000,769000	
<i>Hydrobates castro</i>	Band-rumped Storm-petrel	B	LC		Unknown	13200-4300,13700	
<i>Hydrobates leucorhous</i>	Leach's Storm-petrel	B	VU	B2ab(v)	Unknown	73400-131000,96700	
<i>Hydrobates monteiroi</i>	Monteiro's Storm-petrel	B	VU	D1+2	Unknown	720-770,740	Yes
<i>Hydrobates pelagicus</i>	European Storm-petrel	B	LC		Unknown	270000-308000,285000	
<i>Pelagodroma marina</i>	White-faced Storm-petrel	B	EN	B2ab(iii,v)	Stable	155000-222000,186000	
<i>Pterodroma deserta</i>	Desertas Petrel	B	VU	D1+2	Stable	320-360,340	Yes
<i>Pterodroma madeira</i>	Zino's Petrel	B	EN	D	Increasing	130-160,140	Yes
<i>Puffinus lherminieri</i>	Audubon's Shearwater	B	NT	B2ab(v); C1+2a(i)	Stable	5000-11800,7600	
<i>Puffinus mauretanicus</i>	Balearic Shearwater	B	CR	A4abcde	Decreasing	5800-5900,5800	Yes
<i>Puffinus puffinus</i>	Manx Shearwater	B	LC		Unknown	631000-718000,672000	Yes*
<i>Puffinus yelkouan</i>	Yelkouan Shearwater	B	VU	A2abcde	Decreasing	47000-80700,61400	Yes*
<i>Morus bassanus</i>	Northern Gannet	B	LC		Increasing	721000-722000,722000	Yes*
<i>Pelecanus crispus</i>	Dalmatian Pelican	B	LC		Increasing	4400-5600,5000	
<i>Pelecanus onocrotalus</i>	Great White Pelican	B	LC		Increasing	17200-37900,25500	
<i>Gulosus aristotelis</i>	European Shag	B	LC		Decreasing	70800-88700,80100	
<i>Microcarbo pygmaeus</i>	Pygmy Cormorant	B	LC		Increasing	32300-36600,34200	
<i>Phalacrocorax carbo</i>	Great Cormorant	B	LC		Increasing	444000-533000,489000	
<i>Ardea alba</i>	Great White Egret	B	LC		Increasing	15900-22800,18900	
<i>Ardea cinerea</i>	Grey Heron	B	LC		Increasing	264000-320000,290000	
<i>Ardea purpurea</i>	Purple Heron	B	LC		Decreasing	23500-38100,28600	
<i>Ardeola ralloides</i>	Squacco Heron	B	LC		Decreasing	13100-22000,16700	
<i>Botaurus stellaris</i>	Eurasian Bittern	B	LC		Decreasing	25400-37100,30600	
<i>Bubulcus ibis</i>	Cattle Egret	B	LC		Increasing	128000-156000,142000	
<i>Ciconia ciconia</i>	White Stork	B	LC		Increasing	313000-335000,324000	
<i>Ciconia nigra</i>	Black Stork	B	LC		Increasing	13300-20700,16500	
<i>Egretta garzetta</i>	Little Egret	B	LC		Decreasing	78300-95900,85600	
<i>Ixobrychus minutus</i>	Common Little Bittern	B	LC		Unknown	75600-142000,107000	
<i>Nycticorax nycticorax</i>	Black-crowned Night-heron	B	LC		Decreasing	51900-72100,60200	
<i>Platalea leucorodia</i>	Eurasian Spoonbill	B	LC		Increasing	16800-23900,20300	
<i>Plegadis falcinellus</i>	Glossy Ibis	B	LC		Increasing	22400-63900,34800	
<i>Phoenicopterus roseus</i>	Greater Flamingo	B	LC		Increasing	49200-169000,98300	
<i>Aegypius monachus</i>	Cinereous Vulture	B	LC		Increasing	5200-5400,5300	
<i>Gypaetus barbatus</i>	Bearded Vulture	B	VU	D1	Increasing	450-500,460	
<i>Gyps fulvus</i>	Griffon Vulture	B	LC		Increasing	67400-83800,72400	
<i>Neophron percnopterus</i>	Egyptian Vulture	B	VU	C1	Decreasing	3400-3800,3600	
<i>Aquila adalberti</i>	Spanish Imperial Eagle	B	VU	D1	Increasing	1000-1100,1100	Yes
<i>Aquila chrysaetos</i>	Golden Eagle	B	LC		Increasing	10400-12500,11400	
<i>Aquila fasciata</i>	Bonelli's Eagle	B	LC		Stable	2100-2400,2200	
<i>Aquila heliaca</i>	Eastern Imperial Eagle	B	NT*	D1	Increasing	540-770,640	
<i>Circus gallicus</i>	Short-toed Snake-eagle	B	LC		Increasing	13700-20700,16700	
<i>Clanga clanga</i>	Greater Spotted Eagle	B	CR	C1+2a(i); D	Decreasing	36-70,45	
<i>Clanga pomarina</i>	Lesser Spotted Eagle	B	LC		Increasing	21700-30400,25700	
<i>Haliaeetus albicilla</i>	White-tailed Sea-eagle	B	LC		Increasing	9700-12700,10900	
<i>Hieraetus pennatus</i>	Booted Eagle	B	LC		Increasing	41300-49400,44600	
<i>Pandion haliaetus</i>	Osprey	B	LC		Increasing	12100-15500,14000	
<i>Milvus migrans</i>	Black Kite	B	LC		Increasing	102000-127000,114000	
<i>Milvus milvus</i>	Red Kite	B	LC		Increasing	59400-69600,64000	Yes*
<i>Circus aeruginosus</i>	Western Marsh-harrier	B	LC		Stable	106000-176000,134000	
<i>Circus cyaneus</i>	Hen Harrier	B	VU	A2bcd+3bcd+4bcd	Decreasing	21100-30300,25200	
<i>Circus macrourus</i>	Pallid Harrier	B	EN*	D	Increasing	20-100,40	
<i>Circus pygargus</i>	Montagu's Harrier	B	VU	A2bc+3bc+4bc	Decreasing	28600-40400,33400	
<i>Buteo buteo</i>	Eurasian Buzzard	B	LC		Increasing	1100000-1620000,1380000	
<i>Buteo lagopus</i>	Rough-legged Buzzard	B	VU	A2bcd+3bcd+4bcd; C1	Decreasing	3600-16400,8000	
<i>Buteo rufinus</i>	Long-legged Buzzard	B	LC		Decreasing	2300-4200,3100	
<i>Pernis apivorus</i>	European Honey-buzzard	B	LC		Stable	95600-151000,118000	
<i>Accipiter brevipes</i>	Levant Sparrowhawk	B	LC		Stable	3400-6800,4800	
<i>Accipiter gentilis</i>	Northern Goshawk	B	LC		Decreasing	113000-191000,148000	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (EU28)	IUCN Red List criteria (EU28)	Population trend (EU28)	Population size [mature individuals] min-max, best estimate (EU28)	(Near*) endemic to EU28?
<i>Accipiter nisus</i>	Eurasian Sparrowhawk	B	LC		Stable	397000-684000,546000	
<i>Elanus caeruleus</i>	Black-winged Kite	B	LC		Stable	2600-5600,3800	
<i>Falco biarmicus</i>	Lanner Falcon	B	EN	C1	Decreasing	280-330,300	
<i>Falco cherrug</i>	Saker Falcon	B	EN	C2a(ii)	Decreasing	430-620,540	
<i>Falco columbarius</i>	Merlin	B	VU	A2bc+3bc+4bc	Decreasing	13400-28700,21100	
<i>Falco eleonora</i>	Eleonora's Falcon	B	LC		Stable	28300-28800,28500	Yes*
<i>Falco naumanni</i>	Lesser Kestrel	B	LC		Stable	58000-64900,61200	
<i>Falco peregrinus</i>	Peregrine Falcon	B	LC		Increasing	22500-26600,24300	
<i>Falco rusticolus</i>	Gyr Falcon	B	EN	C1+2a(i)	Decreasing	190-260,220	
<i>Falco subbuteo</i>	Eurasian Hobby	B	LC		Stable	81200-155000,106000	
<i>Falco tinnunculus</i>	Common Kestrel	B	LC		Decreasing	617000-954000,743000	
<i>Falco vespertinus</i>	Red-footed Falcon	B	VU	C1	Decreasing	5000-8000,6300	
<i>Crex crex</i>	Corncrake	B	NT	A2bcde+3bcde+4bcde	Decreasing	254000-537000,365000	
<i>Fulica atra</i>	Common Coot	B	LC		Stable	1080000-1720000,1360000	
<i>Fulica cristata</i>	Red-knobbed Coot	B	CR	C2a(ii)	Decreasing	42-90,60	
<i>Gallinula chloropus</i>	Common Moorhen	B	LC		Decreasing	1410000-1970000,1650000	
<i>Porphyrio porphyrio</i>	Purple Swamphen	B	LC		Decreasing	8100-9500,8700	
<i>Porzana porzana</i>	Spotted Crake	B	LC		Unknown	17100-38300,25000	
<i>Rallus aquaticus</i>	Western Water Rail	B	LC		Unknown	163000-391000,245000	
<i>Zapornia parva</i>	Little Crake	B	LC		Unknown	17400-92700,37200	
<i>Zapornia pusilla</i>	Baillon's Crake	B	NT*	D1	Unknown	230-700,370	
<i>Grus grus</i>	Common Crane	B	LC		Increasing	248000-337000,290000	
<i>Chlamydotis undulata</i>	African Houbara	B	VU	D1	Increasing	610-620,620	
<i>Otis tarda</i>	Great Bustard	B	LC		Increasing	32000-37400,34600	
<i>Tetrax tetrax</i>	Little Bustard	B	EN	A2bce+3bce+4bce	Decreasing	69500-150000,101000	
<i>Actitis hypoleucos</i>	Common Sandpiper	B	LC		Decreasing	521000-789000,623000	
<i>Arenaria interpres</i>	Ruddy Turnstone	B	EN	A2bce+4bce	Decreasing	2800-6800,4900	
<i>Burhinus oedicephalus</i>	Eurasian Thick-knee	B	LC		Unknown	112000-172000,137000	
<i>Calidris alba</i>	Sanderling	W	LC		Increasing	50000-66000,57100	
<i>Calidris alpina</i>	Dunlin	B	LC		Decreasing	61000-102000,81100	
<i>Calidris canutus</i>	Red Knot	W	LC		Decreasing	268000-325000,296000	
<i>Calidris falcinellus</i>	Broad-billed Sandpiper	B	VU	A2bc+3bc+4bc	Decreasing	36700-65800,48300	
<i>Calidris ferruginea</i>	Curlew Sandpiper	W	VU	D1	Increasing	100-1100,420	
<i>Calidris maritima</i>	Purple Sandpiper	W	NT	A2abc	Decreasing	9400-11600,10300	
<i>Calidris minuta</i>	Little Stint	W	LC		Increasing	8500-14200,10900	
<i>Calidris pugnax</i>	Ruff	B	NT	A2bcde	Decreasing	41500-94700,66500	
<i>Calidris temminckii</i>	Temminck's Stint	B	NT	A2bce+3bce+4bce	Decreasing	13200-18000,15600	
<i>Charadrius alexandrinus</i>	Kentish Plover	B	LC		Decreasing	13600-28900,19200	
<i>Charadrius dubius</i>	Little Ringed Plover	B	LC		Unknown	145000-217000,174000	
<i>Charadrius hiaticula</i>	Common Ringed Plover	B	LC		Increasing	54600-80800,68500	
<i>Charadrius leschenaultii</i>	Greater Sandplover	W	CR	C2a(i); D	Decreasing	10-17,13	
<i>Cursorius cursor</i>	Cream-coloured Courser	B	NT	D1	Unknown	1100-1200,1100	
<i>Eudromias morinellus</i>	Eurasian Dotterel	B	LC		Unknown	7500-15600,12100	
<i>Gallinago gallinago</i>	Common Snipe	B	LC		Decreasing	844000-1520000,1130000	
<i>Gallinago media</i>	Great Snipe	B	VU	C1	Decreasing	4400-7600,5900	
<i>Glareola nordmanni</i>	Black-winged Pratincole	B	RE		Decreasing	0-0,0	
<i>Glareola pratincola</i>	Collared Pratincole	B	NT	C1	Decreasing	9200-16500,11700	
<i>Haematopus ostralegus</i>	Eurasian Oystercatcher	B	VU	A4abc	Decreasing	350000-394000,371000	
<i>Himantopus himantopus</i>	Black-winged Stilt	B	LC		Increasing	49400-178000,90100	
<i>Limosa lapponica</i>	Bar-tailed Godwit	B	LC		Stable	2600-9000,3800	
<i>Limosa limosa</i>	Black-tailed Godwit	B	EN	A2bcde+3bcde+4bcde	Decreasing	75800-93400,84000	
<i>Lymnocyptes minimus</i>	Jack Snipe	B	LC		Unknown	17500-46700,31000	
<i>Numenius arquata</i>	Eurasian Curlew	B	NT	A2abcde+4bcde	Decreasing	310000-345000,330000	
<i>Numenius phaeopus</i>	Whimbrel	B	LC		Increasing	94300-136000,115000	
<i>Numenius tenuirostris</i>	Slender-billed Curlew	W	CR (PE)	D	Unknown	0-0,0	
<i>Phalaropus lobatus</i>	Red-necked Phalarope	B	LC		Decreasing	40100-72200,56100	
<i>Pluvialis apricaria</i>	Eurasian Golden Plover	B	LC		Stable	415000-676000,552000	Yes*
<i>Pluvialis squatarola</i>	Grey Plover	W	LC		Decreasing	83300-100000,91300	
<i>Recurvirostra avosetta</i>	Pied Avocet	B	LC		Decreasing	61600-124000,70700	
<i>Scolopax rusticola</i>	Eurasian Woodcock	B	LC		Decreasing	1410000-2940000,2050000	
<i>Tringa erythropus</i>	Spotted Redshank	B	NT	A2bc+3bc+4bc	Decreasing	17900-49000,30900	
<i>Tringa glareola</i>	Wood Sandpiper	B	LC		Stable	845000-1130000,990000	
<i>Tringa nebularia</i>	Common Greenshank	B	LC		Stable	136000-209000,165000	
<i>Tringa ochropus</i>	Green Sandpiper	B	LC		Decreasing	435000-676000,538000	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (EU28)	IUCN Red List criteria (EU28)	Population trend (EU28)	Population size [mature individuals] min-max, best estimate (EU28)	(Near*) endemic to EU28?
<i>Tringa stagnatilis</i>	Marsh Sandpiper	B	VU ^o	D1	Increasing	80-220,130	
<i>Tringa totanus</i>	Common Redshank	B	VU	A2bcde+3bcde+4bcde	Decreasing	185000-233000,204000	
<i>Vanellus spinosus</i>	Spur-winged Lapwing	B	VU ^o	D1	Increasing	160-260,200	
<i>Vanellus vanellus</i>	Northern Lapwing	B	VU	A2bce+4bce	Decreasing	1630000-2250000,1920000	
<i>Xenus cinereus</i>	Terek Sandpiper	B	CR	D	Stable	10-16,12	
<i>Catharacta skua</i>	Great Skua	B	LC		Increasing	19200-19300,19300	
<i>Stercorarius longicaudus</i>	Long-tailed Jaeger	B	LC		Fluctuating	27200-56000,44000	
<i>Stercorarius parasiticus</i>	Arctic Jaeger	B	EN	A2bcd+4bcd	Decreasing	2800-5800,3800	
<i>Hydrocoloeus minutus</i>	Little Gull	B	LC		Stable	23500-31900,27600	
<i>Larus argentatus</i>	European Herring Gull	B	VU	A2bcde+3bcde+4bcde	Decreasing	839000-958000,895000	
<i>Larus armenicus</i>	Armenian Gull	W	VU ^o	D1	Unknown	33-140,70	
<i>Larus audouinii</i>	Audouin's Gull	B	VU	A3bce+4abce	Decreasing	31200-41700,36100	Yes*
<i>Larus cachinnans</i>	Caspian Gull	B	LC		Increasing	11200-19100,14500	
<i>Larus canus</i>	Mew Gull	B	LC		Decreasing	442000-635000,545000	
<i>Larus fuscus</i>	Lesser Black-backed Gull	B	LC		Increasing	617000-675000,647000	
<i>Larus genei</i>	Slender-billed Gull	B	LC		Unknown	10900-17600,13700	
<i>Larus glaucooides</i>	Iceland Gull	W	LC ^{oo}		Unknown	250-300,270	
<i>Larus hyperboreus</i>	Glaucous Gull	W	LC ^{oo}		Unknown	160-220,190	
<i>Larus ichthyaetus</i>	Pallas's Gull	B	NE		N/A	140-240,180	
<i>Larus marinus</i>	Great Black-backed Gull	B	NT	A2bcde+3bcde+4bcde	Decreasing	56300-89900,76100	
<i>Larus melanocephalus</i>	Mediterranean Gull	B	LC		Increasing	33600-50400,40100	
<i>Larus michahellis</i>	Yellow-legged Gull	B	LC		Unknown	767000-1020000,865000	Yes*
<i>Larus ridibundus</i>	Black-headed Gull	B	VU	A2bcde	Decreasing	1630000-2180000,1880000	
<i>Onychoprion fuscatus</i>	Sooty Tern	B	VU ^{oo}	D1	Unknown	2-2,2	
<i>Rissa tridactyla</i>	Black-legged Kittiwake	B	EN	A2abcd+3bcd+4abcd	Decreasing	415000-587000,481000	
<i>Gelochelidon nilotica</i>	Common Gull-billed Tern	B	LC		Increasing	16100-19700,17900	
<i>Hydroprogne caspia</i>	Caspian Tern	B	LC		Decreasing	3400-4100,3800	
<i>Sterna dougallii</i>	Roseate Tern	B	LC		Increasing	4900-6200,5500	
<i>Sterna hirundo</i>	Common Tern	B	LC		Increasing	253000-410000,324000	
<i>Sterna paradisaea</i>	Arctic Tern	B	LC		Increasing	359000-565000,413000	
<i>Sternula albifrons</i>	Little Tern	B	LC		Decreasing	25700-37400,30700	
<i>Thalasseus sandvicensis</i>	Sandwich Tern	B	LC		Increasing	118000-140000,128000	
<i>Chlidonias hybrida</i>	Whiskered Tern	B	LC		Unknown	38300-82900,58200	
<i>Chlidonias leucopterus</i>	White-winged Tern	B	LC		Fluctuating	1000-11700,3400	
<i>Chlidonias niger</i>	Black Tern	B	LC		Stable	23000-35500,28400	
<i>Alca torda</i>	Razorbill	B	LC		Increasing	262000-515000,371000	
<i>Alle alle</i>	Little Auk	W	NE		N/A	0-20,5	
<i>Cephus grylle</i>	Black Guillemot	B	LC		Decreasing	76000-93700,84800	
<i>Fratercula arctica</i>	Atlantic Puffin	B	LC		Increasing	1200000-1210000,1200000	
<i>Uria aalge</i>	Common Murre	B	LC		Increasing	3300000-3340000,3320000	
<i>Pterocles alchata</i>	Pin-tailed Sandgrouse	B	LC		Stable	14900-18000,16300	
<i>Pterocles orientalis</i>	Black-bellied Sandgrouse	B	EN	A2bce+3bce+4bce	Decreasing	3400-8400,6300	
<i>Columba bollii</i>	Dark-tailed Laurel-pigeon	B	LC		Unknown	2500-10000,5000	Yes
<i>Columba junoniae</i>	White-tailed Laurel-pigeon	B	NT	B1ab(iii)+2ab(iii)	Unknown	1000-2500,1600	Yes
<i>Columba livia</i>	Rock Dove	B	LC		Stable	4910000-9470000,6800000	
<i>Columba livia</i>	Rock Dove	B	N/A			20637000-37651000,27875000	
<i>Columba oenas</i>	Stock Dove	B	LC		Increasing	1280000-2110000,1580000	Yes*
<i>Columba palumbus</i>	Common Woodpigeon	B	LC		Increasing	38500000-52700000,44600000	Yes*
<i>Columba trocaz</i>	Madeira Laurel-pigeon	B	LC		Increasing	10000-14000,11800	Yes
<i>Spilopelia senegalensis</i>	Laughing Dove	B	NT ^{oo}	D1	Unknown	130-150,140	
<i>Streptopelia decaocto</i>	Eurasian Collared-dove	B	LC		Increasing	13900000-34900000,19600000	
<i>Streptopelia turtur</i>	European Turtle-dove	B	NT	A2bcde+3bcde+4bcde	Decreasing	3960000-6880000,5260000	
<i>Clamator glandarius</i>	Great Spotted Cuckoo	B	VU	A2bcd+3bcd+4bcd	Decreasing	289000-304000,294000	
<i>Cuculus canorus</i>	Common Cuckoo	B	LC		Decreasing	4490000-7290000,5700000	
<i>Aegolius funereus</i>	Boreal Owl	B	LC		Fluctuating	41900-255000,117000	
<i>Asio flammeus</i>	Short-eared Owl	B	LC		Fluctuating	4300-43200,16600	
<i>Asio otus</i>	Northern Long-eared Owl	B	LC		Unknown	256000-485000,372000	
<i>Athene noctua</i>	Little Owl	B	LC		Unknown	413000-678000,524000	
<i>Bubo bubo</i>	Eurasian Eagle-owl	B	LC		Increasing	26000-36400,30300	
<i>Bubo scandiacus</i>	Snowy Owl	B	CR	D	Fluctuating	0-100,4	
<i>Glaucidium passerinum</i>	Eurasian Pygmy-owl	B	LC		Stable	62100-140000,114000	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (EU28)	IUCN Red List criteria (EU28)	Population trend (EU28)	Population size [mature individuals] min-max, best estimate (EU28)	(Near*) endemic to EU28?
<i>Otus scops</i>	Eurasian Scops-owl	B	LC		Unknown	244000-377000,299000	
<i>Strix aluco</i>	Tawny Owl	B	LC		Stable	874000-1280000,1210000	
<i>Strix nebulosa</i>	Great Grey Owl	B	LC		Increasing	1300-3200,2100	
<i>Strix uralensis</i>	Ural Owl	B	LC		Increasing	23800-42000,73100	
<i>Surnia ulula</i>	Northern Hawk-owl	B	LC		Fluctuating	2800-42400,8600	
<i>Tyto alba</i>	Common Barn-owl	B	LC		Stable	152000-333000,223000	
<i>Caprimulgus europaeus</i>	European Nightjar	B	LC		Unknown	360000-671000,488000	
<i>Caprimulgus ruficollis</i>	Red-necked Nightjar	B	NT	A2bc+3bc+4bc	Decreasing	202000-270000,233000	
<i>Apus affinis</i>	Little Swift	B	VU ^o	D1	Increasing	44-44,44	
<i>Apus apus</i>	Common Swift	B	NT	A2bc+3bc+4bc	Decreasing	24800000-39500000,30900000	
<i>Apus caffer</i>	White-rumped Swift	B	NT ^o	D1	Increasing	500-700,590	
<i>Apus pallidus</i>	Pallid Swift	B	LC		Unknown	322000-2190000,781000	
<i>Apus unicolor</i>	Plain Swift	B	LC		Unknown	15000-40000,24100	Yes
<i>Tachymarptis melba</i>	Alpine Swift	B	LC		Unknown	406000-967000,622000	
<i>Upupa epops</i>	Common Hoopoe	B	LC		Stable	2140000-4990000,3120000	
<i>Alcedo atthis</i>	Common Kingfisher	B	LC		Stable	117000-313000,184000	
<i>Coracias garrulus</i>	European Roller	B	LC		Unknown	26500-44000,32500	
<i>Merops apiaster</i>	European Bee-eater	B	LC		Stable	6670000-9920000,8020000	
<i>Dendrocopos leucotos</i>	White-backed Woodpecker	B	LC		Unknown	72900-176000,111000	
<i>Dendrocopos major</i>	Great Spotted Woodpecker	B	LC		Increasing	9140000-14000000,11100000	
<i>Dendrocopos syriacus</i>	Syrian Woodpecker	B	LC		Unknown	172000-386000,257000	
<i>Dryobates minor</i>	Lesser Spotted Woodpecker	B	NT	A2b+3b+4b	Decreasing	376000-789000,534000	
<i>Dryocopus martius</i>	Black Woodpecker	B	LC		Stable	416000-707000,539000	
<i>Jynx torquilla</i>	Eurasian Wryneck	B	LC		Decreasing	628000-1230000,881000	
<i>Leipicus medius</i>	Middle Spotted Woodpecker	B	LC		Stable	602000-1070000,797000	
<i>Picoides tridactylus</i>	Three-toed Woodpecker	B	LC		Decreasing	86600-184000,122000	
<i>Picus canus</i>	Grey-faced Woodpecker	B	LC		Unknown	170000-278000,216000	
<i>Picus sharpei</i>	Iberian Green Woodpecker	B	LC		Decreasing	488000-938000,671000	Yes*
<i>Picus viridis</i>	Eurasian Green Woodpecker	B	LC		Stable	1050000-1790000,1370000	
<i>Panurus biarmicus</i>	Bearded Reedling	B	LC		Unknown	107000-215000,151000	
<i>Alauda arvensis</i>	Eurasian Skylark	B	LC		Decreasing	56100000-72200000,62800000	
<i>Alaudala rufescens</i>	Lesser Short-toed Lark	B	LC		Decreasing	480000-561000,517000	
<i>Calandrella brachydactyla</i>	Greater Short-toed Lark	B	LC		Increasing	2950000-4680000,3720000	
<i>Chersophilus duponti</i>	Dupont's Lark	B	VU	A2bc+3bc+4bc;C1+2a(i)	Decreasing	4600-4700,4700	
<i>Eremophila alpestris</i>	Horned Lark	B	NT	C1	Decreasing	6300-13300,9100	
<i>Galerida cristata</i>	Crested Lark	B	LC		Decreasing	29800000-38100000,33600000	
<i>Galerida theklae</i>	Thekla's Lark	B	LC		Increasing	3500000-5690000,4380000	
<i>Lullula arborea</i>	Woodlark	B	LC		Stable	3530000-6360000,4660000	
<i>Melanocorypha calandra</i>	Calandra Lark	B	LC		Decreasing	8540000-12900000,10600000	
<i>Cecropis daurica</i>	Red-rumped Swallow	B	LC		Increasing	2740000-6380000,4260000	
<i>Delichon urbicum</i>	Northern House Martin	B	LC		Decreasing	17100000-37200000,25000000	
<i>Hirundo rustica</i>	Barn Swallow	B	LC		Decreasing	38300000-61000000,47200000	
<i>Ptyonoprogne rupestris</i>	Eurasian Crag Martin	B	LC		Stable	265000-493000,343000	
<i>Riparia riparia</i>	Collared Sand Martin	B	LC		Stable	2420000-5600000,3540000	
<i>Anthus berthelotii</i>	Berthelot's Pipit	B	LC		Unknown	60000-300000,134000	Yes
<i>Anthus campestris</i>	Tawny Pipit	B	LC		Stable	1580000-2600000,2030000	
<i>Anthus cervinus</i>	Red-throated Pipit	B	EN	C1	Decreasing	1700-3600,2600	
<i>Anthus petrosus</i>	Rock Pipit	B	LC		Unknown	129000-165000,145000	
<i>Anthus pratensis</i>	Meadow Pipit	B	LC		Stable	9890000-14500000,11600000	
<i>Anthus spinoletta</i>	Water Pipit	B	LC		Unknown	685000-1300000,955000	
<i>Anthus trivialis</i>	Tree Pipit	B	LC		Stable	13900000-23100000,18100000	
<i>Motacilla alba</i>	White Wagtail	B	LC		Stable	13700000-21000000,16700000	
<i>Motacilla cinerea</i>	Grey Wagtail	B	LC		Stable	1930000-3880000,2770000	
<i>Motacilla citreola</i>	Citrine Wagtail	B	LC ^o		Increasing	920-1700,1200	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (EU28)	IUCN Red List criteria (EU28)	Population trend (EU28)	Population size [mature individuals] min-max, best estimate (EU28)	(Near*) endemic to EU28?
<i>Motacilla flava</i>	Western Yellow Wagtail	B	LC		Decreasing	15800000-21200000,18100000	
<i>Cinclus cinclus</i>	White-throated Dipper	B	LC		Unknown	146000-322000,211000	
<i>Bombycilla garrulus</i>	Bohemian Waxwing	B	LC		Increasing	300000-529000,401000	
<i>Prunella collaris</i>	Alpine Accentor	B	LC		Unknown	96800-169000,126000	
<i>Prunella modularis</i>	Dunnock	B	LC		Decreasing	19400000-31900000,23200000	
<i>Cercotrichas galactotes</i>	Rufous-tailed Scrub-robin	B	LC		Decreasing	204000-544000,334000	
<i>Cyanecula svecica</i>	Bluethroat	B	LC		Stable	471000-969000,710000	
<i>Erithacus rubecula</i>	European Robin	B	LC		Stable	78000000-116000000,93400000	
<i>Luscinia luscinia</i>	Thrush Nightingale	B	VU	A2bc+3bc+4bc	Decreasing	958000-1570000,1210000	
<i>Luscinia megarhynchos</i>	Common Nightingale	B	LC		Stable	17300000-24200000,20300000	
<i>Monticola saxatilis</i>	Rufous-tailed Rock-thrush	B	LC		Unknown	42800-76900,56300	
<i>Monticola solitarius</i>	Blue Rock-thrush	B	LC		Stable	86600-179000,122000	
<i>Oenanthe cyprica</i>	Cyprus Wheatear	B	LC		Increasing	100000-200000,141000	Yes
<i>Oenanthe finschii</i>	Finsch's Wheatear	W	NT ^{oo}	D1	Unknown	100-200,140	
<i>Oenanthe hispanica</i>	Black-eared Wheatear	B	LC		Decreasing	902000-2380000,1450000	
<i>Oenanthe isabellina</i>	Isabelline Wheatear	B	LC		Increasing	28400-86000,49400	
<i>Oenanthe leucura</i>	Black Wheatear	B	LC		Increasing	8100-32200,13000	
<i>Oenanthe oenanthe</i>	Northern Wheatear	B	NT	A2bcde+4bcde	Decreasing	4060000-7630000,5600000	
<i>Oenanthe pleschanka</i>	Pied Wheatear	B	LC ^o		Unknown	1200-2400,1700	
<i>Phoenicurus ochruros</i>	Black Redstart	B	LC		Stable	9750000-15500000,12200000	
<i>Phoenicurus phoenicurus</i>	Common Redstart	B	LC		Increasing	4540000-8240000,6290000	
<i>Saxicola dacotiae</i>	Fuerteventura Stonechat	B	NT	B1ab(ii,iii); C2a(ii)	Unknown	13300-15500,14400	Yes
<i>Saxicola rubetra</i>	Whinchat	B	VU	A2bc+3bc+4bc	Decreasing	4890000-7440000,6020000	
<i>Saxicola torquatus</i>	Common Stonechat	B	LC		Decreasing	11000000-17000000,13600000	
<i>Tarsiger cyanurus</i>	Orange-flanked Bush-robin	B	LC		Increasing	1500-11100,3000	
<i>Turdus iliacus</i>	Redwing	B	LC		Decreasing	3920000-4730000,4380000	
<i>Turdus merula</i>	Eurasian Blackbird	B	LC		Increasing	100000000-150000000,120000000	
<i>Turdus philomelos</i>	Song Thrush	B	LC		Increasing	27400000-44400000,34000000	
<i>Turdus pilaris</i>	Fieldfare	B	LC		Increasing	7350000-11400000,9220000	
<i>Turdus torquatus</i>	Ring Ouzel	B	LC		Unknown	329000-636000,448000	
<i>Turdus viscivorus</i>	Mistle Thrush	B	LC		Stable	4990000-9410000,6690000	
<i>Acrocephalus agricola</i>	Paddyfield Warbler	B	LC		Unknown	6100-40400,15700	
<i>Acrocephalus arundinaceus</i>	Great Reed-warbler	B	LC		Decreasing	2510000-3940000,3130000	
<i>Acrocephalus dumetorum</i>	Blyth's Reed-warbler	B	LC		Increasing	240000-626000,381000	
<i>Acrocephalus melanopogon</i>	Moustached Warbler	B	LC		Stable	21600-76300,37200	
<i>Acrocephalus paludicola</i>	Aquatic Warbler	B	NT	C2a(ii)	Increasing	6500-9500,7800	
<i>Acrocephalus palustris</i>	Marsh Warbler	B	LC		Decreasing	5280000-8030000,6490000	
<i>Acrocephalus schoenobaenus</i>	Sedge Warbler	B	LC		Decreasing	3090000-4760000,3700000	
<i>Acrocephalus scirpaceus</i>	Common Reed-warbler	B	LC		Stable	3100000-5290000,4050000	
<i>Cettia cetti</i>	Cetti's Warbler	B	LC		Stable	3580000-5420000,4370000	
<i>Cisticola juncidis</i>	Zitting Cisticola	B	LC		Stable	2270000-3660000,2740000	
<i>Hippolais icterina</i>	Icterine Warbler	B	LC		Stable	2300000-3660000,2880000	
<i>Hippolais olivetorum</i>	Olive-tree Warbler	B	LC		Stable	9500-19000,13400	
<i>Hippolais polyglotta</i>	Melodious Warbler	B	LC		Increasing	4270000-7070000,5210000	Yes*
<i>Iduna caligata</i>	Booted Warbler	B	LC		Increasing	920-3500,1800	
<i>Iduna opaca</i>	Isabelline Warbler	B	LC		Stable	10000-10000,10000	
<i>Iduna pallida</i>	Olivaceous Warbler	B	LC		Increasing	336000-855000,536000	
<i>Locustella fluviatilis</i>	River Warbler	B	VU	A2bc+3bc+4bc	Decreasing	320000-653000,432000	
<i>Locustella luscinioides</i>	Savi's Warbler	B	LC		Unknown	433000-743000,561000	
<i>Locustella naevia</i>	Common Grasshopper-warbler	B	VU	A2bc+3bc+4bc	Decreasing	637000-1140000,823000	
<i>Phylloscopus bonelli</i>	Western Bonelli's Warbler	B	LC		Increasing	4720000-6240000,5390000	
<i>Phylloscopus borealis</i>	Arctic Warbler	B	VU	C1+2a(ii)	Decreasing	320-1700,800	
<i>Phylloscopus canariensis</i>	Canary Islands Chiffchaff	B	LC		Unknown	40000-200000,89400	Yes
<i>Phylloscopus collybita</i>	Common Chiffchaff	B	LC		Increasing	42900000-61900000,50800000	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (EU28)	IUCN Red List criteria (EU28)	Population trend (EU28)	Population size [mature individuals] min-max, best estimate (EU28)	(Near*) endemic to EU28?
<i>Phylloscopus ibericus</i>	Iberian Chiffchaff	B	LC		Stable	924000-1540000,1190000	Yes*
<i>Phylloscopus orientalis</i>	Eastern Bonelli's Warbler	B	LC		Unknown	28000-74100,45300	
<i>Phylloscopus sibilatrix</i>	Wood Warbler	B	LC		Decreasing	6980000-10800000,8640000	
<i>Phylloscopus trochiloides</i>	Greenish Warbler	B	LC		Increasing	82200-249000,141000	
<i>Phylloscopus trochilus</i>	Willow Warbler	B	LC		Decreasing	49300000-83000000,64000000	
<i>Sylvia atricapilla</i>	Eurasian Blackcap	B	LC		Increasing	69900000-108000000,84400000	
<i>Curruca balearica</i>	Balearic Warbler	B	LC		Stable	28000-50000,37400	Yes
<i>Sylvia borin</i>	Garden Warbler	B	LC		Decreasing	9990000-17100000,13200000	
<i>Curruca cantillans</i>	Subalpine Warbler	B	LC		Increasing	6980000-10500000,8380000	
<i>Curruca communis</i>	Common Whitethroat	B	LC		Stable	20400000-28500000,23900000	
<i>Curruca conspicillata</i>	Spectacled Warbler	B	LC		Stable	326000-889000,533000	
<i>Curruca crassirostris</i>	Eastern Orphean Warbler	B	LC		Unknown	32000-124000,61700	
<i>Curruca curruca</i>	Lesser Whitethroat	B	LC		Stable	5960000-8260000,6960000	
<i>Curruca hortensis</i>	Western Orphean Warbler	B	LC		Increasing	247000-492000,349000	
<i>Curruca melanocephala</i>	Sardinian Warbler	B	LC		Stable	13800000-27300000,18600000	
<i>Curruca melanothorax</i>	Cyprus Warbler	B	LC		Stable	120000-180000,147000	Yes
<i>Curruca nisoria</i>	Barred Warbler	B	LC		Unknown	669000-1190000,885000	
<i>Curruca ruppeli</i>	Rüppell's Warbler	B	LC		Stable	6000-20000,11000	
<i>Curruca sarda</i>	Marmora's Warbler	B	LC		Unknown	120000-130000,124000	
<i>Curruca subalpina</i>	Moltoni's Warbler	B	LC		Increasing	102000-265000,164000	Yes
<i>Curruca undata</i>	Dartford Warbler	B	NT	A2bc+3bc+4bc	Decreasing	1190000-2860000,1750000	Yes*
<i>Regulus ignicapilla</i>	Common Firecrest	B	LC		Increasing	8540000-15100000,11300000	Yes*
<i>Regulus madeirensis</i>	Madeira Firecrest	B	LC		Increasing	100000-200000,141000	Yes
<i>Regulus regulus</i>	Goldcrest	B	LC		Stable	20800000-33200000,25700000	
<i>Troglodytes troglodytes</i>	Northern Wren	B	LC		Increasing	60000000-97900000,70500000	
<i>Ficedula albicollis</i>	Collared Flycatcher	B	LC		Unknown	1730000-2850000,2210000	
<i>Ficedula hypoleuca</i>	European Pied Flycatcher	B	LC		Decreasing	4440000-6160000,5180000	
<i>Ficedula parva</i>	Red-breasted Flycatcher	B	LC		Unknown	610000-1250000,872000	
<i>Ficedula semitorquata</i>	Semi-collared Flycatcher	B	LC		Unknown	7800-23000,12700	
<i>Muscicapa striata</i>	Spotted Flycatcher	B	LC		Decreasing	8850000-16800000,12400000	
<i>Aegithalos caudatus</i>	Long-tailed Tit	B	LC		Stable	12000000-19300000,15100000	
<i>Cyanistes caeruleus</i>	Eurasian Blue Tit	B	LC		Stable	51600000-80100000,62500000	
<i>Cyanistes teneriffae</i>	African Blue Tit	B	LC		Unknown	40600-201000,90100	
<i>Lophophanes cristatus</i>	Crested Tit	B	LC		Stable	7070000-12900000,9380000	
<i>Parus major</i>	Great Tit	B	LC		Increasing	94000000-138000000,112000000	
<i>Periparus ater</i>	Coal Tit	B	LC		Increasing	24700000-41000000,30900000	
<i>Poecile cinctus</i>	Siberian Tit	B	NT	A2b	Decreasing	143000-391000,249000	
<i>Poecile lugubris</i>	Sombre Tit	B	LC		Stable	51000-169000,89100	
<i>Poecile montanus</i>	Willow Tit	B	NT	A2b	Decreasing	4070000-6540000,5120000	
<i>Poecile palustris</i>	Marsh Tit	B	LC		Decreasing	6230000-10900000,8150000	
<i>Remiz pendulinus</i>	Eurasian Penduline-tit	B	LC		Unknown	250000-436000,327000	
<i>Sitta europaea</i>	Eurasian Nuthatch	B	LC		Increasing	13500000-23500000,17400000	
<i>Sitta krueperi</i>	Krueper's Nuthatch	B	NT	D1	Increasing	1000-1400,1200	
<i>Sitta neumayer</i>	Western Rock Nuthatch	B	LC		Stable	22400-63000,37300	
<i>Sitta whiteheadi</i>	Corsican Nuthatch	B	VU	C2a(ii)	Decreasing	3000-4400,3800	Yes
<i>Tichodroma muraria</i>	Wallcreeper	B	LC		Unknown	11600-25300,16800	
<i>Certhia brachydactyla</i>	Short-toed Treecreeper	B	LC		Increasing	9570000-15500000,12000000	Yes*

Scientific species name	Common species name	Season of assessment	IUCN Red List category (EU28)	IUCN Red List criteria (EU28)	Population trend (EU28)	Population size [mature individuals] min-max, best estimate (EU28)	(Near*) endemic to EU28?
<i>Certhia familiaris</i>	Eurasian Treecreeper	B	LC		Increasing	7630000-12600000,9800000	
<i>Lanius collurio</i>	Red-backed Shrike	B	LC		Decreasing	10800000-14700000,12600000	
<i>Lanius excubitor</i>	Great Grey Shrike	B	VU	A2bc+3bc+4bc	Decreasing	83500-178000,119000	
<i>Lanius meridionalis</i>	Iberian Grey Shrike	B	VU	A2abc+3bc+4abc	Decreasing	721000-1220000,932000	Yes
<i>Lanius minor</i>	Lesser Grey Shrike	B	LC		Unknown	237000-532000,354000	
<i>Lanius nubicus</i>	Masked Shrike	B	LC		Stable	10400-28400,17100	
<i>Lanius senator</i>	Woodchat Shrike	B	NT	A2bcde+3bcde+4bcde	Decreasing	3780000-5970000,4740000	
<i>Corvus corax</i>	Common Raven	B	LC		Increasing	583000-1110000,791000	
<i>Corvus corone</i>	Carrion Crow	B	LC		Increasing	10800000-16800000,13300000	
<i>Corvus frugilegus</i>	Rook	B	LC		Decreasing	5850000-11100000,7530000	
<i>Corvus monedula</i>	Eurasian Jackdaw	B	LC		Increasing	11500000-20100000,14800000	
<i>Cyanopica cooki</i>	Iberian Azure-winged Magpie	B	LC		Increasing	713000-1800000,1080000	Yes
<i>Garrulus glandarius</i>	Eurasian Jay	B	LC		Stable	8930000-15200000,11500000	
<i>Nucifraga caryocatactes</i>	Northern Nutcracker	B	LC		Unknown	260000-621000,400000	
<i>Perisoreus infaustus</i>	Siberian Jay	B	LC		Fluctuating	141000-310000,231000	
<i>Pica pica</i>	Eurasian Magpie	B	LC		Stable	14100000-20700000,16900000	
<i>Pyrrhoxorax graculus</i>	Yellow-billed Chough	B	LC		Stable	83500-170000,116000	
<i>Pyrrhoxorax pyrrhoxorax</i>	Red-billed Chough	B	LC		Increasing	45200-51500,47800	
<i>Pastor roseus</i>	Rosy Starling	B	LC		Unknown	2000-27000,5100	
<i>Sturnus unicolor</i>	Spotless Starling	B	LC		Decreasing	47700000-61400000,54100000	Yes*
<i>Sturnus vulgaris</i>	Common Starling	B	LC		Decreasing	39000000-64900000,49900000	
<i>Oriolus oriolus</i>	Eurasian Golden Oriole	B	LC		Stable	4750000-7710000,5990000	
<i>Montifringilla nivalis</i>	White-winged Snowfinch	B	LC		Unknown	31800-57800,42700	
<i>Passer domesticus</i>	House Sparrow	B	LC		Stable	228000000-300000000,260000000	
<i>Passer hispaniolensis</i>	Spanish Sparrow	B	LC		Increasing	4170000-12600000,7090000	
<i>Passer italiae</i>	Italian Sparrow	B	VU	A2bce+3bce+4bce	Decreasing	4200000-6850000,5310000	Yes*
<i>Passer montanus</i>	Eurasian Tree Sparrow	B	LC		Stable	24000000-35400000,28800000	
<i>Petronia petronia</i>	Rock Sparrow	B	LC		Increasing	2670000-4360000,3410000	
<i>Acanthis flammea</i>	Redpoll	B	LC		Increasing	1750000-4180000,2820000	
<i>Bucanetes githagineus</i>	Trumpeter Finch	B	LC		Unknown	20600-41200,29100	
<i>Carduelis carduelis</i>	European Goldfinch	B	LC		Stable	48000000-69400000,56800000	
<i>Carduelis citrinella</i>	Citrel Finch	B	LC		Fluctuating	480000-525000,499000	
<i>Carduelis corsicana</i>	Corsican Finch	B	LC		Unknown	26000-51000,36400	Yes
<i>Carpodacus erythrinus</i>	Common Rosefinch	B	NT	A2bce+3bce+4bce	Decreasing	736000-1180000,932000	
<i>Chloris chloris</i>	European Greenfinch	B	LC		Decreasing	33100000-46400000,38800000	
<i>Coccothraustes coccothraustes</i>	Hawfinch	B	LC		Increasing	4920000-8130000,6290000	
<i>Fringilla coelebs</i>	Common Chaffinch	B	LC		Stable	170000000-242000000,201000000	
<i>Fringilla montifringilla</i>	Brambling	B	LC		Stable	4670000-8950000,6850000	
<i>Fringilla polatzeki</i>	Gran Canaria Blue Chaffinch	B	EN	D	Increasing	310-320,320	Yes
<i>Fringilla teydea</i>	Tenerife Blue Chaffinch	B	NT	D2	Increasing	2000-5000,3200	
<i>Linaria cannabina</i>	Common Linnet	B	LC		Increasing	28500000-39800000,33300000	
<i>Linaria flavirostris</i>	Twite	B	VU	A2bc+3bc+4bc	Decreasing	12000-21600,16500	
<i>Pinicola enucleator</i>	Pine Grosbeak	B	LC		Fluctuating	13600-45500,29700	
<i>Pyrrhula murina</i>	Azores Bullfinch	B	VU	D1+2	Stable	510-1700,970	Yes
<i>Pyrrhula pyrrhula</i>	Eurasian Bullfinch	B	LC		Increasing	4780000-8620000,6350000	
<i>Serinus canaria</i>	Island Canary	B	LC		Stable	2380000-4970000,3820000	Yes
<i>Serinus serinus</i>	European Serin	B	LC		Decreasing	33700000-51000000,40300000	Yes*
<i>Spinus spinus</i>	Eurasian Siskin	B	LC		Stable	5390000-9440000,7320000	
<i>Loxia curvirostra</i>	Red Crossbill	B	LC		Increasing	1450000-5460000,3040000	
<i>Loxia leucoptera</i>	Two-barred Crossbill	B	LC		Stable	4400-112000,37500	

Scientific species name	Common species name	Season of assessment	IUCN Red List category (EU28)	IUCN Red List criteria (EU28)	Population trend (EU28)	Population size [mature individuals] min-max, best estimate (EU28)	(Near*) endemic to EU28?
<i>Loxia pytyopsittacus</i>	Parrot Crossbill	B	LC		Increasing	211000-507000,349000	
<i>Loxia scotica</i>	Scottish Crossbill	B	LC		Unknown	8100-22700,13600	Yes
<i>Calcarius lapponicus</i>	Lapland Longspur	B	VU	A2b+3b+4b	Decreasing	326000-716000,494000	
<i>Emberiza aureola</i>	Yellow-breasted Bunting	B	CR(PE)	D	Decreasing	0-0,0	
<i>Emberiza caesia</i>	Cretzschmar's Bunting	B	LC		Increasing	30000-80000,48300	
<i>Emberiza calandra</i>	Corn Bunting	B	LC		Decreasing	35100000-46000000,39900000	
<i>Emberiza cia</i>	Rock Bunting	B	LC		Decreasing	3160000-5870000,4260000	
<i>Emberiza cineracea</i>	Cinereous Bunting	B	EN	C2a(i); D	Decreasing	160-400,250	
<i>Emberiza cirius</i>	Cirl Bunting	B	LC		Stable	4690000-8580000,6220000	Yes*
<i>Emberiza citrinella</i>	Yellowhammer	B	LC		Decreasing	27400000-37700000,31800000	
<i>Emberiza hortulana</i>	Ortolan Bunting	B	NT	A2bcde+3bcde+4bcde	Decreasing	2060000-3500000,2660000	
<i>Emberiza melanocephala</i>	Black-headed Bunting	B	LC		Unknown	561000-1130000,778000	
<i>Emberiza pusilla</i>	Little Bunting	B	LC		Stable	25300-66400,42400	
<i>Emberiza rustica</i>	Rustic Bunting	B	LC		Stable	334000-638000,517000	
<i>Emberiza schoeniclus</i>	Reed Bunting	B	LC		Decreasing	3570000-5760000,4260000	
<i>Plectrophenax nivalis</i>	Snow Bunting	B	LC		Stable	38000-86200,62100	
<i>Lanius meridionalis</i>	Iberian Grey Shrike	B	VU	A2abc+3bc+4abc	Decreasing	721000-1220000,932000	Yes
<i>Lanius minor</i>	Lesser Grey Shrike	B	LC		Decreasing	752000-1610000,1060000	
<i>Lanius nubicus</i>	Masked Shrike	B	LC		Unknown	72400-213000,124000	
<i>Lanius senator</i>	Woodchat Shrike	B	NT	A2bcde+3bcde+4bcde	Decreasing	3860000-6240000,4890000	
<i>Pycnonotus xanthopygos</i>	White-spectacled Bulbul	B	LC		Unknown	12000-36000,20800	
<i>Argya altirostris</i>	Iraq Babbler	B	NT ⁰⁰	D1	Unknown	80-160,110	
<i>Corvus corax</i>	Common Raven	B	LC		Increasing	1170000-3570000,1720000	
<i>Corvus corone</i>	Carrion Crow	B	LC		Stable	16200000-27500000,20800000	
<i>Corvus frugilegus</i>	Rook	B	VU	A2bcde+4bcde	Decreasing	11400000-21300000,15000000	
<i>Corvus monedula</i>	Eurasian Jackdaw	B	LC		Increasing	17800000-36200000,24700000	
<i>Cyanopica cooki</i>	Iberian Azure-winged Magpie	B	LC		Increasing	713000-1800000,1080000	Yes
<i>Garrulus glandarius</i>	Eurasian Jay	B	LC		Stable	12000000-23100000,16400000	
<i>Nucifraga caryocatactes</i>	Northern Nutcracker	B	LC		Stable	596000-1220000,846000	
<i>Perisoreus infaustus</i>	Siberian Jay	B	LC		Unknown	426000-830000,615000	
<i>Pica pica</i>	Eurasian Magpie	B	LC		Stable	19300000-31300000,24100000	
<i>Pyrrhonorax graculus</i>	Yellow-billed Chough	B	LC		Unknown	218000-550000,340000	
<i>Pyrrhonorax pyrrhonorax</i>	Red-billed Chough	B	LC		Unknown	84600-197000,120000	
<i>Pastor roseus</i>	Rosy Starling	B	LC		Fluctuating	101000-403000,187000	
<i>Sturnus unicolor</i>	Spotless Starling	B	LC		Stable	47700000-61400000,54100000	Yes
<i>Sturnus vulgaris</i>	Common Starling	B	LC		Stable	53300000-94400000,69900000	
<i>Oriolus oriolus</i>	Eurasian Golden Oriole	B	LC		Stable	8740000-14600000,11200000	
<i>Carpospiza brachydactyla</i>	Pale Sparrow	B	LC		Stable	12700-24100,17300	
<i>Gymnoris xanthocollis</i>	Chestnut-shouldered Bush-sparrow	B	LC		Unknown	1000-4000,2000	
<i>Montifringilla nivalis</i>	White-winged Snowfinch	B	LC		Unknown	255000-1120000,526000	
<i>Passer domesticus</i>	House Sparrow	B	LC		Decreasing	270000000-378000000,316000000	
<i>Passer hispaniolensis</i>	Spanish Sparrow	B	LC		Unknown	7560000-20000000,12000000	
<i>Passer italiae</i>	Italian Sparrow	B	VU	A2bce+3bce+4bce	Decreasing	4240000-6900000,5360000	Yes
<i>Passer moabiticus</i>	Dead Sea Sparrow	B	LC		Unknown	2000-6000,3500	
<i>Passer montanus</i>	Eurasian Tree Sparrow	B	LC		Decreasing	53700000-76000000,63300000	
<i>Petronia petronia</i>	Rock Sparrow	B	LC		Increasing	4320000-9340000,6270000	
<i>Acanthis flammea</i>	Redpoll	B	LC		Increasing	9010000-19100000,13200000	
<i>Bucanetes githagineus</i>	Trumpeter Finch	B	LC		Unknown	21300-44000,30500	
<i>Bucanetes mongolicus</i>	Mongolian Finch	B	LC		Unknown	1100-3200,1800	
<i>Carduelis carduelis</i>	European Goldfinch	B	LC		Increasing	56100000-88700000,68900000	
<i>Carduelis citrinella</i>	Citril Finch	B	LC		Fluctuating	502000-569000,530000	Yes

Scientific species name	Common species name	Season of assessment	IUCN Red List category (EU28)	IUCN Red List criteria (EU28)	Population trend (EU28)	Population size [mature individuals] min-max, best estimate (EU28)	(Near*) endemic to EU28?
<i>Carduelis corsicana</i>	Corsican Finch	B	LC		Unknown	26000-51000,36400	Yes
<i>Carpodacus erythrinus</i>	Common Rosefinch	B	LC		Decreasing	6830000-14000000,9730000	
<i>Carpodacus rubicilla</i>	Great Rosefinch	B	LC		Unknown	5800-23600,10500	
<i>Chloris chloris</i>	European Greenfinch	B	LC		Decreasing	41500000-64400000,50800000	Yes
<i>Coccothraustes coccothraustes</i>	Hawfinch	B	LC		Increasing	7280000-12100000,9290000	
<i>Fringilla coelebs</i>	Common Chaffinch	B	LC		Stable	308000000-462000000,371000000	
<i>Fringilla montifringilla</i>	Brambling	B	LC		Decreasing	14000000-26000000,19500000	
<i>Fringilla polatzeki</i>	Gran Canaria Blue Chaffinch	B	EN	D	Increasing	430-430,430	Yes
<i>Fringilla teydea</i>	Tenerife Blue Chaffinch	B	NT	D2	Increasing	2000-5000,3200	Yes
<i>Linaria cannabina</i>	Common Linnet	B	LC		Increasing	35800000-64400000,46100000	
<i>Linaria flavirostris</i>	Twite	B	LC		Unknown	159000-548000,285000	
<i>Pinicola enucleator</i>	Pine Grosbeak	B	LC		Stable	254000-448000,341000	
<i>Pyrrhula murina</i>	Azores Bullfinch	B	VU	D1+2	Stable	510-1700,970	Yes
<i>Pyrrhula pyrrhula</i>	Eurasian Bullfinch	B	LC		Increasing	9790000-17300000,12900000	
<i>Rhodopechys sanguineus</i>	Eurasian Crimson-winged Finch	B	LC		Unknown	107000-411000,209000	
<i>Rhodospiza obsoleta</i>	Desert Finch	B	LC		Fluctuating	6100-12400,8700	
<i>Serinus canaria</i>	Island Canary	B	LC		Stable	2380000-4970000,3820000	Yes
<i>Serinus pusillus</i>	Red-fronted Serin	B	LC		Unknown	594000-1920000,1050000	
<i>Serinus serinus</i>	European Serin	B	LC		Decreasing	35200000-55900000,43000000	Yes
<i>Spinus spinus</i>	Eurasian Siskin	B	LC		Stable	13700000-25800000,19000000	
<i>Loxia curvirostra</i>	Red Crossbill	B	LC		Stable	5610000-14200000,9020000	
<i>Loxia leucoptera</i>	Two-barred Crossbill	B	LC		Stable	204000-613000,354000	
<i>Loxia pytyopsittacus</i>	Parrot Crossbill	B	LC		Fluctuating	416000-957000,648000	Yes
<i>Loxia scotica</i>	Scottish Crossbill	B	LC		Unknown	8100-22700,13600	Yes
<i>Calcarius lapponicus</i>	Lapland Longspur	B	LC		Unknown	3120000-5170000,4000000	
<i>Emberiza aureola</i>	Yellow-breasted Bunting	B	CR	A2bcd+3bcd+4bcd	Decreasing	1200-2500,1700	
<i>Emberiza bruniceps</i>	Red-headed Bunting	B	LC		Increasing	8400-25000,14500	
<i>Emberiza buchanani</i>	Grey-necked Bunting	B	LC		Unknown	13800-40300,23400	
<i>Emberiza caesia</i>	Cretzschmar's Bunting	B	LC		Unknown	230000-451000,320000	Yes
<i>Emberiza calandra</i>	Corn Bunting	B	LC		Decreasing	42200000-67800000,52100000	
<i>Emberiza cia</i>	Rock Bunting	B	LC		Decreasing	3880000-8580000,5640000	
<i>Emberiza cineracea</i>	Cinereous Bunting	B	NT	C1	Decreasing	5300-10400,7400	Yes
<i>Emberiza cirrus</i>	Cirl Bunting	B	LC		Decreasing	4910000-9150000,6570000	Yes
<i>Emberiza citrinella</i>	Yellowhammer	B	LC		Decreasing	39000000-54000000,45500000	
<i>Emberiza hortulana</i>	Ortolan Bunting	B	LC		Decreasing	7220000-11300000,8940000	
<i>Emberiza leucocephalos</i>	Pine Bunting	B	RE		N/A	0-0,0	
<i>Emberiza melanocephala</i>	Black-headed Bunting	B	LC		Unknown	4940000-17900000,9080000	
<i>Emberiza pallasi</i>	Pallas's Bunting	B	LC		Unknown	20000-34000,26100	
<i>Emberiza pusilla</i>	Little Bunting	B	LC		Unknown	5020000-7070000,5960000	
<i>Emberiza rustica</i>	Rustic Bunting	B	LC		Decreasing	834000-1640000,1220000	
<i>Emberiza schoeniclus</i>	Reed Bunting	B	LC		Decreasing	9500000-16700000,12300000	
<i>Plectrophenax nivalis</i>	Snow Bunting	B	LC		Unknown	2510000-4940000,3480000	

Appendix 3. Red List status of birds in Europe and the EU28 (in alphabetical order)⁶⁴

Scientific species name	Common species name	IUCN Red List category (Europe)	IUCN Red List category (EU28)
<i>Acanthis flammea</i>	Redpoll	LC	LC
<i>Accipiter badius</i>	Shikra	NT ^o	N/A
<i>Accipiter brevipes</i>	Levant Sparrowhawk	LC	LC
<i>Accipiter gentilis</i>	Northern Goshawk	LC	LC
<i>Accipiter nisus</i>	Eurasian Sparrowhawk	LC	LC
<i>Acrocephalus agricola</i>	Paddyfield Warbler	LC	LC
<i>Acrocephalus arundinaceus</i>	Great Reed-warbler	LC	LC
<i>Acrocephalus dumetorum</i>	Blyth's Reed-warbler	LC	LC
<i>Acrocephalus melanopogon</i>	Moustached Warbler	LC	LC
<i>Acrocephalus paludicola</i>	Aquatic Warbler	VU	NT
<i>Acrocephalus palustris</i>	Marsh Warbler	LC	LC
<i>Acrocephalus schoenobaenus</i>	Sedge Warbler	LC	LC
<i>Acrocephalus scirpaceus</i>	Common Reed-warbler	LC	LC
<i>Actitis hypoleucos</i>	Common Sandpiper	LC	LC
<i>Aegithalos caudatus</i>	Long-tailed Tit	LC	LC
<i>Aegolius funereus</i>	Boreal Owl	LC	LC
<i>Aegypius monachus</i>	Cinereous Vulture	LC	LC
<i>Alauda arvensis</i>	Eurasian Skylark	LC	LC
<i>Alauda leucoptera</i>	White-winged Lark	LC	N/A
<i>Alaudala rufescens</i>	Lesser Short-toed Lark	LC	LC
<i>Alca torda</i>	Razorbill	LC	LC
<i>Alcedo atthis</i>	Common Kingfisher	LC	LC
<i>Alectoris barbara</i>	Barbary Partridge	LC	LC
<i>Alectoris chukar</i>	Chukar	LC	LC
<i>Alectoris graeca</i>	Rock Partridge	NT	NT
<i>Alectoris rufa</i>	Red-legged Partridge	NT	NT
<i>Alle alle</i>	Little Auk	LC	NE
<i>Ammomanes deserti</i>	Desert Lark	EN	N/A
<i>Ammoperdix griseogularis</i>	See-see Partridge	VU	N/A
<i>Anas acuta</i>	Northern Pintail	VU	EN
<i>Anas crecca</i>	Common Teal	LC	LC
<i>Anas platyrhynchos</i>	Mallard	LC	LC
<i>Anhinga rufa</i>	African Darter	RE	N/A
<i>Anser albifrons</i>	Greater White-fronted Goose	LC	LC
<i>Anser anser</i>	Greylag Goose	LC	LC
<i>Anser brachyrhynchus</i>	Pink-footed Goose	LC	LC
<i>Anser caerulescens</i>	Snow Goose	LC	N/A
<i>Anser erythropus</i>	Lesser White-fronted Goose	VU	CR
<i>Anser fabalis</i>	Bean Goose	LC	VU
<i>Anthropoides virgo</i>	Demoiselle Crane	EN	N/A
<i>Anthus berthelotii</i>	Berthelot's Pipit	LC	LC
<i>Anthus campestris</i>	Tawny Pipit	LC	LC
<i>Anthus cervinus</i>	Red-throated Pipit	LC	EN
<i>Anthus gustavi</i>	Pechora Pipit	CR (PE)	N/A
<i>Anthus hodgsoni</i>	Olive-backed Pipit	LC	N/A
<i>Anthus petrosus</i>	Rock Pipit	LC	LC
<i>Anthus pratensis</i>	Meadow Pipit	LC	LC
<i>Anthus spinoletta</i>	Water Pipit	LC	LC
<i>Anthus trivialis</i>	Tree Pipit	LC	LC
<i>Apus affinis</i>	Little Swift	NT	VU ^{oo}
<i>Apus apus</i>	Common Swift	NT	NT
<i>Apus caffer</i>	White-rumped Swift	NT ^o	NT ^o
<i>Apus pallidus</i>	Pallid Swift	LC	LC
<i>Apus unicolor</i>	Plain Swift	LC	LC
<i>Aquila adalberti</i>	Spanish Imperial Eagle	VU	VU
<i>Aquila chrysaetos</i>	Golden Eagle	LC	LC
<i>Aquila fasciata</i>	Bonelli's Eagle	LC	LC
<i>Aquila heliaca</i>	Eastern Imperial Eagle	LC	NT ^o
<i>Aquila nipalensis</i>	Steppe Eagle	CR	N/A
<i>Ardea alba</i>	Great White Egret	LC	LC

⁶⁴ Red List Categories adjusted down by one/two/three Categories because of rescue effect from outside the region are denoted with ^o / ^{oo} / ^{ooo}, respectively.

Scientific species name	Common species name	IUCN Red List category (Europe)	IUCN Red List category (EU28)
<i>Ardea cinerea</i>	Grey Heron	LC	LC
<i>Ardea purpurea</i>	Purple Heron	LC	LC
<i>Ardeola ralloides</i>	Squacco Heron	LC	LC
<i>Arenaria interpres</i>	Ruddy Turnstone	LC	EN
<i>Argya altostris</i>	Iraq Babbler	NT ⁰⁰	N/A
<i>Asio flammeus</i>	Short-eared Owl	LC	LC
<i>Asio otus</i>	Northern Long-eared Owl	LC	LC
<i>Athene noctua</i>	Little Owl	LC	LC
<i>Aythya ferina</i>	Common Pochard	VU	VU
<i>Aythya fuligula</i>	Tufted Duck	NT	VU
<i>Aythya marila</i>	Greater Scaup	LC	EN
<i>Aythya nyroca</i>	Ferruginous Duck	LC	LC
<i>Bombycilla garrulus</i>	Bohemian Waxwing	LC	LC
<i>Bonasa bonasia</i>	Hazel Grouse	LC	VU
<i>Botaurus stellaris</i>	Eurasian Bittern	LC	LC
<i>Branta bernicla</i>	Brent Goose	LC	LC
<i>Branta canadensis</i>	Canada Goose	LC	N/A
<i>Branta leucopsis</i>	Barnacle Goose	LC	LC
<i>Branta ruficollis</i>	Red-breasted Goose	VU	VU
<i>Bubo bubo</i>	Eurasian Eagle-owl	LC	LC
<i>Bubo scandiacus</i>	Snowy Owl	LC	CR
<i>Bubulcus ibis</i>	Cattle Egret	LC	LC
<i>Bucanetes githagineus</i>	Trumpeter Finch	LC	LC
<i>Bucanetes mongolicus</i>	Mongolian Finch	LC	N/A
<i>Bucephala clangula</i>	Common Goldeneye	LC	LC
<i>Bucephala islandica</i>	Barrow's Goldeneye	NT	N/A
<i>Bulweria bulwerii</i>	Bulwer's Petrel	LC	LC
<i>Burhinus oedicephalus</i>	Eurasian Thick-knee	LC	LC
<i>Buteo buteo</i>	Eurasian Buzzard	LC	LC
<i>Buteo lagopus</i>	Rough-legged Buzzard	LC	VU
<i>Buteo rufinus</i>	Long-legged Buzzard	LC	LC
<i>Calandrella brachydactyla</i>	Greater Short-toed Lark	LC	LC
<i>Calcarius lapponicus</i>	Lapland Longspur	LC	VU
<i>Calidris alba</i>	Sanderling	LC	LC
<i>Calidris alpina</i>	Dunlin	LC	LC
<i>Calidris bairdii</i>	Baird's Sandpiper	LC ⁰⁰	N/A
<i>Calidris canutus</i>	Red Knot	LC	LC
<i>Calidris falcinellus</i>	Broad-billed Sandpiper	VU	VU
<i>Calidris ferruginea</i>	Curlew Sandpiper	VU	VU
<i>Calidris maritima</i>	Purple Sandpiper	LC	NT
<i>Calidris minuta</i>	Little Stint	LC	LC
<i>Calidris pugnax</i>	Ruff	NT	NT
<i>Calidris temminckii</i>	Temminck's Stint	LC	NT
<i>Calliope calliope</i>	Siberian Rubythroat	LC	N/A
<i>Calonectris borealis</i>	Cory's Shearwater	LC	LC
<i>Calonectris diomedea</i>	Scopoli's Shearwater	LC	LC
<i>Caprimulgus europaeus</i>	European Nightjar	LC	LC
<i>Caprimulgus ruficollis</i>	Red-necked Nightjar	NT	NT
<i>Carduelis carduelis</i>	European Goldfinch	LC	LC
<i>Carduelis citrinella</i>	Citril Finch	LC	LC
<i>Carduelis corsicana</i>	Corsican Finch	LC	LC
<i>Carpodacus erythrinus</i>	Common Rosefinch	LC	NT
<i>Carpodacus rubicilla</i>	Great Rosefinch	LC	N/A
<i>Carpospiza brachydactyla</i>	Pale Sparrow	LC	N/A
<i>Catharacta skua</i>	Great Skua	LC	LC
<i>Cecropis daurica</i>	Red-rumped Swallow	LC	LC
<i>Cepphus grylle</i>	Black Guillemot	LC	LC
<i>Cercotrichas galactotes</i>	Rufous-tailed Scrub-robin	LC	LC
<i>Certhia brachydactyla</i>	Short-toed Treecreeper	LC	LC
<i>Certhia familiaris</i>	Eurasian Treecreeper	LC	LC
<i>Ceryle rudis</i>	Pied Kingfisher	VU	N/A
<i>Cettia cetti</i>	Cetti's Warbler	LC	LC

Scientific species name	Common species name	IUCN Red List category (Europe)	IUCN Red List category (EU28)
<i>Charadrius alexandrinus</i>	Kentish Plover	LC	LC
<i>Charadrius asiaticus</i>	Caspian Plover	EN ^o	N/A
<i>Charadrius dubius</i>	Little Ringed Plover	LC	LC
<i>Charadrius hiaticula</i>	Common Ringed Plover	LC	LC
<i>Charadrius leschenaultii</i>	Greater Sandplover	NT	CR
<i>Chersophilus duponti</i>	Dupont's Lark	VU	VU
<i>Chlamydotis macqueenii</i>	Asian Houbara	CR (PE)	N/A
<i>Chlamydotis undulata</i>	African Houbara	VU	VU
<i>Chlidonias hybrida</i>	Whiskered Tern	LC	LC
<i>Chlidonias leucopterus</i>	White-winged Tern	LC	LC
<i>Chlidonias niger</i>	Black Tern	LC	LC
<i>Chloris chloris</i>	European Greenfinch	LC	LC
<i>Ciconia ciconia</i>	White Stork	LC	LC
<i>Ciconia nigra</i>	Black Stork	LC	LC
<i>Cinclus cinclus</i>	White-throated Dipper	LC	LC
<i>Circaetus gallicus</i>	Short-toed Snake-eagle	LC	LC
<i>Circus aeruginosus</i>	Western Marsh-harrier	LC	LC
<i>Circus cyaneus</i>	Hen Harrier	LC	VU
<i>Circus macrourus</i>	Pallid Harrier	LC	EN ^o
<i>Circus pygargus</i>	Montagu's Harrier	LC	VU
<i>Cisticola juncidis</i>	Zitting Cisticola	LC	LC
<i>Clamator glandarius</i>	Great Spotted Cuckoo	VU	VU
<i>Clanga clanga</i>	Greater Spotted Eagle	VU	CR
<i>Clanga pomarina</i>	Lesser Spotted Eagle	LC	LC
<i>Clangula hyemalis</i>	Long-tailed Duck	LC	LC
<i>Coccothraustes coccothraustes</i>	Hawfinch	LC	LC
<i>Columba bollii</i>	Dark-tailed Laurel-pigeon	LC	LC
<i>Columba junoniae</i>	White-tailed Laurel-pigeon	NT	NT
<i>Columba livia</i>	Rock Dove	LC	LC
<i>Columba oenas</i>	Stock Dove	LC	LC
<i>Columba palumbus</i>	Common Woodpigeon	LC	LC
<i>Columba trocaz</i>	Madeira Laurel-pigeon	LC	LC
<i>Coracias garrulus</i>	European Roller	LC	LC
<i>Corvus corax</i>	Common Raven	LC	LC
<i>Corvus corone</i>	Carrion Crow	LC	LC
<i>Corvus frugilegus</i>	Rook	VU	LC
<i>Corvus monedula</i>	Eurasian Jackdaw	LC	LC
<i>Coturnix coturnix</i>	Common Quail	NT	LC
<i>Crex crex</i>	Corncrake	LC	NT
<i>Cuculus canorus</i>	Common Cuckoo	LC	LC
<i>Cuculus saturatus</i>	Oriental Cuckoo	LC	N/A
<i>Curruca balearica</i>	Balearic Warbler	LC	LC
<i>Curruca cantillans</i>	Subalpine Warbler	LC	LC
<i>Curruca communis</i>	Common Whitethroat	LC	LC
<i>Curruca conspicillata</i>	Spectacled Warbler	LC	LC
<i>Curruca crassirostris</i>	Eastern Orphean Warbler	LC	LC
<i>Curruca curruca</i>	Lesser Whitethroat	LC	LC
<i>Curruca hortensis</i>	Western Orphean Warbler	LC	LC
<i>Curruca melanocephala</i>	Sardinian Warbler	LC	LC
<i>Curruca melanothorax</i>	Cyprus Warbler	LC	LC
<i>Curruca mystacea</i>	Menetries's Warbler	LC	N/A
<i>Curruca nana</i>	Asian Desert Warbler	VU ^o	N/A
<i>Curruca nisoria</i>	Barred Warbler	LC	LC
<i>Curruca ruppeli</i>	Rüppell's Warbler	LC	LC
<i>Curruca sarda</i>	Marmora's Warbler	LC	LC
<i>Curruca subalpina</i>	Moltoni's Warbler	LC	LC
<i>Curruca undata</i>	Dartford Warbler	NT	NT
<i>Cursorius cursor</i>	Cream-coloured Courser	NT	NT
<i>Cyanecula svecica</i>	Bluethroat	LC	LC
<i>Cyanistes caeruleus</i>	Eurasian Blue Tit	LC	LC
<i>Cyanistes cyanus</i>	Azure Tit	NT	N/A
<i>Cyanistes teneriffae</i>	African Blue Tit	LC	LC

Scientific species name	Common species name	IUCN Red List category (Europe)	IUCN Red List category (EU28)
<i>Cyanopica cooki</i>	Iberian Azure-winged Magpie	LC	LC
<i>Cygnus columbianus</i>	Tundra Swan	VU	VU
<i>Cygnus cygnus</i>	Whooper Swan	LC	LC
<i>Cygnus olor</i>	Mute Swan	LC	LC
<i>Delichon urbicum</i>	Northern House Martin	LC	LC
<i>Dendrocopos leucotos</i>	White-backed Woodpecker	LC	LC
<i>Dendrocopos major</i>	Great Spotted Woodpecker	LC	LC
<i>Dendrocopos syriacus</i>	Syrian Woodpecker	LC	LC
<i>Dryobates minor</i>	Lesser Spotted Woodpecker	LC	NT
<i>Dryocopus martius</i>	Black Woodpecker	LC	LC
<i>Egretta garzetta</i>	Little Egret	LC	LC
<i>Elanus caeruleus</i>	Black-winged Kite	LC	LC
<i>Emberiza aureola</i>	Yellow-breasted Bunting	CR	CR
<i>Emberiza bruniceps</i>	Red-headed Bunting	LC	N/A
<i>Emberiza buchanani</i>	Grey-necked Bunting	LC	N/A
<i>Emberiza caesia</i>	Cretzschmar's Bunting	LC	LC
<i>Emberiza calandra</i>	Corn Bunting	LC	LC
<i>Emberiza cia</i>	Rock Bunting	LC	LC
<i>Emberiza cineracea</i>	Cinereous Bunting	NT	EN
<i>Emberiza cirius</i>	Cirl Bunting	LC	LC
<i>Emberiza citrinella</i>	Yellowhammer	LC	LC
<i>Emberiza hortulana</i>	Ortolan Bunting	LC	NT
<i>Emberiza leucocephalos</i>	Pine Bunting	RE	N/A
<i>Emberiza melanocephala</i>	Black-headed Bunting	LC	LC
<i>Emberiza pallasi</i>	Pallas's Bunting	LC	N/A
<i>Emberiza pusilla</i>	Little Bunting	LC	LC
<i>Emberiza rustica</i>	Rustic Bunting	LC	LC
<i>Emberiza schoeniclus</i>	Reed Bunting	LC	LC
<i>Eremophila alpestris</i>	Horned Lark	LC	NT
<i>Erithacus rubecula</i>	European Robin	LC	LC
<i>Eudromias morinellus</i>	Eurasian Dotterel	LC	LC
<i>Falco biarmicus</i>	Lanner Falcon	NT*	EN
<i>Falco cherrug</i>	Saker Falcon	EN	EN
<i>Falco columbarius</i>	Merlin	VU	VU
<i>Falco eleonora</i>	Eleonora's Falcon	LC	LC
<i>Falco naumanni</i>	Lesser Kestrel	LC	LC
<i>Falco peregrinus</i>	Peregrine Falcon	LC	LC
<i>Falco rusticolus</i>	Gyr Falcon	LC	EN
<i>Falco subbuteo</i>	Eurasian Hobby	LC	LC
<i>Falco tinnunculus</i>	Common Kestrel	LC	LC
<i>Falco vespertinus</i>	Red-footed Falcon	VU	VU
<i>Ficedula albicollis</i>	Collared Flycatcher	LC	LC
<i>Ficedula hypoleuca</i>	European Pied Flycatcher	LC	LC
<i>Ficedula parva</i>	Red-breasted Flycatcher	LC	LC
<i>Ficedula semitorquata</i>	Semi-collared Flycatcher	LC	LC
<i>Francolinus francolinus</i>	Black Francolin	LC	LC
<i>Fratercula arctica</i>	Atlantic Puffin	EN	LC
<i>Fringilla coelebs</i>	Common Chaffinch	LC	LC
<i>Fringilla montifringilla</i>	Brambling	LC	LC
<i>Fringilla polatzeki</i>	Gran Canaria Blue Chaffinch	EN	EN
<i>Fringilla teydea</i>	Tenerife Blue Chaffinch	NT	NT
<i>Fulica atra</i>	Common Coot	NT	LC
<i>Fulica cristata</i>	Red-knobbed Coot	CR	CR
<i>Fulmarus glacialis</i>	Northern Fulmar	VU	EN
<i>Galerida cristata</i>	Crested Lark	LC	LC
<i>Galerida theklae</i>	Thekla's Lark	LC	LC
<i>Gallinago gallinago</i>	Common Snipe	VU	LC
<i>Gallinago media</i>	Great Snipe	LC	VU
<i>Gallinago megala</i>	Swinhoe's Snipe	NT ⁰⁰⁰	N/A
<i>Gallinago stenura</i>	Pintail Snipe	LC	N/A
<i>Gallinula chloropus</i>	Common Moorhen	LC	LC
<i>Garrulus glandarius</i>	Eurasian Jay	LC	LC

*Erratum: Red List status and trend in Europe updated in July 2022 to EN (for more details see: www.iucnredlist.org). This update is not reflected in the graphs and statistics presented in the report.

Scientific species name	Common species name	IUCN Red List category (Europe)	IUCN Red List category (EU28)
<i>Gavia adamsii</i>	Yellow-billed Loon	VU	N/A
<i>Gavia arctica</i>	Arctic Loon	LC	LC
<i>Gavia immer</i>	Common Loon	LC	LC
<i>Gavia stellata</i>	Red-throated Loon	LC	LC
<i>Gelochelidon nilotica</i>	Common Gull-billed Tern	LC	LC
<i>Geronticus eremita</i>	Northern Bald Ibis	RE	N/A
<i>Glareola nordmanni</i>	Black-winged Pratincole	LC	RE
<i>Glareola pratincola</i>	Collared Pratincole	LC	NT
<i>Glaucidium passerinum</i>	Eurasian Pygmy-owl	LC	LC
<i>Grus grus</i>	Common Crane	LC	LC
<i>Gulosus aristotelis</i>	European Shag	LC	LC
<i>Gymnoris xanthocollis</i>	Chestnut-shouldered Bush-sparrow	LC	N/A
<i>Gypaetus barbatus</i>	Bearded Vulture	NT	VU
<i>Gyps fulvus</i>	Griffon Vulture	LC	LC
<i>Haematopus ostralegus</i>	Eurasian Oystercatcher	VU	VU
<i>Halcyon smyrnensis</i>	White-breasted Kingfisher	VU	N/A
<i>Haliaeetus albicilla</i>	White-tailed Sea-eagle	LC	LC
<i>Hieraaetus pennatus</i>	Booted Eagle	LC	LC
<i>Himantopus himantopus</i>	Black-winged Stilt	LC	LC
<i>Hippolais icterina</i>	Icterine Warbler	LC	LC
<i>Hippolais languida</i>	Upcher's Warbler	LC	N/A
<i>Hippolais olivetorum</i>	Olive-tree Warbler	LC	LC
<i>Hippolais polyglotta</i>	Melodious Warbler	LC	LC
<i>Hirundo rustica</i>	Barn Swallow	LC	LC
<i>Histrionicus histrionicus</i>	Harlequin Duck	LC	N/A
<i>Hydrobates castro</i>	Band-rumped Storm-petrel	LC	LC
<i>Hydrobates leucorhous</i>	Leach's Storm-petrel	NT	VU
<i>Hydrobates montei</i>	Monteiro's Storm-petrel	VU	VU
<i>Hydrobates pelagicus</i>	European Storm-petrel	LC	LC
<i>Hydrocoloeus minutus</i>	Little Gull	LC	LC
<i>Hydroprogne caspia</i>	Caspian Tern	LC	LC
<i>Iduna caligata</i>	Booted Warbler	LC	LC
<i>Iduna opaca</i>	Isabelline Warbler	LC	LC
<i>Iduna pallida</i>	Olivaceous Warbler	LC	LC
<i>Iduna rama</i>	Sykes's Warbler	LC	N/A
<i>Irania gutturalis</i>	White-throated Robin	LC	N/A
<i>Ixobrychus minutus</i>	Common Little Bittern	LC	LC
<i>Jynx torquilla</i>	Eurasian Wryneck	LC	LC
<i>Ketupa zeylonensis</i>	Brown Fish-owl	EN	N/A
<i>Lagopus lagopus</i>	Willow Grouse	LC	LC
<i>Lagopus muta</i>	Rock Ptarmigan	LC	LC
<i>Lanius collurio</i>	Red-backed Shrike	LC	LC
<i>Lanius excubitor</i>	Great Grey Shrike	LC	VU
<i>Lanius meridionalis</i>	Iberian Grey Shrike	VU	VU
<i>Lanius minor</i>	Lesser Grey Shrike	LC	LC
<i>Lanius nubicus</i>	Masked Shrike	LC	LC
<i>Lanius senator</i>	Woodchat Shrike	NT	NT
<i>Larus argentatus</i>	European Herring Gull	LC	VU
<i>Larus armenicus</i>	Armenian Gull	LC	VU°
<i>Larus audouinii</i>	Audouin's Gull	VU	VU
<i>Larus cachinnans</i>	Caspian Gull	LC	LC
<i>Larus canus</i>	Mew Gull	LC	LC
<i>Larus fuscus</i>	Lesser Black-backed Gull	LC	LC
<i>Larus genei</i>	Slender-billed Gull	VU	LC
<i>Larus glaucooides</i>	Iceland Gull	LC	LC [∞]
<i>Larus hyperboreus</i>	Glaucaous Gull	LC	LC [∞]
<i>Larus ichthyaetus</i>	Pallas's Gull	LC	NE
<i>Larus marinus</i>	Great Black-backed Gull	LC	NT
<i>Larus melanocephalus</i>	Mediterranean Gull	LC	LC
<i>Larus michahellis</i>	Yellow-legged Gull	LC	LC
<i>Larus ridibundus</i>	Black-headed Gull	LC	VU
<i>Leipopicus medius</i>	Middle Spotted Woodpecker	LC	LC

Scientific species name	Common species name	IUCN Red List category (Europe)	IUCN Red List category (EU28)
<i>Limosa lapponica</i>	Bar-tailed Godwit	LC	LC
<i>Limosa limosa</i>	Black-tailed Godwit	NT	EN
<i>Linaria cannabina</i>	Common Linnet	LC	LC
<i>Linaria flavirostris</i>	Twite	LC	VU
<i>Locustella fluviatilis</i>	River Warbler	LC	VU
<i>Locustella lanceolata</i>	Lanceolated Warbler	LC	N/A
<i>Locustella luscinioides</i>	Savi's Warbler	LC	LC
<i>Locustella naevia</i>	Common Grasshopper-warbler	LC	VU
<i>Lophophanes cristatus</i>	Crested Tit	LC	LC
<i>Loxia curvirostra</i>	Red Crossbill	LC	LC
<i>Loxia leucoptera</i>	Two-barred Crossbill	LC	LC
<i>Loxia pytyopsittacus</i>	Parrot Crossbill	LC	LC
<i>Loxia scotica</i>	Scottish Crossbill	LC	LC
<i>Lullula arborea</i>	Woodlark	LC	LC
<i>Luscinia luscinia</i>	Thrush Nightingale	LC	VU
<i>Luscinia megarhynchos</i>	Common Nightingale	LC	LC
<i>Lymnocyptes minimus</i>	Jack Snipe	LC	LC
<i>Lyrurus mlokosiewiczzi</i>	Caucasian Grouse	NT	N/A
<i>Lyrurus tetrix</i>	Black Grouse	LC	VU
<i>Mareca penelope</i>	Eurasian Wigeon	LC	VU
<i>Mareca strepera</i>	Gadwall	LC	LC
<i>Marmaronetta angustirostris</i>	Marbled Teal	VU	EN
<i>Melanitta fusca</i>	Velvet Scoter	VU	VU
<i>Melanitta nigra</i>	Common Scoter	LC	LC
<i>Melanocorypha bimaculata</i>	Bimaculated Lark	LC	N/A
<i>Melanocorypha calandra</i>	Calandra Lark	LC	LC
<i>Melanocorypha yeltoniensis</i>	Black Lark	EN	N/A
<i>Mergellus albellus</i>	Smew	LC	LC
<i>Mergus merganser</i>	Goosander	LC	LC
<i>Mergus serrator</i>	Red-breasted Merganser	NT	NT
<i>Merops apiaster</i>	European Bee-eater	LC	LC
<i>Merops persicus</i>	Blue-cheeked Bee-eater	LC	N/A
<i>Microcarbo pygmaeus</i>	Pygmy Cormorant	LC	LC
<i>Milvus migrans</i>	Black Kite	LC	LC
<i>Milvus milvus</i>	Red Kite	LC	LC
<i>Monticola saxatilis</i>	Rufous-tailed Rock-thrush	LC	LC
<i>Monticola solitarius</i>	Blue Rock-thrush	LC	LC
<i>Montifringilla nivalis</i>	White-winged Snowfinch	LC	LC
<i>Morus bassanus</i>	Northern Gannet	LC	LC
<i>Motacilla alba</i>	White Wagtail	LC	LC
<i>Motacilla cinerea</i>	Grey Wagtail	LC	LC
<i>Motacilla citreola</i>	Citrine Wagtail	LC	LC°
<i>Motacilla flava</i>	Western Yellow Wagtail	LC	LC
<i>Muscicapa striata</i>	Spotted Flycatcher	LC	LC
<i>Neophron percnopterus</i>	Egyptian Vulture	VU	VU
<i>Netta rufina</i>	Red-crested Pochard	LC	LC
<i>Nucifraga caryocatactes</i>	Northern Nutcracker	LC	LC
<i>Numenius arquata</i>	Eurasian Curlew	NT	NT
<i>Numenius phaeopus</i>	Whimbrel	LC	LC
<i>Numenius tenuirostris</i>	Slender-billed Curlew	CR (PE)	CR (PE)
<i>Nycticorax nycticorax</i>	Black-crowned Night-heron	LC	LC
<i>Oenanthe chrysopygia</i>	Red-tailed Wheatear	VU ⁰	N/A
<i>Oenanthe cypriaca</i>	Cyprus Wheatear	LC	LC
<i>Oenanthe deserti</i>	Desert Wheatear	NT ⁰	N/A
<i>Oenanthe finschii</i>	Finsch's Wheatear	LC	NT ⁰⁰
<i>Oenanthe hispanica</i>	Black-eared Wheatear	LC	LC
<i>Oenanthe isabellina</i>	Isabelline Wheatear	LC	LC
<i>Oenanthe leucura</i>	Black Wheatear	LC	LC
<i>Oenanthe oenanthe</i>	Northern Wheatear	LC	NT
<i>Oenanthe pleschanka</i>	Pied Wheatear	LC	LC°
<i>Oenanthe xanthopyrna</i>	Kurdish Wheatear	LC	N/A
<i>Onychoprion fuscatus</i>	Sooty Tern	VU ⁰⁰	VU ⁰⁰
<i>Oriolus oriolus</i>	Eurasian Golden Oriole	LC	LC

Scientific species name	Common species name	IUCN Red List category (Europe)	IUCN Red List category (EU28)
<i>Otis tarda</i>	Great Bustard	LC	LC
<i>Otus brucei</i>	Pallid Scops-owl	EN	N/A
<i>Otus cypricus</i>	Cyprus Scops-owl	LC	N/A
<i>Otus scops</i>	Eurasian Scops-owl	LC	LC
<i>Oxyura leucocephala</i>	White-headed Duck	VU	VU
<i>Pagophila eburnea</i>	Ivory Gull	VU	N/A
<i>Pandion haliaetus</i>	Osprey	LC	LC
<i>Panurus biarmicus</i>	Bearded Reedling	LC	LC
<i>Parus major</i>	Great Tit	LC	LC
<i>Passer domesticus</i>	House Sparrow	LC	LC
<i>Passer hispaniolensis</i>	Spanish Sparrow	LC	LC
<i>Passer italiae</i>	Italian Sparrow	VU	VU
<i>Passer moabiticus</i>	Dead Sea Sparrow	LC	N/A
<i>Passer montanus</i>	Eurasian Tree Sparrow	LC	LC
<i>Pastor roseus</i>	Rosy Starling	LC	LC
<i>Pelagodroma marina</i>	White-faced Storm-petrel	EN	EN
<i>Pelecanus crispus</i>	Dalmatian Pelican	LC	LC
<i>Pelecanus onocrotalus</i>	Great White Pelican	LC	LC
<i>Perdix perdix</i>	Grey Partridge	LC	VU
<i>Periparus ater</i>	Coal Tit	LC	LC
<i>Perisoreus infaustus</i>	Siberian Jay	LC	LC
<i>Pernis apivorus</i>	European Honey-buzzard	LC	LC
<i>Petronia petronia</i>	Rock Sparrow	LC	LC
<i>Phalacrocorax carbo</i>	Great Cormorant	LC	LC
<i>Phalaropus fulicarius</i>	Red Phalarope	LC	N/A
<i>Phalaropus lobatus</i>	Red-necked Phalarope	LC	LC
<i>Phasianus colchicus</i>	Common Pheasant	LC	EN
<i>Phoenicopterus roseus</i>	Greater Flamingo	LC	LC
<i>Phoenicurus erythrogastrus</i>	White-winged Redstart	NT	N/A
<i>Phoenicurus ochruros</i>	Black Redstart	LC	LC
<i>Phoenicurus phoenicurus</i>	Common Redstart	LC	LC
<i>Phylloscopus bonelli</i>	Western Bonelli's Warbler	LC	LC
<i>Phylloscopus borealis</i>	Arctic Warbler	LC	VU
<i>Phylloscopus canariensis</i>	Canary Islands Chiffchaff	LC	LC
<i>Phylloscopus collybita</i>	Common Chiffchaff	LC	LC
<i>Phylloscopus ibericus</i>	Iberian Chiffchaff	LC	LC
<i>Phylloscopus inornatus</i>	Yellow-browed Warbler	LC	N/A
<i>Phylloscopus nitidus</i>	Green Warbler	LC	N/A
<i>Phylloscopus orientalis</i>	Eastern Bonelli's Warbler	LC	LC
<i>Phylloscopus sibilatrix</i>	Wood Warbler	LC	LC
<i>Phylloscopus sindianus</i>	Mountain Chiffchaff	LC	N/A
<i>Phylloscopus trochiloides</i>	Greenish Warbler	LC	LC
<i>Phylloscopus trochilus</i>	Willow Warbler	LC	LC
<i>Pica pica</i>	Eurasian Magpie	LC	LC
<i>Picoides tridactylus</i>	Three-toed Woodpecker	LC	LC
<i>Picus canus</i>	Grey-faced Woodpecker	LC	LC
<i>Picus sharpei</i>	Iberian Green Woodpecker	LC	LC
<i>Picus viridis</i>	Eurasian Green Woodpecker	LC	LC
<i>Pinicola enucleator</i>	Pine Grosbeak	LC	LC
<i>Platalea leucorodia</i>	Eurasian Spoonbill	LC	LC
<i>Plectrophenax nivalis</i>	Snow Bunting	LC	LC
<i>Plegadis falcinellus</i>	Glossy Ibis	LC	LC
<i>Pluvialis apricaria</i>	Eurasian Golden Plover	LC	LC
<i>Pluvialis squatarola</i>	Grey Plover	LC	LC
<i>Podiceps auritus</i>	Horned Grebe	NT	VU
<i>Podiceps cristatus</i>	Great Crested Grebe	LC	LC
<i>Podiceps grisegena</i>	Red-necked Grebe	VU	VU
<i>Podiceps nigricollis</i>	Black-necked Grebe	VU	LC
<i>Poecile cinctus</i>	Siberian Tit	LC	NT
<i>Poecile hyrcanus</i>	Caspian Tit	LC	N/A
<i>Poecile lugubris</i>	Sombre Tit	LC	LC
<i>Poecile montanus</i>	Willow Tit	LC	NT

Scientific species name	Common species name	IUCN Red List category (Europe)	IUCN Red List category (EU28)
<i>Poecile palustris</i>	Marsh Tit	LC	LC
<i>Polysticta stelleri</i>	Steller's Eider	LC	EN
<i>Porphyrio porphyrio</i>	Purple Swamphen	LC	LC
<i>Porzana porzana</i>	Spotted Crake	LC	LC
<i>Prinia gracilis</i>	Graceful Prinia	LC	N/A
<i>Prunella atrogularis</i>	Black-throated Accentor	LC	N/A
<i>Prunella collaris</i>	Alpine Accentor	LC	LC
<i>Prunella modularis</i>	Dunnock	LC	LC
<i>Prunella montanella</i>	Siberian Accentor	LC	N/A
<i>Prunella ocularis</i>	Radde's Accentor	LC	N/A
<i>Pterocles alchata</i>	Pin-tailed Sandgrouse	LC	LC
<i>Pterocles orientalis</i>	Black-bellied Sandgrouse	EN	EN
<i>Pterodroma deserta</i>	Desertas Petrel	VU	VU
<i>Pterodroma madeira</i>	Zino's Petrel	EN	EN
<i>Ptyonoprogne rupestris</i>	Eurasian Crag Martin	LC	LC
<i>Puffinus lherminieri</i>	Audubon's Shearwater	NT	NT
<i>Puffinus mauretanicus</i>	Balearic Shearwater	CR	CR
<i>Puffinus puffinus</i>	Manx Shearwater	LC	LC
<i>Puffinus yelkouan</i>	Yelkouan Shearwater	VU	VU
<i>Pycnonotus xanthopygos</i>	White-spectacled Bulbul	LC	N/A
<i>Pyrrhonorax graculus</i>	Yellow-billed Chough	LC	LC
<i>Pyrrhonorax pyrrhonorax</i>	Red-billed Chough	LC	LC
<i>Pyrrhula murina</i>	Azores Bullfinch	VU	VU
<i>Pyrrhula pyrrhula</i>	Eurasian Bullfinch	LC	LC
<i>Rallus aquaticus</i>	Western Water Rail	LC	LC
<i>Recurvirostra avosetta</i>	Pied Avocet	LC	LC
<i>Regulus ignicapilla</i>	Common Firecrest	LC	LC
<i>Regulus madeirensis</i>	Madeira Firecrest	LC	LC
<i>Regulus regulus</i>	Goldcrest	LC	LC
<i>Remiz pendulinus</i>	Eurasian Penduline-tit	LC	LC
<i>Rhodopechys sanguineus</i>	Eurasian Crimson-winged Finch	LC	N/A
<i>Rhodospiza obsoleta</i>	Desert Finch	LC	N/A
<i>Rhodostethia rosea</i>	Ross's Gull	EN ⁰	N/A
<i>Riparia riparia</i>	Collared Sand Martin	LC	LC
<i>Rissa tridactyla</i>	Black-legged Kittiwake	VU	EN
<i>Saxicola dacotiae</i>	Fuerteventura Stonechat	NT	NT
<i>Saxicola rubetra</i>	Whinchat	LC	VU
<i>Saxicola torquatus</i>	Common Stonechat	LC	LC
<i>Scolopax rusticola</i>	Eurasian Woodcock	LC	LC
<i>Serinus canaria</i>	Island Canary	LC	LC
<i>Serinus pusillus</i>	Red-fronted Serin	LC	N/A
<i>Serinus serinus</i>	European Serin	LC	LC
<i>Sitta europaea</i>	Eurasian Nuthatch	LC	LC
<i>Sitta krueperi</i>	Krueper's Nuthatch	LC	NT
<i>Sitta neumayer</i>	Western Rock Nuthatch	LC	LC
<i>Sitta tephronota</i>	Eastern Rock Nuthatch	LC	N/A
<i>Sitta whiteheadi</i>	Corsican Nuthatch	VU	VU
<i>Somateria mollissima</i>	Common Eider	EN	VU
<i>Somateria spectabilis</i>	King Eider	LC	N/A
<i>Spatula clypeata</i>	Northern Shoveler	LC	NT
<i>Spatula querquedula</i>	Garganey	LC	VU
<i>Spilopelia senegalensis</i>	Laughing Dove	LC	NT ^{oo}
<i>Spinus spinus</i>	Eurasian Siskin	LC	LC
<i>Stercorarius longicaudus</i>	Long-tailed Jaeger	LC	LC
<i>Stercorarius parasiticus</i>	Arctic Jaeger	EN	EN
<i>Stercorarius pomarinus</i>	Pomarine Jaeger	LC	N/A
<i>Sterna dougallii</i>	Roseate Tern	LC	LC
<i>Sterna hirundo</i>	Common Tern	LC	LC
<i>Sterna paradisaea</i>	Arctic Tern	LC	LC
<i>Sternula albifrons</i>	Little Tern	LC	LC
<i>Streptopelia decaocto</i>	Eurasian Collared-dove	LC	LC
<i>Streptopelia turtur</i>	European Turtle-dove	VU	NT

Scientific species name	Common species name	IUCN Red List category (Europe)	IUCN Red List category (EU28)
<i>Strix aluco</i>	Tawny Owl	LC	LC
<i>Strix nebulosa</i>	Great Grey Owl	LC	LC
<i>Strix uralensis</i>	Ural Owl	LC	LC
<i>Sturnus unicolor</i>	Spotless Starling	LC	LC
<i>Sturnus vulgaris</i>	Common Starling	LC	LC
<i>Surnia ulula</i>	Northern Hawk-owl	LC	LC
<i>Sylvia atricapilla</i>	Eurasian Blackcap	LC	LC
<i>Sylvia borin</i>	Garden Warbler	LC	LC
<i>Syrrhaptes paradoxus</i>	Pallas's Sandgrouse	RE	N/A
<i>Tachybaptus ruficollis</i>	Little Grebe	LC	LC
<i>Tachymarptis melba</i>	Alpine Swift	LC	LC
<i>Tadorna ferruginea</i>	Ruddy Shelduck	LC	NT°
<i>Tadorna tadorna</i>	Common Shelduck	LC	LC
<i>Tarsiger cyanurus</i>	Orange-flanked Bush-robin	LC	LC
<i>Tetrao urogallus</i>	Western Capercaillie	LC	LC
<i>Tetraogallus caspius</i>	Caspian Snowcock	LC	N/A
<i>Tetraogallus caucasicus</i>	Caucasian Snowcock	LC	N/A
<i>Tetrax tetrax</i>	Little Bustard	VU	EN
<i>Thalasseus sandvicensis</i>	Sandwich Tern	LC	LC
<i>Tichodroma muraria</i>	Wallcreeper	LC	LC
<i>Tringa erythropus</i>	Spotted Redshank	LC	NT
<i>Tringa glareola</i>	Wood Sandpiper	LC	LC
<i>Tringa nebularia</i>	Common Greenshank	LC	LC
<i>Tringa ochropus</i>	Green Sandpiper	LC	LC
<i>Tringa stagnatilis</i>	Marsh Sandpiper	LC	VU°
<i>Tringa totanus</i>	Common Redshank	VU	VU
<i>Troglodytes troglodytes</i>	Northern Wren	LC	LC
<i>Turdus atrogularis</i>	Black-throated Thrush	LC	N/A
<i>Turdus iliacus</i>	Redwing	LC	LC
<i>Turdus merula</i>	Eurasian Blackbird	LC	LC
<i>Turdus philomelos</i>	Song Thrush	LC	LC
<i>Turdus pilaris</i>	Fieldfare	LC	LC
<i>Turdus torquatus</i>	Ring Ouzel	LC	LC
<i>Turdus viscivorus</i>	Mistle Thrush	LC	LC
<i>Turnix sylvaticus</i>	Common Buttonquail	RE	RE
<i>Tyto alba</i>	Common Barn-owl	LC	LC
<i>Upupa epops</i>	Common Hoopoe	LC	LC
<i>Uria aalge</i>	Common Murre	LC	LC
<i>Uria lomvia</i>	Thick-billed Murre	LC	N/A
<i>Vanellus gregarius</i>	Sociable Lapwing	CR	N/A
<i>Vanellus indicus</i>	Red-wattled Lapwing	VU ⁰⁰	N/A
<i>Vanellus leucurus</i>	White-tailed Lapwing	LC	N/A
<i>Vanellus spinosus</i>	Spur-winged Lapwing	LC	VU°
<i>Vanellus vanellus</i>	Northern Lapwing	VU	VU
<i>Xema sabini</i>	Sabine's Gull	LC	N/A
<i>Xenus cinereus</i>	Terek Sandpiper	LC	CR
<i>Zapornia parva</i>	Little Crake	LC	LC
<i>Zapornia pusilla</i>	Baillon's Crake	LC	NT°
<i>Zoothera aurea</i>	White's Thrush	LC	N/A

Appendix 4. Countries and territories covered by the European Red List of Birds (countries and territories included in the EU28 level of assessment shown in bold)

COUNTRIES AND TERRITORIES	
Albania	Italy
Andorra	Kosovo
Armenia	Latvia
Austria	Liechtenstein
Azerbaijan	Lithuania
The Azores	Luxembourg
Belarus	Northern Macedonia
Belgium	Madeira
Bosnia and Herzegovina	Malta
Bulgaria	Moldova
The Canary Islands	Montenegro
Croatia	The Netherlands
Cyprus	Norway
Czechia	Poland
Denmark	Portugal
Estonia	Romania
The Faroe Islands	European Russia
Finland	Serbia
France	Slovakia
Georgia	Slovenia
Germany	Spain
Gibraltar	Svalbard and Jan Mayen
Greece	Sweden
Greenland	Switzerland
Hungary	Turkey
Iceland	Ukraine
Ireland	The United Kingdom





Azure Tit (*Cynaistes cyanus*)
Photo: Istock



Atlantic Puffin (*Fratercula arctica*)
Photo: Martha de Jong-Lantink



Sooty Tern (*Onychoprion fuscatus*)
Photo: Istock

Photo credits:

Cover: Demoiselle Crane (*Anthropoides virgo*) - Photo: Istock

Page 9: Kingfisher (*Alcedo atthis*) – Photo: Jan Veber; Steppe Eagle (*Aquila nipalensis*) – Photo: 4028mdk09; Black-necked Grebe (*Podiceps nigricollis*) – Photo: Martin Mecnarowski; Yellow-breasted Bunting (*Emberiza aureola*) – Photo: Hiyashi Haka; Atlantic Puffin (*Fratercula arctica*) – Photo: Sue Cro; Woodchat Shrike (*Lanius senator*) - Photo: Juan Emilio; Demoiselle Crane (*Anthropoides virgo*) - Photo: Istock; Northern Pintail (*Anas acuta*) – Photo: Ivan Dudáček; Greater Spotted Eagle (*Clanga clanga*) – Photo: Koshy Koshy; Rook (*Corvus frugilegus*) – Photo: Jiřina Chalupská; Atlantic Puffin (*Fratercula arctica*) – Photo: Martha de Jong-Lantink; Common Eider (*Somateria mollissima*) – Photo: Ondřej Prosícký; Common Redshank (*Tringa totanus*) – Photo: Britt-Marie Sohlström; Red-necked Grebe (*Podiceps grisegena*) – Photo: Becky Matsubara; Balearic Shearwater (*Puffinus mauretanicus*) – Photo: Cabrera Natura; Sooty Tern (*Onychoprion fuscatus*) - Photo: Istock; Greater Spotted Eagle (*Clanga clanga*) – Photo: Peter Steward; Common Quail (*Coturnix coturnix*) – Photo: Petr Šaj; Atlantic Puffin (*Fratercula arctica*) – Photo: Sue Cro; Gran Canaria Blue Chaffinch (*Fringilla polatzeki*) – Photo: Miguel Angel Peña Estévez; Arctic Skua (*Stercorarius parasiticus*) – Photo: Tony Morris; Red-knobbed Coot (*Fulica cristata*) – Photo: Muchaxo; Eurasian Curlew (*Numenius arquata*) – Photo: Petr Šaj; Azure Tit (*Cynaistes cyanus*) - Photo: Istock; Red-necked Grebe (*Podiceps grisegena*) – Photo: Becky Matsubara; Common Snipe (*Gallinago gallinago*) – Photo: Martin Mecnarowski; Gran Canaria Blue Chaffinch (*Fringilla polatzeki*) – Photo: Miguel Angel Peña Estévez; Red-necked Grebe (*Podiceps grisegena*) – Photo: Kim Taylor; Greater Spotted Eagle (*Clanga clanga*) – Photo: Swati Kulkarni; Black-tailed Godwit (*Limosa limosa*) – Photo: Petr Šaj; Red-necked Grebe (*Podiceps grisegena*) – Photo: Mah Connor; Demoiselle Crane (*Anthropoides virgo*) – Photo: Tim Strater; Redshank (*Tringa totanus*) – Photo: Frans Vandewalle; Sociable Lapwing (*Vanellus gregarius*) – Photo: Rob Llewellyn; Sociable Lapwing (*Vanellus gregarius*) – Photo: Adrian Drummond-Hill; Common Redshank (*Tringa totanus*) – Photo: Britt-Marie Sohlström

Page 21: Kingfisher (*Alcedo atthis*) – Photo: Jan Veber; Greater Spotted Eagle (*Clanga clanga*) – Photo: Koshy Koshy; Black-tailed Godwit (*Limosa limosa*) - Photo: Petr Šaj

Page 22: Eurasian Curlew (*Numenius arquata*) – Photo: Petr Šaj; Common Snipe (*Gallinago gallinago*) – Photo: Martin Mecnarowski;

Common Redshank (*Tringa totanus*) – Photo: Britt-Marie Sohlström; Northern Pintail (*Anas acuta*) – Photo: Ivan Dudáček;

Page 23: Common Eider (*Somateria mollissima*) - Photo: Ondřej Prosícký; Common Quail (*Coturnix coturnix*) – Photo: Petr Šaj;

Red-necked Grebe (*Podiceps grisegena*) – Photo: Becky Matsubara; Black-necked Grebe (*Podiceps nigricollis*) – Photo: Martin Mecnarowski



Stichting BirdLife Europe gratefully acknowledges financial support from the LIFE programme of the European Union.

European Red List of Birds

Compiled by BirdLife International



Publications Office
of the European Union