

1 **2019 REFINEMENT TO THE 2006**
2 **IPCC GUIDELINES FOR**
3 **NATIONAL GREENHOUSE GAS**
4 **INVENTORIES**

5 **OVERVIEW**

Final Draft

6 **Authors**

7 Eduardo Calvo Buendia (Peru), Sabin Guendehou (Benin), Bundit Limmeechokchai (Thailand), Riitta Pipatti
8 (Finland), Yasna Rojas (Chile), Rob Sturgiss (Australia), Kiyoto Tanabe (Japan), Tom Wirth (USA)

9 Daniela Romano (Italy), Jongikhaya Witi (South Africa), Amit Garg (India), Melissa M. Weitz (USA), Bofeng
10 Cai (China), Deborah A. Ottinger (USA), Hongmin Dong (China), James Douglas MacDonald (Canada),
11 Stephen Michael Ogle (USA), Marcelo Theoto Rocha (Brazil), Maria José Sanz Sanchez (Spain), Deborah M.
12 Bartram (USA), Sirintornthep Towprayoon (Thailand)

13 **Review Editors**

14 Dario Gómez (Argentina) and William Irving (USA)

15

16

Contents

17 Overview

18	1	Introduction	4
19	2	Background	5
20	3	Coverage of the 2019 Refinement	6
21	4	Relationship with the 2006 IPCC Guidelines	8
22	5	Specific developments in the 2019 Refinement	10
23	6	Clarification on key concepts in the 2019 Refinement.....	14
24			

25

Tables

26	Table 1	Types of refinement from inventory compliers' perspective.....	8
----	---------	--	---

27

Boxes

28	Box 1	Significance and prioritization criteria	6
29	Box 2	Example of mapping table for the 2019 Refinement to the 2006 IPCC Guidelines (Vol.4)....	9
30			

Final Draft

31 1 INTRODUCTION

32 The *2006 IPCC Guidelines for National Greenhouse Gas Inventories (2006 IPCC Guidelines)* which were
33 published in 2006 provide methodologies for estimating national inventories of anthropogenic emissions by
34 sources and removals by sinks of greenhouse gases. This IPCC Methodology Report titled the *2019 Refinement to*
35 *the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (2019 Refinement)* was published in 2019 to
36 refine the *2006 IPCC Guidelines* with the aim to provide an updated and sound scientific basis for supporting the
37 preparation and continuous improvement of national greenhouse gas inventories.

38 In order to achieve this overall aim, the *2019 Refinement*:

- 39 • Provides supplementary methodologies for sources or sinks of greenhouse gases only where currently there
40 are gaps or where new technologies and production processes have emerged requiring elaborated
41 methodologies or for sources or sinks that are not well covered by the *2006 IPCC Guidelines*;
- 42 • Provides updated default values of emission factors and other parameters based on the latest available
43 scientific information only where significant differences from default values presented in the *2006 IPCC*
44 *Guidelines* are identified;
- 45 • Provides additional or alternative up-to-date information and guidance, where possible, as clarification or
46 elaboration of existing guidance in the *2006 IPCC Guidelines*.

47 The *2019 Refinement* does not revise the *2006 IPCC Guidelines*, but updates, supplements and/or elaborates the
48 *2006 IPCC Guidelines* where gaps or out-of-date science have been identified. It does not replace the *2006 IPCC*
49 *Guidelines*, but should be used in conjunction with the *2006 IPCC Guidelines* and the *2013 Supplement to the*
50 *2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands (Wetlands Supplement)*.

51

52 2 BACKGROUND

53 The Bureau of IPCC Task Force on National Greenhouse Gas Inventories (TFB), at its 26th meeting held in Ottawa,
54 Canada in August 2014, concluded that:

55 • The *2006 IPCC Guidelines* provide a technically sound methodological basis of national greenhouse gas
56 inventories, and therefore fundamental revision was unnecessary.

57 • To maintain the scientific validity of the *2006 IPCC Guidelines*, certain refinements might be required, taking
58 into account scientific and other technical advances that had matured sufficiently since 2006.

59 Following these conclusions by the TFB and also in accordance with the approval by the IPCC at its 40th Session
60 held in Berlin, Germany in April 2014, the Task Force on National Greenhouse Gas Inventories (TFI) started a
61 technical assessment of IPCC inventory guidelines through an on-line questionnaire survey and four expert
62 meetings in 2015 and 2016^{1,2,3,4}.

63 The technical assessment revealed that there had been abundant new scientific and empirical knowledge published
64 since 2006 that the IPCC should take into account, particularly with respect to data for emission factor development
65 for some categories and gases. Consequently, the TFB recognized the necessity and usefulness of refining the *2006*
66 *IPCC Guidelines* (e.g. updating default emission factors), and concluded that the refinement should be made as
67 early as possible so as to help all the Parties to the United Nations Framework Convention on Climate Change
68 (UNFCCC) use *good practice* inventory methodologies based on up-to-date scientific knowledge.

69 The IPCC, at its 43rd Session in Nairobi, Kenya in April 2016, approved the proposal made by the Co-Chairs of
70 TFI on “Refinement of 2006 IPCC Guidelines for National Greenhouse Gas Inventories, including production of
71 a Methodology Report(s)” as contained in the Decision IPCC/XLIII-8 “Update of methodologies on National
72 Greenhouse Gas Inventories”. At the same session, the IPCC decided to consider the outline of the new
73 Methodology Report(s) at the 44th Session of the IPCC in October 2016 and to consider the draft Methodology
74 Report(s) at a Plenary session of the IPCC in May 2019 (Decision IPCC/XLIII-7 “Sixth Assessment Report (AR6)
75 Products. Strategic Planning”). Following this decision, a scoping meeting for a Methodology Report(s) to refine
76 the *2006 IPCC Guidelines* was held in Minsk, Belarus in August 2016. The outcome of the scoping meeting was
77 approved by the TFB at its 28th meeting which was held immediately after the scoping meeting, and submitted to
78 the IPCC for consideration. Finally the IPCC, at its 44th Session in Bangkok, Thailand in October 2016, Thailand,
79 decided to prepare the *2019 Refinement* and adopted its scope and outline (Decision IPCC/XLIV-5 “Sixth
80 Assessment Report (AR6) Products, Outline of the Methodology Report(s) to refine the 2006 Guidelines for
81 National Greenhouse Gas Inventories”).

¹ Expert Meeting for Technical Assessment of IPCC Inventory Guidelines (Energy, IPPU, Waste Sectors), 29 June - 1 July 2015, Geneva, Switzerland

² Expert Meeting for Technical Assessment of IPCC Inventory Guidelines (AFOLU Sector), 13-16 July 2015, São Paulo, Brazil

³ Expert meeting for Technical Assessment of IPCC Inventory Guidelines: follow-up on specified issues from the 2015 expert meetings, 25-26 April 2016, Wollongong, Australia

⁴ Expert meeting for Technical Assessment of IPCC Inventory Guidelines (Cross-sectoral issues), 27-29 April 2016, Wollongong, Australia

Final Draft

82 3 COVERAGE OF THE 2019 REFINEMENT

83 The *2019 Refinement* covers all IPCC inventory sectors but refinements are included for only those categories
 84 where the science was considered to have sufficiently advanced since 2006 or where new or additional guidance
 85 was required. The specific categories that have been refined through this process were selected through a technical
 86 assessment carried out in 2015 and 2016 and a subsequent scoping meeting held in August 2016 using the
 87 significance and prioritization criteria shown in Box 1⁵.

88

89
90

Box 1
SIGNIFICANCE AND PRIORITIZATION CRITERIA

- 91 • Significance of the source/sink and the gas within the sector on a global scale. Sources significant only for a limited number of particular countries, currently or in the foreseeable future, may not meet this criterion. The adequacy of the existing guidance for a particular category should be considered, as should the likelihood that new information would lead to a definite improvement in the IPCC Guidelines.
- 92
- 93
- 94
- 95
- 96 • Availability of relevant new scientific results.
- 97 • Sufficient data availability and maturity of scientific advances since 2006 to provide a basis for methodological development or refinement, including (1) ability to develop new or updated default emission/removal factors; and (2) feasibility of obtaining the necessary data to implement the methods.
- 98
- 99
- 100 • Emergence of new sources or gases meeting these criteria.
- 101

102

103 The outline of the *2019 Refinement* adopted by the IPCC was developed on the basis of the categories selected
 104 through the process mentioned above. However, the IPCC also agreed on the following principles when the outline
 105 was adopted.

- 106 • Authors should develop modifications even for those chapters/sections/subsections where “No refinement” is
 107 indicated in the approved table of contents, if deemed necessary to ensure consistency with the refinements
 108 made in the other chapters/sections/subsections.
- 109 • Authors may conclude no refinement should be made even for the chapters/sections/subsections where
 110 refinement is expected in this approved table of contents, after comprehensive review of available literature.
 111

112 In addition to the greenhouse gases included in the *2006 IPCC Guidelines*, the *2019 Refinement* includes gases for
 113 which global warming potential (GWP) values are given in one of the subsequent IPCC Assessment Reports (e.g.,
 114 the Fourth or Fifth Assessment Report), unless the gases are covered by Annexes A through E of the Montreal
 115 Protocol. (Annex F of the Montreal Protocol lists hydrofluorocarbons, which are included in the *2019 Refinement*.)
 116 The *2019 Refinement* also provides estimation methods for halogenated greenhouse gases for which GWP values
 117 were not available from IPCC Assessment Reports at the time the *2019 Refinement* was developed. (See Volume
 118 1, Chapter 8 and Volume 3, Chapters 6, 7, and 8 for examples of both sets of gases.)

119 The *2006 IPCC Guidelines* contain links to information on methods used under other agreements and conventions,
 120 for the estimation of emissions of tropospheric precursors which may be used to supplement the reporting of
 121 emissions and removals of greenhouse gases. The *2019 Refinement* follows this approach, and does not provide
 122 methods for the estimation of emissions of tropospheric precursors.

123 The structure of the *2019 Refinement* is the same as that of the *2006 IPCC Guidelines* so as to make it easier for
 124 inventory compilers to use the *2019 Refinement* with the *2006 IPCC Guidelines*. It comprises an Overview Chapter
 125 (this chapter) and the following five volumes.

- 126 • Volume 1: General Guidance and Reporting
- 127 • Volume 2: Energy

⁵ There are potential issues for future revision or refinement that are not addressed in the *2019 Refinement*, including those identified during the Technical Assessment in 2015-2016 but considered too early to work on, as well as those identified after the approval of table of contents of the *2019 Refinement* by the IPCC at its 44th Session in Bangkok, Thailand in October 2016.

128 • Volume 3: Industrial Processes and Product Use (IPPU)

129 • Volume 4: Agriculture, Forestry and Other Land Use (AFOLU)

130 • Volume 5: Waste

131 Across all the volumes, some additional sections have been included. The guidance focuses on inventory
132 methodologies rather than on scientific discussions of the background material, for which references are provided.

133

Final Draft

4 RELATIONSHIP WITH THE 2006 IPCC GUIDELINES

As stated in Section 1 above, the *2019 Refinement* does not revise the *2006 IPCC Guidelines*, but updates, supplements and/or elaborates the *2006 IPCC Guidelines* where gaps or out-of-date science have been identified. It does not replace the *2006 IPCC Guidelines*, but should be used in conjunction with the *2006 IPCC Guidelines* and the *Wetlands Supplement*.⁶

In line with this principle, the *2019 Refinement* was developed by:

- updating existing guidance in the *2006 IPCC Guidelines* to address the needs explained in the first or second bullet in Section 1 (Introduction) of this chapter. When updating a section or an entire chapter, in some cases these sections or chapters have been entirely rewritten with new structure.
- elaborating existing guidance in the *2006 IPCC Guidelines* to address the needs explained in the first or third bullet of Section 1 (Introduction) of this chapter.
- adding completely new guidance on issues for which there is essentially no guidance in the *2006 IPCC Guidelines* to address the needs explained in the first bullet in Section 1 (Introduction) of this chapter.

With a view to helping inventory compilers understand the relationship between the *2019 Refinement* and the *2006 IPCC Guidelines*, each volume has an annex titled “Mapping table for the *2019 Refinement* to the *2006 IPCC Guidelines*”. This annex in each volume provides a road map for relating sections, equations, tables, figures and boxes in the *2019 Refinement* to the *2006 IPCC Guidelines*. In these annexes, types of refinement (update, new guidance, no refinement or removed) are clarified for chapters/sections/subsections from inventory compilers’ perspective in order to help them understand how they should use the *2019 Refinement*. The types of refinement used there are explained in Table 1 below.

Type of refinement	Explanation
Update	Inventory compilers should use the chapter/section/subsection in the <i>2019 Refinement</i> instead of the corresponding chapter/section/subsection in the <i>2006 IPCC Guidelines</i> . (Inventory compilers should read the chapter/section/subsection in the <i>2019 Refinement</i> only.)
New Guidance	Inventory compilers should use the section/chapter in the <i>2019 Refinement</i> without reference to specific chapter/section/subsection in the <i>2006 IPCC Guidelines</i> , recognizing that there is essentially no corresponding guidance in the <i>2006 IPCC Guidelines</i> . (Inventory compilers should read the chapter/section/subsection in the <i>2019 Refinement</i> only.)
No Refinement	Inventory compilers should read the corresponding chapter/section/subsection in the <i>2006 IPCC Guidelines</i> , because no refinement has been made in that chapter/section/subsection.
Removed	There were few cases where guidance/sections were removed because they were no longer relevant.

Every chapter/section/subsection in the *2019 Refinement* which is explained as “Update” or “New Guidance” contains all necessary paragraphs, tables, figures, etc. including those unchanged from the *2006 IPCC Guidelines*, so that inventory compilers do not have to look back to the *2006 IPCC Guidelines*.

In addition, with a view to ensure consistency with the *2006 IPCC Guidelines* to the extent possible, efforts were made to keep the numbering of tables, figures, boxes, equations the same as in the *2006 IPCC Guidelines*. As a consequence, the numbering of new/updated tables, figures, boxes, equations in the *2019 Refinement* is not necessarily in sequence.

⁶ The IPCC decided to produce the *2019 Refinement* as a separate Methodology Report which should be used in conjunction with the *2006 IPCC Guidelines*. Consolidating all methodological guidance into a single Methodology Report would require a new decision by the IPCC.

163
164

Section Title	Type of Refinement	2006 Guidelines Section Number	2019 Refinement Section Number
Introduction	NR	2.1	2.1
Inventory Framework	U	2.2	2.2
Overview of carbon stock change estimation	NR	2.2.1	2.2.1
Overview of non-CO ₂ emission estimation	NR	2.2.2	2.2.2
Conversion of C stock changes to CO ₂ emissions	NR	2.2.3	2.2.3
Generic methods for CO ₂ emissions and removals	NR	2.3	2.3
Change in biomass carbon stocks (above-ground biomass and below-ground biomass)	NR	2.3.1	2.3.1
Land remaining in a land-use category	NR	2.3.1.1	2.3.1.1
Land converted to a new land-use category	NR	2.3.1.2	2.3.1.2
Additional generic guidance for tier 2 methods	NG	-	2.3.1.3
Change in dead organic matter	NR	2.3.2	2.3.2
Land remaining in a land-use category	U	2.3.2.1	2.3.2.1
Land converted to a new land-use category	U	2.3.2.2	2.3.2.2

U = Update, NG = New Guidance, NR = No Refinement

165

166

167

168 "Good practice" is a key concept for inventory compilers to follow in preparing national greenhouse gas
 169 inventories. The key concept does not change in the *2019 Refinement*. The term "good practice" has been defined,
 170 since 2000 when this concept was introduced⁷, as "a set of procedures intended to ensure that greenhouse gas
 171 inventories are accurate in the sense that they are systematically neither over- nor underestimates so far as can
 172 be judged, and that uncertainties are reduced so far as practicable". This definition has gained general acceptance
 173 amongst countries as the basis for inventory development. This can be also interpreted as "a set of procedures
 174 intended to ensure that greenhouse gas inventories are accurate in the sense that they are systematically neither
 175 over- nor underestimates so far as can be judged, and that they are precise so far as practicable" in the context
 176 of refinement of Chapter 3 of Volume 1⁸.

177 These requirements are intended to ensure that estimates of anthropogenic emissions by sources and removals by
 178 sinks, even if uncertain, are *bona fide* estimates, in the sense of not containing any biases that could have been
 179 identified and eliminated, and that random errors have been reduced as far as practicable, given national
 180 circumstances. Estimates of this type are presumably the best attainable, given current scientific knowledge and
 181 available resources. The *2019 Refinement* also recognises the principles of transparency, accuracy, completeness,
 182 consistency and comparability defined in the *2006 IPCC Guidelines*.

183 The *2006 IPCC Guidelines* are intended to help prepare national inventories of anthropogenic emissions by sources
 184 and removals by sinks. Nonetheless, the Guidelines can also be relevant for estimating actual emissions or removals
 185 at the subnational, entity or project level. This is the case also with the *2019 Refinement*. Some examples of the use
 186 of the guidance in the *2006 IPCC Guidelines* and/or the *2019 Refinement* for other purposes than national inventory
 187 preparation are provided in some boxes in the *2019 Refinement*.

188

⁷ The definition was originally introduced with the *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* published in 2000.

⁸ In the *2006 IPCC Guidelines*, the term "uncertainty" was used with two meanings – a general one which is associated with both accuracy and precision, and a specific one only as the inverse of precision. This definition text of "good practice" which uses the term "precise" instead of "uncertainties" overcomes inconsistencies with general IPCC definition for uncertainty, without changing the original concept of "good practice".

Final Draft

189 5 SPECIFIC DEVELOPMENTS IN THE 2019 190 REFINEMENT

191 The *2019 Refinement* contains the following specific developments, among others, since the *2006 IPCC Guidelines*:

192 *Volume 1 (General Guidance and Reporting)*

- 193 • *National greenhouse gas inventory arrangements and management tools*: The *2019 Refinement* elaborates
194 guidance on establishing greenhouse gas inventory arrangements to support the development, improvement
195 and maintenance of national greenhouse gas inventories. This guidance is not designed to be prescriptive
196 given that the shape and form of greenhouse gas inventory arrangements depend on national circumstances.
197 Instead, the guidance provides approaches and examples of national greenhouse gas inventory arrangements
198 that could be useful. In addition, institutional arrangements include the interactions between organisations that
199 are involved with the greenhouse gas inventory inputs, compilation processes, and outputs. The *2019*
200 *Refinement* provides new guidance on generic inventory management tools such as workplans, improvement
201 plans, data management systems, quality systems, training and capacity building and documentation
202 procedures. The tools presented in this new guidance should not be considered prescriptive. However, they
203 provide some suggested approaches and examples demonstrated to be useful when developing greenhouse gas
204 inventory systems. (Chapter 1)
- 205 • *Data collection strategy*: General guidance for collecting existing national/international data and new data is
206 elaborated. The material can be used by both countries establishing a data collection strategy for the first time
207 and countries with established data collection procedures. It has also the advantage to be applicable to emission
208 factor, activity, and uncertainty data collection. (Chapter 2)
- 209 • *Use of facility-level data in inventories*: Detailed industrial facility-level data that are increasingly collected
210 for various goals such as tracking the progress of emission trading programmes or climate change policies has
211 the potential to be utilized in national greenhouse gas inventories. The challenge for inventory compilers is
212 assessing how best to integrate facility reported data to achieve improvements, especially if there are some
213 outstanding coverage and completeness issues. The *2019 Refinement* includes new guidance on how best to
214 use facility-level data that are not originally designed for national greenhouse gas inventory compilation. A
215 new decision tree for selecting facility-level data is provided as well as *good practice* reporting considerations
216 associated with facility-level data used in the national greenhouse gas inventory. (Chapter 2)
- 217 • *Uncertainty analysis*: The *2019 Refinement* provides an update on uncertainties associated with activity data.
218 It also incorporates guidance on how to derive uncertainty estimates from activity data generated based on
219 random samples. This elaborated guidance has useful applications particularly in the AFOLU sector in dealing
220 with uncertainty estimates from land use surveys or forest cover surveys. The updated guidance also includes
221 key requirements for use of Approach 1 uncertainty analysis with examples. A practical step-by-step example
222 demonstrating the use of Approach 2 uncertainty analysis (Monte-Carlo analysis) is also provided to guide
223 inventory compilers. (Chapter 3)
- 224 • *Key category analysis*: No major modifications with respect to the *2006 IPCC Guidelines* have occurred but
225 a simplification of the equation to perform *key category analysis* using trend assessment (Approach 1) has
226 been implemented in the *2019 Refinement*. General principles and guidance have been updated and an updated
227 trend approach is described. Priorities for maintenance and improvement of the inventory are addressed, and
228 new guidance in determining the appropriate level of disaggregation of greenhouse gas estimates to identify
229 key categories is provided. (Chapter 4)
- 230 • *Non-linear interpolation*: New methodology with an example has been added in the *2019 Refinement* on non-
231 linear interpolation analysis. This is relevant in cases where time series consistency is best represented by
232 multiplicative (exponential) rather than additive (linear) relationships. (Chapter 5)
- 233 • *Comparison of greenhouse gas emission estimates with atmospheric measurements*: Guidance on comparison
234 of greenhouse gas emission estimates with atmospheric measurement has been updated and elaborated to
235 reflect the state of science for atmospheric measurements and their application to verifying national emissions.
236 The most notable advances were achieved in the application of inverse models of atmospheric transport for
237 emission estimates at the national scale. Thus, atmospheric measurements are being used to provide useful
238 quality assurance of the national greenhouse gas emission estimates. The guidance highlights key components
239 and necessary steps needed for national greenhouse gas inventory verification using atmospheric
240 measurements. (Chapter 6)
- 241 • *Use and reporting of models*: The *2019 Refinement* provides new guidance on the use and reporting of models.
242 This new guidance applies to complex models, generally Tier 3 approaches. A step-by-step approach to report

- 243 on the use of models in emission inventories is presented along with a checklist for ensuring *good practice* in
244 the use of complex, higher tier models in national greenhouse gas inventories. (Chapter 6)
- 245 • *Indirect greenhouse gas emissions*: The *2019 Refinement* includes a series of updates ranging from
246 improvements in the explanation of the methodology for indirect nitrous oxide (N₂O) emissions from the
247 atmospheric deposition of nitrogen in nitrogen oxides (NO_x) and ammonia (NH₃), detailed methodological
248 guidance on treatment of carbon dioxide (CO₂) inputs to the atmosphere from emissions of carbon-containing
249 compounds, an update on the background science on precursors and indirect emissions. The *2019 Refinement*
250 also incorporates new guidance on non-biogenic sources of CO₂ from the atmospheric oxidation of methane
251 (CH₄), carbon monoxide (CO), and non-methane volatile organic compounds (NMVOCs), examples of
252 NMVOCs from the different source categories as well as guidance on carbon content of various materials
253 (percent carbon by mass) and percent of total solvent NMVOC emissions (by mass) (Chapter 7).
- 254 **Volume 2 (Energy)**
- 255 • *Fugitive emissions from mining, processing, storage and transportation of coal*: The *2019 Refinement*
256 includes guidance on emissions from exploration, and on CO₂ emissions from underground mines. (Chapter
257 4)
 - 258 • *Fugitive emissions from oil and natural gas systems*: The *2019 Refinement* includes emission factors
259 representative of current practice, including for unconventional oil and gas exploration, and methods and
260 emission factors for abandoned wells. (Chapter 4)
 - 261 • *Fuel transformation*: The *2019 Refinement* includes a new section on fugitive emissions from fuel
262 transformation, including methods for fugitive emissions from charcoal production, coke production, coal to
263 liquids and gas to liquids. (Chapter 4)
- 264 **Volume 3 (Industrial Processes and Product Use)**
- 265 • *New categories and new gases*: The *2019 Refinement* expands the scope of the *2006 IPCC Guidelines* to
266 include more manufacturing sectors identified as sources of greenhouse gases. These include production of
267 hydrogen, rare earth metals, and alumina, and waterproofing of circuit boards. In addition, a basis for future
268 methodological development is provided for fluorinated treatment of textiles, carpet, leather and paper.
269 Additional greenhouse gases identified in the IPCC Fourth and Fifth Assessment Reports, as well as other
270 references, are also included where anthropogenic sources have been identified. Greenhouse gases identified
271 in the IPCC Fourth and Fifth Assessment Reports include, for example, additional hydrofluorocarbons,
272 perfluorocarbons, and halogenated ethers, such as PPFMIE (a perfluoropolyether widely used as a heat
273 transfer fluid in electronics manufacturing). (Chapters 3, 4, 6, 8)
 - 274 • *Updates and elaborations*: The guidance for several source categories has been updated and elaborated. This
275 includes the guidance for production of nitric acid, fluorochemicals, iron and steel, aluminum, and electronics,
276 and for the production and use of refrigeration and air-conditioning equipment. (Chapters 3, 4, 6, 7)
- 277 **Volume 4 (Agriculture, Forestry and Other Land Use)**
- 278 • *Interannual variability (IAV)*: A new section has been introduced to provide an option that may be used to
279 disaggregate Managed Land Proxy (MLP) emissions and removals into those that are considered to result
280 from human effects and those that are considered to result from natural effects. These approaches may be of
281 interest to countries with AFOLU sector emissions that have high IAV due to natural effects. The section first
282 addresses definitional issues, followed by a description of whether or not different methodological approaches
283 used to estimate carbon stock changes quantify the interannual variability of emissions and removals. A
284 generic methodology to estimate, disaggregate and report the contribution of natural disturbances to the
285 emissions and removals on managed lands is then provided, along with country-specific examples of
286 approaches to disaggregating anthropogenic and natural effects on managed lands. The purpose of this
287 guidance is to support countries that wish to increase the transparency of anthropogenic greenhouse gas flux
288 estimates on managed lands. (Chapter 2)
 - 289 • *Biomass estimates*: Biomass Tier 1 factors have been updated for Forest Land, Cropland and Settlements.
290 Guidance on Tier 1 methods for Cropland and Settlements has been refined and clarified. Guidance for time
291 series consistency for Forest Land has also been updated. Factors for dead organic matter have been updated,
292 and new sections on Tier 2 guidance for the use of allometric models and biomass maps have been introduced.
293 (Chapters 2, 4, 5, 8)
 - 294 • *Soil carbon*: Tier 1 carbon stock change factors have been updated for tillage management, grassland
295 management and land use based on evolving understanding of management impacts on soils. Many of the
296 updated factors reflect a smaller impact of anthropogenic activity on soil carbon than default factors provided
297 in the *2006 IPCC Guidelines*. Reference C stocks have also been updated based on an analysis of a global
298 dataset that produces more representative reference stocks for different soil types by climate regions. The
299 default method has also been refined to estimate the impact of biochar amendments on soil carbon stocks in

Final Draft

- 300 mineral soils for cropland and grassland. More guidance is provided for developing Tier 2 stock change factors
 301 and Tier 3 methods. In addition, an alternative Tier 2 approach, i.e., steady-state method, been provided in the
 302 *Cropland Remaining Cropland* section of the report. (Chapters 2, 4, 5, 6, 7, 8)
- 303 • *Flooded Lands*: New guidance is provided for CO₂ and non-CO₂ emissions from *Land Converted to Flooded*
 304 *Lands* and *Flooded Land Remaining Flooded Land*. Methods for future development associated with these
 305 sources were included in Appendix 2 and Appendix 3 of Volume 4 of the *2006 IPCC Guidelines*. The science
 306 has matured over the past decade and these sources are now included in the main guidance (instead of
 307 appendices) of Chapter 7, Volume 4 of the *2019 Refinement* for a more complete inventory of greenhouse gas
 308 emissions from managed lands. The methods include approaches for factoring-out emissions that would have
 309 occurred otherwise if the land remained unmanaged⁹. (Chapter 7)
 - 310 • *Livestock and manure management*: Tier 1 emission factors have been updated considering current
 311 productivity data and integrating differential emission factors and for high and low productivity systems.
 312 Further, for major animal categories, Tier 1 parameters such as enteric fermentation EFs, volatile solids and
 313 nitrogen excretion are derived based on consistent data sources. The Tier 1 method to estimate CH₄ emissions
 314 from manure management has been updated for consistency with N₂O emissions. Certain Tier 2 parameters
 315 have been refined. The methane conversion rate (Y_m) for cattle and buffalo, varies based on animal diet and
 316 level of productivity. The methane conversion factor (MCF) for animal waste management systems are
 317 presented based on climatic regions, as opposed to annual temperatures and a simple calculation model for
 318 deriving the MCF based on monthly temperature regimes has been presented. Finally, improved guidance has
 319 been developed for the treatment of nitrogen transfers among livestock emission source categories and
 320 transfers to agricultural soils. (Chapter 10)
 - 321 • *Soil N₂O*: Tier 1 estimates have been updated based on the latest science for direct and indirect emission factors.
 322 A key development is the disaggregation of emission factors by climate region. (Chapter 11)
 - 323 • *Harvested wood products (HWPs)*: The methods and equations in the *2006 IPCC Guidelines* have been
 324 updated. The updated methods and equations better help inventory compilers to include the HWP pool
 325 estimates in greenhouse gas inventories using any of the three approaches: 'stock-change' approach,
 326 'production' approach and 'atmospheric flow' approach¹⁰. (Chapter 12)
- 327 **Volume 5 (Waste)**
- 328 • *Waste generation, composition and management*: The *2019 Refinement* updates key parameters used in the
 329 first order decay (FOD) method include waste generation rate and waste composition by countries and region
 330 using UN classification were updated in the comparable year (2010) and provides default value and
 331 uncertainty of carbon content, nitrogen content and degradable organic carbon (DOC) of domestic and
 332 industrial sludge in percent of dry matter. (Chapter 2)
 - 333 • *Estimation of CH₄ emission from landfill*: Guidance on the use of methane correction factor (MCF) in different
 334 management conditions of solid waste disposal sites (SWDS) has been updated. New default values for the
 335 MCF to estimate CH₄ emissions from active aeration landfill have been provided by level of landfill
 336 management (poorly and well managed). The IPCC Waste Model has been updated according to the
 337 refinement. Default values for the fraction of degradable organic carbon which decomposes (DOC_f) for
 338 different waste components and their uncertainties have been updated, and relevant guidance has been added.
 339 (Chapter 3)
 - 340 • *Incineration and open burning of waste*: Guidance on emission estimation from new technologies including
 341 gasification and pyrolysis has been elaborated with provision of CH₄ and N₂O emission factors to ensure a
 342 more complete coverage of sources. Oxidation factor of municipal solid waste (MSW) open burning has been
 343 updated. (Chapter 5)

⁹ In the outline of the *2019 Refinement* approved by the IPCC at its 44th Session in Bangkok, Thailand in October 2016, refinement was requested to update CO₂ emission factors for land converted to flooded land (Wetlands) and to develop, on the basis of comprehensive review of available literature, consistent methodologies that take into account factoring out of emissions and removals that would otherwise occur in the absence of the flooded area for estimating CO₂ and CH₄ emissions from flooded lands (both land converted to flooded land and flooded land remaining flooded land). Methods have been developed to comply with the direction from the 44th Session of the IPCC. These methods provide transparency in terms of the total emissions from Flooded Land following the Managed Land Proxy, which is consistent with other methodologies for the AFOLU sector, in addition to the net emissions based on factoring-out the amount that would have otherwise occurred if the land remained unmanaged for both *Flooded Land Remaining Flooded Land* and *Land Converted to Flooded Land*.

¹⁰ At the time the *2019 Refinement* was developed, these approaches were under discussion within the UNFCCC process in the context of revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention.

- 344 • *CH₄ emissions from wastewater treatment*: Updated guidance is provided for the estimation of CH₄ from
345 wastewater treatment, and updated emission factors for septic systems and centralised wastewater treatment
346 plants are provided. Updated emission factors are also provided for CH₄ emissions from wastewater after
347 disposal of untreated wastewater or wastewater treatment effluent into aquatic environments. (Chapter 6)
- 348 • *N₂O emissions from wastewater treatment*: New guidance and emission factors are provided for N₂O
349 emissions from domestic and industrial wastewater treatment plants, and updated emission factors are
350 provided for N₂O emissions from wastewater after disposal of untreated wastewater or wastewater treatment
351 effluent into aquatic environments. (Chapter 6)
- 352 • *Abiogenic (fossil) CO₂ emissions from wastewater treatment and discharge*: A discussion of abiogenic (fossil)
353 CO₂ emissions from wastewater treatment and discharge, where fossil organic carbon is present in wastewater
354 or treatment sludge, is presented as an appendix as a basis for future methodological development. (Chapter
355 6)
- 356 • *Discharge into aquatic environments*: An alternate set of emission factors is provided for CH₄ and N₂O
357 emissions from wastewater after disposal of untreated wastewater or wastewater treatment effluent into aquatic
358 environments when the country has activity data to differentiate the conditions of the waterbody receiving the
359 discharge. (Chapter 6)

Final Draft

360 **6 CLARIFICATION ON KEY CONCEPTS IN THE** 361 **2019 REFINEMENT**

362 The following key concepts should be noted when using the *2019 Refinement*. These are consistent with the *2006*
363 *IPCC Guidelines*.

- 364 • The *2019 Refinement* provides guidance for data gathering, compilation, and reporting. Reporting refers to the
365 presentation of emission inventory estimates in tables or other formats used to communicate inventory
366 information. The *2019 Refinement* does not provide guidance for reporting of information used to assess
367 compliance with commitments.
- 368 • Reporting tables are provided in Volume 1 as part of general guidance for reporting. They are not intended to
369 prescribe specific reporting formats under the UNFCCC that should be developed and agreed by the Parties
370 to the UNFCCC.
- 371 • The *2019 Refinement* provides methods for estimating emissions (and removals as appropriate) for each gas
372 in mass units. It does not recommend any specific metrics (e.g., GWP values) to calculate emission estimates
373 in CO₂ equivalent units. Some guidance included in the *2019 Refinement* (e.g., key category analysis) suggests
374 calculation of emission estimates in CO₂ equivalent, for which any metrics can be used.
- 375 • A *tier* represents a level of methodological complexity. Usually three tiers are provided. Tier 1 is the basic
376 method, Tier 2 intermediate and Tier 3 most demanding in terms of complexity and data requirements. Tiers
377 2 and 3 are sometimes referred to as *higher tier* methods and are generally considered to be more accurate on
378 condition that adequate data are available to develop, evaluate and apply a higher tier method.

379