

## BOUNDARY ELEMENT METHODS FOR HELMHOLTZ PROBLEMS WITH WEAKLY IMPOSED BOUNDARY CONDITIONS\*

TIMO BETCKE<sup>†</sup>, ERIK BURMAN<sup>†</sup>, AND MATTHEW W. SCROGGS<sup>‡</sup>

**Abstract.** We consider boundary element methods where the Calderón projector is used for the system matrix and boundary conditions are weakly imposed using a particular variational boundary operator designed using techniques from augmented Lagrangian methods. Regardless of the boundary conditions, both the primal trace variable and the flux are approximated. We focus on the imposition of Dirichlet conditions on the Helmholtz equation and extend the analysis of the Laplace problem from *Boundary element methods with weakly imposed boundary conditions* [Betcke, Burman, and Scroggs, *SIAM J. Sci. Comput.*, 41 (2019), pp. A1357–A1384] to this case. The theory is illustrated by a series of numerical examples.

**Key words.** boundary element methods, Helmholtz equation, wave scattering, Nitsche’s method

**MSC codes.** 65N38, 65R20

**DOI.** 10.1137/20M1334802

**1. Introduction.** In a previous paper [3], we introduced a method of weakly imposing boundary conditions on the boundary element method, inspired by Nitsche’s method [17] and Babuška’s penalty method [1] for the finite element method. Weak imposition of boundary conditions here means that neither the Dirichlet trace nor the Neumann trace is imposed exactly; instead an  $h$ -dependent boundary condition is imposed that is weighted in such a way that optimal error estimates may be derived and the exact boundary condition is recovered in the asymptotic limit.

In [3], we introduced the weak imposition of Dirichlet, Neumann, and Robin boundary conditions on Laplace’s equation; in [7], we applied this method to Signorini contact conditions, again for Laplace’s equation. In this paper, we look at how this method and its analysis can be extended to be used for the Helmholtz equation, focusing on the exterior Helmholtz Dirichlet problem: Find  $u = u^{\text{inc}} + u^{\text{scat}} \in H_{\text{loc}}^1(\Delta, \Omega^+)$  such that

$$(1.1a) \quad -\Delta u - k^2 u = 0 \quad \text{in } \Omega^+,$$

$$(1.1b) \quad \frac{\partial u^{\text{scat}}}{\partial |\mathbf{x}|} - iku^{\text{scat}} = o(|\mathbf{x}|^{-1}) \quad \text{as } |\mathbf{x}| \rightarrow \infty,$$

$$(1.1c) \quad u = g_D \quad \text{on } \Gamma,$$

where  $\Omega^- \subset \mathbb{R}^3$  is a bounded interior domain with polyhedral boundary  $\Gamma$ ,  $\Omega^+ = \mathbb{R}^3 \setminus \Omega^-$  is the domain exterior to  $\Omega^-$ ,  $\boldsymbol{\nu}$  is the unit normal to the surface  $\Gamma$  pointing outwards into  $\Omega^+$ ,  $u^{\text{inc}}$  is a known incident wave, and  $k \in \mathbb{R}$  is the wavenumber of the problem. We assume that  $g_D \in H^{1/2}(\Gamma)$ , and  $u \in H^{3/2+\epsilon}(\Gamma)$  for some  $\epsilon > 0$ .

\*Submitted to the journal’s Methods and Algorithms for Scientific Computing section April 28, 2020; accepted for publication (in revised form) June 21, 2022; published electronically September 6, 2022.

<https://doi.org/10.1137/20M1334802>

**Funding:** The work of the first and second authors was supported by EPSRC grants EP/P01576X/1 and EP/P012434/1.

<sup>†</sup>Department of Mathematics, University College London, London, WC1E 6BT, UK (t.betcke@ucl.ac.uk, e.burman@ucl.ac.uk).

<sup>‡</sup>Department of Engineering, University of Cambridge, Cambridge, CB2 1PZ, UK (mws48@cam.ac.uk, <https://www.msccroggs.co.uk>).

Due to the Sommerfeld radiation condition (1.1b), (1.1) has a unique solution [20]. The formulation of Helmholtz problems using boundary integral equations are covered in detail in [16], and their discretization is examined in [23, section 7.6].

The use of discretization and the boundary element method to solve Helmholtz problems has been well studied. For sufficiently small wavenumbers  $k$  and sufficiently smooth boundaries, the operators involved are coercive, and hence a priori error bounds can be derived [22, 21, 8]. For values of  $k$  and domains for which coercivity cannot be shown, error bounds have been shown that involve both the mesh size  $h$  and the wavenumber  $k$  [2, 19]. If the wavenumber is varied, then the mesh must be refined to keep the value of  $hk$  constant in order to maintain a low error [10]. The use of  $hp$ -BEM for Helmholtz has also been studied and analyzed [14, 15].

The use of blocked operator formulations to solve Helmholtz problems is common for domain decomposition problems, where the boundary element method is used in multiple domains with different wavenumbers [18, 12], or a combination of finite and boundary element methods can be used [11]. To avoid the appearance of spurious resonances in solutions, coupled stabilized formulations can be solved [9]. The formulations presented in this paper are, in general, more expensive than standard formulations, as they require the assembly of the full Calderón system. In these cases, however, larger blocked systems are already being assembled, and so it may be possible to impose boundary conditions on them weakly with little additional cost.

The method proposed in this paper is applicable to low and medium frequency problems. In practice, preconditioning limits the method's effectiveness for higher frequency problems. However, one main advantage of this method is its immunity to eigenvalues of the interior problem: the solution to (1.1) can be found for any wavenumber using our method without any modification to stabilize against eigenvalues being necessary.

Although we do not present an analysis of problems with mixed boundary conditions, we discuss our method's potential application to such problems in section 6. As we discussed in [3] for Laplace problems, our method of weakly imposing boundary conditions is advantageous when solving mixed problems, as they can be implemented by assembling different sparse terms on different parts of the mesh without needing to adapt the Calderón part of the formulation.

In section 2, we define the boundary operators used in our formulations and present some of their important properties. In section 3, we derive our formulation for Dirichlet Helmholtz problems. In section 4, we analyze this formulation and prove a priori error bounds. In section 5, we present some numerical experiments, and in section 6 we give some concluding remarks.

**2. Boundary operators.** We define the Green's function for the Helmholtz operator in  $\mathbb{R}^3$  by

$$(2.1) \quad G(\mathbf{x}, \mathbf{y}) = \frac{e^{ik|\mathbf{x}-\mathbf{y}|}}{4\pi|\mathbf{x}-\mathbf{y}|}.$$

In this paper, we focus on the problem in  $\mathbb{R}^3$ . Similar analysis can be used for problems in  $\mathbb{R}^2$ , in which case this definition should be replaced by  $G(\mathbf{x}, \mathbf{y}) = \frac{i}{4}H_0^{(1)}(k|\mathbf{x}-\mathbf{y}|)$ , where  $H_0^{(1)}$  is a Hankel function of the first kind.

In the standard fashion (see, e.g., [23, Chapter 6]), we define the single layer potential operator,  $\mathcal{V} : H^{-1/2}(\Gamma) \rightarrow H_{\text{loc}}^1(\Omega^+)$ , and the double layer potential operator,  $\mathcal{K} : H^{1/2}(\Gamma) \rightarrow H_{\text{loc}}^1(\Omega^+)$  for  $v \in H^{1/2}(\Gamma)$ ,  $\mu \in H^{-1/2}(\Gamma)$ , and  $\mathbf{x} \in \Omega^+ \setminus \Gamma$  by

$$(2.2) \quad (\mathcal{V}\mu)(\mathbf{x}) := \int_{\Gamma} G(\mathbf{x}, \mathbf{y})\mu(\mathbf{y}) \, d\mathbf{y},$$

$$(2.3) \quad (\mathcal{K}v)(\mathbf{x}) := \int_{\Gamma} \frac{\partial G(\mathbf{x}, \mathbf{y})}{\partial \nu_{\mathbf{y}}} v(\mathbf{y}) \, d\mathbf{y}.$$

We define the space  $H^1_{\text{loc}}(\Delta, \Omega^+) := \{v \in H^1_{\text{loc}}(\Omega^+) : \Delta v \in L^2(\Omega^+)\}$ , and then we define the Dirichlet and Neumann traces,  $\gamma^{\pm}_D : H^1_{\text{loc}}(\Omega^+) \rightarrow H^{1/2}(\Gamma)$  and  $\gamma^{\pm}_N : H^1_{\text{loc}}(\Delta, \Omega^+) \rightarrow H^{-1/2}(\Gamma)$ , by

$$(2.4) \quad \gamma^+_D f(\mathbf{x}) := \lim_{\Omega^+ \ni \mathbf{y} \rightarrow \mathbf{x} \in \Gamma} f(\mathbf{y}),$$

$$(2.5) \quad \gamma^+_N f(\mathbf{x}) := \lim_{\Omega^+ \ni \mathbf{y} \rightarrow \mathbf{x} \in \Gamma} \nu_{\mathbf{x}} \cdot \nabla f(\mathbf{y}).$$

We recall that if the Dirichlet and Neumann traces of a solution of (1.1a) are known, then the potentials (2.2) and (2.3) may be used to reconstruct the function in  $\Omega^+$  using the following relation.

$$(2.6) \quad u = \mathcal{K}(\gamma^+_D u) - \mathcal{V}(\gamma^+_N u).$$

It is also known [23, Lemma 6.6] that for all  $\mu \in H^{-1/2}(\Gamma)$ , the function

$$(2.7) \quad u^{\mathcal{V}}_{\mu} := \mathcal{V}\mu$$

satisfies  $-\Delta u^{\mathcal{V}}_{\mu} - k^2 u^{\mathcal{V}}_{\mu} = 0$ . Similarly, for the double layer potential there holds [23, Lemma 6.10] that for all  $v \in H^{1/2}(\Gamma)$ , the function

$$(2.8) \quad u^{\mathcal{K}}_v := \mathcal{K}v$$

satisfies  $-\Delta u^{\mathcal{K}}_v - k^2 u^{\mathcal{K}}_v = 0$ .

We define  $\{\gamma_D f\}_{\Gamma}$  and  $\{\gamma_N f\}_{\Gamma}$  to be the averages of the interior and exterior Dirichlet and Neumann traces of  $f$ . We define the single layer, double layer, adjoint double layer, and hypersingular boundary integral operators,  $\mathbb{V} : H^{-1/2}(\Gamma) \rightarrow H^{1/2}(\Gamma)$ ,  $\mathbb{K} : H^{1/2}(\Gamma) \rightarrow H^{1/2}(\Gamma)$ ,  $\mathbb{K}' : H^{-1/2}(\Gamma) \rightarrow H^{-1/2}(\Gamma)$ , and  $\mathbb{W} : H^{1/2}(\Gamma) \rightarrow H^{-1/2}(\Gamma)$ , by

$$(2.9a) \quad (\mathbb{K}v)(\mathbf{x}) := \{\gamma_D \mathcal{K}v\}_{\Gamma}(\mathbf{x}), \quad (\mathbb{V}\mu)(\mathbf{x}) := \{\gamma_D \mathcal{V}\mu\}_{\Gamma}(\mathbf{x}),$$

$$(2.9b) \quad (\mathbb{W}v)(\mathbf{x}) := -\{\gamma_N \mathcal{K}v\}_{\Gamma}(\mathbf{x}), \quad (\mathbb{K}'\mu)(\mathbf{x}) := \{\gamma_N \mathcal{V}\mu\}_{\Gamma}(\mathbf{x}),$$

where  $\mathbf{x} \in \Gamma$ ,  $v \in H^{1/2}(\Gamma)$  and  $\mu \in H^{-1/2}(\Gamma)$  [23, Chapter 6].

We define  $\llbracket \gamma_D \rrbracket_{\Gamma} := \gamma^+_D - \gamma^-_D$  and  $\llbracket \gamma_N \rrbracket_{\Gamma} := \gamma^+_N - \gamma^-_N$  to be the jumps of the interior and exterior Dirichlet and Neumann traces across the boundary. In [23, Chapter 6], the following jump conditions are shown:

$$(2.10) \quad \llbracket \gamma_D \rrbracket_{\Gamma} \mathcal{V} = \llbracket \gamma_N \rrbracket_{\Gamma} \mathcal{K} = 0, \quad \llbracket \gamma_N \rrbracket_{\Gamma} \mathcal{V} = -\llbracket \gamma_D \rrbracket_{\Gamma} \mathcal{K} = -\text{Id},$$

where  $\text{Id}$  is the identity operator.

It follows from (2.9) and (2.10) that

$$(2.11a) \quad \gamma^+_D \mathcal{V} = \mathbb{V}, \quad \gamma^+_N \mathcal{V} = -\sigma \text{Id} + \mathbb{K}',$$

$$(2.11b) \quad \gamma^+_D \mathcal{K} = (1 - \sigma) \text{Id} + \mathbb{K}, \quad \gamma^+_N \mathcal{K} = -\mathbb{W},$$

$$(2.11c) \quad \gamma^-_D \mathcal{V} = \mathbb{V}, \quad \gamma^-_N \mathcal{V} = \sigma \text{Id} + \mathbb{K}',$$

$$(2.11d) \quad \gamma^-_D \mathcal{K} = (1 - \sigma) \text{Id} + \mathbb{K}, \quad \gamma^-_N \mathcal{K} = -\mathbb{W},$$

where  $\sigma$  is defined as in [23, equation 6.11]. We note that  $\sigma = \frac{1}{2}$  almost everywhere.

The following coercivity results are known for the single layer and hypersingular operators in  $\mathbb{R}^3$ , where  $\langle \cdot, \cdot \rangle_\Gamma$  denotes the  $H^{1/2}(\Gamma)$ – $H^{-1/2}(\Gamma)$  duality pairing.

LEMMA 2.1 (Gårding's inequality for  $\mathbf{V}$ ). *There exists a compact operator  $\mathbf{T}_\mathbf{V} : H^{-1/2}(\Gamma) \rightarrow H^{1/2}(\Gamma)$  and  $\alpha_\mathbf{V} > 0$  such that*

$$(2.12) \quad \alpha_\mathbf{V} \|\mu\|_{H^{-1/2}(\Gamma)}^2 \leq \langle \mathbf{V}\mu, \mu \rangle_\Gamma + \langle \mathbf{T}_\mathbf{V}\mu, \mu \rangle_\Gamma \quad \forall \mu \in H^{-1/2}(\Gamma).$$

*Proof.* [23, Theorem 6.40].

LEMMA 2.2 (Gårding's inequality for  $\mathbf{W}$ ). *There exists a compact operator  $\mathbf{T}_\mathbf{W} : H^{1/2}(\Gamma) \rightarrow H^{-1/2}(\Gamma)$  and  $\alpha_\mathbf{W} > 0$  such that*

$$(2.13) \quad \alpha_\mathbf{W} \|v\|_{H^{1/2}(\Gamma)}^2 \leq \langle \mathbf{W}v, v \rangle_\Gamma + \langle \mathbf{T}_\mathbf{W}v, v \rangle_\Gamma \quad \forall v \in H_*^{1/2}(\Gamma),$$

where  $H_*^{1/2}(\Gamma)$  denotes the set of functions  $v \in H^{1/2}(\Gamma)$  such that  $\bar{v} = 0$ , where  $\bar{v} := \frac{\langle v, \mathbf{1} \rangle_\Gamma}{\langle \mathbf{1}, \mathbf{1} \rangle_\Gamma}$  is the average value of  $v$ .

*Proof.* This follows by applying the proof of [23, Theorem 6.40] to the hypersingular operator.  $\square$

The following boundedness results are also known.

LEMMA 2.3 (boundedness). *There exist  $C_\mathbf{V}, C_\mathbf{K}, C'_\mathbf{K}, C_\mathbf{W} > 0$  such that*

$$(2.14) \quad i) \quad \|\mathbf{V}\mu\|_{H^{1/2}(\Gamma)} \leq C_\mathbf{V} \|\mu\|_{H^{-1/2}(\Gamma)} \quad \forall \mu \in H^{-1/2}(\Gamma),$$

$$(2.15) \quad ii) \quad \|\mathbf{K}v\|_{H^{1/2}(\Gamma)} \leq C_\mathbf{K} \|v\|_{H^{1/2}(\Gamma)} \quad \forall v \in H^{1/2}(\Gamma),$$

$$(2.16) \quad iii) \quad \|\mathbf{K}'\mu\|_{H^{-1/2}(\Gamma)} \leq C'_\mathbf{K} \|\mu\|_{H^{-1/2}(\Gamma)} \quad \forall \mu \in H^{-1/2}(\Gamma),$$

$$(2.17) \quad iv) \quad \|\mathbf{W}v\|_{H^{-1/2}(\Gamma)} \leq C_\mathbf{W} \|v\|_{H^{1/2}(\Gamma)} \quad \forall v \in H^{1/2}(\Gamma).$$

*Proof.* [23, sections 6.2–6.5 and 6.9].  $\square$

We define the exterior Calderón projector by

$$(2.18) \quad \mathbf{C}^+ := \begin{pmatrix} (1 - \sigma)\text{Id} + \mathbf{K} & -\mathbf{V} \\ -\mathbf{W} & \sigma\text{Id} - \mathbf{K}' \end{pmatrix},$$

and recall that if  $u$  is a solution of (1.1a), then it satisfies

$$(2.19) \quad \mathbf{C}^+ \begin{pmatrix} \gamma_\mathbf{D}^+ u \\ \gamma_\mathbf{N}^+ u \end{pmatrix} = \begin{pmatrix} \gamma_\mathbf{D}^+ u \\ \gamma_\mathbf{N}^+ u \end{pmatrix}.$$

When considering eigenvalues of the Laplacian, we will make use of the interior Calderón projector. This is defined by

$$(2.20) \quad \mathbf{C}^- := \begin{pmatrix} (1 - \sigma)\text{Id} - \mathbf{K} & \mathbf{V} \\ \mathbf{W} & \sigma\text{Id} + \mathbf{K}' \end{pmatrix}.$$

If  $u^-$  is a solution of an interior Helmholtz problem, then it satisfies

$$(2.21) \quad \mathbf{C}^- \begin{pmatrix} \gamma_\mathbf{D}^- u^- \\ \gamma_\mathbf{N}^- u^- \end{pmatrix} = \begin{pmatrix} \gamma_\mathbf{D}^- u^- \\ \gamma_\mathbf{N}^- u^- \end{pmatrix}.$$

Taking the product of the exterior identity (2.19) with two test functions and using the fact that  $\sigma = \frac{1}{2}$  almost everywhere, we arrive at the following equations:

$$(2.22) \quad \langle \gamma_D^+ u, \mu \rangle_\Gamma = \langle (\frac{1}{2} \text{Id} + \mathbf{K}) \gamma_D^+ u, \mu \rangle_\Gamma - \langle \mathbf{V} \gamma_N^+ u, \mu \rangle_\Gamma \quad \forall \mu \in H^{-1/2}(\Gamma),$$

$$(2.23) \quad \langle \gamma_N^+ u, v \rangle_\Gamma = \langle (\frac{1}{2} \text{Id} - \mathbf{K}') \gamma_N^+ u, v \rangle_\Gamma - \langle \mathbf{W} \gamma_D^+ u, v \rangle_\Gamma \quad \forall v \in H^{1/2}(\Gamma).$$

For a more compact notation, we write  $u$  in the place of  $\gamma_D^+ u$  and introduce  $\lambda = \gamma_N^+ u$  and the exterior Calderón form

$$(2.24) \quad \mathcal{C}^+[(u, \lambda), (v, \mu)] := \langle (\frac{1}{2} \text{Id} + \mathbf{K})u, \mu \rangle_\Gamma - \langle \mathbf{V}\lambda, \mu \rangle_\Gamma + \langle (\frac{1}{2} \text{Id} - \mathbf{K}')\lambda, v \rangle_\Gamma - \langle \mathbf{W}u, v \rangle_\Gamma.$$

We may then rewrite (2.22) and (2.23) as

$$(2.25) \quad \mathcal{C}^+[(u, \lambda), (v, \mu)] = \langle u, \mu \rangle_\Gamma + \langle \lambda, v \rangle_\Gamma.$$

We will also frequently use the multitrace form, defined by

$$(2.26) \quad \mathcal{A}[(u, \lambda), (v, \mu)] := - \langle \mathbf{K}u, \mu \rangle_\Gamma + \langle \mathbf{V}\lambda, \mu \rangle_\Gamma + \langle \mathbf{K}'\lambda, v \rangle_\Gamma + \langle \mathbf{W}u, v \rangle_\Gamma,$$

and the multitrace form with compact perturbation,

$$(2.27) \quad \mathcal{A}_\Gamma[(u, \lambda), (v, \mu)] := - \langle \mathbf{K}u, \mu \rangle_\Gamma + \langle (\mathbf{V} + \mathbf{T}_V)\lambda, \mu \rangle_\Gamma + \langle \mathbf{K}'\lambda, v \rangle_\Gamma + \langle (\mathbf{W} + \mathbf{T}_W)u, v \rangle_\Gamma.$$

Using (2.26), we may rewrite (2.25) as

$$(2.28) \quad \mathcal{A}[(u, \lambda), (v, \mu)] = -\frac{1}{2} \langle u, \mu \rangle_\Gamma - \frac{1}{2} \langle \lambda, v \rangle_\Gamma.$$

To quantify the two traces we introduce the product space

$$\mathbb{V} := H^{1/2}(\Gamma) \times H^{-1/2}(\Gamma).$$

We also introduce the associated norm

$$\|(v, \mu)\|_{\mathbb{V}} := \|v\|_{H^{1/2}(\Gamma)} + \|\mu\|_{H^{-1/2}(\Gamma)}.$$

Using the results in Lemmas 2.1–2.3, we obtain the continuity and coercivity of  $\mathcal{A}$ .

LEMMA 2.4 (continuity). *There exists  $C > 0$  such that*

$$|\mathcal{A}[(w, \eta), (v, \mu)]| \leq C \|(w, \eta)\|_{\mathbb{V}} \|(v, \mu)\|_{\mathbb{V}} \quad \forall (w, \eta), (v, \mu) \in \mathbb{V}.$$

*Proof.* Use Lemma 2.3. □

LEMMA 2.5 (coercivity). *There exists  $\alpha > 0$  and compact operators  $\mathbf{T}_V$  and  $\mathbf{T}_W$  such that*

$$\alpha \left( \|v\|_{H^{1/2}(\Gamma)}^2 + \|\mu\|_{H^{-1/2}(\Gamma)}^2 \right) \leq \mathcal{A}[(v, \mu), (v, \mu)] + \langle \mathbf{T}_W v, v \rangle_\Gamma + \langle \mathbf{T}_V \mu, \mu \rangle_\Gamma \quad \forall (v, \mu) \in \mathbb{V}.$$

*Proof.* Use the coercivity of  $\mathbf{V}$  and  $\mathbf{W}$  from Lemmas 2.1 and 2.2, and let  $\alpha = \min(\alpha_W, \alpha_V)$ . □

**3. Derivation of a formulation for Helmholtz Dirichlet problems.** In this section, we derive a formulation for the exterior Helmholtz problem with non-homogeneous Dirichlet conditions.

As in [3], we write the boundary condition as

$$(3.1) \quad R_\Gamma(u, \lambda) = 0$$

and look to solve

$$(3.2) \quad \mathcal{A}[(u, \lambda), (v, \mu)] = -\frac{1}{2} \langle u, \mu \rangle_\Gamma - \frac{1}{2} \langle \lambda, v \rangle_\Gamma + \langle R_\Gamma(u, \lambda), \beta_1 v + \beta_2 \mu \rangle_\Gamma.$$

**3.1. Dirichlet boundary condition.** To impose a Dirichlet boundary condition, we choose  $\beta_1 = -i\beta_D^{1/2}$ ,  $\beta_2 = i\beta_D^{-1/2}$ , where  $\beta_D$  will be identified with a mesh-dependent penalty parameter, and

$$(3.3) \quad R_D(u, \lambda) := i\beta_D^{1/2}(g_D - u),$$

where  $g_D \in H^{1/2}(\Gamma)$  is the Dirichlet data.

Inserting this into (3.2), we obtain the formulation

$$(3.4) \quad \mathcal{A}[(u, \lambda), (v, \mu)] - \frac{1}{2} \langle u, \mu \rangle_\Gamma + \frac{1}{2} \langle \lambda, v \rangle_\Gamma + \langle \beta_D u, v \rangle_\Gamma = \langle g_D, \beta_D v - \mu \rangle_\Gamma.$$

This leads us to the following formulation for the Helmholtz Dirichlet problem: Find  $(u, \lambda) \in \mathbb{V}$  such that

$$(3.5) \quad \mathcal{A}[(u, \lambda), (v, \mu)] + \mathcal{B}_D^+[(u, \lambda), (v, \mu)] = \mathcal{L}_D^+(v, \mu) \quad \forall (v, \mu) \in \mathbb{V},$$

where

$$(3.6) \quad \mathcal{B}_D^+[(u, \lambda), (v, \mu)] := \frac{1}{2} \langle \lambda, v \rangle_\Gamma - \frac{1}{2} \langle u, \mu \rangle_\Gamma + \langle \beta_D u, v \rangle_\Gamma,$$

$$(3.7) \quad \mathcal{L}_D^+(v, \mu) := \langle g_D, \beta_D v - \mu \rangle_\Gamma.$$

We now show that a solution of the Helmholtz problem (1.1) is also a solution of this weak formulation.

**PROPOSITION 3.1.** *If  $u$  is a solution of (1.1), then  $(\gamma_D^+ u, \gamma_N^+ u)$  is a solution of (3.5).*

*Proof.* Let  $(v, \mu) \in \mathbb{V}$ . By (2.28), we see that

$$\begin{aligned} & (\mathcal{A} + \mathcal{B}_D^+)[(\gamma_D^+ u, \gamma_N^+ u), (v, \mu)] \\ &= -\frac{1}{2} \langle \gamma_D^+ u, \mu \rangle_\Gamma - \frac{1}{2} \langle \gamma_N^+ u, v \rangle_\Gamma + \frac{1}{2} \langle \gamma_N^+ u, v \rangle_\Gamma - \frac{1}{2} \langle \gamma_D^+ u, \mu \rangle_\Gamma + \langle \beta_D \gamma_D^+ u, v \rangle_\Gamma \\ &= -\langle \gamma_D^+ u, \mu \rangle_\Gamma + \langle \beta_D \gamma_D^+ u, v \rangle_\Gamma \\ &= \langle \gamma_D^+ u, \beta_D v - \mu \rangle_\Gamma. \end{aligned}$$

Using (1.1c), we see that

$$\begin{aligned} (\mathcal{A} + \mathcal{B}_D^+)[(\gamma_D^+ u, \gamma_N^+ u), (v, \mu)] &= \langle g_D, \beta_D v - \mu \rangle_\Gamma \\ &= \mathcal{L}_D^+(v, \mu). \quad \square \end{aligned}$$

To discretize (3.5), we introduce a family of conforming, shape regular triangulations of  $\Gamma$ ,  $\{\mathcal{T}_h\}_{h>0}$ , indexed by the largest element diameter of the mesh,  $h$ . We then consider the following finite element spaces:

$$\begin{aligned} \mathbb{P}_h^p &:= \{v_h \in C^0(\Gamma) : v_h|_T \in \mathbb{P}_p(T), \text{ for every } T \in \mathcal{T}_h\}, \\ \text{DP}_h^q &:= \{v_h \in L^2(\Gamma) : v_h|_T \in \mathbb{P}_q(T), \text{ for every } T \in \mathcal{T}_h\}, \end{aligned}$$

where  $\mathbb{P}_p(T)$  denotes the space of polynomials of order less than or equal to  $p$  on a triangle  $T$ , and  $\{\Gamma_i\}_{i=1}^M$  are the polygonal faces of  $\Gamma$ . We observe that  $\mathbb{P}_h^p \subset H^{1/2}(\Gamma)$  and  $\text{DP}_h^q \subset L^2(\Gamma)$ .

We let  $\mathbb{V}_h$  be a discrete product space: in our case, we take this to be equal to either  $\mathbb{P}_h^p(\Gamma) \times \text{DP}_h^q(\Gamma)$  or  $\mathbb{P}_h^p(\Gamma) \times \mathbb{P}_h^q(\Gamma)$ , although a wide range of other choices are possible. We let  $\mathbb{W}$  be a product Hilbert space, such that  $\mathbb{V}_h \subset \mathbb{W} \subset \mathbb{V}$ , and we let  $\|\cdot\|_{\mathbb{B}}$  be a norm defined on  $\mathbb{W}$ , such that for all  $(v, \mu) \in \mathbb{W}$ ,  $\|(v, \mu)\|_{\mathbb{B}} \geq \|(v, \mu)\|_{\mathbb{V}}$ .

Using the space  $\mathbb{V}_h$ , we look to solve the discrete problem: Find  $(u_h, \lambda_h) \in \mathbb{V}_h$  such that

$$(3.8) \quad \mathcal{A}[(u_h, \lambda_h), (v_h, \mu_h)] + \mathcal{B}_D^-[u_h, \lambda_h], (v_h, \mu_h)] = \mathcal{L}_D^-(v_h, \mu_h) \quad \forall (v_h, \mu_h) \in \mathbb{V}_h.$$

We define the norm

$$\|(v, \mu)\|_{\mathcal{B}_D} := \|(v, \mu)\|_{\mathbb{V}} + |\beta_D|^{1/2} \|v\|_{L^2(\Gamma)}.$$

**4. Analysis.** In this section, we analyze the formulation derived in the previous section.

**4.1. Analysis of the continuous problem.** We begin by analyzing the continuous problem (3.5). In the same way as we did in [3], we prove that the form  $\mathcal{A} + \mathcal{B}_D^+$  is continuous.

**PROPOSITION 4.1 (continuity).** *There exists  $M > 0$  such that  $\forall (w, \eta), (v, \mu) \in \mathbb{W}$ ,*

$$|\mathcal{A}[(w, \eta), (v, \mu)] + \mathcal{B}_D^+[(w, \eta), (v, \mu)]| \leq M \|(w, \eta)\|_{\mathcal{B}_D} \|(v, \mu)\|_{\mathcal{B}_D}.$$

*Proof.* This can be proved in the same way as [3, Proposition 4.13] but with  $|\beta_D|$  in the place of  $\beta_D$ . □

Eigenvalues of the interior Laplacian have an effect on the boundary integral formulation of the exterior problem. These eigenvalues are defined as follows.

**DEFINITION 4.2 (eigenvalues of the Laplacian).** *If the interior Laplace problem (find  $u$  such that*

$$(4.1a) \quad -\Delta u = l_D u \quad \text{in } \Omega^-,$$

$$(4.1b) \quad u = 0 \quad \text{on } \Gamma)$$

*has multiple solutions, then  $l_D$  is called a Dirichlet eigenvalue of the interior Laplacian.*

*If the interior Laplace problem (find  $u$  such that*

$$(4.2a) \quad -\Delta u = l_N u \quad \text{in } \Omega^-,$$

$$(4.2b) \quad \frac{\partial u}{\partial \nu} = 0 \quad \text{on } \Gamma)$$

*has multiple solutions that differ by more than a constant, then  $l_N$  is called a Neumann eigenvalue of the interior Laplacian.*

*If the interior Laplace problem (find  $u$  such that*

$$(4.3a) \quad -\Delta u = l_R u \quad \text{in } \Omega^-,$$

$$(4.3b) \quad \frac{\partial u}{\partial \nu} + \beta_D u = 0 \quad \text{on } \Gamma$$

has multiple solutions, then we call  $l_R$  a Robin eigenvalue of the interior Laplacian with Robin parameter  $\beta_D$ .

We now prove some important properties of these eigenvalues.

LEMMA 4.3. *For  $l \in \mathbb{C}$ , at most one of the following statements is true:*

- $l$  is a Dirichlet eigenvalue of the interior Laplacian;
- $l$  is a Neumann eigenvalue of the interior Laplacian;
- $l$  is a Robin eigenvalue of the interior Laplacian (for some  $\beta_D \neq 0$ ).

*Proof.* By [23, section 7.6], we know that a value cannot be both a Dirichlet and a Neumann eigenvalue.

If  $l$  is both a Dirichlet and a Robin eigenvalue, then there exist  $u_1$  and  $u_2$  satisfying (4.1a) whose values on  $\Gamma$  satisfy  $u_1 = u_2 = 0$  and

$$\begin{aligned} 0 &= \frac{\partial u_1}{\partial \nu} + \beta_D u_1 = \frac{\partial u_1}{\partial \nu}, \\ 0 &= \frac{\partial u_2}{\partial \nu} + \beta_D u_2 = \frac{\partial u_2}{\partial \nu}. \end{aligned}$$

If  $u_1$  and  $u_2$  differed by a constant, their values on  $\Gamma$  would differ by the same constant, contradicting the boundary conditions  $u_1 = u_2 = 0$ . Therefore  $l$  is also a Neumann eigenvalue, which is a contradiction.

Similarly, if  $l$  is both a Neumann and a Robin eigenvalue, then in the same way we see that  $l$  is also a Dirichlet eigenvalue, leading to a similar contradiction.  $\square$

LEMMA 4.4. *If  $\text{Im}(\beta_D) \neq 0$ , then the interior Laplacian with Robin parameter  $\beta_D$  has no nontrivial real Robin eigenvalues.*

*Proof.* Let  $\beta_D \in \mathbb{C}$  with  $\text{Im}(\beta_D) \neq 0$ . Suppose that  $l_R \in \mathbb{R} \setminus \{0\}$  is a Robin eigenvalue of the interior Laplacian with corresponding eigenfunction  $u_R$ , i.e.,

$$\begin{aligned} -\Delta u_R &= l_R u_R && \text{in } \Omega^-, \\ \frac{\partial u_R}{\partial \nu} + \beta_D u_R &= 0 && \text{on } \Gamma. \end{aligned}$$

Consider the weak formulation of this problem: find  $u_R \in H^1(\Omega)$  such that

$$\langle \nabla u_R, \nabla v \rangle_\Omega + \beta_D \langle u_R, v \rangle_\Gamma = l_R \langle u_R, v \rangle_\Omega \quad \forall v \in H^1(\Omega).$$

Taking  $v = u_R$  leads to

$$\|\nabla u_R\|_{L^2(\Omega)}^2 + \beta_D \|u_R\|_{L^2(\Gamma)}^2 = l_R \langle u_R, u_R \rangle_\Omega.$$

Taking the imaginary part of this gives

$$\text{Im}(\beta_D) \|u_R\|_{L^2(\Gamma)}^2 = 0.$$

As  $\text{Im}(\beta_D) \neq 0$ , this implies that  $\|u_R\|_{L^2(\Gamma)} = 0$ , and so  $u_R = 0$  on  $\Gamma$ .



This, however, implies that  $u_R$  is also a Dirichlet eigenfunction of the Laplacian with eigenvalue  $l_R$ , contradicting Lemma 4.3. Hence no such real eigenvalue exists.  $\square$

We now proceed to prove that the form  $\mathcal{A} + \mathcal{B}_D^+$  is injective. First we prove this when  $k^2$  is not a Dirichlet eigenvalue. Note that by Lemma 4.4, the assumption that  $k^2$  is not a Robin eigenvalue of the interior Laplacian holds whenever  $\text{Im}(\beta_D) \neq 0$ .

LEMMA 4.5 (injectivity, part one). *Let  $(v, \mu) \in \mathbb{W}$ . If  $k^2$  is not a Dirichlet eigenvalue of the interior Laplacian and  $k^2$  is not a Robin eigenvalue of the interior Laplacian with Robin parameter  $\beta_D$ , then  $\forall (w, \eta) \in \mathbb{W}$*

$$\mathcal{A}[(v, \mu), (w, \eta)] + \mathcal{B}_D^+[(v, \mu), (w, \eta)] = 0$$

implies that  $(v, \mu) = 0$ .

*Proof.* Suppose that  $(v, \mu) \in \mathbb{W}$  such that

$$(4.4) \quad (\mathcal{A} + \mathcal{B}_D^+)[(v, \mu), (w, \eta)] = 0 \quad \forall (w, \eta) \in \mathbb{W}.$$

Taking  $w = 0$  in (4.4), we see that,  $\forall \eta \in H^{-1/2}(\Gamma)$ ,

$$(4.5) \quad \langle \mathbf{V}\mu, \eta \rangle_\Gamma = \langle (\frac{1}{2}\text{Id} + \mathbf{K})v, \eta \rangle_\Gamma.$$

$k^2$  is not a Dirichlet eigenvalue of the interior Laplacian, so by [23, section 7.6] we see that  $\mathbf{V}$  is invertible, and (as (4.5) is a direct boundary integral formulation) there exists a solution to the interior Helmholtz equation  $\tilde{p} \in H^1(\Omega^-)$  such that

$$\begin{aligned} \gamma_D^- \tilde{p} &= v, \\ \gamma_N^- \tilde{p} &= \mu. \end{aligned}$$

Taking  $\eta = 0$  in (4.4), we see that,  $\forall w \in H^{1/2}(\Gamma)$ ,

$$(4.6) \quad 0 = \langle \mathbf{W}v, w \rangle_\Gamma + \langle \beta_D v, w \rangle_\Gamma + \langle (\frac{1}{2}\text{Id} + \mathbf{K}')\mu, w \rangle_\Gamma.$$

We also know from the second line of the interior Calderón identity (2.21) that

$$(4.7) \quad \langle \mathbf{W}v, w \rangle_\Gamma + \langle (\frac{1}{2}\text{Id} + \mathbf{K}')\mu, w \rangle_\Gamma = \langle \mathbf{W}\gamma_D^- \tilde{p}, w \rangle_\Gamma + \langle (\frac{1}{2}\text{Id} + \mathbf{K}')\gamma_N^- \tilde{p}, w \rangle_\Gamma$$

$$(4.8) \quad = \langle \gamma_N^- \tilde{p}, w \rangle_\Gamma$$

$$(4.9) \quad = \langle \mu, w \rangle_\Gamma,$$

and so using (4.6) we see that

$$(4.10) \quad 0 = \langle \mu, w \rangle_\Gamma + \langle \beta_D v, w \rangle_\Gamma,$$

and so  $\mu = -\beta_D v$ .

This means that  $\tilde{p}$  is the solution (4.3) with  $l_R = k^2$ . If  $k^2$  is not an eigenvalue of (4.3), then the unique solution of this problem is  $\tilde{p} = 0$ , and so  $v = \mu = 0$ .  $\square$

Next, we prove that the  $\mathcal{A} + \mathcal{B}_D^+$  is injective when  $k^2$  is a Dirichlet eigenvalue.

LEMMA 4.6 (injectivity, part two). *Let  $(v, \mu) \in \mathbb{W}$ . If  $k^2$  is a Dirichlet eigenvalue of the interior Laplacian, then  $\forall (w, \eta) \in \mathbb{W}$*

$$\mathcal{A}[(v, \mu), (w, \eta)] + \mathcal{B}_D^+[(v, \mu), (w, \eta)] = 0$$

implies that  $(v, \mu) = 0$ .

*Proof.* Suppose that  $(v, \mu) \in \mathbb{W}$  such that

$$(4.11) \quad (\mathcal{A} + \mathcal{B}_D^+)[(v, \mu), (w, \eta)] = 0 \quad \forall (w, \eta) \in \mathbb{W}.$$

Taking  $w = 0$  in (4.11), we see that,  $\forall \eta \in H^{-1/2}(\Gamma)$ ,

$$(4.12) \quad \langle \mathbb{V}\mu, \eta \rangle_\Gamma = \langle (\frac{1}{2}\text{Id} + \mathbf{K})v, \eta \rangle_\Gamma.$$

As  $k^2$  is a Dirichlet eigenvalue of the interior Laplacian, then we see by Lemma 4.3 that  $k^2$  is not a Neumann eigenvalue or a Robin eigenvalue.

As  $k^2$  is not a Neumann eigenvalue, we see by [23, Lemma 7.6] that the equation

$$\langle (\frac{1}{2}\text{Id} + \mathbf{K})u, \eta \rangle_\Gamma = \langle g, \eta \rangle_\Gamma \quad \forall \eta \in H^{-1/2}(\Gamma)$$

has a unique solution. By (4.12), we see that  $v$  is the solution of this equation with  $g = \mathbb{V}\mu$ .

Let  $\tilde{p} = \mathcal{V}\mu - \mathcal{K}v$ . By (2.7) and (2.8), we see that

$$(4.13) \quad -\Delta\tilde{p} - k^2\tilde{p} = 0 \quad \text{in } \Omega^-.$$

Taking the Dirichlet trace of  $\tilde{p}$  and applying (4.12), we see that on  $\Gamma$

$$(4.14) \quad \begin{aligned} \gamma_D^-\tilde{p} &= \mathbb{V}\mu + (\frac{1}{2}\text{Id} - \mathbf{K})v \\ &= (\frac{1}{2}\text{Id} + \mathbf{K})v + (\frac{1}{2}\text{Id} - \mathbf{K})v \\ &= v. \end{aligned}$$

We therefore conclude, using the first line of the interior Calderón identity (2.21) and (4.12), that

$$(4.15) \quad \begin{aligned} \mathbb{V}\gamma_N^-\tilde{p} &= (\frac{1}{2}\text{Id} + \mathbf{K})\gamma_D^-\tilde{p} \\ &= (\frac{1}{2}\text{Id} + \mathbf{K})v \\ &= \mathbb{V}\mu. \end{aligned}$$

Note that as  $k^2$  is a Dirichlet eigenvalue,  $\mathbb{V}$  is not injective so this does not necessarily imply that  $\gamma_N^-\tilde{p} = \mu$ .

By [23, Lemma 7.6], the boundary integral equation

$$(4.16) \quad (\frac{1}{2}\text{Id} + \mathbf{K})f = \mu - \gamma_N^-\tilde{p}$$

has a unique solution. Let  $\tilde{q} = \mathcal{V}f$ . By (2.7), we see that

$$(4.17) \quad -\Delta\tilde{q} - k^2\tilde{q} = 0 \quad \text{in } \Omega^-.$$

Taking the Neumann trace of  $\tilde{q}$  and using (2.11c) and (4.16), we see that on  $\Gamma$

$$(4.18) \quad \begin{aligned} \gamma_N^-\tilde{q} &= \gamma_N^-(\mathcal{V}f) \\ &= (\frac{1}{2}\text{Id} + \mathbf{K})f \\ &= \mu - \gamma_N^-\tilde{p}. \end{aligned}$$

We know from the first line of the interior Calderón identity (2.21) that

$$(\frac{1}{2}\text{Id} + \mathbf{K})\gamma_D^-\tilde{q} = \mathbb{V}\gamma_N^-\tilde{q}.$$

Applying (4.18) and then using (4.15) gives

$$(4.19) \quad \begin{aligned} (\tfrac{1}{2}\text{Id} + \mathbf{K})\gamma_{\bar{\mathbf{D}}}\tilde{q} &= \mathbf{V}(\mu - \gamma_{\bar{\mathbf{N}}}\tilde{p}) \\ &= 0. \end{aligned}$$

By [23, Lemma 7.6],  $\tfrac{1}{2}\text{Id} + \mathbf{K}$  is injective, so  $\gamma_{\bar{\mathbf{D}}}\tilde{q} = 0$ . By [23, Lemma 7.5], this implies that  $(\tfrac{1}{2}\text{Id} - \mathbf{K}')\gamma_{\bar{\mathbf{N}}}\tilde{q} = 0$ , and so by (4.18),

$$(4.20) \quad (\tfrac{1}{2}\text{Id} - \mathbf{K}')\gamma_{\bar{\mathbf{N}}}\tilde{p} = (\tfrac{1}{2}\text{Id} - \mathbf{K}')\mu.$$

Using (4.20) and the second line of the interior Calderón identity (2.21), we see that,  $\forall w \in H^{1/2}(\Gamma)$ ,

$$(4.21) \quad \begin{aligned} \langle \mathbf{W}v, w \rangle_{\Gamma} + \langle (\tfrac{1}{2}\text{Id} + \mathbf{K}')\mu, w \rangle_{\Gamma} &= \langle \mathbf{W}v, w \rangle_{\Gamma} + \langle \mu, w \rangle_{\Gamma} + \langle (-\tfrac{1}{2}\text{Id} + \mathbf{K}')\mu, w \rangle_{\Gamma} \\ &= \langle \mathbf{W}\gamma_{\bar{\mathbf{D}}}\tilde{p}, w \rangle_{\Gamma} + \langle \mu, w \rangle_{\Gamma} + \langle (-\tfrac{1}{2}\text{Id} + \mathbf{K}')\gamma_{\bar{\mathbf{N}}}\tilde{p}, w \rangle_{\Gamma} \\ &= \langle \mu, w \rangle_{\Gamma}. \end{aligned}$$

Taking  $\eta = 0$  in (4.4) and applying (4.21), we see that

$$(4.22) \quad \begin{aligned} 0 &= \langle \mathbf{W}v, w \rangle_{\Gamma} + \langle \beta_{\mathbf{D}}v, w \rangle_{\Gamma} + \langle (\tfrac{1}{2}\text{Id} + \mathbf{K}')\mu, w \rangle_{\Gamma} \\ &= \langle \mu, w \rangle_{\Gamma} + \langle \beta_{\mathbf{D}}v, w \rangle_{\Gamma}, \end{aligned}$$

and so  $\mu = -\beta_{\mathbf{D}}v$ .

Consider  $\tilde{p} + \tilde{q}$ . From (4.13) and (4.17), we see that

$$(4.23) \quad -\Delta(\tilde{p} + \tilde{q}) - k^2(\tilde{p} + \tilde{q}) = 0 \quad \text{in } \Omega^-.$$

Using the fact that  $\gamma_{\bar{\mathbf{D}}}\tilde{q} = 0$  (as we concluded from (4.19)) and (4.14) and (4.18), we see that on  $\Gamma$

$$(4.24) \quad \begin{aligned} \gamma_{\bar{\mathbf{D}}}(\tilde{p} + \tilde{q}) &= \gamma_{\bar{\mathbf{D}}}\tilde{p} + \gamma_{\bar{\mathbf{D}}}\tilde{q} \\ &= v, \end{aligned}$$

$$(4.25) \quad \begin{aligned} \gamma_{\bar{\mathbf{N}}}(\tilde{p} + \tilde{q}) &= \gamma_{\bar{\mathbf{N}}}\tilde{p} + \gamma_{\bar{\mathbf{N}}}\tilde{q} \\ &= \mu, \end{aligned}$$

and so, using (4.22), we see that

$$(4.26) \quad \begin{aligned} \frac{\partial(\tilde{p} + \tilde{q})}{\partial \nu} + \beta_{\mathbf{D}}(\tilde{p} + \tilde{q}) &= \mu + \beta_{\mathbf{D}}v \\ &= 0. \end{aligned}$$

Therefore  $\tilde{p} + \tilde{q}$  is solution of the Robin problem (4.3). Since  $k^2$  is not a Robin eigenvalue, we see that  $\tilde{p} + \tilde{q} = 0$  in  $\Omega^-$ , and so by (4.24) and (4.25)  $v = \mu = 0$ .  $\square$

We now combine the previous two results.

**PROPOSITION 4.7 (injectivity).** *Let  $(v, \mu) \in \mathbb{W}$ . If  $k^2$  is not a Robin eigenvalue of the interior Laplacian with Robin parameter  $\beta_{\mathbf{D}}$ , then  $\forall (w, \eta) \in \mathbb{W}$*

$$\mathcal{A}[(v, \mu), (w, \eta)] + \mathcal{B}_{\mathbf{D}}^+[(v, \mu), (w, \eta)] = 0$$

*implies that  $(v, \mu) = 0$ .*

*Proof.* Combine Lemmas 4.5 and 4.6.  $\square$

Using Lemma 4.4, we see that the following corollary holds.

**COROLLARY 4.8.** *If  $\text{Im}(\beta_D) \neq 0$  and  $\forall (w, \eta) \in \mathbb{W}$*

$$\mathcal{A}[(v, \mu), (w, \eta)] + \mathcal{B}_D^+[(v, \mu), (w, \eta)] = 0,$$

then  $(v, \mu) = 0$ .

We now prove that the form  $\mathcal{A}_T + \mathcal{B}_D^+$  is coercive, where  $\mathcal{A}_T$  is the multitrace form with compact perturbation as defined in (2.27).

**LEMMA 4.9.** *If  $\text{Re}(\beta_D) \geq 0$ , then there exists  $\alpha > 0$  such that  $\forall (v, \mu) \in \mathbb{V}$ ,*

$$\alpha \|(v, \mu)\|_{\mathbb{V}}^2 \leq |(\mathcal{A}_T + \mathcal{B}_D^+)[(v, \mu), (v, \mu)]|.$$

*Proof.* Using the definitions of  $\mathcal{A}_T$  and  $\mathcal{B}_D^+$ , we see that

$$\begin{aligned} \text{Re}((\mathcal{A}_T + \mathcal{B}_D^+)[(v, \mu), (v, \mu)]) &= \text{Re}(-\langle \mathbf{K}v, \mu \rangle_{\Gamma} + \langle (\mathbf{V} + \mathbf{T}_V)\mu, \mu \rangle_{\Gamma} + \langle \mathbf{K}'\mu, v \rangle_{\Gamma} \\ &\quad + \langle (\mathbf{W} + \mathbf{T}_W)v, v \rangle_{\Gamma} + \frac{1}{2} \langle \mu, v \rangle_{\Gamma} - \frac{1}{2} \langle v, \mu \rangle_{\Gamma} + \langle \beta_D v, v \rangle_{\Gamma}) \\ &= \langle (\mathbf{V} + \mathbf{T}_V)\mu, \mu \rangle_{\Gamma} + \langle (\mathbf{W} + \mathbf{T}_W)v, v \rangle_{\Gamma} + \langle \text{Re}(\beta_D)v, v \rangle_{\Gamma}. \end{aligned}$$

Applying Lemmas 2.1 and 2.2 and using the assumption that  $\text{Re}(\beta_D) \geq 0$ , we see that

$$\begin{aligned} \text{Re}((\mathcal{A}_T + \mathcal{B}_D^+)[(v, \mu), (v, \mu)]) &\geq \alpha_V \|\mu\|_{H^{-1/2}(\Gamma)}^2 + \alpha_W \|v\|_{H^{1/2}(\Gamma)}^2 + \text{Re}(\beta_D) \|v\|_{L^2(\Gamma)}^2 \\ &\geq \alpha_V \|\mu\|_{H^{-1/2}(\Gamma)}^2 + \alpha_W \|v\|_{H^{1/2}(\Gamma)}^2 \\ &\geq \alpha \|(v, \mu)\|_{\mathbb{V}}^2 \end{aligned}$$

for some  $\alpha > 0$ .

The result then follows from that fact that for any  $c \in \mathbb{C}$ ,  $|c| \geq \text{Re}(c)$ .  $\square$

Using the previous lemma, we can now prove that the form  $\mathcal{A} + \mathcal{B}_D^+$  is coercive.

**PROPOSITION 4.10 (coercivity).** *If  $\text{Re}(\beta_D) > 0$  and  $k^2$  is not a Robin eigenvalue of the interior Laplacian with Robin parameter  $\beta_D$ , then there exists  $\alpha > 0$  such that  $\forall (v, \mu)$*

$$\alpha \|(v, \mu)\|_{\mathbb{V}} \leq \sup_{(w, \eta) \in \mathbb{V}} \frac{|(\mathcal{A} + \mathcal{B}_D^+)[(v, \mu), (w, \eta)]|}{\|(w, \eta)\|_{\mathbb{V}}}.$$

*Proof.* Following the proof of [13, Theorem 2], we suppose (for a contradiction) that the result does not hold. This means that there exists a sequence  $(v_n, \mu_n)$  such that  $\|(v_n, \mu_n)\|_{\mathbb{V}} = 1$  ( $\forall n$ ) and

$$\sup_{(w, \eta) \in \mathbb{V}} \frac{|(\mathcal{A} + \mathcal{B}_D^+)[(v_n, \mu_n), (w, \eta)]|}{\|(w, \eta)\|_{\mathbb{V}}} \leq \frac{1}{n}.$$

This implies that for every  $(w, \eta) \in \mathbb{V}$ ,

$$|(\mathcal{A} + \mathcal{B}_D^+)[(v_n, \mu_n), (w, \eta)]| \leq \frac{1}{n} \|(w, \eta)\|_{\mathbb{V}},$$

and so

$$(4.27) \quad \lim_{n \rightarrow \infty} (\mathcal{A} + \mathcal{B}_D^+)[(v_n, \mu_n), (w, \eta)] = 0.$$

$H^{1/2}(\Gamma)$  and  $H^{-1/2}(\Gamma)$  are Hilbert spaces, so  $\mathbb{V}$  is a Hilbert space. The bounded sequence  $(v_n, \mu_n)$  therefore contains a weakly convergent subsequence  $(\tilde{v}_n, \tilde{\mu}_n)$  that weakly converges to  $(v_\infty, \mu_\infty) \in \mathbb{V}$ .

By weak convergence and (4.27), we see that, for any  $(w, \eta) \in \mathbb{V}$ ,

$$\begin{aligned} 0 &= \lim_{j \rightarrow \infty} (\mathcal{A} + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (w, \eta)] \\ &= (\mathcal{A} + \mathcal{B}_D^+)[(v_\infty, \mu_\infty), (w, \eta)]. \end{aligned}$$

By Proposition 4.7, it follows that  $(v_\infty, \mu_\infty) = 0$ .

Using Lemma 4.9 and the fact that  $\|(\tilde{v}_n, \tilde{\mu}_n)\|_{\mathbb{V}} = 1$ , we see that there is  $\alpha > 0$  such that  $(\forall n)$

$$\begin{aligned} &(\mathcal{A} + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (\tilde{v}_n, \tilde{\mu}_n)] \\ &= (\mathcal{A}_T + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (\tilde{v}_n, \tilde{\mu}_n)] - \langle T_V \tilde{\mu}_n, \tilde{\mu}_n \rangle_\Gamma - \langle T_W \tilde{v}_n, \tilde{v}_n \rangle_\Gamma \\ &\geq \alpha \|(\tilde{v}_n, \tilde{\mu}_n)\|_{\mathbb{V}}^2 - \langle T_V \tilde{\mu}_n, \tilde{\mu}_n \rangle_\Gamma - \langle T_W \tilde{v}_n, \tilde{v}_n \rangle_\Gamma \\ (4.28) \quad &= \alpha - \langle T_V \tilde{\mu}_n, \tilde{\mu}_n \rangle_\Gamma - \langle T_W \tilde{v}_n, \tilde{v}_n \rangle_\Gamma. \end{aligned}$$

We also know that

$$\begin{aligned} &\langle T_V \tilde{\mu}_n, \tilde{\mu}_n \rangle_\Gamma + \langle T_W \tilde{v}_n, \tilde{v}_n \rangle_\Gamma \\ &\leq \|T_V \tilde{\mu}_n\|_{H^{1/2}(\Gamma)} \|\tilde{\mu}_n\|_{H^{-1/2}(\Gamma)} + \|T_W \tilde{v}_n\|_{H^{-1/2}(\Gamma)} \|\tilde{v}_n\|_{H^{1/2}(\Gamma)} \\ &\leq \|(T_W \tilde{v}_n, T_V \tilde{\mu}_n)\|_{\mathbb{V}} \|(\tilde{v}_n, \tilde{\mu}_n)\|_{\mathbb{V}} \\ (4.29) \quad &= \|(T_W \tilde{v}_n, T_V \tilde{\mu}_n)\|_{\mathbb{V}}. \end{aligned}$$

But since  $T_W$  and  $T_V$  are compact operators and  $(\tilde{v}_n, \tilde{\mu}_n)$  converge weakly to zero we have

$$(4.30) \quad \lim_{n \rightarrow 0} \|(T_W \tilde{v}_n, T_V \tilde{\mu}_n)\|_{\mathbb{V}} = 0.$$

Combining (4.27)–(4.30), we see that

$$0 < \alpha \leq \lim_{n \rightarrow 0} (\mathcal{A} + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (\tilde{v}_n, \tilde{\mu}_n)] = 0,$$

which is a contradiction. □

We conclude our analysis of the continuous problem by showing that it is well-posed.

**PROPOSITION 4.11.** *If  $\text{Re}(\beta_D) > 0$  and  $k^2$  is not a Robin eigenvalue of the interior Laplacian with Robin parameter  $\beta_D$ , then the continuous problem (3.5) has a unique solution.*

*Proof.* As shown in [23, Theorem 3.15], this is a direct consequence of Propositions 4.7 and 4.10. □

**4.2. Analysis of the discrete problem.** We now proceed to analyze the discrete problem (3.8). In this section we will use the following notation.

**DEFINITION 4.12.** *For two quantities  $a$  and  $b$  that may vary with  $h$ , we write  $a \lesssim b$  if there is a constant  $h_0 > 0$  and an  $h$ -independent constant  $c \in \mathbb{R}$  such that  $a \leq cb \forall h < h_0$ . We write  $a \approx b$  if  $a \lesssim b$  and  $b \lesssim a$ .*

We start by proving that coercivity holds in the discrete case.

PROPOSITION 4.13 (discrete coercivity). *If  $|\beta_D| \lesssim 1$  and  $k^2$  is not a Robin eigenvalue of the interior Laplacian with Robin parameter  $\beta_D$ , then there exist  $h_0 \in \mathbb{R}$  and  $\alpha > 0$  such that  $\forall h < h_0$  and  $\forall (v_h, \mu_h) \in \mathbb{V}_h$ ,*

$$\alpha \|(v_h, \mu_h)\|_{\mathcal{B}_D} \leq \sup_{(w_h, \eta_h) \in \mathbb{V}_h \setminus \{0\}} \frac{|\mathcal{A}[(v_h, \mu_h), (w_h, \eta_h)] + \mathcal{B}_D^+[(v_h, \mu_h), (w_h, \eta_h)]|}{\|(w_h, \eta_h)\|_{\mathcal{B}_D}}.$$

*Proof.* Following the proof of [6, Theorem 2.2], we suppose (for a contradiction) that the result does not hold. This implies that there exists a sequence  $(h_n)_{n \in \mathbb{N}}$  such that  $\lim_{n \rightarrow \infty} h_n = 0$ , and for each  $n$  there exists  $(v_{h_n}, \mu_{h_n}) \in \mathbb{V}_{h_n}$  such that

$$(4.31) \quad \|(v_{h_n}, \mu_{h_n})\|_{\mathcal{B}_D} = 1,$$

$$(4.32) \quad \sup_{(w_{h_n}, \eta_{h_n}) \in \mathbb{V}_{h_n}} \frac{|(\mathcal{A} + \mathcal{B}_D^+)[(v_{h_n}, \mu_{h_n}), (w_{h_n}, \eta_{h_n})]|}{\|(w_{h_n}, \eta_{h_n})\|_{\mathcal{B}_D}} < \tau_n,$$

where  $\tau_n > 0$  and  $\lim_{n \rightarrow \infty} \tau_n = 0$ .

We define  $F : \mathbb{V}_{h_n} \rightarrow \mathbb{W}$ , for  $(t_{h_n}, \kappa_{h_n}) \in \mathbb{V}_{h_n}$ , by

$$\langle F(t_{h_n}, \kappa_{h_n}), (w_{h_n}, \eta_{h_n}) \rangle_{\Gamma} = (\mathcal{A} + \mathcal{B}_D^+)[(t_{h_n}, \kappa_{h_n}), (w_{h_n}, \eta_{h_n})] \quad \forall (w_{h_n}, \eta_{h_n}) \in \mathbb{V}_{h_n}.$$

By Lemmas 2.1 and 2.2, there exists a compact operator  $T : \mathbb{V}_{h_n} \rightarrow \mathbb{V}_{h_n}$  such that the real part of the operator  $F + T$  is elliptic. Following the proofs for the Laplace problem in [3], we see that there exists  $\alpha > 0$  such that

$$\begin{aligned} \alpha \|(v_{h_n}, \mu_{h_n})\|_{\mathcal{B}_D} &\leq \sup_{(w_{h_n}, \eta_{h_n}) \in \mathbb{V}_{h_n}} \frac{\operatorname{Re}(\langle (F + T)(v_{h_n}, \mu_{h_n}), (w_{h_n}, \eta_{h_n}) \rangle_{\Gamma})}{\|(w_{h_n}, \eta_{h_n})\|_{\mathcal{B}_D}} \\ &\leq \sup_{(w_{h_n}, \eta_{h_n}) \in \mathbb{V}_{h_n}} \frac{|\langle (F + T)(v_{h_n}, \mu_{h_n}), (w_{h_n}, \eta_{h_n}) \rangle_{\Gamma}|}{\|(w_{h_n}, \eta_{h_n})\|_{\mathcal{B}_D}}. \end{aligned}$$

Using (4.32), we see that

$$\begin{aligned} \alpha \|(v_{h_n}, \mu_{h_n})\|_{\mathcal{B}_D} &\leq \sup_{(w_{h_n}, \eta_{h_n}) \in \mathbb{V}_{h_n}} \frac{|\langle F(v_{h_n}, \mu_{h_n}), (w_{h_n}, \eta_{h_n}) \rangle_{\Gamma}|}{\|(w_{h_n}, \eta_{h_n})\|_{\mathcal{B}_D}} \\ &\quad + \sup_{(w_{h_n}, \eta_{h_n}) \in \mathbb{V}_{h_n}} \frac{|\langle T(v_{h_n}, \mu_{h_n}), (w_{h_n}, \eta_{h_n}) \rangle_{\Gamma}|}{\|(w_{h_n}, \eta_{h_n})\|_{\mathcal{B}_D}} \\ &\leq \tau_n + \sup_{(w_{h_n}, \eta_{h_n}) \in \mathbb{V}_{h_n}} \frac{|\langle T(v_{h_n}, \mu_{h_n}), (w_{h_n}, \eta_{h_n}) \rangle_{\Gamma}|}{\|(w_{h_n}, \eta_{h_n})\|_{\mathcal{B}_D}}. \end{aligned}$$

By the Cauchy–Schwarz inequality,

$$\begin{aligned} |\langle T(v_{h_n}, \mu_{h_n}), (w_{h_n}, \eta_{h_n}) \rangle_{\Gamma}| &\leq \|T(v_{h_n}, \mu_{h_n})\|_{\mathbb{V}} \|(w_{h_n}, \eta_{h_n})\|_{\mathbb{V}} \\ &\leq \|T(v_{h_n}, \mu_{h_n})\|_{\mathbb{V}} \|(w_{h_n}, \eta_{h_n})\|_{\mathcal{B}_D}. \end{aligned}$$

Let  $0 < \epsilon < 1$ . Taking  $N_1 \in \mathbb{N}$  large enough so that  $\forall n > N_1$ ,  $\tau_n < \frac{\epsilon}{2}$ , we see that

$$(4.33) \quad \alpha \|(v_{h_n}, \mu_{h_n})\|_{\mathcal{B}_D} < \frac{\epsilon}{2} + \|T(v_{h_n}, \mu_{h_n})\|_{\mathbb{V}}.$$

The bounded sequence  $(v_{h_n}, \mu_{h_n})$  contains a weakly convergent subsequence  $(\tilde{v}_n, \tilde{\mu}_n)$  that converges to  $(v_0, \mu_0) \in \mathbb{V}$ . This means that there exists  $N_2 \in \mathbb{N}$  such that  $\forall n > N_2$ ,

$$(4.34) \quad \|\mathbb{T}[(\tilde{v}_n, \tilde{\mu}_n) - (v_0, \mu_0)]\|_{\mathbb{V}} \leq \frac{\epsilon}{2}.$$

Let  $\delta > 0$ . Let  $(w, \eta) \in \mathbb{V}$ . The spaces  $\mathbb{V}_h$  are asymptotically dense in  $\mathbb{V}$ , so there exists  $h_0$  such that  $\forall h < h_0$ , there exists  $(w_h, \eta_h) \in \mathbb{V}_h$  such that

$$(4.35) \quad \|(w, \eta) - (w_h, \eta_h)\|_{\mathbb{V}} < \delta.$$

Suppose that  $n$  is large enough that  $(\tilde{v}_n, \tilde{\mu}_n) \in \mathbb{V}_{\tilde{h}_n}$  with  $\tilde{h}_n < h_0$ .

Using (4.32), we see that

$$(4.36) \quad \begin{aligned} & (\mathcal{A} + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (w, \eta)] \\ &= (\mathcal{A} + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (w, \eta) - (w_h, \eta_h)] + (\mathcal{A} + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (w_h, \eta_h)] \\ &< (\mathcal{A} + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (w, \eta) - (w_h, \eta_h)] + \tilde{\tau}_n \|(w_h, \eta_h)\|_{\mathcal{B}_D}, \end{aligned}$$

where  $\tilde{\tau}_n$  represents the values in (4.32) corresponding to the subsequence  $(\tilde{v}_n, \tilde{\mu}_n)$ .

Using Proposition 4.1 and (4.31), we see that

$$(4.37) \quad \begin{aligned} & (\mathcal{A} + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (w, \eta) - (w_h, \eta_h)] \leq M \|(\tilde{v}_n, \tilde{\mu}_n)\|_{\mathcal{B}_D} \|(w, \eta) - (w_h, \eta_h)\|_{\mathcal{B}_D} \\ &= M \|(w, \eta) - (w_h, \eta_h)\|_{\mathcal{B}_D}. \end{aligned}$$

Using the definition of  $\|\cdot\|_{\mathbb{V}}$ , we see that

$$\begin{aligned} \|(w, \eta) - (w_h, \eta_h)\|_{\mathcal{B}_D} &= \|(w, \eta) - (w_h, \eta_h)\|_{\mathbb{V}} + |\beta_D|^{1/2} \|w - w_h\|_{L^2(\Gamma)} \\ &\leq \|(w, \eta) - (w_h, \eta_h)\|_{\mathbb{V}} + |\beta_D|^{1/2} \|(w, \eta) - (w_h, \eta_h)\|_{\mathbb{V}} \\ &= (1 + |\beta_D|^{1/2}) \|(w, \eta) - (w_h, \eta_h)\|_{\mathbb{V}}. \end{aligned}$$

Using our assumption that  $|\beta_D| \lesssim 1$ , i.e.,  $|\beta_D| \leq c$  for some constant  $c \in \mathbb{R}$ , and (4.35), we see that

$$(4.38) \quad \begin{aligned} \|(w, \eta) - (w_h, \eta_h)\|_{\mathcal{B}_D} &\leq c \|(w, \eta) - (w_h, \eta_h)\|_{\mathbb{V}} \\ &< c\delta. \end{aligned}$$

Combining (4.36)–(4.38), we see that

$$(4.39) \quad (\mathcal{A} + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (w, \eta)] < M'\delta + \tilde{\tau}_n \|(w_h, \eta_h)\|_{\mathcal{B}_D},$$

where the  $M' = Mc$  is constant.

As  $|\beta_D| \leq c$ ,

$$\begin{aligned} \|(w_h, \eta_h)\|_{\mathcal{B}_D} &= \|(w_h, \eta_h)\|_{\mathbb{V}} + |\beta_D|^{1/2} \|w_h\|_{L^2(\Gamma)} \\ &\leq \|(w_h, \eta_h)\|_{\mathbb{V}} + c^{1/2} \|w_h\|_{L^2(\Gamma)} \\ &\leq c' \|(w_h, \eta_h)\|_{\mathbb{V}} \end{aligned}$$

for some constant  $c' \in \mathbb{R}$ . Using the triangle inequality and (4.35), we obtain

$$(4.40) \quad \begin{aligned} \|(w_h, \eta_h)\|_{\mathbb{V}} &\leq \|(w, \eta) - (w_h, \eta_h)\|_{\mathbb{V}} + \|(w, \eta)\|_{\mathbb{V}} \\ &< \delta + \|(w, \eta)\|_{\mathbb{V}}, \end{aligned}$$

and so

$$(4.41) \quad (\mathcal{A} + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (w, \eta)] < M''\delta + c'\tilde{\tau}_n \|(w, \eta)\|_{\mathbb{V}},$$

where  $M'' = M' + c'\tilde{\tau}_n$ . As  $\|(w, \eta)\|_{\mathbb{V}}$  is constant, it is possible to take  $n$  large enough that  $\tilde{\tau}_n < \delta/\|(w, \eta)\|_{\mathbb{V}}$ . Hence, we conclude that

$$\lim_{h \rightarrow 0} ((\mathcal{A} + \mathcal{B}_D^+)[(\tilde{v}_n, \tilde{\mu}_n), (w, \eta)]) = 0.$$

As this is true  $\forall (w, \eta) \in \mathbb{V}$ , it follows that

$$(\mathcal{A} + \mathcal{B}_D^+)[(v_0, \mu_0), (w, \eta)] = 0,$$

and so, by Proposition 4.7, we have that  $(v_0, \mu_0) = 0$  whenever  $|\beta_D| \lesssim 1$ .

We therefore conclude using (4.33) and (4.34) that for any  $n > \max(N_1, N_2)$ ,

$$\alpha \|(\tilde{v}_n, \tilde{\mu}_n)\|_{\mathcal{B}_D} < \epsilon.$$

This contradicts (4.31).  $\square$

Based on the results in [3] and the Nitsche formulations for other problems, we would expect this coercivity result to hold for  $|\beta_D| \lesssim h^{-1}$ , which is a wider range of values than we have proven it for. When  $|\beta_D| \lesssim h^{-1}$  and  $|\beta_D| \not\lesssim 1$ , the proof can be modified to show that (4.39) still holds. However, when  $|\beta_D| \approx h^{-1}$  the term  $\|(w_h, \eta_h)\|_{\mathcal{B}_D}$  depends on  $h$ , and it is not necessarily possible to take  $\tilde{\tau}_n$  small enough to counter this, so this prevents us from completing the proof in this case. Note, however, that the method is optimally convergent using the uniformly bounded  $\beta_D$ , so in practice there is no reason to use a stronger penalty parameter.

The coercivity result allows us to prove that the discrete problem is well-posed.

**PROPOSITION 4.14.** *If  $|\beta_D| \lesssim 1$  and  $k^2$  is not a Robin eigenvalue of the interior Laplacian with Robin parameter  $\beta_D$ , then there exists  $h_0 \in \mathbb{R}$  such that  $\forall h < h_0$ , the discrete problem (3.8) has a unique solution.*

*Proof.* Suppose that  $(u_h, \lambda_h)$  and  $(u'_h, \lambda'_h)$  are two solutions of (3.8). Applying Proposition 4.13 to  $(u_h, \lambda_h) - (u'_h, \lambda'_h)$ , we see that

$$\alpha \|(u_h, \lambda_h) - (u'_h, \lambda'_h)\|_{\mathcal{B}_D} \leq \sup_{(w_h, \eta_h) \in \mathbb{V}_h \setminus \{0\}} \frac{|(\mathcal{A} + \mathcal{B}_D^+)[(u_h, \lambda_h) - (u'_h, \lambda'_h), (w_h, \eta_h)]|}{\|(w_h, \eta_h)\|_{\mathcal{B}_D}}.$$

As  $(u_h, \lambda_h)$  and  $(u'_h, \lambda'_h)$  are both solutions of (3.8),  $(\mathcal{A} + \mathcal{B}_D^+)[(u_h, \lambda_h), (w_h, \eta_h)] = (\mathcal{A} + \mathcal{B}_D^+)[(u'_h, \lambda'_h), (w_h, \eta_h)]$ , and so

$$\alpha \|(u_h, \lambda_h) - (u'_h, \lambda'_h)\|_{\mathcal{B}_D} \leq 0.$$

It follows from Proposition 4.7 that  $(u_h, \lambda_h) = (u'_h, \lambda'_h)$ , and so (3.8) has at most one solution.

However, as (3.8) is a linear system with a square matrix, having at most one solution implies that it has exactly one solution.  $\square$

**4.3. A priori error bounds.** To prove our a priori error bounds, we will use the following lemma.

**LEMMA 4.15.** *Assume that  $\operatorname{Re}(\beta_D) > 0$ ,  $h < h_0$ , and  $k^2$  is not a Robin eigenvalue of the interior Laplacian with Robin parameter  $\beta_D$ . Let  $(u, \lambda)$  be the solution of (3.5), and let  $(u_h, \lambda_h)$  be the solution of (3.8). If  $|\beta_D| \lesssim 1$ , then*



$$\|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}_D} \lesssim \inf_{(v_h, \mu_h) \in \mathbb{V}_h} \|(u - v_h, \lambda - \mu_h)\|_{\mathcal{B}_D}.$$

*Proof.* Let  $(v_h, \mu_h) \in \mathbb{V}_h$ . By the triangle inequality, we obtain

$$(4.42) \quad \|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}_D} \leq \|(u - v_h, \lambda - \mu_h)\|_{\mathcal{B}_D} + \|(u_h - v_h, \lambda_h - \mu_h)\|_{\mathcal{B}_D}.$$

Applying Proposition 4.13 to the second term on the right gives

$$\|(u_h - v_h, \lambda_h - \mu_h)\|_{\mathcal{B}_D} \leq \frac{1}{\alpha} \sup_{(w_h, \eta_h) \in \mathbb{V}_h \setminus \{0\}} \frac{|(\mathcal{A} + \mathcal{B}_D^+)[(u_h - v_h, \lambda_h - \mu_h), (w_h, \eta_h)]|}{\|(w_h, \eta_h)\|_{\mathcal{B}_D}}.$$

It follows from (3.5) and (3.8) that  $(\mathcal{A} + \mathcal{B}_D^+)[(u_h, \lambda_h), (w_h, \eta_h)] = (\mathcal{A} + \mathcal{B}_D^+)[(u, \lambda), (w_h, \eta_h)]$ , and so

$$\|(u_h - v_h, \lambda_h - \mu_h)\|_{\mathcal{B}_D} \leq \frac{1}{\alpha} \sup_{(w_h, \eta_h) \in \mathbb{V}_h \setminus \{0\}} \frac{|(\mathcal{A} + \mathcal{B}_D^+)[(u - v_h, \lambda - \mu_h), (w_h, \eta_h)]|}{\|(w_h, \eta_h)\|_{\mathcal{B}_D}}.$$

Applying Proposition 4.1 gives

$$(4.43) \quad \begin{aligned} \|(u_h - v_h, \lambda_h - \mu_h)\|_{\mathcal{B}_D} &\leq \frac{M}{\alpha} \sup_{(w_h, \eta_h) \in \mathbb{V}_h \setminus \{0\}} \frac{\|(u - v_h, \lambda - \mu_h)\|_{\mathcal{B}_D} \|(w_h, \eta_h)\|_{\mathcal{B}_D}}{\|(w_h, \eta_h)\|_{\mathcal{B}_D}} \\ &= \frac{M}{\alpha} \|(u - v_h, \lambda - \mu_h)\|_{\mathcal{B}_D}. \end{aligned}$$

Combining (4.42) and (4.43) and taking the infimum yields the desired result. □

If  $\Gamma$  is a discretization of a smooth surface, then we can take  $\mathbb{V}_h = P_h^p(\Gamma) \times P_h^q(\Gamma)$  as our discrete space. The following result gives the approximation properties of this space.

LEMMA 4.16 (approximation in  $P_h^p(\Gamma) \times P_h^q(\Gamma)$ ). *If  $|\beta_D| \lesssim h^{-1}$ , then  $\forall (v, \mu) \in H^s(\Gamma) \times H^r(\Gamma)$ ,*

$$\inf_{(w_h, \eta_h) \in P_h^p(\Gamma) \times P_h^q(\Gamma)} \|(v - w_h, \mu - \eta_h)\|_{\mathcal{B}_D} \lesssim h^{\zeta-1/2} |v|_{H_{pw}^\zeta(\Gamma)} + h^{\xi+1/2} |\mu|_{H_{pw}^\xi(\Gamma)},$$

where  $\zeta = \min(p + 1, s)$ ,  $\xi = \min(q + 1, r)$ ,  $s \geq \frac{1}{2}$  and  $r \geq -\frac{1}{2}$ .

*Proof.* This can be proved in the same way as [3, Proposition 4.14]. □

We can now prove the a priori error bound for  $\mathbb{V}_h = P_h^p(\Gamma) \times P_h^q(\Gamma)$ .

THEOREM 4.17. *Assume that  $h < h_0$ ,  $\text{Re}(\beta_D) > 0$ ,  $|\beta_D| \lesssim 1$ , and  $k^2$  is not a Robin eigenvalue of the interior Laplacian with Robin parameter  $\beta_D$ . The solution,  $(u, \lambda) \in H^s(\Gamma) \times H^r(\Gamma)$ , of (3.5), and the solution,  $(u_h, \lambda_h) \in P_h^p(\Gamma) \times P_h^q(\Gamma)$ , of (3.8) satisfy*

$$\|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}_D} \lesssim h^{\zeta-1/2} |u|_{H_{pw}^\zeta(\Gamma)} + h^{\xi+1/2} |\lambda|_{H_{pw}^\xi(\Gamma)},$$

where  $\zeta = \min(p + 1, s)$  and  $\xi = \min(q + 1, r)$ .

*Proof.* Combine Lemma 4.16 and Lemma 4.15. □

When  $\Gamma$  is not smooth, the flux space must be adapted so that it can represent jumps in the normal derivative between cells. In this case, we take  $\mathbb{V}_h = P_h^p(\Gamma) \times DP_h^q(\Gamma)$ . The following result gives the approximation properties of this space.

LEMMA 4.18 (approximation in  $P_h^p(\Gamma) \times DP_h^q(\Gamma)$ ). *If  $|\beta_D| \lesssim h^{-1}$ , then  $\forall (v, \mu) \in H^s(\Gamma) \times H^r(\Gamma)$ ,*

$$\inf_{(w_h, \eta_h) \in P_h^p(\Gamma) \times DP_h^q(\Gamma)} \|(v - w_h, \mu - \eta_h)\|_{\mathcal{B}_D} \lesssim h^{\zeta-1/2} |v|_{H_{pw}^\zeta(\Gamma)} + h^{\xi+1/2} |\mu|_{H_{pw}^\xi(\Gamma)},$$

where  $\zeta = \min(p + 1, s)$ ,  $\xi = \min(q + 1, r)$ ,  $s \geq \frac{1}{2}$  and  $r \geq -\frac{1}{2}$ .

*Proof.* See [3, Proposition 4.14]. □

We can now prove the a priori error bound for  $\mathbb{V}_h = P_h^p(\Gamma) \times DP_h^q(\Gamma)$ .

THEOREM 4.19. *Assume that  $h < h_0$ ,  $\text{Re}(\beta_D) > 0$ ,  $|\beta_D| \lesssim 1$ , and  $k^2$  is not a Robin eigenvalue of the interior Laplacian with Robin parameter  $\beta_D$ . The solution,  $(u, \lambda) \in H^s(\Gamma) \times H^r(\Gamma)$ , of (3.5), and the solution,  $(u_h, \lambda_h) \in P_h^p(\Gamma) \times DP_h^q(\Gamma)$ , of (3.8) satisfy*

$$\|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}_D} \lesssim h^{\zeta-1/2} |u|_{H_{pw}^\zeta(\Gamma)} + h^{\xi+1/2} |\lambda|_{H_{pw}^\xi(\Gamma)},$$

where  $\zeta = \min(p + 1, s)$  and  $\xi = \min(q + 1, r)$ .

*Proof.* Combine Lemma 4.18 and Lemma 4.15. □

Recall that due to Lemma 4.4, the Robin eigenvalue assumption in the results in this section holds whenever  $\text{Im}(\beta_D) \neq 0$ .

**5. Numerical results.** In this section, we demonstrate the theory with a series of numerical examples. All the results presented were computed using version 0.2.4 of Bempp-cl, an open source Python BEM library [4]. We precondition all linear systems in this section with blocked mass matrix preconditioners applied from the left. We take  $\mathbb{V}_h = P_h^1(\Gamma) \times P_h^1(\Gamma)$  throughout this section, so we use the preconditioner

$$\begin{pmatrix} M & 0 \\ 0 & M \end{pmatrix},$$

where  $M = (m_{ij})$  is defined by

$$m_{ij} = \langle \phi_j, \phi_i \rangle_\Gamma,$$

where  $\{\phi_0, \phi_1, \dots\}$  is the basis of the space  $P_h^1(\Gamma)$ . The preconditioning corresponds to taking the discrete strong form of the operator, as described in [5].

Define

$$(5.1) \quad g_D(\mathbf{x}) = \frac{e^{ik|\mathbf{r}_1|}}{|\mathbf{r}_1|} + \frac{e^{ik|\mathbf{r}_2|}}{|\mathbf{r}_2|},$$

where  $\mathbf{r}_1 = \mathbf{x} - (\frac{1}{10}, \frac{1}{2}, \frac{1}{2})$  and  $\mathbf{r}_2 = \mathbf{x} - (\frac{1}{10}, \frac{1}{4}, \frac{1}{4})$ . Let  $\Omega^-$  be a bounded domain such that  $(\frac{1}{10}, \frac{1}{2}, \frac{1}{2}) \in \Omega^-$  and  $(\frac{1}{10}, \frac{1}{4}, \frac{1}{4}) \in \Omega^-$ , and let  $\Omega^+ := \mathbb{R}^3 \setminus \Omega^-$ . It is easy to check that for any wavenumber  $k > 0$ ,  $u(\mathbf{x}) = \frac{e^{ik|\mathbf{r}_1|}}{|\mathbf{r}_1|} + \frac{e^{ik|\mathbf{r}_2|}}{|\mathbf{r}_2|}$  is the solution of the exterior Helmholtz problem

$$(5.2a) \quad -\Delta u - k^2 u = 0 \quad \text{in } \Omega^+,$$

$$(5.2b) \quad \frac{\partial u^{\text{scat}}}{\partial |\mathbf{x}|} - iku^{\text{scat}} = o(|\mathbf{x}|^{-1}) \quad \text{as } |\mathbf{x}| \rightarrow \infty,$$

$$(5.2c) \quad u = g_D \quad \text{on } \Gamma,$$




FIG. 1. The error (left) and GMRES iteration counts (right) of the penalty method with  $\beta_D = 1$  for the Helmholtz Dirichlet problem with varying  $k$  on the unit sphere with  $h = 2^{-2}$ . Here we take  $(u_h, \lambda_h), (v_h, \mu_h) \in P_h^1(\Gamma) \times P_h^1(\Gamma)$  and solve to a GMRES tolerance of  $10^{-5}$ .




FIG. 2. The error (left) and GMRES iteration counts (right) of the penalty method with varying real  $\beta_D$  for the Helmholtz Dirichlet problem with  $k = 2.759$  on the unit sphere with  $h = 2^{-2}$  (red triangles),  $h = 2^{-3}$  (red diamonds), and  $h = 2^{-4}$  (red pentagons). Here we take  $(u_h, \lambda_h), (v_h, \mu_h) \in P_h^1(\Gamma) \times P_h^1(\Gamma)$  and solve to a GMRES tolerance of  $10^{-5}$ .

with  $u^{\text{inc}} = 0$  (and so  $u = u^{\text{scat}}$ ). It can also be seen that

$$(5.3) \quad \frac{\partial u}{\partial \nu} = \frac{(ik|\mathbf{r}_1| - 1)e^{ik|\mathbf{r}_1|}}{|\mathbf{r}_1|^3} \mathbf{r}_1 \cdot \nu + \frac{(ik|\mathbf{r}_2| - 1)e^{ik|\mathbf{r}_2|}}{|\mathbf{r}_2|^3} \mathbf{r}_2 \cdot \nu \quad \text{on } \Gamma_N.$$

Using (5.1) and (5.3), we can compute the error of the solutions obtained in this section.

Figure 1 shows how the error and number of GMRES iterations vary as  $k$  is increased using  $\beta_D = 1$ . It can be seen that at various points the number of iterations and error grows as  $k$  approaches a problematic Robin eigenvalue of the problem: one of these is near 2.759, as indicated by the dashed lines.

In section 4, we saw that our formulation can fail to have a unique solution when  $k^2$  is Robin eigenvalues of the Laplacian with Robin parameter  $\beta_D$ . This implies that adjusting the parameter  $\beta_D$  will adjust the locations of these eigenvalues. This can be observed in Figure 2, where the error and number of iterations are shown for the problem with  $k = 2.759$  and varying real  $\beta_D$ . The spike in the error and number




FIG. 3. The error (top) and GMRES iteration counts (bottom) of the penalty method with varying  $\beta_D$  with  $\text{Re}(\beta_D) = 1$  for the Helmholtz Dirichlet problem with  $k = 2.759$  on the unit sphere with  $h = 2^{-2}$  (red triangles),  $h = 2^{-3}$  (red diamonds), and  $h = 2^{-4}$  (red pentagons). Here we take  $(u_h, \lambda_h), (v_h, \mu_h) \in P_h^1(\Gamma) \times P_h^1(\Gamma)$  and solve to a GMRES tolerance of  $10^{-5}$ .

of iterations at  $\beta_D = 1$  (as shown by the dashed lines) is due to the approaching same eigenvalue as we saw in Figure 1. We observe that the increase in the error is less pronounced for meshes with a lower value of  $h$ . This is due to 2.759 being near an eigenvalue of a given discretization of the sphere: the exact approximation of the sphere changes a little as we increase the number of cells in the mesh, and so the eigenvalues differ on the different meshes used.

As we saw in [3] for Laplace problems, we observe in Figure 2 that the error and iteration counts for the problem increase once  $\beta_D$  is increased above a certain level.

Figure 3 shows how the error and iteration counts vary as we adjust the imaginary part of  $\beta_D$  when the real part of  $\beta_D$  is fixed as 1, again taking  $k = 2.759$  so that we hit an eigenvalue when  $\text{Im}(\beta_D) = 0$ . We see that once  $|\text{Im}(\beta_D)|$  is greater than around  $10^{-2}$ , the error and iteration count drop. Once  $|\text{Im}(\beta_D)|$  is too large, the error and iteration count rise in a similar way to that we observed when taking a large real  $\beta_D$ . We observe that the iteration count is slightly lower for a small range of values when  $\text{Im}(\beta_D)$  is negative.

Motivated by our observations in Figure 3, we fix  $\beta_D = 1 - i$ . Figure 4 shows how the error and iteration counts change as we increase  $k$  with this value of  $\beta_D$ . In agreement with Lemma 4.4, we observe that (in contrast to Figure 1) there is no vulnerability to eigenvalues in this case, and the iteration count remains steady as we increase  $k$ .




FIG. 4. The error (left) and GMRES iteration counts (right) of the penalty method with  $\beta_D = 1 - i$  for the Helmholtz Dirichlet problem with varying  $k$  on the unit sphere with  $h = 2^{-2}$ . Here we take  $(u_h, \lambda_h), (v_h, \mu_h) \in P_h^1(\Gamma) \times P_h^1(\Gamma)$  and solve to a GMRES tolerance of  $10^{-5}$ .




FIG. 5. The error (left) and GMRES iteration counts (right) of the penalty method with  $\beta_D = 1 - i$  for the Helmholtz Dirichlet problem with  $k = 3$  on the unit sphere as we reduce  $h$ . The dashed line shows order 2 convergence. Here we take  $(u_h, \lambda_h), (v_h, \mu_h) \in P_h^1(\Gamma) \times P_h^1(\Gamma)$  and solve to a GMRES tolerance of  $10^{-5}$ .

Figure 5 shows the error and number of iterations as we reduce  $h$ . We observe order 2 convergence, and see that the number of iterations remains the same as  $h$  is reduced, demonstrating the effectiveness of the mass matrix preconditioner.

**6. Conclusions.** In this paper, we have derived and analyzed a formulation for weak imposition of Dirichlet boundary conditions on the Helmholtz equation. By taking a parameter with a nonzero imaginary part, Helmholtz problems can be solved at any wavenumber without any difficulties caused by resonances of the interior problem.

The formulation derived in this paper bears a close resemblance to the formulations for Laplace that we derived and analyzed in [3]. Formulations for Helmholtz problems with mixed Dirichlet–Neumann or Robin boundary conditions could be derived in the same way. We expect that these formulations could be analyzed following a similar method as used here, although this analysis appears to pose some additional challenges. The weak imposition of mixed boundary conditions is demonstrated in Figure 6, where we have plotted the scattering of an incident wave colliding with a collection of sound-hard (a 0 Neumann boundary condition) and sound-soft (a 0 Dirichlet boundary condition) spheres.




FIG. 6. The incident wave  $u^{\text{inc}} = e^{ik\mathbf{x}\cdot\mathbf{d}}$ , where  $\mathbf{d} = (0, 1, 0)$  and  $k = 2$ , scattering off 25 spheres. The white spheres are sound-hard and the black spheres are sound-soft.

One benefit of formulating mixed problems in this way is that the boundary conditions are imposed by adding sparse terms to the full Calderón system: for mixed boundary conditions, sparse terms assembled on parts of the boundary can be added without any need to adjust the dense Calderón term. When solving an inverse problem—for example, when looking to find the material properties that should be used to give a scatterer a certain desired property—the Calderón term (which is the most expensive part to assemble) can be reused and different sparse terms added to solve the same problem with different boundary conditions.

One avenue of interest for further interest would be the weak imposition of boundary condition on Maxwell problems. In the experiments we have run to explore this, however, we have been unable to obtain good solutions in a reasonable amount of time. Maxwell problems are prone to being strongly ill-conditioned, and it appears that mass matrix preconditioning is not enough to achieve good performance in the Maxwell case. Therefore, we believe that it is necessary to design more powerful preconditioners for these weak formulations in order to make this method feasible for Maxwell problems.

#### REFERENCES

- [1] I. BABUŠKA, *The finite element method with penalty*, Math. Comput., 27 (1973), pp. 221–228, <https://doi.org/10.2307/2005611>.
- [2] L. BANJAI AND S. SAUTER, *A refined Galerkin error and stability analysis for highly indefinite variational problems*, SIAM J. Numer. Anal., 45 (2007), pp. 37–53, <https://doi.org/10.1137/060654177>.
- [3] T. BETCKE, E. BURMAN, AND M. W. SCROGGS, *Boundary element methods with weakly imposed boundary conditions*, SIAM J. Sci. Comput., 41 (2019), pp. A1357–A1384, <https://doi.org/10.1137/18M119625X>.
- [4] T. BETCKE AND M. W. SCROGGS, *Bempp-cl: A fast Python based just-in-time compiling boundary element library*, J. Open Source Softw., (2021), p. 2879, <https://doi.org/10.21105/joss.02879>.
- [5] T. BETCKE, M. W. SCROGGS, AND W. ŚMIGAJ, *Product algebras for Galerkin discretisations of boundary integral operators and their applications*, ACM Trans. Math. Softw., 46 (2020), pp. 4:1–4:22, <https://doi.org/10.1145/3368618>.

- [6] A. S. BONNET-BEN DHIA, P. CIARLET JR., AND C. M. ZWÖLF, *Time harmonic wave diffraction problems in materials with sign-shifting coefficients*, J. Comput. App. Math., 234 (2010), pp. 1912–1919, <https://doi.org/10.1016/j.cam.2009.08.041>.
- [7] E. BURMAN, S. FREI, AND M. W. SCROGGS, *Weak imposition of Signorini boundary conditions on the boundary element method*, SIAM J. Numer. Anal., 58 (2020), pp. 2334–2350, <https://doi.org/10.1137/19M1281721>.
- [8] V. DOMINGUEZ, I. GRAHAM, AND V. SMYSHLYAEV, *A hybrid numerical-asymptotic boundary integral method for high-frequency acoustic scattering*, Numer. Math., 106 (2007), pp. 471–510, <https://doi.org/10.1007/s00211-007-0071-4>.
- [9] S. ENGLER AND O. STEINBACH, *Stabilized boundary element methods for exterior Helmholtz problems*, Numer. Math., 110 (2008), pp. 145–160, <https://doi.org/10.1007/s00211-008-0161-y>.
- [10] I. GRAHAM, M. LÖHNDORF, J. MELENK, AND E. SPENCE, *When is the error in the h-BEM for solving the Helmholtz equation bounded independently of  $k$ ?*, BIT Numer. Math., 55 (2015), pp. 171–214, <https://doi.org/10.1007/s10543-014-0501-5>.
- [11] R. HIPTMAIR AND P. MEURY, *Stabilized FEM-BEM coupling for Helmholtz transmission problems*, SIAM J. Numer. Anal., 44 (2006), pp. 2107–2130, <https://doi.org/10.1137/050639958>.
- [12] A. LALIENA, M.-L. RAPÚN, AND F.-J. SAYAS, *Symmetric boundary integral formulations for Helmholtz transmission problems*, Appl. Numer. Math., 59 (2009), pp. 2814–2823, <https://doi.org/10.1016/j.apnum.2008.12.030>.
- [13] R. LAZAROV, *Inf-sup condition and Banach–Nečas–Babuška theorem*, [http://www.math.tamu.edu/raytcho.lazarov/access/Sofia\\_L.04.pdf](http://www.math.tamu.edu/raytcho.lazarov/access/Sofia_L.04.pdf), 2013, accessed 2019-05-10.
- [14] M. LÖHNDORF AND J. M. MELENK, *Wavenumber-explicit hp-BEM for high frequency scattering*, SIAM J. Numer. Anal., 49 (2011), pp. 2340–2363, <https://doi.org/10.1137/100786034>.
- [15] J. M. MELENK, *Mapping properties of combined field Helmholtz boundary integral operators*, SIAM J. Math. Anal., 44 (2012), pp. 2599–2636, <https://doi.org/10.1137/100784072>.
- [16] J.-C. NÉDÉLEC, *Acoustic and Electromagnetic Equations*, Appl. Math. Sci. 144, Springer-Verlag, New York, 2001, <https://doi.org/10.1007/978-1-4757-4393-7>.
- [17] J. NITSCHKE, *Über ein Variationsprinzip zur Lösung von Dirichlet-Problemen bei Verwendung von Teilräumen, die keinen Randbedingungen unterworfen sind*, Abh. Math. Semin. Univ. Hamburg, 36 (1971), pp. 9–15, <https://doi.org/10.1007/BF02995904>.
- [18] M.-L. RAPÚN AND F.-J. SAYAS, *Mixed boundary integral methods for Helmholtz transmission problems*, J. Comput. Appl. Math., 214 (2008), pp. 238–258, <https://doi.org/10.1016/j.cam.2007.02.028>.
- [19] A. H. SCHATZ, *An observation concerning Ritz–Galerkin methods with indefinite bilinear forms*, Math. Comput., 28 (1974), pp. 959–962, <https://doi.org/10.1090/S0025-5718-1974-0373326-0>.
- [20] S. H. SCHOT, *Eighty years of Sommerfeld’s radiation condition*, Historia Math., 19 (1992), pp. 385–401, [https://doi.org/10.1016/0315-0860\(92\)90004-U](https://doi.org/10.1016/0315-0860(92)90004-U).
- [21] E. A. SPENCE, S. N. CHANDLER-WILDE, I. G. GRAHAM, AND V. P. SMYSHLYAEV, *A new frequency-uniform coercive boundary integral equation for acoustic scattering*, Commun. Pure Appl. Math., 64 (2011), pp. 1384–1415, <https://doi.org/10.1002/cpa.20378>.
- [22] E. A. SPENCE, I. V. KAMOTSKI, AND V. P. SMYSHLYAEV, *Coercivity of combined boundary integral equations in high-frequency scattering*, Commun. Pure Appl. Math., 68 (2015), pp. 1587–1639, <https://doi.org/10.1002/cpa.21543>.
- [23] O. STEINBACH, *Numerical Approximation Methods for Elliptic Boundary Value Problems*, Springer-Verlag, New York, 2008, <https://doi.org/10.1007/978-0-387-68805-3>.